



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

LANE MEDICAL LIBRARY STANFORD STOR
D205 C29v 1896
The microscope and its revelations / by



24503387380



LANE

MEDICAL



LIBRARY


LEVI COOPER LANE FUND


LIBRARY
OF
Cooper Medical College

DATE..... *Oct 1894*

NO. *3301* CLASS.....

GIFT OF
H. Ellinwood





THE
MICROSCOPE:
AND ITS
REVELATIONS.

WORKS BY DR. CARPENTER.

I.

PRINCIPLES OF COMPARATIVE PHYSIOLOGY.

A NEW AMERICAN, FROM THE FOURTH AND REVISED LONDON EDITION.

With more than three hundred beautiful illustrations. In one large and handsome octavo volume of 752 pages.

Without pretending to it, it is an Encyclopædia of the subject, accurate and complete in all respects—a truthful reflection of the advanced state at which the science has now arrived.—*Dublin Quarterly Journal of Medical Science.*

A truly magnificent work—in itself a perfect physiological study.—*Ranking's Abstract.*

This work stands without its fellow. It is one, few men in Europe could have undertaken; it is one, no man, we believe,

could have brought to so successful an issue as Dr. Carpenter. We feel that this abstract can give the reader a very imperfect idea of the fulness of this work, and no idea of its unity, of the admirable manner in which material has been brought, from the most various sources, to conduce to its completeness, of the lucidity of the reasoning it contains, or of the clearness of language in which the whole is clothed.—*Medical Times.*

II.

PRINCIPLES OF HUMAN PHYSIOLOGY.

WITH THEIR CHIEF APPLICATIONS TO

PSYCHOLOGY, PATHOLOGY, THERAPEUTICS, HYGIENE, AND FORENSIC MEDICINE.

A new American, from the last and revised London Edition, with nearly three hundred illustrations. Edited, with additions, by F. Gurney Smith, M.D. In one large and handsome volume of about nine hundred pages.

The most complete work on the science in our language.—*Am. Med. Journal.*

The most complete work now extant in our language.—*N. O. Med. Register.*

The best text-book in the language on this extensive subject.—*London Med. Times.*

A complete cyclopædia of this branch of science.—*N. Y. Med. Times.*

The most complete exposition of physiology which any language can at present give.—*Brit. and For. Med.-Chirurg. Review.*

III.

ELEMENTS (OR MANUAL) OF PHYSIOLOGY.

INCLUDING PHYSIOLOGICAL ANATOMY.

Second American, from the last and revised London Edition, with one hundred and ninety illustrations. In one handsome octavo volume of 566 pages.

In his former works it would seem that he had exhausted the subject of Physiology. In the present, he gives the essence, as it were, of the whole.—*N. Y. Journal of Medicine.*

Those who have occasion for an elementary treatise on Physiology, cannot do better than to possess themselves of the manual of Dr. Carpenter.—*Medical Examiner.*

PREPARING.

PRINCIPLES OF GENERAL PHYSIOLOGY, including ORGANIC CHEMISTRY, and HISTOLOGY. With a general sketch of the Vegetable and Animal Kingdoms. With several hundred illustrations. In one large and handsome octavo volume.

In his last edition of the "Comparative Physiology" and "Human Physiology," the author found it desirable to omit the chapters connected with "General Physiology." He has therefore undertaken to prepare a volume devoted exclusively to that subject, forming an introduction to the other works, or, taken in conjunction with them, constituting a complete and extended system of Physiology, in all its branches.

THE
MICROSCOPE:
AND ITS
REVELATIONS.

BY

WILLIAM B. CARPENTER, M.D., F.R.S., F.G.S.,
EXAMINER IN PHYSIOLOGY AND COMPARATIVE ANATOMY IN THE UNIVERSITY OF LONDON;
PROFESSOR OF MEDICAL JURISPRUDENCE IN UNIVERSITY COLLEGE;
PRESIDENT OF THE MICROSCOPICAL SOCIETY OF LONDON; ETC.

WITH AN APPENDIX

CONTAINING THE

APPLICATIONS OF THE MICROSCOPE TO CLINICAL MEDICINE, ETC.

BY

FRANCIS GURNEY SMITH, M.D.,
PROFESSOR OF THE INSTITUTES OF MEDICINE
IN THE MEDICAL DEPARTMENT OF PENNSYLVANIA COLLEGE, ETC.

ILLUSTRATED BY FOUR HUNDRED AND THIRTY-FOUR
Engravings on Wood.



PHILADELPHIA:
BLANCHARD AND LEA.

1856.

At

Entered, according to Act of Congress, in the year 1856.

BY BLANCHARD AND CO.,

In the Clerk's Office of the District Court for the Eastern District of Pennsylvania.

C. SHERMAN & SON, PRINTERS,

19 St. James Street.

□ 205
C 293
1856

PREFACE TO THE AMERICAN EDITION.

THE American Edition of Dr. Carpenter's Work has been reprinted with the Author's sanction, from advance sheets furnished by him to the American publishers. In assuming the supervision of the press, the Editor has been careful to leave the work as it came from the Author's hands. Such additions as seemed most required by the Students of this country have been made in the form of an Appendix. The reader will find Dr. Carpenter's reasons for omitting the Clinical Applications of the Microscope, fully detailed in his Preface; but as the various works on this subject are not readily accessible on this side of the Atlantic, it was thought that a selection from them, in a compendious form, might enhance the usefulness of the work. Free use has been made, therefore, of the excellent manuals of Beale and Bennett; and the various kindred treatises and journals have also been drawn upon. All that this portion of the work claims is to present a general view of those subjects which seem to be most required by the student of the Microscope. The growing interest in this important field of inquiry will, it is hoped, afford sufficient apology for its introduction.

A short account of American Microscopes, their modifications and accessories, has also been added, and the whole Appendix has been fully illustrated, through the liberality of the Publishers, by the addition of upwards of one hundred wood-engravings.

FRANCIS GURNEY SMITH, M.D.

428 WALNUT STREET,
June, 1856.

P R E F A C E .

THE rapid increase which has recently taken place in the use of the Microscope,—both as an instrument of scientific research, and as a means of gratifying a laudable curiosity and of obtaining a healthful recreation,—has naturally led to a demand for information, both as to the mode of employing the instrument and its appurtenances, and as to the objects for whose minute examination it is most appropriate. And as none of the existing Treatises, either British or Foreign, on the Microscope and its Uses, have seemed to the Author fully adapted to meet this demand (some of them being almost exclusively concerned with the Microscope itself, and others with some special branch or branches of Microscopic study), he has felt encouraged to carry out a plan which he had formed several years since; by endeavoring to combine, within a moderate compass, that information in regard to the use of his “tools” which is most essential to the working Microscopist, with such an account of the objects best fitted for his study, as might qualify him to comprehend what he observes, and might thus prepare him to benefit science, whilst expanding and refreshing his own mind.

In his account of the various forms of Microscopes and Accessory Apparatus, the Author has not attempted to describe everything which is in use in this country; still less, to go into details respecting the construction of foreign instruments. He is satisfied that in all which relates both to the mechanical and the optical arrangements of their instruments, the chief English Microscope-makers are decidedly in advance of their Continental rivals; and on this point he speaks not only from his own conviction, but from the authority of a highly accomplished German Microscopist, who has recently visited London for the

express purpose of making the comparison. Even among the products of English skill, it was necessary for him to make a selection; and he trusts that he will be found to have done adequate justice to all those who have most claim to an honorable mention.

The great objection to English Microscopes, especially on the western side of the Atlantic, seems to be their costliness; and as it can be affirmed with truth, that the instruments of Nacet, Oberhauser, and other Continental makers, are adequate for all essential purposes, a general preference is given to the latter (as the Author understands) among the Microscopists of the United States. He feels sure, however, that no one who has ever been accustomed to work with a well-constructed English Microscope will ever give the preference to a foreign instrument; and he is glad to be able to announce that one of the best London firms is now prepared to supply a Microscope of excellent quality at a price very little exceeding that paid for Continental instruments, of far superior capabilities. (See p. 103, note.)

In treating of the Applications of the Microscope, the Author has constantly endeavored to meet the wants of those who come to the study of the minute forms of Animal and Vegetable life with little or no previous scientific preparation, but who desire to gain something more than a mere *sight* of the objects to which their observation may be directed. Some of these may perhaps object to the general tone of his work as too highly pitched, and may think that he might have rendered his descriptions simpler by employing fewer scientific terms. But he would reply to such, that he has had much opportunity of observing, among the votaries of the Microscope, a desire for such information as he has attempted to convey (of the extent of which desire, the success of the "Quarterly Journal of Microscopical Science" is a very gratifying evidence); and that the use of scientific terms cannot be dispensed with, since there are no others in which the facts can be expressed. As he has made a point of explaining these, in the places where they are first introduced, he cannot think that any of his readers need find much difficulty in apprehending their meaning.

The proportion of space allotted to the various departments, has been determined, not so much by their Physiological importance, as by their special interest to the Microscopist; and the remembrance of this consideration will serve to account for

much that might otherwise appear strange. The Author has thought it particularly needful to restrain himself, in treating of certain very important subjects which are fully discussed in treatises expressly devoted to them (such, for example, as the structure of Insects, and the Primary Tissues of Vertebrata), in order that he might give more space to those on which no such sources of information are readily accessible. For the same reason he has omitted all reference to the applications of the Microscope to Pathological inquiry; a subject which would interest only one division of his readers, and on which it would have been impossible for him to compress, within a sufficiently narrow compass, a really useful summary of what such readers can readily learn elsewhere. So again, the application of the Microscope to the detection of Adulterations in Food, &c., is a topic of such a purely special character, and must be so entirely based on detailed descriptions of the substances in question, that he has thought it better to leave this also untouched.

It has been the Author's object throughout, to guide the possessor of a Microscope to the *intelligent* study of any department of Natural History, that his individual tastes may lead him to follow out, and his particular circumstances may give him facilities for pursuing. And he has particularly aimed to show, under each head, how small is the amount of reliable knowledge already acquired, compared with that which remains to be attained by the zealous and persevering student. Being satisfied that there is a large quantity of valuable *Microscope-power* at present running to waste in this country,—being applied in such desultory observations as are of no service whatever to science, and of very little to the mind of the observer,—he will consider the labor he has bestowed upon the production of this Manual as well repaid, if it should tend to direct this power to more systematic labors, in those fertile fields which only await the cultivator to bear abundant fruit.

In all that concerns the *working* of the Microscope, the Author has mainly drawn upon his own experience, which dates back almost to the time when Achromatic Object-glasses were first constructed in this country. He would be ungrateful, however, if he were not to acknowledge that he has derived many valuable hints from the Practical Treatises of Mr. Quekett and Dr. Beale, and from the Micrographic Dictionary of Messrs. Griffith and Henfrey. And among the works by which he has been

specially aided in treating of the Applications of the instrument, he would especially name Mr. Quekett's valuable Lectures on Histology (Vegetable and Animal), Mr. Ralfs's beautiful Monograph on the British Desmidiæ, Prof. W. Smith's on the Diatomaceæ (which will, when complete, be quite worthy to take rank with the preceding), and the Micrographic Dictionary.

All the Illustrations have been drawn by Mr. W. Bagg, and have been engraved under his superintendence; and the Author ventures to affirm, that for fidelity as well as for beauty of execution, they will bear comparison with any Microscopic delineations yet executed on wood. A large proportion of the subjects are original; the sources of all that are not so, are specified in the list (p. xvii).

The Author feels that some apology is due for the long delay which has attended the appearance of this work. When it was first announced as forthcoming, his full intention was to apply himself immediately to its production; but the unexpected demand for new editions of his two large Treatises on Physiology, required that the whole of his disposable time and attention should be given up during two years to carrying these through the press. When he at last found himself free to apply himself to the "Microscope," he fully expected that the forward state of his preparations would enable him to complete it by October, 1855. But in this expectation he has been disappointed by the occurrence of two severe attacks of indisposition, which compelled him for a time to suspend all mental exertion, and have rendered it necessary for him carefully to abstain from overtasking himself; so that he feels assured that those who have kindly waited for the appearance of this volume, will not, when acquainted with these circumstances, blame him for a delay, the causes of which have lain so completely beyond his control.

UNIVERSITY HALL, LONDON,
Feb. 9, 1856.

TABLE OF CONTENTS.

INTRODUCTION.

	PAGE
History of the Microscope and Microscopic Research,	33

CHAPTER I.

Optical Principles of the Microscope,	65
---	----

CHAPTER II.

CONSTRUCTION OF THE MICROSCOPE.

General Principles,	86
Simple Microscopes,	90
Ross's,	91
Gairdner's,	92
Field's,	93
Quekett's,	94
Compound Microscopes,	96
Field's,	97
Highley's,	99
Nachet's,	100
Smith and Beck's Student's,	102
Do. Dissecting,	103
Warrington's Universal,	105
Ross's,	108
Powell and Lealand's,	111
Smith and Beck's,	113
Nachet's Binocular,	114

CHAPTER III.

ACCESSORY APPARATUS.

Draw-Tube,	116
Erector,	119
Micrometer,	120
Goniometer,	124
Indicator,	124
Camera Lucida,	124
Object-Glass Holder,	127

	PAGE
Object-Marker,	127
Lever Stage,	128
Object-Finder,	129
Magnetic Stage,	130
Diaphragm-Plate,	131
Achromatic Condenser,	131
Reflecting Prisms,	133
White Cloud Illuminator,	134
Oblique Illuminators,	135
Black-Ground Illuminators,	138
Polarizing Apparatus,	140
Illuminators for Opaque Objects,	143
Stage-Forceps,	146
Glass Stage-Plate,	147
Aquatic Box,	148
Zoophyte Trough,	149
Compressorium,	150
Dipping-Tubes,	152
Forceps,	153

CHAPTER IV.

MANAGEMENT OF THE MICROSCOPE.

Support,	154
Light,	154
Position of Light,	157
Care of the Eyes,	158
Care of the Microscope,	159
General Arrangements,	160
Arrangement for Transparent Objects,	168
Arrangement for Opaque Objects,	175
Errors of Interpretation,	180
Comparative Values of Object-Glasses ; Test-Objects,	192
Determination of Magnifying Power,	199

CHAPTER V.

PREPARATION, MOUNTING, AND COLLECTION OF OBJECTS.

Microscopic Dissection,	201
Cutting Sections of Soft Substances,	204
Cutting Sections of Harder Substances,	205
Grinding and Polishing of Sections,	207
Chemical Actions,	211
Glass Slides,	214
Thin Glass,	214
Varnishes and Cements,	217
Mounting Objects Dry,	221
Mounting Objects in Canada Balsam,	224
Preservative Fluids,	232
Mounting Objects in Fluid,	234
Cement-Cells,	235
Thin-Glass Cells,	236

CONTENTS.

xiii

	PAGE
Plate-Glass and Shallow Cells,	237
Deep and Built-up Cells,	239
Mounting Objects in Cells,	240
Importance of Cleanliness,	242
Labelling and Preserving of Objects,	242
Collection of Objects,	243

CHAPTER VI.

MICROSCOPIC FORMS OF VEGETABLE LIFE—PROTOPHYTES.

Boundary between Animal and Vegetable Kingdoms,	247
Characters of Vegetable Cell,	248
Life-History of Simplest Protophytes,	251
Volvocinæ,	259
Desmidiaceæ,	265
Diatomaceæ,	277
Palmellaceæ,	307
Ulvaceæ,	309
Oscillatoriaceæ,	311
Siphonaceæ,	313
Confervaceæ,	317
Conjugatæ,	319
Chætophoraceæ,	321
Batrachospermeæ,	321
Characeæ,	322

CHAPTER VII.

MICROSCOPIC STRUCTURE OF HIGHER CRYPTOGAMIA.

Algæ,	327
Lichens,	333
Fungi,	334
Hepaticæ,	344
Mossee,	348
Ferns,	352
Equisetaceæ,	358

CHAPTER VIII.

MICROSCOPIC STRUCTURE OF PHANEROGAMIC PLANTS.

Elementary Tissues,	360
Structure of Stem and Root,	376
Structure of Cuticle and Leaves,	388
Structure of Flowers,	395

CHAPTER IX.

MICROSCOPIC FORMS OF ANIMAL LIFE:—PROTOZOA; ANIMALCULES.

Protozoa,	404
Rhizopoda,	410
Infusoria,	412
Rotifera,	426

CHAPTER X.

FORAMINIFERA, POLYCYSTINA, AND SPONGES.

	PAGE
Foraminifera,	436
Polycystina,	449
Sponges,	452

CHAPTER XI.

ZOOPHYTES.

Hydra,	457
Hydrozoa,	461
Acalephæ,	467
Anthozoa,	472

CHAPTER XII.

ECHINODERMATA.

Structure of Shell and Spines,	477
Echinoderm Larvæ,	486

CHAPTER XIII.

POLYZOA AND COMPOUND TUNICATA.

Polyzoa,	492
Compound Tunicata,	500

CHAPTER XIV.

MOLLUSCOUS ANIMALS GENERALLY.

Shells of Mollusca,	506
Tongues of Gasteropods,	517
Development of Gasteropods,	521
Ciliary motion on Gills,	526
Organs of Sense of Mollusks,	527
Chromatophores of Cephalopods,	528

CHAPTER XV.

ANNULOSA OR WORMS.

Entozoa,	527
Turbellaria,	533
Annelida,	535

CHAPTER XVI.

CRUSTACEA.

Pycnogonidæ,	539
Entomostraca,	541
Suctoria,	547
Cirripeda,	548
Shell of Decapoda,	550
Metamorphosis of Decapoda,	551

CONTENTS.

XV

CHAPTER XVII.

INSECTS AND ARACHNIDA.

	PAGE
Great number and variety of Objects afforded by Insects,	554
Structure of Integument,	555
Tegumentary Appendages,	556
Parts of the Head,	561
Circulation of the Blood,	569
Respiratory Apparatus,	570
Wings,	574
Feet,	576
Stings and Ovipositors,	578
Eggs,	579
Acarida,	581
Parts of Spiders,	583

CHAPTER XVIII.

VERTEBRATED ANIMALS.

Primary Tissues,	584
Bone,	584
Teeth,	588
Scales of Fish,	591
Hairs,	594
Feathers,	597
Hoofs, Horns, &c.,	598
Blood,	598
White and Yellow Fibres,	602
Skin, Mucous, and Serous Membranes,	604
Epidermis,	605
Pigment-Cells,	605
Epithelium,	606
Fat,	607
Cartilage,	608
Glands,	609
Muscle,	611
Nerve,	614
Circulation of the Blood,	616
Injected Preparations,	619
Vessels of Respiratory Organs,	624

CHAPTER XIX.

APPLICATIONS OF THE MICROSCOPE TO GEOLOGY.

Fossilized Wood, Coal,	625
Fossil Foraminifera, Chalk,	631
Nummulites,	634
Orbitoides,	637
Organic materials of Rocks,	639
Structure of Fossil Bones, Teeth, &c.,	641

CHAPTER XX.

INORGANIC OR MINERAL KINGDOM.—POLARIZATION.

	PAGE
Mineral objects,	644
Crystallization of Salts,	645
Objects suitable for Polariscope,	645

APPENDIX.

Microscope as a means of Diagnosis,	650
Examination of the Nervous System,	653
Muscular System,	655
Respiratory System,	657
Morbid Lung,	658
Examination of the Glandular System,	662
Liver,	662
Kidney,	663
Morbid Kidney,	664
Salivary Glands,	664
Thymus and Thyroid Glands,	664
Adipose Tissue,	664
Fatty Degeneration,	665
Examination of Vascular and Absorbent Systems,	666
Skin, Mucous and Serous Membrane,	667
The Eye,	671
The Hard Tissues,	672
Morbid Growths,	674
Animal Fluids,	681
Serous and Dropsical Fluids,	699
Injections,	700
Microscopes of American Manufacture,	704

LIST OF ILLUSTRATIONS.

	PAGE
1. Refraction of Parallel Rays by Plano-convex Lens,	67
2. Ditto by Double-convex Lens,	67
3. Refraction of Converging Rays,	68
4. Refraction of Diverging Rays,	68
5. Formation of Images by Convex Lenses,	70
6. Spherical Aberration,	70
7. Chromatic Aberration,	72
8. Section of the Achromatic Object-glass,	75
9. Effect of Covering-glass,	76
10. Action of Simple Microscope,	78
11. Simplest form of Compound Microscope,	82
12. Complete Compound Microscope,	82
13. Huyghenian Eye-piece,	83
14. Ross's Simple Microscope,	91
15. Gairdner's Simple Microscope,	92
16. Field's Simple Microscope,	94
17. Quekett's Dissecting Microscope,	95
18. Field's Compound Microscope,	98
19. Highley's Hospital Microscope,	99
20. Nacet's Compound Microscope,	101
21. Smith and Beck's Student's Microscope,	102
22. Ditto Dissecting Microscope,	104
23. Warrington's Universal Microscope,	105
24. Ditto Ditto,	106
25. Ditto Ditto,	106
26. Ditto Ditto,	107
27. Ross's Large Compound Microscope,	108
28. Powell and Lealand's Ditto,	111
29. Smith and Beck's Ditto,	113
30. Arrangement of Binocular Prisms,	115
31. Nacet's Binocular Microscope,	116
32. Draw-tube with Erector,	119
33. Jackson's Eye-piece Micrometer,	122
34. Microscope arranged for Drawing,	125
35. Ross's Achromatic Condenser,	133
36. Smith and Beck's Ditto,	133
37. Powell and Lealand's Ditto,	134
38. White-Cloud Illuminator,	135
39. Amici's Prism,	137
40. Parabolic Illuminator,	139
41. Fitting of Polarizing Prism,	141
42. Fitting of Analyzing Prism,	141
43. Selenite Object-Carrier,	142
44. Bull's-Eye Condenser,	143
45. Ordinary Condensing Lens,	144
46. Side Reflector,	145

	PAGE
47. Stage-Forceps,	147
48. Aquatic Box,	148
49. Zoophyte-Trough,	150
50. Compressorium,	151
51. Dipping-Tubes,	152
52. Forceps,	153
53. Section of Adjusting Object-Glass,	165
54. Arrangement of Microscope for Transparent Objects,	169
55. Ditto Ditto Opaque Objects,	177
56. Spring Scissors,	203
57. Curved Scissors,	204
58. Valentin's Knife,	205
59. Section-Instrument,	206
60. Lever of Contact,	216
61. Slider-Forceps,	226
62. Spring-Press,	227
63. Turn-Table,	236
64. Plate-glass Cells,	238
65. Tube-Cells,	239
66. Built-up Cells,	240
67. <i>Palmoglaea macrococca</i> , after Braun,	251
68. <i>Protococcus pluviialis</i> , after Cohn,	254
69. <i>Volvox globator</i> , after Ehrenberg,	259
70. Development of <i>Volvox</i> , after Williamson,	262
71. Various species of <i>Staurastrum</i> , after Ralfs,	266
72. <i>Closterium lunula</i> , after S. G. Osborne,	267
73. Development of <i>Pediastrum granulatum</i> , after Braun,	271
74. Various forms of <i>Pediastrum</i> , after Ralfs,	272
75. Conjugation of <i>Cosmarium botrytis</i> , after Ralfs,	274
76. Ditto of <i>Closterium</i> , after Ralfs,	275
77. <i>Didymoprium Grevillii</i> , after Ralfs,	275
78. Portion of <i>Isthmia nervosa</i> , after Smith,	280
79. <i>Triceratium favus</i> , after Smith,	281
80. <i>Pleurosigma angulatum</i> , after Wenham,	282
81. <i>Biddulphia pulchella</i> , after Smith,	283
82. Conjugation of <i>Epithemia</i> , after Thwaites,	285
83. Conjugation of <i>Aulacoseira</i> , after Thwaites,	286
84. <i>Actinocyclus undulatus</i> , after Smith,	289
85. <i>Heliopecta</i> ,	290
86. <i>Arachnoidiscus Ehrenbergii</i> , after Smith,	291
87. <i>Campylodiscus costatus</i> , after Smith,	292
88. <i>Surirella constricta</i> , after Smith,	293
89. <i>Gomphonema geminatum</i> , after Smith,	294
90. Ditto, more highly magnified, after Smith,	295
91. <i>Licmophora stbellata</i> , after Smith,	296
92. <i>Meridion circulare</i> and <i>Bacillaria paradoxa</i> , after Smith,	296
93. <i>Achnanthes longipes</i> , after Smith,	297
94. <i>Diatoma vulgare</i> , after Smith,	298
95. <i>Grammatophora serpentina</i> , after Smith,	298
96. <i>Isthmia nervosa</i> , after Smith,	298
97. <i>Meloseira subflexilis</i> , after Smith,	299
98. <i>Meloseira varians</i> , after Smith,	299
99. <i>Mastogloia Smithii</i> , after Smith,	300
100. <i>Mastogloia lanceolata</i> , after Smith,	300
101. Fossil <i>Diatomaceæ</i> from Oran, after Ehrenberg,	302
102. Fossil <i>Diatomaceæ</i> from Mourné Mountain, after Ehrenberg,	304
103. <i>Hematococcus sanguineus</i> , after Hassal,	308
104. Development of <i>Ulva</i> , after Kützing,	309
105. Zoospores of <i>Ulva</i> , after Thuret,	310
106. Zoospores of <i>Achlya</i> , after Unger,	316
107. Cell-multiplication of <i>Conferva</i> , after Mohl,	318

LIST OF ILLUSTRATIONS.

xix

	PAGE
108. <i>Zygnema quininum</i> , after Kützing,	320
109. <i>Chaetophora elegans</i> , after Thuret,	321
110. <i>Batrachospermum moniliforme</i> ,	322
111. <i>Nitella flexilis</i> , after Slack,	323
112. Antheridia of <i>Chara</i> , after Thuret,	325
113. <i>Mesogloia vermicularis</i> , after Payer,	327
114. <i>Sphacelaria cirrhosa</i> ,	329
115. Receptacle of <i>Fucus</i> , after Thuret,	329
116. Antheridia and antherozoids of <i>Fucus</i> , after Thuret,	330
117. Tetraspores of <i>Carpocaulon</i> , after Kützing,	332
118. <i>Torula Cerevisiæ</i> , after Mandl,	335
119. <i>Sarcina ventriculi</i> , after Robin,	336
120. <i>Botrytis bassiana</i> , after Robin,	337
121. <i>Enterobryus spiralis</i> , after Leidy,	338
122. Structure of <i>Enterobryus</i> , after Leidy,	339
123. Fungoid Vegetation, from <i>Pussulus</i> , after Leidy,	340
124. <i>Sysanus caput-medusæ</i> , after Payer,	341
125. <i>Puccinia graminis</i> ,	342
126. <i>Æcidium tussilaginis</i> , after Payer,	343
127. <i>Clavaria crispula</i> , after Payer,	344
128. Fructification of <i>Marchantia</i> , after Payer,	345
129. Stomata of <i>Marchantia</i> , after Mirbel,	345
130. Conceptacles of <i>Marchantia</i> , after Mirbel,	346
131. Archegonia of <i>Marchantia</i> , after Payer,	347
132. Elater and spores of <i>Marchantia</i> , after Payer,	348
133. Portion of Leaf of <i>Sphagnum</i> ,	349
134. Structure of <i>Mosses</i> , after Jussieu,	349
135. Antheridia and antherozoids of <i>Polytrichum</i> , after Thuret,	350
136. Mouth of Capsule of <i>Funaria</i> ,	351
137. Peristome of <i>Fontinalis</i> , after Payer,	351
138. Ditto of <i>Bryum</i> , ditto,	351
139. Ditto of <i>Cinclidium</i> , ditto,	351
140. Petiole of <i>Fern</i> ,	352
141. Sori of <i>Polypodium</i> , after Payer,	353
142. Ditto of <i>Hæmionitis</i> , ditto,	353
143. Sorus and Indusium of <i>Aspidium</i> ,	354
144. Ditto of <i>Deparia</i> , after Payer,	354
145. Development of Prothallium of <i>Pteris</i> , after Suminski,	355
146. Antheridia and antherozoids of <i>Pteris</i> , after Suminski,	356
147. Archegonium of <i>Pteris</i> , after Suminski,	357
148. Spores of <i>Equisetum</i> , after Payer,	358
149. Section of leaf of <i>Agave</i> , after Hartig,	361
150. Section of <i>Aralia</i> (rice-paper),	362
151. Stellate Parenchyma of <i>Rush</i> ,	362
152. Cubical parenchyma of <i>Nuphar</i> ,	362
153. Circulation in hairs of <i>Tradescantia</i> , after Slack,	366
154. Testa of <i>Star-Anise</i> ,	367
155. Section of <i>Cherry-stone</i> ,	368
156. Section of <i>Coquilla-nut</i> ,	368
157. Spiral cells of <i>Oncidium</i> ,	369
158. Spiral fibres of <i>Collomia</i> ,	369
159. Cells of <i>Pæony</i> , filled with starch,	370
160. Starch-grains, under polarized light,	370
161. Glandular fibres of <i>Coniferous Wood</i> ,	373
162. Vascular tissue of <i>Italian Reed</i> , after Schleiden,	374
163. Transverse Section of stem of <i>Palm</i> ,	377
164. Ditto of <i>Wunghie Cane</i> ,	377
165. Ditto of <i>Clematis</i> ,	379
166. Ditto of <i>Cedar</i> ,	379
167. Ditto of <i>Buckthorn</i> ,	380
168. Ditto Ditto more highly magnified,	380

	PAGE
169. Transverse Section of <i>Hazel</i> ,	380
170. Ditto of <i>Fossil Conifer</i> ,	381
171. Vertical section of <i>Fossil Conifer</i> , radial,	382
172. Ditto Ditto, tangential,	382
173. Ditto of <i>Mahogany</i> ,	382
174. Transverse Section of Fossil-wood,	383
175. Vertical Section of Ditto,	383
176. Transverse Section of Fossil-wood,	383
177. Vertical Section Ditto,	383
178. Transverse Section of <i>Aristolochia</i> , (?)	385
179. Ditto of <i>Burdock</i> ,	385
180. Cuticle of <i>Yucca</i> ,	388
181. Ditto of <i>Indian Corn</i> ,	388
182. Ditto of <i>Apple</i> , after Brongniart,	389
183. Ditto of <i>Rochea</i> ,	390
184. Vertical Section of leaf of <i>Rochea</i> , after Brongniart,	390
185. Cuticle of <i>Iris</i> , Ditto,	392
186. Vertical Section of leaf of <i>Iris</i> , Ditto,	393
187. Longitudinal Section of ditto, Ditto,	393
188. Cuticle of Petal of <i>Geranium</i> ,	395
189. Pollen-grains of <i>Althæa</i> , &c.,	398
190. Seeds of <i>Poppy</i> , &c.,	402
191. <i>Anaba princeps</i> , after Ehrenberg,	405
192. <i>Actinophrys sol</i> , after Clarapède,	407
193. Various forms of <i>Rhizopods</i> , after Ehrenberg,	411
194. <i>Kerona silurus</i> , after Milne-Edwards,	414
195. <i>Paramecium caudatum</i> , Ditto,	414
196. Group of <i>Vorticella</i> , after Ehrenberg,	416
197. Fissiparous Multiplication of <i>Chilodon</i> , after Ehrenberg,	418
198. Metamorphoses of <i>Vorticella</i> , after Stein,	419
199. Ditto of <i>Trichoda</i> , after Haime,	420
200. <i>Brachionus pala</i> , after Milne-Edwards,	427
201. <i>Rotifer vulgaris</i> , after Ehrenberg,	429
202. <i>Stephanoceros Eichornii</i> , Ditto,	432
203. <i>Noteus quadricornis</i> , Ditto,	434
204. <i>Gromia oviformis</i> , after Schulze,	437
205. <i>Rosalina ornata</i> , after Schulze,	439
206. <i>Orbitolites complanatus</i> ,	441
207. Animal of simple type of ditto,	442
208. Animal of complex type of ditto,	444
209. Section of <i>Faujasina</i> , after Williamson,	447
210. <i>Podocyrthis Schomburgkii</i> , after Ehrenberg,	450
211. <i>Rhopalocanium ornatum</i> , Ditto,	450
212. <i>Haliomma Humboldtii</i> , Ditto,	450
213. <i>Perichlamyidium prætextum</i> , Ditto,	450
214. <i>Polycystina</i> from Barbadoes, Ditto,	451
215. <i>Stylodictya gracilis</i> , Ditto,	452
216. <i>Astromma Aristotelis</i> , Ditto,	452
217. Structure of <i>Grantia</i> , after Dobie,	453
218. Portion of <i>Halichondria</i> ,	454
219. Siliceous spicules of <i>Pachymatisma</i> ,	454
220. <i>Hydra fusca</i> , after Milne-Edwards,	458
221. Ditto in gemmation, after Trembley,	460
222. Medusa-buds of <i>Syncoryna</i> , after Sars,	460
223. <i>Campanularia gelatinosa</i> , after Van Beneden,	464
224. <i>Sertularia cupressina</i> , after Johnston,	466
225. <i>Thaumantias pilosella</i> , after E. Forbes,	467
226. Development of <i>Medusa-buds</i> , after Dalyell,	468
227. Development of <i>Medusæ</i> , Ditto,	469
228. <i>Cydippe</i> and <i>Beroë</i> , after Milne-Edwards,	470
229. <i>Noctiluca miliaris</i> , after Quatrefages,	571

LIST OF ILLUSTRATIONS.

xxi

	PAGE
230. Spicules of <i>Alcyonium</i> and <i>Gorgonia</i> ,	472
231. Ditto of <i>Gorgonia guttata</i> ,	472
232. Spicules of <i>Muricea elongata</i> ,	472
233. Filiferous capsules of <i>Actinia</i> , &c., after Gosse,	475
234. Section of Shell of <i>Echinus</i> ,	478
235. Calcareous reticulation of Spine of <i>Echinus</i> ,	478
236. Ambulacral disk of <i>Echinus</i> ,	478
237. Transverse Section of Spine of <i>Acrocladia</i> ,	479
238. Spines of <i>Spatangus</i> ,	481
239. Calcareous skeleton of <i>Astrophyton</i> ,	482
240. Ditto of <i>Holothuria</i> ,	485
241. Ditto of <i>Synapta</i> ,	486
242. Ditto of <i>Chirodota</i> ,	486
243. Bipinnarian larva of <i>Star-fish</i> , after Müller,	487
244. Pluteus-larva of <i>Echinus</i> , after Müller,	489
245. Cells of <i>Leprælia</i> , after Johnston,	492
246. <i>Laguncula repens</i> , after Van Beneden,	493
247. Bird's-head processes of <i>Cellularia</i> and <i>Bugula</i> , after Johnston and Busk,	498
248. <i>Amaroucium proliferum</i> , after Milne-Edwards,	501
249. <i>Botryllus violaceus</i> , Ditto	502
250. <i>Perophora</i> , after Lister,	503
251. Transverse Section of <i>Pinna</i> ,	507
252. Membranous basis of Ditto	507
253. Vertical Section of Ditto	507
254. Oblique Section of Ditto	508
255. Section of hinge-tooth of <i>Mya</i> ,	509
256. Nacre of <i>Arricula</i> ,	510
257. Tubular shell-structure of <i>Anomia</i> ,	512
258. Vertical Section of shell of <i>Unio</i> ,	512
259. Internal surface of shell of <i>Terebratula</i> ,	513
260. External Ditto Ditto	513
261. Vertical Sections of Ditto	514
262. Horizontal Section of shell of <i>Terebratula bullata</i> ,	514
263. Ditto Ditto of <i>Megerlia lima</i> ,	514
264. Ditto Ditto of <i>Spiriferina rostrata</i> ,	514
265. Palate of <i>Helix hortensis</i> ,	518
266. Ditto of <i>Zonites cellarius</i> ,	518
267. Ditto of <i>Trochus zizyphinus</i> ,	519
268. Ditto of <i>Doris tuberculata</i> ,	519
269. Ditto of <i>Buccinum</i> , under polarized light,	521
270. Embryoes of <i>Nudibranchs</i> , after Alder and Hancock,	522
271. Embryonic development of <i>Purpura</i> ,	523
272. Latter stages of the same,	525
273. Structure of <i>Polycelis</i> , after Quatrefages,	534
274. Circulation of <i>Terebella</i> , after Milne-Edwards,	535
275. <i>Ammothea pycnogonoides</i> , after Quatrefages,	540
276. <i>Cyclops quadricornis</i> , after Baird,	543
277. Development of <i>Balanus</i> , after Bate,	549
278. Metamorphoses of <i>Carcinus</i> , after Couch,	551
279. Scale of <i>Morpho Menelaus</i> ,	557
280. Battledoor scale of <i>Polyommatus argus</i> , after Quekett,	557
281. Scales of <i>Podura plumbea</i> ,	558
282. Hairs of <i>Myriapod</i> and <i>Dermeestes</i> ,	560
283. Head and eyes of <i>Bee</i> ,	561
284. Section of Eye of <i>Melolontha</i> , after Strauss-Durckheim,	562
285. Antenna of <i>Cockchafer</i> ,	565
286. Portions of ditto more highly magnified,	565
287. Tongue of <i>Fly</i> ,	566
288. Tongue, &c. of <i>Honey Bee</i> ,	567
289. Proboscis of <i>Vanessa</i> ,	568
290. Tracheal system of <i>Nepa</i> , after Milne-Edwards,	571

	PAGE
291. Trachæe of <i>Dytiscus</i> ,	571
292. Spiracle of <i>Fly</i> ,	572
293. Spiracle of Larva of <i>Cockchaffer</i> ,	572
294. Foot of <i>Fly</i> , after Hepworth,	577
295. Foot of <i>Dytiscus</i> ,	577
296. Eggs of Insects, after Burmeister,	580
297. Foot, with combs, of <i>Spider</i> ,	583
298. Ordinary and glutinous threads of <i>Spider</i> ,	583
299. Minute Structure of Bone, after Wilson,	585
300. Lacunæ of Ditto, highly magnified, after Mandl,	586
301. Section of Bony scale of <i>Lepidosteus</i> ,	587
302. Vertical Section of Tooth of <i>Lamna</i> , after Owen,	588
303. Transverse Ditto of <i>Pristis</i> , ditto,	588
304. Ditto Ditto of <i>Myliobates</i> ,	589
305. Vertical Section of Human Tooth, after Mandl,	590
306. Portion of Skin of <i>Sole</i> ,	592
307. Scale of <i>Sole</i> ,	592
308. Hair of <i>Musk-Deer</i> ,	595
309. Hair of <i>Sable</i> ,	595
310. Hairs of <i>Squirrel</i> and <i>Indian Bat</i> ,	595
311. Transverse section of Hair of <i>Pecari</i> ,	595
312. Structure of <i>Human Hair</i> , after Wilson,	596
313. Transverse section of Horn of <i>Rhinoceros</i> ,	598
314. Blood-corpuscles of <i>Frog</i> , after Donnè,	599
315. Ditto of <i>Man</i> , Ditto,	599
316. Fibrous Membrane of Egg-shell,	603
317. White Fibrous Tissue,	603
318. Yellow Fibrous Tissue,	603
319. Pigment-cells of Choroid, after Henle,	605
320. Pigment-cells of <i>Tadpole</i> , after Schwann,	606
321. Ciliated Epithelium, after Mandl,	607
322. Areolar and Adipose Tissue, after Mandl,	608
323. Cartilage of Ear of <i>Mouse</i> ,	609
324. Cartilage of <i>Tadpole</i> , after Schwann,	609
325. Striated Muscular Fibre,	612
326. Ultimate fibrillæ of Ditto,	612
327. Capillary Circulation in Web of <i>Frog's</i> foot, after Wagner,	618
328. Intestinal villi of <i>Monkey</i> ,	622
329. Various forms of Capillary network, after Berres,	623
330. Portion of Gill of <i>Eel</i> ,	624
331. Interior of Lung of <i>Frog</i> ,	624
332. Section of Lung of <i>Fowl</i> ,	625
333. Section of <i>Human</i> Lung,	626
334. Microscopic organisms in Levant Mud, after Williamson,	630
335. Ditto in Chalk from Gravesend, after Ehrenberg,	631
336. Ditto Ditto from Meudon, Ditto	632
337. Vertical section of <i>Nummulite</i> ,	635
338. Portion of Ditto, more highly magnified,	636
339. Horizontal section of Ditto,	637
340. Section of <i>Orbitoides Prattii</i> parallel to its surface,	637
341. Portions of Ditto, more highly magnified,	638
342. Vertical section of Ditto,	638
343. Eye of <i>Trilobite</i> , after Buckland,	640
344. Section of Tooth of <i>Labyrinthodon</i> , after Owen,	641
345. Crystallized Silver,	645
346. Oily Matter in Fibres of Brain (Todd and Bowman),	653
347. Corpora Amylacea of Pineal Gland (Sieveking and Jones),	654
348. Fatty Degeneration of Nerve-Fibre,	654
349. Striated Muscular Fibre,	654
350. Elongated Muscle-Cells (Non-Striated),	656
351. Sarcolemma,	656

LIST OF ILLUSTRATIONS.

xxiii

	PAGE
352. Structure of Lung,	657
353. Capillaries of Lung,	658
354. Tubercle Corpuscles,	658
355. Yellow Tubercle (Bennett),	658
356. } Scrofulous Pus (Bennett),	658
357. }	
358. Section of Gray Granulations,	659
359. Cretaceous and Cheesy Variety of Tubercle (Bennett),	659
360. Pigmentary Matter mixed with Tubercle (Bennett),	659
361. } Structure of Tubercular Mass from Cerebellum,	659
362. }	
363. Tubercle Corpuscle from Lung,	660
364. Plastic or Pyoid Corpuscle,	660
365. Granular Corpuscle from Cerebral Swelling,	660
366. Corpuscles from Reticulum of Cancer,	660
367. Lung in first Stage of Pneumonia (Da Costa),	661
368. Ditto 2d Ditto Ditto	661
369. Ditto Ditto Ditto Ditto	661
370. Ditto 3d Ditto Ditto	662
371. Transverse Section of Lobule of Human Liver (Leidy),	662
372. Hepatic Cells (Leidy),	663
373. Urine from tube and Epithelial Cell,	663
374. Adipose Tissue,	665
375. Fatty Degeneration of Vessels of Brain,	666
376. } Atheromatous Deposits in Vessels,	667
377. }	
378. Fibre-Cells passing into Fibres,	669
379. Fibrous Tissue formed from Fibre-Cell,	669
380. Altered Epithelial Cells from Ulcer of Lip,	669
381. Epithelial and Fibre-Cells from same,	669
382. Pus Corpuscles,	677
383. Ditto after Acetic Acid,	678
384. Free Cancer Nuclei,	678
385. } Polygonal Cancer-Cells (Donaldson),	679
386. }	
387. Caudate Cancer-Cells (Donaldson),	680
388. Fusiform Cells,	680
389. Concentric Cancer-Cells,	681
390. Compound, a Mother Cell,	681
391. Agglomerated Nuclei (Donaldson),	681
392. Corpuscles of Blood,	681
393. Same after exposure to Air,	681
394. Colorless Corpuscles,	682
395. Blood in Leucocythæmia,	682
396. Caudate Blood-Corpuscles,	683
397. Milk and Colostrum-Granules,	683
398. Healthy Milk-Globules,	684
399. Salivary Corpuscles, Epithelial Scales and Granules,	684
400. Tubercle-Corpuscles (Bennett),	686
401. Sputum of Calcareous Matter,	686
402. Sputum in Acute Pneumonia,	686
403. Pus-Globules,	687
404. Scrofulous Pus,	687
405. Mucous Corpuscles and Epithelium in Urine,	690
406. Secreting Canal and Epithelium from Kidney,	690
407. Mucus, Pus, Blood, and Epithelium in Leucorrhœa,	691
408. Fibrinous Casts from Tubuli Uriniferi,	691
409. Ditto Ditto Ditto in Bright's Disease,	692
410. Tube containing a Homogeneous Cast,	692
411. Spermatozoa,	692
412. Torula from Urine of Diabetes,	693

	PAGE
413. Oil-Globules from Urine,	693
414. Lithic Acid Crystals,	693
415. Various forms of Lithates,	694
416. Crystal of Cystine,	694
417. Oxalate of Lime,	695
418. Same (various forms),	695
419. Dumb-Bell Crystals,	695
420. Urate of Ammonia,	696
421. Triple Phosphate,	696
422. Stellar form of Triple Phosphate,	696
423. Starch Corpuscles after Partial Digestion,	698
424. Microscopic Appearance of Cancerous Juice from Urine,	699
425. Fibres and Corpuscles from Ovarian Fluid,	699
426. Spencer's Trunnion Microscope,	705
427. Queen's Student's Microscope,	706
428. Grunow's Student's Microscope,	707
429. Ditto Ditto smaller size,	708
430. Ditto first-class Microscope,	709
431. Diagram of Smith's Inverted Microscope,	710
432. Dr. J. L. Smith's Inverted Microscope,	711
433. } Dr. J. L. Smith's Goniometer and Micrometer,	712-13
434. }	

THE MICROSCOPE.

INTRODUCTION.

No one who attentively examines the progress of any department of Science, save such as are (like Mathematics or Metaphysics) of a purely abstract character, can fail to perceive how much it is dependent upon the perfection of its *instruments*. There are few instances, in fact, in which the invention of a new instrument, or the improvement of an old one, has not given a fresh stimulus to investigation; even where it has done no more than afford that degree of precision to the results of inquiries already in progress, which alone could enable them to be made available as data for philosophical reasoning. But there are many cases in which such inventions or improvements have opened out entirely new paths of scientific research, leading to fertile fields of investigation whose very existence had been previously unknown, to rich mines of discovery whose treasures had lain uncared for because entirely unsuspected. A few examples of this general truth may not be inappropriate, by way of preface to the brief notice which it is intended to give in the present Introduction, of the most important epochs in the history, as well of the Microscope itself, as of its application to the purposes of scientific research.

Thus in taking a retrospective survey of the history of *Astronomy*, we find that every great advance in our knowledge of the Celestial Universe, has been preceded by improvements, either in those instruments for *measuring space and time*, by which the places of the Heavenly Bodies are determined, the rate of their movements estimated, and a basis for the computation of their distances ascertained; or, again, in the *telescope*, by which our power of sight is so wonderfully augmented, that we are enabled, when gazing through it into the unfathomable depths of space, to take cognizance of world beyond world and system beyond system, whose remoteness cannot be expressed by any form of words that shall convey a distinct idea to the mind, and to bring the members of our own group within such *visual* proximity to

ourselves, that we can scrutinize their appearance nearly as well as if they had *actually* been brought a thousand times nearer to us. For it was the increased precision of celestial observations on the places and movements of the Planets, which furnished the data whereon Kepler was enabled to base his statement of the laws of their motion. It was the application of the pendulum to the measurement of short intervals of time, that enabled Galileo to ascertain the law of Falling Bodies. And it was not until the precise measurement of a degree upon the surface of the Earth had furnished the means of determining both its own diameter and its distance from the Moon, that Newton was enabled to verify and establish his grand conception, of the identity of that force which deflects the planets from a rectilinear course into elliptical orbits, with that which draws a stone to the ground; and thus to establish that Law of Universal Gravitation, which still remains the most comprehensive, as well as the most simple, of all the generalizations, within which the intellect of man has been able to comprehend the phenomena of Nature. So, again, it was only when the elder Herschel had developed new powers in the telescope, that Sidereal Astronomy could be pursued with any view much higher than that of mapping the distribution of the stars in the celestial sphere; and the present state of our knowledge of double, triple, and other combinations of stars, with their mutually adjusted movements, of the multiform clusters of luminous points which seem like repetitions of our own firmament in remote depths of space, and of those nebulous films which may be conceived to be new worlds and systems in process of formation, has only been rendered attainable by the improvements which have been subsequently made in the construction of that majestic instrument.

If we glance at the mode in which the fabric of our existing *Chemistry* has been upreared, we at once see that it could not have attained its present elevation and stability, but for the instrumentality of the perfected *balance*; by whose unerring indications it was that the first decisive blow was given to the old "phlogistic" theory, that the foundation was laid for true ideas of chemical combination, that the Laws of that Combination were determined, and that the Combining Equivalents of different elementary substances were ascertained; and by whose means alone can any of those analytical researches be prosecuted, which are not only daily adding to our knowledge of the composition of the bodies which surround us, and suggesting the most important applications of that knowledge to almost every department of the Arts of Life, but which are preparing a broad and secure foundation for a loftier and more comprehensive system of Chemical Philosophy.

So, again, the *balance of torsion*, the ingenious invention of Cavendish and Coulomb, enables the Physical philosopher not merely to render sensible, but to subject to precise measurement

and subdivision, degrees of force that are far too feeble to affect the nicest balance of the ordinary construction, even if it were possible to bring them to act upon it; and strange as it may seem, it has been in such a balance that the Earth itself has been weighed, and that a basis has been thus afforded for the computation of the weights of the different Planets and even of the Sun; whilst in the opposite direction it is employed to furnish those data in regard to the intensity of the electric and magnetic forces, on which alone can any valid theory of their operation be constructed.

The *galvanometer*, again, in which the minutest Electric disturbances are rendered sensible by the deflection of the magnetic needle, has not only brought to light a vast class of most interesting electric changes which were previously unsuspected (one of the most remarkable of these being the existence of electric currents in the nerves of living animals, first ascertained by M. du Bois-Reymond), but has enabled those changes to be estimated with a marvellous amount of exactness; thus furnishing to observations made by its means, a precision which is quite unattainable in any other mode, and which is absolutely essential to the establishment of any valid theory of electric action. And this same instrument is scarcely less valuable, as serving, by a particular modification of it, for the detection and estimation of changes of Temperature far too minute to be measured by the ordinary thermometer; thus affording the requisite means of exactness to observation, in a department of science to which at first sight it appeared to have no relation.

“What an important influence,” says Sir John Herschel, “may be exercised over the progress of a single branch of science, by the invention of a ready and convenient mode of executing a definite measurement, and the construction and common introduction of an instrument adapted for it, cannot be better exemplified than by the instance of the *reflecting goniometer*; this simple, cheap, and portable little instrument has changed the whole face of Mineralogy, and given it all the characters of one of the exact sciences.”

Of all the instruments which have been yet applied to scientific research, there is perhaps not one which has undergone such important improvements within so brief a space of time, as the *Microscope* has received during the second quarter of the present century; and there is certainly none whose use, under its improved form has been more largely or more rapidly productive of most valuable results. As an optical instrument, the Microscope is now at least as perfect as the Telescope; for the 6-foot parabolic speculum of Lord Rosse's gigantic instrument, is not more completely adapted to the Astronomical survey of the heavenly bodies, than the achromatic combination of lenses so minute that they can scarcely be themselves discerned by the unaided eye, is to the scrutiny of the Physiologist into the mysteries of life

and organization. Nor are the revelations of the one less surprising to those who find their greatest charm in novelty, or less interesting to those who apply themselves to the study of their scientific bearings, than are those of the other. The universe which the Microscope brings under our ken, seems as unbounded in its limit as that whose remotest depths the Telescope still vainly attempts to fathom. Wonders as great are disclosed in a speck of whose minuteness the mind can scarcely form any distinct conception, as in the most mysterious of those nebulae whose incalculable distance baffles our hopes of attaining a more intimate knowledge of their constitution. And the general doctrines to which the labors of Microscopists are manifestly tending, in regard to the laws of Organization and the nature of Vital Action, seem fully deserving to take rank in comprehensiveness and importance with the highest principles yet attained in Physical or Chemical Science.

As the primary object of this treatise is to promote the *use* of the Microscope, by explaining its construction, by instructing the learner in the best methods of employing it, and by pointing out the principal directions in which these may be turned to good account, any detailed review of its *history* would be misplaced. It will suffice to state, that whilst the *simple* microscope or magnifying-glass was known at a very remote period, the *compound* microscope,—the powers of which, like those of the telescope, depend upon the combination of two or more lenses,—was not invented until about the end of the sixteenth century; the earlier microscopes having been little else than modified telescopes, and the essential distinction between the two not having been at first appreciated. Still, even in the very imperfect form which the instrument originally possessed, the attention of scientific men was early attracted to the Microscope; for it opened to them a field of research altogether new, and promised to add largely to their information concerning the structure of every kind of organized body. The Transactions of the Royal Society contain the most striking evidence of the interest taken in microscopic investigations two centuries ago. Their early volumes, as Mr. Quekett truly remarks, “literally teem” with improvements in the construction of the Microscope, and with discoveries made by its means. The *Micrographia* of Robert Hooke, published in 1667, was, for its time, a most wonderful production; but this was soon surpassed by the researches of Leeuwenhoek, whose name first appears in the Philosophical Transactions, in the year 1673. That with such imperfect instruments at his command, this accurate and pains-taking observer should have seen *so much* and *so well*, as to make it dangerous for any one even now to announce a discovery, without having first consulted his works, in order to see whether some anticipation of it may not be found there, must ever remain a marvel to the microscopist. This is partly to be explained by the fact, that he

trusted less to the compound microscope, than to single lenses of high power, the use of which is attended with difficulty, but which are comparatively free from the errors inseparable from the first-named instrument in its original form. The names of Grew and Malpighi, also, appear as frequent contributors to the early volumes of the Philosophical Transactions; the researches of the former having been chiefly directed to the minute structure of Plants, and those of the latter to that of Animals. Both were attended with great success. The former laid the foundation of our anatomical knowledge of the Vegetable tissues, and described their disposition in the roots and stems of a great variety of plants and trees; besides making out many important facts in regard to their physiological actions. The latter did the same for the Animal body; and seems to have been the first to witness the marvellous spectacle of the movement of Blood in the capillary vessels of the Frog's foot,—thus verifying by ocular demonstration that doctrine of the passage of blood from the smallest arteries to the smallest veins, which had been propounded as a rational probability by the sagacious Harvey.

Glimpses of the invisible world of Animalcular life were occasionally revealed to the earlier Microscopists, by which their curiosity must have been strongly excited; yet they do not appear to have entered on this class of investigations, with any large portion of that persevering zeal which they devoted to the analysis of the higher forms of organic structure. Its wonders, however, were gradually unfolded; so that in the various treatises on the Microscope published during the eighteenth century, an account of the plants and animals (but especially of the latter) too minute to be seen by the unaided eye, occupies a conspicuous place. It was towards the middle of that period, that M. Trembley of Geneva first gave to the world his researches on the "Fresh-water Polype" or *Hydra*; the publication of which may be considered to have marked a most important epoch in the history of microscopic inquiry. For it presented to the naturalist the first known example of a class of animals (of which the more delicate and flexible *Zoophytes* are, so to speak, the skeletons) whose claim to that designation had been previously doubted or even denied, the terms "sea-mosses," "sea ferns," &c., having been applied to them, not merely as appropriately indicating their form and aspect, but as expressive of what even the most eminent Zoologists, as well as Botanists, considered to be their vegetable nature. And it presented to the Physiologist an entirely new type of *animal life*; the wonderful nature of which was fitted not only to excite the liveliest interest, but also to effect a vast extension in the range of the ideas entertained up to that time regarding its nature and capacities. For what animal previously known, could propagate itself by buds like a plant,—could produce afresh any part that might be cut away,—could form any number of new heads by the completion of the halves

into which the previous heads had been slit (thus realizing the ancient fable of the Hydra),—could even regenerate the whole from a minute portion, so that when the body of one individual was positively minced into fragments, each of these should grow into a new and complete polype,—could endure being turned inside out, so that what was previously the external surface should become the lining of the stomach, and *vice versâ*,—and could sustain various other kinds of treatment not less strange (such as the grafting of two individuals together, head to head, or tail to tail, or the head of the one to the tail of another), not only without any apparent injury, but with every indication, in the vigor of its life, of being entirely free from suffering or damage? (See Chap. XI).—It was by our own countryman, Ellis, that the discoveries of Trembley were first applied to the elucidation of the real animal nature of the so-called Corallines;¹ the structure of which was so carefully investigated by him, that subsequent observers added little to our knowledge of it, until a comparatively recent period.

The true *animalcules* were first systematically studied, in the latter part of the last century, by Gleichen, a German microscopist, who devised the ingenious plan of feeding them with particles of coloring matter, so as to make apparent the form and position of their digestive cavities; and this study was afterwards zealously pursued by the eminent Danish naturalist, Otho Fred. Müller, to the results of whose labors in this field but little was added by others, until Professor Ehrenberg entered upon the investigation with the advantage of greatly improved instruments. It was at about the same period with Müller, that Vaucher, a Genevese botanist, systematically applied the Microscope to the investigation of the lower forms of Vegetable life; and made many curious discoveries in regard both to their structure and to the history of their lives. He was the first to notice the extraordinary phenomenon of the spontaneous movement of the *Zoospores* of the humbler aquatic plants, which is known to be the means provided by nature for the dispersion of the race (Chaps. VI, VII); but being possessed with the idea (common to all Naturalists of that period and still very generally prevalent) that *spontaneous motion* evinces *Animal life*, he interpreted the facts which he observed, as indicating the existence of a class of beings which are Plants at one phase of their lives, and animals at another,—a doctrine which has since been completely set aside by the advance of physiological knowledge. Notwithstanding this and other errors of *interpretation*, however, the work of Vaucher on the “Fresh-water Confervæ” contains such a vast body of accurate *observation* on the growth and reproduction of the Microscopic Plants to the study of which he devoted himself, that it is quite worthy to take rank with that of Trembley, as having laid

¹ The structures to which this term is now *scientifically* restricted, are really vegetable.

the foundation of all our scientific knowledge of these very interesting forms. Although the curious phenomenon of "conjugation" had been previously observed by Müller, yet its connection with the function of Reproduction had not been even suspected by him; and it was by Vaucher that its real import was first discerned, and that its occurrence (which had been regarded by Müller as an isolated phenomenon, peculiar to a single species) was found to be common to a large number of humble aquatic forms of vegetation. But little advance was made upon the discoveries of Vaucher in regard to these, save by addition to the number of their specific forms, until a fresh stimulus had been given to such investigations by the improvement of the instrument itself. At present, they are among the most favorite objects of study among a large number of observers, both in this country and on the Continent; and are well deserving of the attention which they receive.

Less real progress seems to have been made in Microscopic inquiry, during the first quarter of the present century, than during any similar period since the invention of the instrument. The defects inseparable from its original construction, formed a bar to all discovery beyond certain limits; and although we are now continually meeting with new wonders, which patient and sagacious observation would have detected at any time and with any of the instruments then in use, yet it is not surprising that the impression should have become general, that almost everything which it *could* accomplish *had* already been done. The instrument fell under a temporary cloud from another cause; for having been applied by Anatomists and Physiologists to the determination of the elementary structure of the animal body, their results were found to be so discordant, as to give rise to a general suspicion of a want of trustworthiness in the Microscope, and in everything announced upon its authority. Thus both the instrument and its advocates were brought into more or less discredit; and as they continue to lie under this, in the estimation of many, to the present day, it will be desirable to pause here for awhile, to inquire into the sources of that discrepancy, to consider whether it is avoidable, and to inquire how far it should lead to a distrust of Microscopic observations, carefully and sagaciously made, and accurately recorded.

It is a tendency common to *all* observers, and not by any means peculiar to Microscopists, to describe what they *believe* and *infer*, rather than what they actually *witness*. The older Microscopic observers were especially liable to fall into this error; since the want of definiteness in the images presented to their eyes, left a great deal to be completed by the imagination. And when, as frequently happened, Physiologists began with *theorizing* on the elementary structure of the body, and allowed themselves to twist their imperfect observations into accordance with their theories, it was not surprising that their accounts of what

they professed to have seen should be extremely discordant. But from the moment that the visual image presented by a well-constructed Microscope, gave almost as perfect an idea of the object, as we could have obtained from the sight of the object itself, if enlarged to the same size and viewed with the unassisted eye, Microscopic observations admitted of nearly the same certainty as observations of any other class; it being only in a comparatively small number of cases, that a doubt can fairly remain about any question of *fact*, as to which the Microscope can be expected to inform us.¹

Another fallacy, common like the last to all observations, but with which the Microscopic observations of former times were perhaps especially chargeable, arises from a want of due attention to the *conditions* under which the observations are made. Thus one observer described the Human Blood-corpuscles as flattened disks resembling pieces of money, another as slightly concave on each surface, a third as slightly convex, a fourth as highly convex, and a fifth as globular; and the former prevalence of the last opinion, is marked by the habit which still lingers in popular phraseology, of designating these bodies as "blood-globules." Yet all microscopists are now agreed, that their real form, when examined in freshly drawn blood, is that of circular disks, with slightly concave surfaces; and the diversity in previous statements was simply due to the alteration effected in the shape of these disks, by the action of water or other liquids added for the sake of dilution; the effect of this being to render their surfaces first flat, then slightly convex, then more highly convex, at last changing their form to that of perfect spheres. But microscopical inquiries are not in themselves more liable to fallacies of this description, than are any other kinds of scientific investigation; and it will always be found here, as well as elsewhere, that—good instruments and competent observers being presupposed—the accordance in *results* will be precisely proportional to the accordance of *conditions*, that is, to the similarity of the objects, the similarity of the treatment to which they may be subjected, and the similarity of the mode in which they may be viewed.²

The more completely, therefore, the statements of Microscopic observers are kept free from those fallacies, to which observations of *any* kind are liable, wherein due care has not been taken

¹ One of the most remarkable of the *questiones vexatæ* at present agitated, is the nature of the markings on the siliceous valves of *Diatomacæ* (Chap. VI); some observers affirming those spots of the surface to be *elevations*, which others consider to be *depressions*. The difference is here one of *interpretation*, rather than of direct *observation*; the nature of the case preventing that kind of view of the object, which could leave no doubt as to the fact; and the conclusion formed being one of inference from a variety of appearances, which will differently impress the minds of different individuals.

² In objects of the most difficult class, such as the *Diatomacæ*, this last point is one of fundamental importance; very different appearances being presented by the *same object*, according to the mode in which it is illuminated, and the focal adjustment of the object-glass under which it is examined.—See Chap. VI.

to guard against them, the more completely will it be found that an essential agreement exists among them all, in regard to the *facts* which they record. And although the influence of *preconceived theories* still too greatly modifies, in the minds of some, the descriptions they profess to give of the facts actually presented to their visual sense, yet on the whole it is remarkable to what a unity of doctrine the best Microscopists of all countries are converging, in regard to all such subjects of this kind of inquiry, as have been studied by them with adequate care and under similar conditions. Hence it is neither fair to charge upon the Microscopists of the present day the errors of their predecessors; nor is it just to lay to the account of the instrument, what entirely proceeds from the fault of the observer, in recording, not what he sees in it, but what he fancies he can see.

It was at the commencement of the second quarter of the present century, that the principle of *Achromatic correction*, which had long before been applied to the Telescope, was first brought into efficient operation in the construction of the Microscope; for although its theoretical possibility was well known, insuperable difficulties were believed to exist in its practical application. The nature of this most important improvement will be explained in its proper place (Chap. I); and at present it will be sufficient to say, that within eight or ten years from the date of its first introduction, the character of the Microscope had been in effect so completely transformed, that it became an altogether new instrument; and from being considered but little better than a scientific toy, it soon acquired the deserved reputation of being one of the most perfect instruments ever devised by Art for the investigation of Nature. To this reputation it has a still greater claim at the present time; and though it would be hazardous to deny the possibility of any further improvement, yet the statements of theorists as to what *may* be accomplished, are so nearly equalled by what *has* been effected, that little room for improvement can be considered to remain, unless an entirely new theory shall be devised, which shall create a new set of possibilities.

Neither Botanists or Zoologists, Anatomists or Physiologists, were slow to avail themselves of the means of perfecting and extending their knowledge, thus unexpectedly put into their hands; and the records of Scientific Societies, and the pages of Scientific Journals, have ever since teemed, like the early Transactions of the Royal Society, with discoveries made by its instrumentality. All really philosophic inquirers soon came to feel, how vastly the use of the improved Microscope must add to their insight into every department of Organic Nature; and numbers forthwith applied themselves diligently to the labor of investigation. Old lines of research, which had been abandoned as unlikely to lead to any satisfactory issue, were taken up again with the confident expectation of success, which the result has shown to have

been well grounded; and new paths were soon struck out, each of which, leading into some region previously unexplored, soon cleared the way to others which became alike productive; thus laying open an almost unlimited range of inquiry, which the time that has since elapsed has served rather to extend than to contract, and which the labor that has been devoted to it has rather amplified than exhausted. A slight sketch of what has already been accomplished by the assistance of the Microscope, in the investigation of the phenomena of Life, seems an appropriate Introduction to the more detailed account of the instrument and its uses, which the present Treatise is designed to embrace.

The comparative simplicity of the structure of Plants, and the relatively large scale of their elementary parts, had allowed the Vegetable Anatomist, as we have seen, to elucidate some of its most important features, without any better assistance than the earlier Microscopes were capable of supplying. And many of those humbler forms of Cryptogamic vegetation, which only manifest themselves to the unaided eye when by their multiplication they aggregate into large masses, had been made the objects of careful study, which had yielded some most important results. Hence there seemed comparatively little to be done by the Microscopist in Botanical research; and it was not immediately perceived what was the direction in which his labors were likely to be most productive. Many valuable memoirs had been published, from time to time, on various points of vegetable structure; the increased precision and greater completeness of which, bore testimony to the importance of the aid which had been afforded by the greater efficiency of the instrument employed in such researches. But it was when the attention of Vegetable Physiologists first began to be prominently directed to the history of *development*, as the most important of all the subjects which presented themselves for investigation, that the greatest impulse was given to Scientific Botany; and its subsequent progress has been largely influenced by that impulse, both in the accelerated rate at which it has advanced, and in the direction which it has taken. Although Robert Brown had previously observed and recorded certain phenomena of great importance, yet it is in the Memoir of Prof. Schleiden, first published in 1837, that this new movement may be considered to have had its real origin; so that, whatever may be the errors with which his statements (whether on that occasion, or subsequently) are chargeable, there cannot be any reasonable question as to the essential service he has rendered to science, in pointing out the way to others, on whose results greater reliance may be placed. It was by Schleiden that the fundamental truth was first broadly enunciated, that as there are as many among the lowest orders of Plants, in which a single cell constitutes the entire individual, each living *for* and *by* itself alone, so each of the cells, by the aggregation of

which any individual among the higher Plants consists, has an independent life of its own, besides the "incidental" life which it possesses as a part of the organism at large: and that the doctrine was first proclaimed, that the *life-history of the individual cell* is therefore the very first and absolutely indispensable basis, not only for Vegetable Physiology, but (as was even then foreseen by his far-reaching mental vision) for Comparative Physiology in general. The first problem, therefore, which he set himself to investigate, was—*how does the cell itself originate?* It is unfortunate that he should have had recourse for its solution, to some of those cases in which the investigation is attended with peculiar difficulty, instead of making more use of the means and opportunities which the "single-celled" plants afford; and it is doubtless in great part to this cause, that we are to attribute certain fallacies in his results, of which subsequent researches have furnished the correction.

In no department of Botany, has recent Microscopy been more fertile in curious and important results, than in that which relates to the humblest forms of *Cryptogamia* that abound not only in our seas, rivers, and lakes, but even more in our marshes, pools, and ditches. For, in the first place, these present us with a number of most beautiful and most varied *forms*, such as on that account alone are objects of great interest to the Microscopist; this is especially the case with the curious group (ranked among Animalcules by Prof. Ehrenberg), which, from the bipartite form of their cells, has received the designation of *Desmidiæ*. In another group, that of *Diatomaceæ* (also still regarded as Animalcules, not only by Ehrenberg, but by many other Naturalists), not only are the forms of the plants often very remarkable, but their surfaces exhibit *markings* of extraordinary beauty and symmetry, which are among the best "test-objects" that can be employed for the higher powers of the instrument (Chap. IV): moreover, the membrane of each cell being coated externally with a film of silica, which not only takes its form, but receives the impress of its minutest markings, the siliceous skeletons remain unchanged after the death of the plants which formed them, sometimes accumulating to such an amount, as to give rise to deposits of considerable thickness at the bottoms of the lakes or pools which they inhabit; and similar deposits, commonly designated as beds of "fossil animalcules," are not unfrequently found at a considerable distance from the surface of the ground, on the site of what must have probably once been a lake or estuary, occasionally extending over such an area, and reaching to such a depth, as to constitute no insignificant part of the crust of the globe.—It is not only in these particulars, however, that the foregoing and other humble Plants have special attractions for the Microscopist; for the study of their *living actions* brings to view many phenomena, which are not only well calculated to excite the interest of those who find their chief pleasure

in the act of observing, but are also of the highest value to the Physiologist, who seeks to determine from the study of them what are the acts wherein Vitality may be said essentially to consist, and what are the fundamental distinctions between Animal and Vegetable life. Thus it is among these plants, that we can best study the history of the multiplication of cells by "binary subdivision," which seems to be the most general mode of growth and increase throughout the Vegetable kingdom; and it is in these, again, that the process of sexual generation is presented to us under its simplest aspect, in that curious act of "conjugation" to which reference has already been made (p. 39). But further, nearly all these Plants have at some period or other of their lives, a power of spontaneous *movement*; which in many instances so much resembles that of Animalcules, as to *seem* unmistakably to indicate their animal nature, more especially as this movement is usually accomplished by the agency of visible *cilia*: and the determination of the conditions under which it occurs, and of the purposes it is intended to fulfil, is only likely to be accomplished after a far more extensive as well as more minute study of their entire history, than has yet been prosecuted, save in a small number of instances. It is not a little remarkable, that in several of the cases in which the life-history of these plants has been most completely elucidated, they have been found to present a great variety of forms and aspects at different periods of their existence, and also to possess several different methods of reproduction; and hence it can be very little doubted, that numerous forms which are commonly reputed to be distinct and unrelated species, will prove in the end to be nothing else than successive stages of one and the same type. One of the most curious results attained by Microscopic inquiry of late years, has been the successive transfer of one group of reputed Animalcules after another, from the Animal to the Vegetable side of the line of demarcation between the two kingdoms; and although, as to the precise points across which this line should be drawn, there is not yet a unanimous agreement, yet there is now an increasing accord as to its general situation, which, even a few years since, was energetically canvassed. Those who are acquainted with the well-known *Volvox* (commonly termed the "globe-animalcule") will be surprised to learn that this, with its allies, constituting the family *Volvocineæ*, is now to be considered as on the *Vegetable* side of the boundary. (On the subjects of this paragraph, see Chap. VI.)

Not only this lowest type of Vegetable existence, but the *Cryptogamic* series as a whole, has undergone of late years a very close scrutiny, which has yielded results of the highest importance; many new and curious forms having been brought to light (some of them in situations in which their existence might have been least anticipated), and some of the most obscure portions of their history having received an unexpectedly clear eluci-

dation. Thus the discovery was announced by M. Audouin in 1837, that the disease termed *muscardine*, which annually carried off large numbers of the silkworms bred in the south of France, really consists in the growth of a fungous vegetation in the interior of their bodies, the further propagation of which may be almost entirely prevented by appropriate means; in the succeeding year, the fact was brought forward by several Microscopists, that *yeast* also is composed of vegetable cells, which grow and multiply during the process of fermentation; and subsequent researches have shown that the bodies of almost all animals, not even excepting Man himself, are occasionally infested by Vegetable as well as by animal Parasites, many of them remarkable for their beauty of configuration, and others for the variety of the forms they assume. The various parasites which attack our cultivated plants, again,—such as the “blights” of corn, the potato fungus, and the vine fungus,—have received a large measure of attention from Microscopists, and much valuable information has been collected in regard to them. It is still a question, however, which has to be decided upon other than microscopic evidence, how far the attacks of these fungi are to be considered as the *causes* of the diseases to which they stand related, or whether their presence (as is undoubtedly the case in many parallel instances) is the *effect* of the previously unhealthy condition of the plants which they infest; the general evidence appears to the author to incline to the latter view.

Of all the additions which our knowledge of the structure and life-history of the higher types of Cryptogamic vegetation has received, since the achromatic microscope has been brought to bear upon them, there is none so remarkable as that which relates to their Reproductive function. For the existence in that group of anything at all corresponding to the *sexual* generation of Flowering Plants, was scarcely admitted by any Botanists; and those few who did affirm it were unable to substantiate their views by any satisfactory proof, and were (as the event has shown) quite wrong as to the grounds on which they based them. Various isolated facts, the true meaning of which was quite unrecognized, had been discovered from time to time,—such as the existence of the moving filaments now termed “antherozoids,” in the “globules” of the *Chara* (first demonstrated by Mr. Varley in 1834), and in the “antheridia” of *Mosses* and *Liverworts* (as shown by Unger and Meyen in 1837), and the presence of “antheridia” upon what had been always previously considered the embryo-frond of the *Ferns* (first detected by Nageli in 1844); but of the connection of these with the generative function, no valid evidence could be produced, and the sexual reproduction of the Cryptogamia was treated by many Botanists of the greatest eminence, as a doctrine not less chimerical, than the doctrine of the sexuality of Flowering Plants had appeared to be to the opponents of Linnæus. It was by the admirable researches

of Count Suminski upon the development of the Ferns (1848), that the way was first opened to the right comprehension of the reproductive process in that group; and the doctrine of the fertilizing powers of the "antherozoids," once established in a single case, was soon proved to apply equally well to many others. Not a year has since elapsed, without the production of new evidence of the like sexuality in the several groups of the Cryptogamic series; this having been especially furnished by Hofmeister in regard to the higher types, by Thuret and Decaisne as to the marine Algæ, and by Tulasne with respect to Lichens and Fungi; and the doctrine may now be considered as established beyond the reach of cavil from any but those, who, having early committed themselves dogmatically to the negative opinion, have not the candor to allow due weight to the evidence on the affirmative side. With the study of the Reproduction of these plants, that of the history of their *Development* has naturally been connected; and some of the facts already brought to light, especially by the study of certain forms of Fungous vegetation, demonstrate the extreme importance of this inquiry in settling the foundations of Classification. For whereas the arrangement of Fungi, as of other Plants, has been based upon the characters furnished by their fructification, these characters have been found by Tulasne to be frequently subject to variations so wide, that one and the same *individual* shall present two or more kinds of fructification, such as had been previously considered to be peculiar to distinct *orders*. In this department of study, which has been scarcely at all cultivated by Microscopists of our own country, there is a peculiarly wide field for careful and painstaking research, and a sure prospect of an ample harvest of discovery. (On the subjects of the two preceding paragraphs, see Chap. VII.)

Although it has been in Cryptogamic Botany, that the zealous pursuit of Microscopic inquiry has been most conducive to scientific progress, yet the attention of Vegetable Anatomists and Physiologists has been also largely and productively directed to the minute structure and life-history of Flowering Plants. For although some of the general features of that structure had been made out by the earlier observers, and successive additions had been made to the knowledge of them, previously to the new era to which reference has so often been made, yet all this knowledge required to be completed and made exact, by a more refined examination of the Elementary Tissues than was before possible; and little was certainly known in regard to those processes of growth, development, and reproduction, in which their activity as living organisms consists. All the researches which have been made upon this point, tend most completely to bear out the general doctrine so clearly set forth by Schleiden, as to the independent vitality of each integral part of the fabric; and among the most curious results of the inquiries which have been

prosecuted in this direction, may be mentioned the discovery, that the movement of "rotation" of the *protoplasma* (or the viscid granular fluid at the expense of which the nutritive act seems to take place) within the cells, which was first observed by the Abbé Corti in the *Chara* in 1776, is by no means a unique or exceptional case; for that it may be detected in so large a number of instances, among Phanerogamia no less than among Cryptogamia, as apparently to justify the conclusion that it takes place in Vegetable cells generally, at some period or other of their evolution. In studying the phenomena of Vegetable Nutrition, the Microscope has been most effectually applied, not merely to the determination of changes in the form and arrangement of the elementary parts, but also to the detection of such changes in their composition, as ordinary Chemistry would be quite at fault to discover; each individual cell being (so to speak) a laboratory in itself, within which a transformation of organic compounds is continually taking place, not only for its own requirements, but for those of the economy at large; and these changes being at once made apparent by the application of chemical reagents to microscopic specimens whilst actually under observation. Hence the Vegetable Physiologist finds, in this Microscopic Chemistry, one of his most valuable means of tracing the succession of the changes in which Nutrition consists, as well as of establishing the chemical nature of particles far too minute to be analyzed in the ordinary way; and he derives further assistance in the same kind of investigation, from the application of Polarized Light (§ 63), which immediately enables him to detect the presence of mineral deposits, of starch-granules, and of certain other substances which are peculiarly affected by it. One of the most interesting among the general results of such researches, has been the discovery that the *true* cell-wall of the Plant (the "primordial utricle" of Mohl) has the same *albuminous* composition as that of the Animal; the external *cellulose* envelope, which had been previously considered as the distinctive attribute of the Vegetable cell, being in reality but a secretion from its surface. Of all the applications of the Microscope, however, to the study of the life-history of the Flowering plant, there is none which has excited so much interest, or given rise to so much discussion, as the nature of the process by which the *Ovule* is *fecundated* by the penetration of the pollen-tube. This question, in the opinion of the author, may be considered as now determined; and the conclusion arrived at is one so strictly in harmony with the general results obtained by the study of the (apparently) very different phenomena presented by the Generative process of the Cryptogamia, that it justifies the Physiologist in advancing a general doctrine as to the nature of the function, which proves to be no less applicable to the Animal kingdom than it is to the Vegetable. (See Chap. VIII.)

Among the objects of interest so abundantly offered by the

Animal Kingdom to the observation of Microscopists furnished with vastly improved instruments of research, it was natural that those minuter forms of Animal life which teem in almost every stationary collection of water, should engage their early attention; and among those Naturalists who applied themselves to this study, the foremost rank must undoubtedly be assigned to the celebrated German Microscopist, Prof. Ehrenberg. For although it is now unquestionable that he has committed numerous errors,—many doctrines which at first gained considerable currency on the strength of his high reputation, having now been abandoned by almost every one save their originator,—yet when we look at the vast advances which he unquestionably made in our knowledge of Animalcular life, the untiring industry which he has displayed in the study of it, the impulse which he has given to the investigations of others, and the broad foundation which he has laid for their inquiries in the magnificent works in which his own observations are recorded, we cannot but feel that his services have been almost invaluable, since, but for him, this department of microscopic inquiry would certainly have been in a position far behind that to which it has now advanced. Yet, great as has been the labor bestowed by him and by his followers in the same line of pursuit, it has become increasingly evident of late years that our knowledge of Infusory Animalcules is still in its infancy; that the great fabric erected by Prof. Ehrenberg rests upon a most insecure foundation; and that the Anatomy, Physiology, and systematic arrangement of these beings need to be restudied completely, *ab initio*. For, in the first place, there can be no doubt whatever, that a considerable section of the so-called Animalcules belongs to the Vegetable kingdom; consisting, as already pointed out, of the *motile* forms of the humbler Plants, of which a very large proportion pass, at some period of their existence, through a stage of activity that serves for their diffusion. Moreover, in another group, whose character has been entirely misconceived by the great German Microscopist, and was first clearly discriminated by M. Dujardin, there is neither mouth nor stomach of any kind; the minute plants and animals which serve it as food, being incorporated, as it were, with the soft animal jelly, which constitutes the almost homogeneous body; and this jelly further extending itself into “pseudopodial” prolongations, whereby these alimentary particles are laid hold of and drawn in. It was by the same distinguished French Microscopist, that the important fact was first discovered, that animals of this *Rhizopod* type are really the fabricators of those minute shells, which, from their Nautilus-like aspect, had been previously regarded as belonging to the highest class of the Molluscous Sub-Kingdom; and the whole of this most interesting group, which had received from M. D’Orbigny (who first perceived the speciality of its nature, and made a particular study of it) the designation of *Foraminifera*, has thus had its place in

the Animal scale most strangely reversed; being at once degraded from a position but little removed from Vertebrated animals, to a level in some respects even lower than that of the ordinary Animalcules.

But even when Prof. Ehrenberg's class of *Polygastrica* has been thus reduced, by the removal of those forms which are true Plants, and by the detachment of such as belong to the Rhizopod group, we find that our knowledge of its real nature is almost wholly *to be gained*; since little else has yet been accomplished than a description of a multitude of *forms*, of whose *history* as living beings scarcely anything else is known, than that they take food into the interior of their bodies by means of an oral orifice, that they digest this food and appropriate it to their own growth, and that they multiply themselves by binary subdivision. Now there is a very strong analogical probability, that many even of the most dissimilar forms of these Animalcules will prove to be different states of one and the same; for their multiplication by binary subdivision being not a true generative process, but being merely (so to speak) the growth of the individual, we may be almost certain that sooner or later a new phase will present itself, consisting in the evolution of proper sexual bodies, which will perform a true generative act, the products of which may be very probably quite different from the forms we are accustomed to regard as peculiar to each species. The attention of several eminent Microscopists at the present time is strongly fixed upon this part of the inquiry; which can only be efficiently prosecuted, by *limiting* the range of observation for a time to a *small number* of forms, and pursuing these through all the phases of their existence.

Among the most important of Prof. Ehrenberg's unquestioned discoveries, we are undoubtedly to place that of the comparatively high organization of the *Rotifera*, or Wheel-Animalcules and their allies; for which, though previously confounded with a similar Infusoria, he asserted and vindicated a claim to a far more elevated rank. For although in this instance, too, some of his descriptions have been shown to be incorrect, and many of his inferences to be erroneous, and although subsequent observers are not agreed among themselves as to many important particulars, yet all assent to the general accuracy of Prof. Ehrenberg's statements, and recognize the title of the *Rotifera* to a place not far removed from that of the Vermiform tribes. A parallel discovery was made about the same time by MM. Audouin and Milne Edwards, in regard to the *Flustra* and their allies, which had previously ranked among those flexible Zoophytes popularly known as "corallines," and are often scarcely to be distinguished from them in mode of growth or general aspect;¹ but which

¹ "You go down," says Mr. Kingsley, "to any shore after a gale of wind, and pick up a few delicate little sea-ferns. You have two in your hand (*Sertularia operculata* and *Genellaria loriculata*), which probably look to you, even under a good pocket magnifier, identical or nearly so. But you are told, to your surprise, that however like the

were separated as a distinct order by these observers, on account of their possession of a second orifice to the alimentary canal, and the general tendency of their plan of organization to that which characterizes the inferior Mollusca. The importance of this distinction was at once recognized; and the group received the designation of *Polyzoa* from Mr. J. V. Thompson, and of *Bryozoa* from Prof. Ehrenberg. The organization of this very interesting group was further elucidated, some years subsequently, by the admirable observations of Dr. Arthur Farre upon a newly-discovered form (named by him *Bowerbankia*), the transparency of whose envelopes allowed its internal structure to be distinctly made out; and the additional features which he detected, were all such as to strengthen the idea already entertained of its essentially Molluscan character. This idea received its final and complete confirmation from the admirable researches of M. Milne Edwards on the *Compound Ascidians*, which are the lowest animals whose Molluscan nature had been previously acknowledged; these having been discovered by him to agree with Zoophytes in their plant-like attribute of extension by "gemmation" or budding, and to present, in all the most important features of their organization, an extremely close approximation to the Bryozoa. Thus whilst Microscopic research has degraded the Foraminifera from their supposed rank with the Nautilus and Cuttle-fish, to the level of the Sponge, it has raised the Wheel-Animalcules into proximity with the aquatic Worms, and the humble "Sea-Mat," formerly supposed to be a Plant, to a position not much below that of the Oyster and Mussel.¹

Another most curious and most important field of microscopic inquiry has been opened up in the discovery of the *Transformations* which a large proportion of the lower animals undergo, during the early stages of their existence; and notwithstanding that it has even yet been very imperfectly cultivated, the unexpected result has been already attained, that the fact of "metamorphosis,"—previously known only in the cases of Insects and Tadpoles, and commonly considered as an altogether *exceptional* phenomenon,—is almost *universal* among the inferior tribes; it being a rare occurrence for the offspring to come forth from the egg in a condition bearing any resemblance to that which characterizes the adult, and the latter being in general attained only after a long series of changes, in the course of which many curious phases are presented. One of the earliest and most remarkable discoveries which was made in this direction,—that of the metamorphosis of the *Cirrhipeds* (Barnacles and their allies) by Mr. J. V. Thompson,—proved of most important assistance in the determination of the true place of that group, which had previously been a matter of controversy; for although in their outward

dead horny polypidoms which you hold may be, the two species of animal which have formed them, are at least as far apart in the scale of creation as a quadruped is from a fish."

¹ With reference to the subjects of the three preceding paragraphs, see Chaps. IX, X, XI.

characters they bear such a resemblance to Mollusks, that the Barnacles which attach themselves to floating timber, and the Acorn-shells which incrust the surfaces of rocks, are unhesitatingly ranked by Shell-collectors among their "multivalves," yet the close resemblance which exists between their early forms and the little Water-Fleas which swarm in our pools, makes it quite certain that the Barnacles not only belong to the Articulated instead of to the Molluscous series, but that they must be ranked in close proximity to the Entomostracous division of the Crustacea, if not actually as members of it. To the same discoverer, moreover, we owe the knowledge that even the common *Crab* undergoes metamorphoses scarcely less strange, its earliest form being a little creature of most grotesque shape, which had been previously described as an adult and perfect Entomostracan; so that, although scarcely any two creatures can apparently be more unlike than a Barnacle and a Crab, they have (so to speak) the same starting-point; the difference in their ultimate aspect chiefly arising from the difference in the proportionate development of parts which are common to both. A still more remarkable series of Metamorphoses has recently been shown by Prof. Müller to exist among the *Echinoderms* (Star-fish, Sea-urchins, &c.); whose development he has studied with great perseverance and sagacity. Thus the larva of the Star-fish is an active free-swimming animal, having a long body with six slender arms on each side, from one end of which the young star-fish is (so to speak) budded off; and when this has attained a certain stage of development, the long twelve-armed body separates from it and dies off, its chief function having apparently been, to carry the young Star-fish to a distance from its fellows, and thus to prevent overcrowding by the accumulation of individuals in particular spots, which would be liable to occur if they never had any more active powers of locomotion than they possess in their adult state. Scarcely less remarkable are the changes which are to be witnessed in the greater number of aquatic *Mollusks*, almost all of which, however inert in their adult condition, possess active powers of locomotion in their larval state; some being propelled by the vibratile movement of cilia disposed upon the head somewhat after the fashion of those of Wheel-animalcules, and others by the lateral strokes of a sort of tail which afterwards disappears like that of a tadpole. Among the *Annelids* or marine worms, again, there is found to be an extraordinary dissimilarity, though of a somewhat different nature, between the larval and the adult forms: for they commonly come forth from the egg in a condition but little advanced beyond that of Animalcules; and, although they do not undergo any metamorphosis comparable to that of Insects, they pass through a long series of phases of development (chiefly consisting in the successive production of new joints or segments, and of the organs appertaining to these) before they acquire their complete type. In nearly all the foregoing cases it may be re-

marked, that the larval forms of different species bear to one another a far stronger resemblance than exists among adults, the distinguishing characters of the latter being only evolved in the course of their development; and every new discovery in this direction only gives fresh confirmation to the great law of development early detected by the sagacity of Von Baer, that *the more special forms of structure arise out of the more general, and this by a gradual change*. The meaning of this law will become obvious hereafter, when some of the principal cases to which it applies shall have been brought in illustration of it (Chap. XII).

A still more curious series of discoveries has been made, by means of the Microscope, in regard to the early development of the Medusan *Acalephs* (jelly-fish, &c.), and the relationship that exists between them and the *Hydraform Zoophytes*;—two groups of animals, which had been previously ranked in different classes, and had not been supposed to possess anything in common. For it has been clearly made out by the careful observations of Sars, Siebold, Dalyell, and others, that those delicate arborescent Zoophytes, each polype of which is essentially a Hydra, not only grow by extending themselves into new branches, like plants, sometimes also budding off detached *gemmæ*, which multiply their kind by developing themselves into Zoophytic forms like those whence they sprang; but also produce peculiar buds having all the characters of *Medusæ*, which contain the proper generative organs of the Zoophyte, but which, usually detaching themselves from the stock that bore them, swim freely through the ocean as minute jelly-fish, without exhibiting the slightest trace of their originally attached condition. The *Medusæ* in due time produce fertile eggs; and each egg develops itself, not into the form of its immediate progenitor, but into that of the Zoophyte from which the *Medusa* was budded off. And thus a most extraordinary alteration of forms is presented, between the Zoophyte, which may be compared to the *growing* or *vegetating* stage of a Plant (its polypes representing the leaf-buds), and the *Medusa*, the development of which marks its *flowering* stage. So again, from the investigation of the early history of those larger forms of “jelly-fish” with which every visitor to the sea-coast is familiar, it has been rendered certain that they too are developed from Polype larvæ, usually of very minute size, which give off *Medusa* buds; so that whilst *they* are best known to us in their *Medusan* state, and the *Hydraform Zoophytes* in their polypoid state, each of these groups is the representative of a certain stage in the life-history of one and the same tribe of these curious beings, which, when complete, includes both states, as will hereafter be shown in more detail (Chap. XI). Changes very similar in kind, and in many respects even more remarkable, have been found by microscopic inquiry to take place among the *Entozoa* (intestinal worms); but being interesting only to professed Naturalists and scientific Physiologists, they scarcely call for particular notice in a treatise like the present.

It has not been among the least important results of the new turn which Zoological inquiry has thus taken, that a far higher spirit has been introduced into the cultivation of this science, than previously pervaded it. Formerly it was thought, both in Zoology and in Botany, that Classification might be adequately based on external characters alone; and the scientific acquirements of a Naturalist were estimated rather by the extent of his acquaintance with these, than by any knowledge he might possess of their internal organization. The great system of Cuvier, it is true, professed to rest upon organization as its basis; but the acquaintance with this which was considered requisite for the purpose, was very limited in its amount and superficial in its character; and no Naturalist formerly thought of studying the history of Development as a necessary adjunct to the Science of Classification. How essential a knowledge of it has now become, however, if only as a basis for any truly natural arrangement of Animals, must have become apparent from the preceding sketch; and it has thus come to be felt and admitted amongst all truly philosophic Naturalists, that the complete study of any particular group, even for the purposes of classification, involves the acquirement of a knowledge, not only of its intimate structure, but of its entire life-history. And thus Natural History and Physiology,—two departments of the great Science of Life, which the Creator inextricably blended, but which Man has foolishly striven to separate,—are now again being brought into their original and essential harmony; and it is coming to be thought more credible, to give a complete elucidation of the history of even a single species, than to describe any number of new forms, about which nothing else is made out save what shows itself on the surface.

Thus every Microscopist, however limited may be his opportunities, has a wide range of observation presented to him in the study of the lower forms of Animal life; with the strongest incitement to persevering and well-directed inquiry, that the anticipation of novelty, and the expectation of valuable results, can afford. For, notwithstanding the large number of admirable records which have been already published (chiefly, we must admit with regret, by Continental Naturalists) upon the developmental history of the lower tribes of Animals, there is no one of the subjects that have been just passed in review, of which the knowledge hitherto gained can be regarded as more than a sample of that which remains to be acquired. Records like those already referred to, might easily be multiplied a hundred fold, with infinite advantage to Science; if those Microscopists who spend their time in desultory observation, and in looking at some favorite objects over and over again, would but concentrate their attention upon some particular species or group, and work out its entire history with patience and determination. And the observer himself would find this great advantage in so doing,—that an inquiry

thus pursued gradually becomes to him an object of such attractive interest, that he experiences a zest in its pursuit to which the mere *dilettante* is an entire stranger,—besides enjoying all the mental profit, which is the almost necessary result of the *thorough* performance of *any* task that is not in itself unworthy. And what *can* be a more worthy occupation, than the attempt to gain an insight, however limited, into the operations of Creative Wisdom?—these being not less wonderfully displayed among the forms of Animal life which are accounted the simplest and least attractive, than in those which more conspicuously solicit the attention of the Student of Nature, by the beauty of their aspect or the elaborateness of their organization.

It has not been, however, in the study of the minuter forms of Animal life alone, that the Microscope has been turned to valuable account; for the Anatomists and Physiologists who had made the Human fabric the especial object of their study, and who had been led to believe that the knowledge accumulated by their repeated and persevering scrutiny into every portion accessible to their vision, was all which it lay within their power to attain, have found in this new instrument of research, the means of advancing far nearer towards the *penetralia* of Organization, and of gaining a much deeper insight into the mysteries of Life, than had ever before been conceived possible. For every part of the entire organism has been, so to speak, decomposed into its *elementary tissues*, the structure and actions of each of which have been separately and minutely investigated; and thus a new department of study, which is known as *Histology* (or Science of the Tissues), has not only been marked out, but has already made great advances towards completeness. In the pursuit of this inquiry, the Microscopists of our day have not limited themselves to the fabric of Man, but have extended their researches through the entire range of the Animal kingdom; and in so doing, have found, as in every other department of Nature, a combination of endless variety in detail, with a marvellous simplicity and uniformity of general plan. Thus the *bones* which constitute the skeleton of the Vertebrated animal,—however different from each other in their external configuration, in the arrangement of their compact and their cancellated portions, and other such particulars as specially adapt them for the purposes they have to perform in each organism,—all consist of a certain kind of tissue, distinguished under the microscope by features of a most peculiar and interesting kind; and these features, whilst presenting (like those of the Human countenance) a certain general conformity to a common plan, exhibit (as Prof. Quekett has shown) such distinctive modifications of that plan in the different classes and orders of the Vertebrated series, that it is generally possible by the microscopic examination of the merest fragment of a bone, to pronounce with great probability as to the *natural family* to which it has belonged. Still

more is this the case in regard to the *Teeth*, whose organic structure (originally detected by Leeuwenhoek) has been newly and far more completely elucidated by Profrs. Purkinje, Retzius, Owen, and Tomes; for the inquiry into the comparative structure of these organs, which has been prosecuted by Prof. Owen, in particular, through the entire range of the Vertebrated series, has shown that, with an equally close conformity to a certain *general* plan of structure, there are at the same time still wider diversities in detail, which are so characteristic of their respective groups, that it is often possible to discriminate, not only families, but even the genera and species, by careful attention to the minute features of their structure. Similar inquiries, with results in many respects analogous, have been carried out by the Author, in regard to the *Shells* of Mollusks, Crustaceans, and Echinoderms; his researches having not only demonstrated the regularly-organized structure of these protective envelopes (which had been previously affirmed to be mere inorganic exudations, presenting in many instances a crystalline texture), but having shown that many natural groups are so distinctively characterized by the microscopic peculiarities they present, that the inspection of a minute fragment of Shell will often serve to determine, no less surely than in the case of bones and teeth, the position of the animal of which it formed a part. The soft parts of the Animal body, moreover, such as the *cartilages* which cover the extremities of the bones and the *ligaments* which hold them together at the joints, the *muscles* whose contraction develops motion and the *tendons* which communicate that motion, the *nervous ganglia* which generate nervous force and the *nerve-fibres* which convey it, the *skin* which clothes the body and the *mucous* and *serous* membranes which line its cavities, the *assimilating glands* which make the blood and the *secreting glands* which keep it in a state of purity,—these, and many other tissues that might be enumerated, are severally found to present characteristic peculiarities of structure, which are more or less distinctly recognizable throughout the Animal series, and which bear the strongest testimony to the Unity of the Design in which they all originated. As we descend to the lower forms of Animal life, however, we find these distinctions less and less obvious; and we at last come to fabrics of such extreme simplicity and *homogeneousness*, that every part seems to resemble every other in structure and action; no provision being made for that “division of labor” which marks the higher types of organization, and which, being the consequence of the development of separate *organs* each having its special work to do, can only be effected where there is a “differentiation” of parts, that gives to the entire fabric a character of *heterogeneousness*.

The Microscopic investigation whose nature has thus been sketched, has not only been most fruitful in the discovery of individual facts, but has led to certain general results, of great

value in Physiological Science. Among the most important of these, is the complete metamorphosis which has been effected in the ideas previously entertained regarding *living action*; such having been essentially based on the Circulation of the blood, as the only vital phenomenon of which any direct cognizance could be gained through the medium of the senses. For it gradually came to be clearly perceived, that in the Animal as in the Plant, *each integral portion of the Organism possesses an independent Life of its own*, in virtue of which it performs a series of actions peculiar to itself, provided that the conditions requisite for those actions be supplied to it; and that the Life of the body as a whole (like a symphony performed by a full orchestra) consists in the harmonious combination of its separate instrumental acts, —the circulation of the blood, instead of *making the tissues*, simply affording the supply of prepared nutriment, at the expense of which they *evolve themselves* from germs previously existing. This general doctrine was first put prominently forwards by Schwann, whose “Microscopical Researches into the Accordance in the Structure and Growth of Animals and Plants,” published in 1839, marks the commencement of a new era in all that department of Animal Physiology, which comprises the simply vegetative life of the organized fabric. These researches, avowedly based upon the ideas advanced by Schleiden, were prosecuted in the same direction as his had been; the object which this admirable observer and philosophic reasoner specially proposed to himself, being the study of the *development* of the Animal tissues. He found that although their evolution cannot be watched while in actual progress, its history may be traced out by the comparison of the successive stages brought to light by Microscopic research; and in so far as this has been accomplished for each separate part of the organism, the structure and actions of its several components, however diverse in their fully developed condition, are found to resemble each other more and more closely, the more nearly these parts are traced back to their earliest appearance. Thus we arrive in our retrospective survey, at a period in the early history of Man, at which the whole embryonic mass is but *a congeries of cells*, all apparently similar and equal to each other; and going still further back, it is found that all these have had their origin in the subdivision of *a single primordial cell*, which is the first defined product of the generative act. On this single cell, the Physiologist bases his idea of the most elementary type of Organization; whilst his actions present him with all that is essential to the notion of Life. And in pursuing the history of the germ, from this, its simplest and most *homogeneous* form, to the assumption of that completed and perfected type which is marked by the extreme *heterogeneousness* of its different parts, he has another illustration of that law of *progress from the general to the special*, which is one of the highest principles yet attained in the science of Vitality.

But, further, the Physiologist, not confining his inquiries to Man, pursues the like researches into the developmental history of other living beings, and is soon led to the conclusion that the same is true of them also; each Animal, as well as each Plant, having the same starting-point in the single cell; and the distinctive features by which its perfected form is characterized, how striking and important soever these may be, arising in the course of its development towards the condition it is ultimately to present. In the progress of that evolution, those fundamental differences which mark out the great natural divisions of the Animal and the Vegetable kingdoms respectively, are the first to manifest themselves; and the subordinate peculiarities which distinguish classes, orders, families, genera, and species, successively make their appearance, usually (but not by any means constantly) in the order of importance which Systematists have assigned to them. And it is in thus pursuing, by the aid which the Microscope alone can afford to his visual power, the history of the Organic Germ, from that simple and homogeneous form which seems common to every kind of living being, either to that complex and most heterogeneous organism which is the mortal tenement of Man's immortal spirit, or only to that humble Protophyte or Protozoon, which lives and grows and multiplies without showing any essential advance upon its embryonic type,—that the Physiologist is led to his grandest conception of the Unity and All-Comprehensive nature of that Creative Design, of which the development of every individual Organism, from the lowest to the highest, is a separate exemplification, at once perfect in itself, and harmonious with every other.

It has been the purpose of the foregoing sketch, to convey an idea, not merely of the services which the Microscope has already rendered to the *collector of facts* in every department of the Science of Life, but also of the value of these facts as a foundation for *philosophical reasoning*. For it is when thus utilized, that observations, whether made with the Microscope or with the Telescope, or by any other instrumentality, acquire their highest value, and excite the strongest interest in the mind. But as it is not every one who is prepared by his previous acquirements to appreciate such researches, according to the scientific estimate of their importance, it may be well now to address ourselves to that large and increasing number, who are disposed to apply themselves to Microscopic research as *amateurs*, following the pursuit rather as a means of wholesome recreation to their own minds, than with a view to the extension of the boundaries of existing knowledge; and to those in particular who are charged, whether as parents or as instructors, with the direction and training of the youthful mind.

All the advantages which have been urged at various times,

with so much sense and vigor,¹ in favor of the study of Natural History, apply with full force to Microscopical inquiry. What better encouragement and direction can possibly be given to the exercise of the *observing* powers of a child, than to habituate him to the employment of this instrument upon the objects which immediately surround him, and then to teach him to search out novelties among those less immediately accessible? The more we limit the natural exercise of these powers, by the use of those methods of education which are generally considered to be specially advantageous for the development of the Intellect,—the more we take him from fields and woods, from hills and moors, from river-side and sea-shore, and shut him up in close school-rooms and narrow play-grounds, limiting his attention to abstractions, and cutting him off even in his hours of sport from those sights and sounds of Nature which seem to be the appointed food of the youthful spirit,—the more does it seem important that he should in some way be brought into contact with her, that he should have his thoughts sometimes turned from the pages of books to those of Creation, from the teachings of Man to those of God. Now if we attempt to give this direction to the thoughts and feelings in a merely *didactic* mode, it loses that spontaneousness which is one of its most valuable features. But if we place before the young a set of objects which can scarcely fail to excite their healthful curiosity, satisfying this only so far as to leave them still inquirers, and stimulating their interest from time to time by the disclosure of such new wonders as arouse new feelings of delight, they come to look upon the pursuit as an ever-fresh fountain of happiness and enjoyment, and to seek every opportunity of following it for themselves.

There are no circumstances or conditions of life, which need be altogether cut off from these sources of interest and improvement. Those who are brought up amidst the wholesome influences of a country life, have, it is true, the greatest direct opportunities of thus drawing from the Natural Creation the appropriate nurture for their own spiritual life. But the very familiarity of the objects around them, prevents these from exerting their most wholesome influence, unless they be led to see how much there is *beneath the surface* even of what they seem to know best; and in rightly training them to look for this, how many educational objects,—physical, intellectual, and moral,—may be answered at the same time! “A walk without an object,” says Mr. Kingsley, “unless in the most lovely and novel scenery, is a poor exercise; and as a recreation utterly nil. If we wish rural walks to do our children any good, we must give them a love for rural sights, an object in every walk; we must teach them—and we can teach them—to find wonder in every insect, sublimity in every hedge-row, the records of past worlds in every pebble, and

¹ By none more forcibly than by Mr. Kingsley, in his recent little volume entitled “Glaucus, or the Wonders of the Shore.”

boundless fertility upon the barren shore; and so, by teaching them to make full use of that limited sphere in which they now are, make them faithful in a few things, that they may be fit hereafter to be rulers over much." What can be a more effectual means of turning such opportunities to the best account, than the employment of an aid which not only multiplies almost infinitely the sources of interest presented by the objects with which our eyes are most familiar, but finds inexhaustible life where all seems lifeless, ceaseless activity where all seems motionless, perpetual change where all seems inert? Turn, on the other hand, to the young who are growing up in our great towns, in the heart of the vast Metropolis, whose range of vision is limited on every side by bricks and mortar, who rarely see a green leaf or a fresh blade of grass, and whose knowledge of animal life is practically limited to the dozen or two of creatures that everywhere attach themselves to the companionship of Man, and shape their habits by his. To attempt to inspire a real love of Nature by books and pictures, in those who have never felt her influences, is almost hopeless. A child may be interested by accounts of her wonders, as by any other instructive narrative; but they have little of *life* or *reality* in his mind,—far less than has the story of adventure which appeals to his own sympathies, or even than the fairy tale which charms and fixes his imagination. But here the Microscope may be introduced with all the more advantage, as being almost the only means accessible under such circumstances, for supplying what is needed. A single rural or even suburban walk will afford stores of pleasurable occupation for weeks, in the examination of its collected treasures. A large glass jar may be easily made to teem with life, in almost as many and as varied forms as could be found by the unaided eye in long and toilsome voyages over the wide ocean; and a never-ending source of amusement is afforded by the observation of their growth, their changes, their movements, their habits. The school-boy thus trained, looks forward to the holiday which shall enable him to search afresh in some favorite pool, or to explore the wonders of some stagnant basin, with as much zest as the keenest sportsman longs for a day's shooting on the moors, or a day's fishing in the best trout stream; and with this great advantage over him,—that his excursion is only the beginning of a fresh stock of enjoyment, instead of being in itself the whole.

This is no imaginary picture, but one which we have constantly under our eyes; and no argument can be needed to show the value of such a taste, to such, at least, as have set clearly before their minds the objects at which they should aim in the great work of Education. For we have not merely to train the intellectual powers and to develop the moral sense; but to form those tastes—those "likes and dislikes"—which exercise a more abiding and a more cogent influence on the conduct, than either the reason or the mere knowledge of duty. It is our object to foster

all the higher aspirations, to keep in check all that is low and degrading. But the mind *must* have recreation and amusement; and the more closely it is kept by the system of education adopted, to the exercise of any one set of powers, the more potent will be that reaction which will urge it, when restraint is removed, to activity of some other kind; and the more important is it, that this reaction should receive a direction to what is healthful and elevating, instead of to what is weakening and degrading. It is quite a mistake to imagine that those evil habits which result from a wrong exercise of the natural powers, a wrong direction of the natural tendencies, can be effectually antagonized by the simple effort at *repression*. The constant exercise either of external coercion or of internal restraint, tends to keep the attention directed towards the forbidden object of gratification; the malady is only held in check, not cured; and it will break out, perhaps with augmented force, whenever the perpetually-present impulses shall derive more than ordinary strength from some casual occurrence, or the restraining power shall have been temporarily weakened. The only effectual mode of keeping in check the *wrong*, is by making use of these same powers and tendencies in a *right* mode; by finding out objects whereon they may be beneficially exercised; and by giving them such a direction and encouragement, as may lead them to *expend themselves* upon these, instead of fretting and chafing under restraint, ready to break loose at the first opportunity. There is no object on which the youthful energy can be employed more worthily, than in the pursuit of Knowledge; no kind of knowledge can be made more attractive, than that which is presented by the Works of Creation; no source is more accessible, no fountain more inexhaustible; and there is none which affords, both in the mode of pursuing it, and in its own nature, so complete and beneficial a diversion from the ordinary scholastic pursuits.

If there be one class more than another, which especially needs to have its attention thus awakened to such objects of interest, as, by drawing its better nature into exercise, shall keep it free from the grovelling sensuality in which it too frequently loses itself, it is our Laboring population; the elevation of which is one of the great social problems of the day. On those who are actively concerned in promoting and conducting its education, the claims and advantages of the Study of Nature can scarcely be too strongly urged; since experience has fully proved,—what might have been *a priori* anticipated,—that where the taste for this pursuit has been *early* fostered by judicious training, it becomes so completely a part of the mind, that it rarely leaves the individual, however unfavorable his circumstances may be to its exercise, but continues to exert a refining and elevating influence through his whole subsequent course of life. Now for the reasons already stated, the Microscope is not merely a most valuable adjunct in such instruction, but its assistance is essential

in giving to almost every Natural object its highest educational value; and whilst the *country* Schoolmaster has the best opportunities of turning it to useful account, it is to the *city* Schoolmaster that, in default of other opportunities, its importance as an educational instrument should be the greatest. It was from feeling very strongly how much advantage would accrue from the introduction of a form of Microscope, which should be at once *good* enough for Educational purposes, and *cheap* enough to find its way into every well-supported School in town and country, that the Author suggested to the Society of Arts in the summer of 1854, that it should endeavor to carry out an object so strictly in accordance with the enlightened purposes which it is aiming to effect; and this suggestion having been considered worthy of adoption, a Committee, chiefly consisting of experienced Microscopists, was appointed to carry it into effect. It was determined to aim at obtaining two instruments;—a *simple* and low-priced microscope for the use of Scholars, to whom it might be appropriately given as a reward for zeal and proficiency in the pursuit of Natural History, not in books, but in the field;—and a *compound* Microscope for the use of Teachers, of capacity sufficient to afford a good view of every kind of object most likely to interest the pupil or to be within the reach of the instructor. Notwithstanding the apprehensions generally expressed, that no instruments at all likely to answer the intended purpose could possibly be produced at the prices specified, the result has proved the fallacy; for among several instruments of greater or less efficiency, sent in competition for the award, the Committee was able to select a Simple and a Compound Microscope fully answering their expectations, and henceforth to be supplied to the public at a cost so low as to place these instruments (it may be hoped) within the reach of almost every one to whom they are likely to be of service. An account of these two Microscopes will be given hereafter. (Chap. II, §§ 29, 31.)

It is not alone, however, as furnishing an attractive object of pursuit for the young—fitted at once to excite a wholesome taste for novelty, ever growing with what it feeds on, and to call forth the healthful exercise of all those powers, both physical and mental, which can minister to its gratification,—that Natural History studies in general, and Microscopic inquiry in particular, are to be specially commended as a means of intellectual and moral discipline; for there is no capacity, however elevated, to which they do not furnish ample material for the exercise of all its best powers, no period of life which may not draw from them its purest pleasures. Even to *observe well* is not so easy a thing as some persons imagine. Some are too hasty, imagining that they can take in everything at a glance, and hence often forming very erroneous or imperfect notions, which may give an entirely wrong direction not only to their own views but to those of others, and may thus render necessary an amount

of labor for the ultimate determination of the truth, many times as great as that which would have sufficed in the first instance, had the original observations been accurately made and faithfully recorded. Others, again, are too slow and hesitating; and fix their attention too much upon details, to be able to enter into the real significance of what may be presented to the vision. Although ignorance has doubtless much to do in producing both these faults, yet they both have their source in mental tendencies which are not corrected by the mere acquisition of knowledge, and which are very inimical, not merely to its fair reception, but also to the formation of a sound judgment upon any subject whatever. The habit of guarding against them, therefore, once acquired in regard to Microscopic observation, will be of invaluable service in every walk of life. Not less important is it (as has been already shown), to keep our observations free alike from the bias of preconceived ideas, and from the suggestive influence of superficial resemblances; and here, too, we find the training which Microscopical study affords, especially when it is prosecuted under the direction of an experienced guide, of the highest value in forming judicious habits of thought and action. To set the young observer to examine and investigate for himself, to tell him merely *where* to look and (in general terms) *what* to look for, to require from him a careful account of what he sees, and then to lead him to compare this with the descriptions of similar objects by Microscopists of large experience and unquestionable accuracy, is not only the best training he can receive as a Microscopist, but one of the best means of preparing his mind for the exercise of its powers in any sphere whatever.

It cannot be too strongly or too constantly kept in view, that the value of the results of Microscopic inquiry will depend far more upon the sagacity, perseverance, and accuracy of the observer, than upon the elaborateness of his instrument. The most perfect Microscope ever made, in the hands of one who knows not how to turn it to account, is valueless; in the hands of a careless, a hasty, or a prejudiced observer, it is worse than valueless, as furnishing new contributions to the already large stock of errors that pass under the guise of scientific truths. On the other hand, the least costly Microscope that has ever been constructed, how limited soever its powers, provided that it gives no *false* appearances; shall furnish to him who knows what *may* be done with it, a means of turning to an account, profitable alike to science and to his own immortal spirit, those hours which might otherwise be passed in languid *ennui*, or in frivolous or degrading amusements,¹ and even of immortalizing his name by the discovery of secrets in Nature as yet undreamed of. A very

¹ "I have seen," says Mr. Kingsley, "the cultivated man, craving for travel and success in life, pent up in the drudgery of London work, and yet keeping his spirit calm, and his morals perhaps all the more righteous, by spending over his Microscope evenings which would too probably have gradually been wasted at the theatre."

large proportion of the great achievements of Microscopic research that have been noticed in the preceding outline, have been made by the instrumentality of microscopes which would be generally condemned in the present days as utterly unfit for any scientific purpose; and it cannot for a moment be supposed, that the field which Nature presents for the prosecution of inquiries with instruments of comparatively limited capacity, has been in any appreciable degree exhausted. On the contrary, what *has been* done by these and scarcely superior instruments, only shows how much there is *to be* done. The author may be excused for citing, as an apposite example of his meaning, the curious results he has recently obtained from the study of the development of the *Purpura lapillus* (rockwhelk), which will be detailed in their appropriate place (Chap. XII); for these were obtained almost entirely by the aid of *single lenses*, the Compound Microscope having been only occasionally applied to, for the verification of what had been previously worked out, or for the examination of such minute details as the power employed did not suffice to reveal.

But it should be urged upon such as are anxious to do service to science, by the publication of discoveries which they suppose themselves to have made with comparatively imperfect instruments, that they will do well to refrain from bringing these forward, until they shall have obtained the opportunity of verifying them with better. It is, as already remarked, when an object is *least* clearly seen, that there is *most* room for the exercise of the imagination; and there was sound sense in the reply once made by a veteran observer, to one who had been telling him of wonderful discoveries which another was said to have made "*in spite of the badness of his Microscope*,"—"No, sir, it was *in consequence of the badness of his Microscope*." If those who observe, with however humble an instrument, will but rigidly observe the rule of recording only what they can *clearly see*, they can neither go far astray themselves, nor seriously mislead others.

Among the erroneous tendencies which Microscopic inquiry seems especially fitted to correct, is that which leads to the estimation of things by their merely sensuous or material greatness, instead of by their value in extending our ideas and elevating our aspirations. For we cannot long scrutinize the "world of small" to which we thus find access, without having the conviction forced upon us, that all *size* is but relative, and that *mass* has nothing to do with real grandeur. There is something in the extreme of minuteness, which is no less wonderful,—might it not almost be said, no less majestic?—than the extreme of vastness. If the mind loses itself in the contemplation of the immeasurable depths of space, and of the innumerable multitudes of stars and systems by which they are peopled, it is equally lost in wonder and admiration, when the eye is turned to those countless multitudes of living beings which a single drop of

water may contain, and when the attention is given to the wondrous succession of phenomena which the life-history of every individual among them exhibits, and to the order and constancy which this presents. Still more is this the case, when we direct our scrutiny to the penetration of that universe which may be said to be included in the body of Man, or of any one of the higher forms of organized being, and survey the innumerable assemblage of elementary parts, each having its own independent life, yet each working in perfect harmony with the rest, for the completion of the wondrous aggregate which the life of the whole presents. In the study of the one class of phenomena, no less than in the survey of the other, we are led towards that Infinity, in comparison with which the greatest and the least among the objects of Man's regard are equally insignificant; and in that Infinity alone can we seek for a Wisdom to design, or a Power to execute, results so vast and so varied, by the orderly co-operation of the most simple means.

CHAPTER I.

OPTICAL PRINCIPLES OF THE MICROSCOPE.

1. ALL Microscopes in ordinary use, whether *simple* or *compound*, depend for their magnifying power on that influence exerted by lenses in altering the course of the rays of light passing through them, which is termed *refraction*.¹ This influence takes place in accordance with the two following laws, which are fully explained and illustrated in every elementary treatise on optics.²

I. A ray of light passing from a rarer into a denser medium, is refracted *towards* a line drawn perpendicularly to the plane which divides them; and *vice versâ*.

II. The *sines* of the angles of *incidence* and *refraction* (that is, of the angles which the ray makes with the perpendicular *before* and *after* its refraction) bear to one another a constant ratio for each substance, which is known as its *index of refraction*.

It follows from the first of these laws, that a ray of light entering any denser medium *perpendicularly*, undergoes *no* refraction, but continues in its straight course; and from the second, that the rays nearest the perpendicular are refracted less than those more distant from it. The "index of refraction" is determined for different substances, by the amount of the refractive influence which they exert upon rays passing into them, not from air, but from a vacuum; and in expressing it, the sine of the angle of refraction is considered as the *unit*, to which that of the angle of incidence bears a fixed relation. Thus when we say that the "index of refraction" of Water is 1.336, we mean that the sine of the angle of incidence of a ray passing into water from a vacuum, is to that of the angle of refraction, as 1.336 to 1, or almost exactly as $1\frac{1}{3}$ to 1, or as 4 to 3. And thus, the angle of incidence being given, that of the angle of refraction may be found by *dividing* it by the index of refraction.

2. On the other hand, when a ray emerges from a dense medium into a rare one, it is bent *from* the perpendicular, accord-

¹ It is not considered necessary in the present Treatise, to describe the *reflecting* Microscope of Amici; since this, although superior to the Microscopes in use previously to its introduction, has been completely superseded by the application of the Achromatic principle to the ordinary Microscope.

² See especially Dr. Golding Bird's "Manual of Natural Philosophy," Chap. XXII.

ing to the same ratio; and to find the course of the emergent ray, the sine of the angle of incidence must be *multiplied* by the "index of refraction," which will give the sine of the angle of refraction. Now when an emergent ray falls very obliquely upon the surface, the refraction which it would sustain in passing forth, tending as it does to deflect it still farther from the perpendicular, becomes so great that the ray cannot pass out at all, and is reflected back from the plane which separates the two media, into the one from which it was emerging. This internal reflection will take place, whenever the product of the sine of the angle of incidence, multiplied by the index of refraction, exceeds the sine of 90° , which is the radius of the circle; and therefore the "limiting angle," beyond which an oblique ray suffers internal reflection, varies for different substances in proportion to their respective indices of refraction. Thus, the index of refraction of water being 1.336, no ray can pass out of it into a vacuum,¹ if its angle of incidence exceed $48^\circ 28'$, since the sine of that angle, multiplied by 1.336, equals the radius; and in like manner, the "limiting angle" for flint-glass, its index of refraction being 1.60, is $38^\circ 41'$. This fact imposes certain limits upon the performance of microscopic Lenses; whilst at the same time it enables the optician to make most advantageous use of glass Prisms for the purpose of *reflection*; the proportion of the light which they throw back being much greater than that returned from the best polished metallic surfaces, and the brilliancy of the reflected image being consequently higher. Such prisms are of great value to the Microscopist for particular purposes, as will hereafter appear (§§ 40, 41, 57, 60).

3. The lenses employed in the construction of Microscopes are chiefly *convex*; those of the opposite kind, or *concave*, being only used to make certain modifications in the course of the rays passing through convex lenses, whereby their performance is rendered more exact (§§ 10, 12). It is easily shown to be in accordance with the laws of refraction already cited, that when a "pencil" of parallel rays, passing through air, impinges upon a *convex* surface of glass, the rays will be made to converge; for they will be bent towards the centre of the circle, the radius being the perpendicular to each point of curvature. The central or axial ray, as it coincides with the perpendicular, will undergo no refraction; the others will be bent from their original course in an increasing degree, in proportion as they fall at a distance from the centre of the lens; and the effect upon the whole will be such, that they will be caused to meet at a point, called the *focus*, some distance beyond the centre of curvature. This effect will not be materi-

¹ The reader may easily make evident to himself the internal reflection of water, by nearly filling a wineglass with water, and holding it at a higher level than his eye, so that he sees the surface of the fluid obliquely from beneath; no object held above the water will then be visible through it, if the eye be placed beyond the limiting angle; whilst the surface itself will appear as if silvered, through its reflecting back to the eye the light which falls upon it from beneath.

ally changed by allowing rays to pass into air again through a *plane* surface of glass, perpendicular to the axial ray (Fig. 1); a lens of this description is called a *plano-convex* lens; and it will hereafter be shown to possess properties, which render it very useful in the construction of microscopes. But if, instead of passing through a plane surface, the rays re-enter the air through a second *convex* surface, turned in the opposite direction, as in a *double-convex* lens, they will be made to converge still more. This will be readily comprehended, when it is borne in mind that the contrary direction of the second surface, and the

FIG. 1.





FIG. 2.



Parallel rays, falling on a *plano-convex* lens, brought to a focus at the distance of its diameter; and conversely, rays diverging from that point, rendered parallel. ● Parallel rays, falling on a *double-convex* lens, brought to a focus in its centre; conversely, rays diverging from that point, rendered parallel. ●


contrary direction of its refraction (this being *from* the denser medium instead of *into* it), antagonize each other; so that the second convex surface exerts an influence on the course of the rays passing through it, which is almost exactly equivalent to that of the first. Hence the focus of a *double-convex* lens will be just half the distance, or (as commonly expressed) will be at half the length, of the focus of a *plano-convex* lens having the same curvature on one side (Fig. 2).

4. The distance of the focus from the lens will depend, not merely upon its degree of curvature, but also upon the refracting power of the substance of which it may be formed; since the *lower* the index of refraction, the *less* will the oblique rays be deflected towards the axial ray, and the *more remote* will be their point of meeting; and conversely; the *greater* the refractive index, the *more* will the oblique rays be deflected towards the axial ray, and the *nearer* will be their point of convergence. A lens made of any substance whose index of refraction is 1.5, will bring parallel rays to a focus at the distance of its *diameter* of curvature, after they have passed through *one* convex surface (Fig. 1), and at the distance of its *radius* of curvature, after they have passed through *two* convex surfaces (Fig. 2); and as this ratio almost exactly expresses the refractive power of ordinary Glass, we may for all practical purposes consider the "principal focus" (as the focus for parallel rays is termed), of a *double-convex* lens

to be at the distance of its radius, that is, in its centre of curvature, and that of a *plano-convex* lens to be at the distance of twice its radius, that is, at the other end of the diameter of its sphere of curvature.


5. It is evident from what has preceded, that as a double-convex lens brings parallel rays to a focus in its centre of curvature, it will on the other hand cause those rays to assume a parallel direction, which are diverging from that centre before they impinge upon it (Fig. 2); so that, if a luminous body be placed in the principal focus of a double-convex lens, its divergent rays, falling on one surface of the lens as a *cone*, will pass forth from its other side as a *cylinder*. Again, if rays *already converging* fall upon a double-convex lens, they will be brought together at a point nearer to it than its centre of curvature (Fig. 3); whilst, if the incident rays be *diverging* from a distant point, their focus will be more distant from the lens than its principal focus (Fig. 4).

FIG. 3.



Rays already converging, brought together at a point nearer than the principal focus; and rays diverging from a point within the principal focus, still diverging, though in a diminished degree.

FIG. 4.



Rays diverging from points more distant than the principal focus on either side, brought to a focus beyond it; if the point of divergence be within the circle of curvature, the focus of convergence will be beyond it; and *vice versa*.

The further from the point from which they diverge, the more nearly will the rays approach the parallel direction; until, at length, when the object is very distant, its rays in effect become parallel, and are brought together in the principal focus. If the rays which fall upon a double-convex lens, be diverging from the farther extremity of the diameter of its sphere of curvature, they will be brought to a focus at an equal distance on the other side of the lens; but the more the point of divergence is approximated to the centre or principal focus, the further removed on the other side will be the point of convergence, until, the point of divergence being *at the centre*, there is no convergence at all, the rays being merely rendered parallel. If the point of divergence be *within* the principal focus, they will neither be brought to converge nor be rendered parallel, but will diverge in a diminished degree (Fig. 3). The same principles apply equally to a *plano-convex* lens; allowance being made for the double distance of its principal focus. They also apply to a lens whose surfaces have

different curvatures; the principal focus of such a lens is found by multiplying the radius of one surface by the radius of the other, and dividing this product by half the sum of the same radii. The rules by which the foci of convex lenses may be found, for rays of different degrees of convergence and divergence, will be found in works on Optics.


6. The refracting influence of *concave* lenses will evidently be precisely the opposite of that of convex. Rays which fall upon them in a parallel direction, will be made to *diverge* as if from the principal focus, which is here called the *negative* focus. This will be, for a plano-concave lens, at the distance of the diameter of the sphere of curvature; and for a double-concave, in the centre of that sphere. In the same manner, rays which are converging to such a degree, that, if uninterrupted, they would have met in the principal focus, will be rendered parallel; if converging more, they will still meet, but at a greater distance; and if converging less, they will diverge as from a negative focus at a greater distance than that for parallel rays. If already diverging, they will diverge still more, as from a negative focus nearer than the principal focus; but this will approach the principal focus, in proportion as the distance of the point of divergence is such, that the direction of the rays approaches the parallel.

7. If a lens be convex on one side and concave on the other, forming what is called a *meniscus*, its effect will depend upon the proportion between the two curvatures. If they are equal, as in a watch glass, no perceptible effect will be produced; if the convex curvature be the greater, the effect will be that of a less powerful *convex* lens: and if the concave curvature be the more considerable, it will be that of a less powerful *concave* lens. The focus of convergence for parallel rays in the first case, and of divergence in the second, may be found by dividing the product of the two radii by half their difference.

8. Hitherto we have considered only the effects of lenses upon a "pencil" of rays issuing from a single luminous point, and that point situated in the line of its axis. If the point be situated above the line of its axis, the focus will be below it, and *vice versa*. The surface of every luminous body may be regarded as comprehending an infinite number of such points, from every one of which a pencil of rays proceeds, and is refracted according to the laws already specified; so that a perfect but inverted image or picture of the object is formed upon any surface placed in the focus, and adapted to receive the rays. It will be evident from what has gone before, that if the object be placed at twice the distance of the principal focus, the image, being formed at an equal distance on the other side of the lens (§ 5), will be of the same dimensions with the object: whilst, on the other hand, if the object (Fig. 5, *a b*) be nearer the lens, the image *A B* will be farther from it, and of larger dimensions; but if the object *A B* be farther from the lens, the image *a b* will be nearer to it, and

smaller than itself. Further, it is to be remarked, that the larger the image in proportion to the object, the less bright it

FIG. 5.



Formation of Images by convex lenses.

will be, because the same amount of light has to be spread over a greater surface; whilst an image that is smaller than the object, will be more brilliant in the same proportion. 9. The knowledge of these general facts will enable us readily to understand the ordinary operation of the Microscope; but the instrument is subject to certain optical imperfections, the mode of remedying which cannot be comprehended without an acquaintance with their nature. One of these imperfections results from the unequal refraction of the rays which have passed through lenses, whose curvatures are equal over their whole surfaces. If the course of the rays passing through an ordinary convex lens be carefully laid down (Fig. 6), it will be found that

FIG. 6.




Diagram illustrating Spherical Aberration.

they do not all meet exactly in the foci already stated, but that the focus F of the rays $A B, A B$, which have passed through the peripheral portion of the lens, is much closer to it than that of the rays ab, ab , which are nearer the line of its axis; so that, if a screen be held in the former, the rays which have passed through the central portion of the lens will be stopped by it before they have come to a focus; and if the screen be carried back into the focus of the latter, the rays which were most distant from the axis will have previously met and crossed, so that they will come to it in a state of divergence, and will pass to c and d . In either case, therefore, the image will have a certain degree of indistinctness; and there is no one point to which all the rays can be brought by a single lens of spherical curvature. The difference between the focal points of the central and of the peripheral rays, is termed the *Spherical Aberration*. It is obvious that, to produce the desired effect, the curvature requires to be increased around the centre of the lens, so as to bring the rays which pass through it more speedily to a focus; and to be diminished towards the circumference, so as to throw the focus of the rays influenced by it to a greater distance. The requisite conditions may be theoretically fulfilled by a lens, one of whose surfaces, instead of being spherical, should be a portion of an ellipsoid or hyperboloid of

certain proportions; but the difficulties in the way of the mechanical execution of lenses of this description are such, that, for practical purposes, this plan of construction is altogether unavailable.

9 a. Various means have been devised for reducing the Aberration of lenses of spherical curvature. It may be considerably diminished, by making the most advantageous use of ordinary lenses. Thus, the aberration of a plano-convex lens, whose convex side is turned towards parallel rays, is only $\frac{1}{100}$ ths of its thickness; whilst, if its plane side be turned towards them, the aberration is $4\frac{1}{2}$ times the thickness of the lens. Hence, in the employment of a plano-convex lens, its convex surface should be turned towards a distant object, when it is used to form an image by bringing to a focus parallel or slightly diverging rays; but it should be turned towards the eye, when it is used to render parallel the rays which are diverging from a very near object. The single lens having the least spherical aberration, is a double-convex whose radii are as one to six: when its flattest face is turned towards parallel rays, the aberration is nearly $3\frac{1}{2}$ times its thickness; but when its most convex side receives or transmits them, the aberration is only $\frac{1}{20}$ ths of its thickness. The aberration is further diminished, by reducing the aperture or working-surface of the lens, so as to employ only the rays that pass through the central part, which, if sufficiently small in proportion to the whole sphere, will bring them all to nearly the same focus. The use of this may be particularly noticed in the object-glasses of common (non-achromatic) Microscopes; in which, whatever be the size of the lens itself, the greater portion of its surface is rendered inoperative by a *stop*, which is a plate with a circular aperture interposed between the lens and the rest of the instrument. If this aperture be gradually enlarged, it will be seen that, although the image becomes more and more illuminated, it is at the same time becoming more and more indistinct; and that, in order to gain *defining power*, the aperture must be reduced again. Now this reduction is attended with two great inconveniences; in the first place, the loss of intensity of light, the degree of which will depend upon the quantity transmitted by the lens, and will vary, therefore, with its aperture; and, secondly, the diminution of the "angle of aperture," that is, of the angle (*abc*, Fig. 8) made by the most diverging of the rays of the pencil issuing from any point of an object, which can enter the lens; on the extent of which angle depend some of the most important qualities of a Microscope (§ 100).

10. The Spherical Aberration may be got rid of altogether, however, by making use of *combinations* of lenses, so disposed that their opposite aberrations shall correct each other, whilst magnifying power is still gained. For it is easily seen that, as the aberration of a concave lens is just the opposite of that of a convex lens, the aberration of a convex lens placed in its most

favorable position may be corrected by a concave lens of much less power in its most unfavorable position; so that, although the power of the convex lens is weakened, all the rays which pass through this combination will be brought to one focus. It is by a method of this kind, that the Optician aims to correct the spherical aberration, in the construction of those combinations of lenses which are now employed as object-glasses, in all Compound Microscopes that are of any real value as instruments of observation. But it sometimes happens that this correction is not perfectly made; and the want of it becomes evident, in the *fog* by which the distinctness of the image of the object, and especially the precision of its outlines, is obscured.

11. But the spherical aberration is not the only imperfection with which the optician has to contend in the construction of microscopes. A difficulty equally serious arises from the unequal refrangibility of the several colored rays, which together make up white or colorless light,¹ so that they are not all brought to the same focus, even by a lens free from spherical aberration. It is this difference in their refrangibility, which causes their complete separation by the prism into a spectrum; and it manifests itself, though in a less degree, in the image formed by a convex lens. For if parallel rays of white light fall upon a convex surface, the *most* refrangible of its component rays, namely, the *violet*, will be brought to a focus at a point somewhat nearer to the lens than the principal focus, which is the mean of the whole; and the converse will be true of the *red* rays, which are the *least* refrangible, and whose focus will therefore be more distant. Thus in Fig. 7, the rays of white light, *A B*, *A B*, which fall

Fig. 7.




Diagram illustrating Chromatic Aberration.

on the peripheral portion of the lens, are so far decomposed, that the violet rays are brought to a focus at *c*, and crossing there, diverge again and pass on towards *F F*. On the other hand, the red rays are not brought to a focus until *d*, and cross the diverging violet rays at *E E*. The foci of the intermediate rays of the spectrum (indigo, blue, green, yellow, and orange) are intermediate between these two extremes. If the image be received upon a screen placed at *c*, the focus of the violet rays, violet will predominate in its own color, and it will be surrounded by a prismatic fringe in which blue, green, yellow, orange, and red may be successively distinguished. If, on the other hand, the screen be placed at *d*, the focus of the red rays, the image will have a

¹ It has been deemed better to adhere to the ordinary phraseology, when speaking of this fact, as more generally intelligible than the language in which it might be more scientifically described, and at the same time leading to no practical error.

predominantly red tint, and will be surrounded by a series of colored fringes in inverted order, formed by the other rays of the spectrum, which have met and crossed.¹ The line $\epsilon \epsilon$, which joins the points of intersection between the red and the violet rays, marks the "mean focus," that is the situation in which the colored fringes will be narrowest, the "dispersion" of the colored rays being the least; whilst the interval $c d$, which separates the foci of the extreme rays, is termed the *Chromatic Aberration* of the lens. As the axial ray $A' B'$ undergoes no refraction, neither does it sustain any dispersion; and the nearer the rays are to the axial ray, the less dispersion do they suffer. Again, the more oblique the direction of the rays, whether they pass through the central or the peripheral portion of the lens, the greater will be the refraction they undergo, and the greater also will be their dispersion; and thus it happens that when, by using only the central part of a lens (§ 12), the chromatic aberration is reduced to its minimum, the central part of a picture may be tolerably free from false colors, whilst its marginal portion shall exhibit broad fringes.²

12. The Chromatic Aberration of a lens, like the Spherical, may be diminished by the contraction of its aperture, so that only its central portion is employed. But the error cannot be got rid of entirely by any such reduction, which, for the reasons already mentioned, is in itself extremely undesirable. Hence it is of the first importance in the construction of a really efficient Microscope, that the chromatic aberration of its "object-glasses" (in which the principal dispersion is liable to occur) should be entirely *corrected*, so that the largest possible aperture should be given to these lenses, without the production of any false colors. No such correction can be accomplished even theoretically in a single lens; but it may be effected by the combination of two or more, advantage being taken of the different relations which the *refractive* and the *dispersive* powers bear to each other in different substances. For if we can unite with a *convex* lens, whose dispersive power is *low* as compared with its refractive power, a *concave* of lower curvature, whose dispersive power is relatively *high*, it is obvious that the dispersion of the rays occasioned by the convex lens may be effectually *neutralized* by the opposite dispersion of the concave (§ 6); whilst the refracting power of the convex is only *lowered* by the opposite refraction of the concave, in virtue of the longer focus of the latter. No difficulty stands in the way of carrying this theoretical correction into practice. For the "dispersive" power of *flint-glass* bears so much larger a ratio to its refractive power than does that of

¹ This experiment is best tried with a lens of long focus, of which the central part is covered with an opaque stop, so that the light passes only through a peripheral ring; since, if its whole aperture be in use, the regular formation of the fringes is interfered with by the *spherical* aberration, which gives a different focus to the rays passing through each annular zone.

² This is well seen in the large pictures exhibited by Oxy-hydrogen Microscopes.


crown-glass, that a convex lens of the former, the focal length of which is $7\frac{1}{4}$ inches, will produce the same degree of color as a convex lens of *crown-glass*, whose focal length is $4\frac{1}{4}$ inches. Hence a concave lens of the former material and curvature, will fully correct the dispersion of a convex lens of the latter; whilst it diminishes its refractive power only to such an extent as to make its focus 10 inches. The correction for chromatic aberration in such a lens would be perfect, if it were not that, although the extreme rays, violet and red, are thus brought to the same focus, the dispersion of the rest is not equally compensated; so that what is termed a *secondary spectrum* is produced, the images of objects seen through such a lens being bordered on one side with a purple fringe, and on the other with a green fringe. Moreover such a lens is not corrected for spherical aberration; and it must of course be rendered free from this, to be of any real service, however complete may be the freedom of its image from false colors. The double correction may be accomplished theoretically by the combination of *three* lenses, namely, a double-concave of flint placed between two double-convex of crown, ground to certain curvatures; and this method has long been employed in the construction of the large object-glasses of Telescopes, which are, by means of it, rendered *Achromatic*,—that is, are enabled to exert their refractive power without producing either chromatic or spherical aberration.

13. It has only been of late years, however, that the construction of Achromatic object-glasses for Microscopes has been considered practicable; their extremely minute size having been thought to forbid the attainment of that accuracy which is necessary in the adjustment of the several curvatures, in order that the errors of each separate lens which enters into the combination, may be effectually balanced by the opposite errors of the rest. The first successful attempt was made in this direction, in the year 1823, by M. Selligues of Paris; the plan which he adopted being that of the combination of two or more pairs of lenses, each pair consisting of a double-convex of crown-glass, and a plano-concave of flint. In the next year, Mr. Tulley of London, without any knowledge of what M. Selligues had accomplished, applied himself (at the suggestion of Dr. Goring) to the construction of achromatic object-glasses for the microscope; and succeeded in producing a single combination of three lenses (on the telescope plan), the corrections of which were extremely complete. This combination, however, was not of high power, nor of large angular aperture; and it was found that these advantages could not be gained, without the addition of a second combination. Prof. Amici at Modena, also, who attempted the construction of microscopic object-glasses as early as 1812, but, despairing of success, had turned his attention to the application of the *reflecting* principle to the Microscope, resumed his original labors on hearing of the success of M. Selligues; and, by work-

ing on his plan, he produced, in 1827, an achromatic combination of three pairs of lenses, which surpassed anything of the same kind that had been previously executed. From that time, the superiority of the plan of combining *three pairs* of lenses (Fig. 8; 1, 2, 3), which should be so adjusted as to correct each other's errors, to the telescopic combinations adopted by Mr. Tulley, may be considered to have been completely established; and English opticians, working on this method, soon rivalled the best productions of Continental skill.

14. It was in this country that the next important improvements originated; these being the result of the theoretical investigations of Mr. J. J. Lister,¹ which led him to the discovery of certain properties in achromatic combinations, that had not been previously detected. Acting upon the rules which he laid down, practical opticians at once succeeded in producing combinations far superior to any which had been previously executed, both in wideness of aperture, flatness of field, and perfectness of correction; and continued progress has been since made in the same direction, by the like combination of theoretical acumen with manipulative skill.² For the subsequent investigations of Mr. Lister have led him to suggest new combinations, which have been speedily carried into practical execution; and there is good reason to believe that the limit of perfection has now been nearly reached, since almost everything which seems theoretically possible has been actually accomplished. The most perfect combinations at present in use for high powers, consist of as many as *eight* distinct lenses; namely, in front, a triplet composed of two plano-convex lenses of crown-glass, with a plano-concave of dense flint between them; next, a doublet, composed of a double-convex of crown, and a double-

FIG. 8.



Section of an Achromatic Object-glass.


¹ See his Memoir in the "Philosophical Transactions," for 1829.

² The first British Opticians (after Mr. Tulley) who applied themselves to the construction of Achromatic object-glasses for microscopes, were Mr. Ross and Mr. Powell. Mr. James Smith did not enter the field until some time afterwards; but, having the advantage of Mr. Lister's special superintendence, he soon equalled, in the lower powers at least, the best productions of his predecessors. With Mr. Ross, his son has been subsequently associated: with Mr. Powell, his brother-in-law, Mr. Lealand; and with Mr. Smith, Mr. Beck, a nephew of Mr. Lister. These three firms have constantly kept up an honorable rivalry, which has been very advantageous to the *perfectionment* of the Microscope; and have maintained a position which is still far in advance of that of all other manufacturing Opticians in this country or the Continent. The lenses produced by each are distinguished by excellencies of their own; and it would be scarcely possible fairly to assign an absolute preference to either above the others. Among the amateurs who have occupied themselves in the construction of microscopic Achromatics, Mr. Wenham has been the most successful. An American rival has recently been announced, in the person of Mr. Spencer; who, taking advantage of all that had been previously accomplished, is said to have produced combinations not only equalling, but, in some important particulars, surpassing those of English makers. Only one of these, however, has found its way (the author believes) to this country; and not having had the opportunity of seeing it himself, he can only judge of it by report. (See Appendix.)

concave of flint; and at the back, another triplet, consisting of two double-convex lenses of crown, with a double-concave of flint interposed between them. By the use of this combination, an angular aperture of no less than 170° has been obtained with an objective of 1-12th inch focus; and it is obvious that as an increase of divergence of no more than 10° would bring the extreme rays into a straight line with each other, they would not enter the lens at all; so that no further enlargement of the aperture can be practically useful.

15. The enlargement of the angle of aperture, and the greater completeness of the corrections, first obtained by the adoption of Mr. Lister's principles, soon rendered sensible an imperfection in the performance of these lenses under certain circumstances

Fig. 9.



which had previously passed unnoticed; and the important discovery was made by Mr. Ross, that a very obvious difference existed in the precision of the image, according as the object is viewed *with* or *without* a covering of talc or thin glass; an object-glass which is perfectly adapted to either of these conditions, being sensibly defective under the other. The mode in which this difference arises, is explained by Mr. Ross as follows.¹ Let o , Fig. 9, be any point of an object; $o p$ the axial ray of the pencil that diverges from it; and $o t$, $o t'$, two diverging rays, the one near to, the other remote from, the axial ray. Now if $g g g g$ represent the section of a piece of thin glass, intervening between the object and the object-glass, the rays $o t$ and $o t'$ will be refracted in their passage through it, in the directions $t r$, $t' r'$; and on emerging from it again, they will pass on towards e and e' . Now if the course of these emergent rays be traced backwards, as by the dotted lines, the ray $e r$ will seem to have issued from x , and the ray $e' r'$ from y ; and the distance $x y$ is an aberration quite sufficient to disturb the previous balance of the aberrations of the lens composing the object-glass. The requisite correction may be effected, as Mr. Ross pointed out, by giving to the *front* pair (Fig. 8, 1) of the three of which the objective is composed, an excess of positive aberration (*i. e.* by under-correcting it), and by giving to the other two pairs (2, 3) an excess of negative aberration (*i. e.* by over-correcting them), and by making the distance between the former and the latter susceptible of alteration. For when the front pair is approximated most nearly to the other two, and its distance from the object is increased,

¹ "Transactions of the Society of Arts," vol. li.

its positive aberration is more strongly exerted upon the other pairs, than it is when the distance between the lenses is increased, and the distance between the front pair and the object is diminished. Consequently, if the lenses be so adjusted that their correction is perfect for an uncovered object, the front pair being removed to a certain distance from the others, its approximation to them will give to the whole combination an excess of positive aberration, which will neutralize the negative aberration occasioned by covering the object with a thin plate of glass.¹ It is obvious that this correction will be more important to the perfect performance of the combination, the larger is its angle of aperture; since, the wider the divergence of the oblique rays from the axial ray, the greater will be the refraction which they will sustain in passing through a plate of glass, and the greater therefore will be the negative aberration produced, which will, if uncorrected, seriously impair the distinctness of the image. And it is consequently not required for *low* powers, whose angle of aperture is comparatively small; nor even for the higher, so long as their angle of aperture does not exceed 50°. As a large proportion of the lenses made by foreign Opticians do not range beyond this, the adjustment in question may be dispensed with; and even where the angle is much larger, if the corrections be made perfect for a thickness of glass of 1-100th of an inch (which is about an average of that with which objects of the finer kind are usually covered), they will not be much deranged by a difference of a few hundredths of an inch, more or less, in that amount.

16. We are now prepared to enter upon the application of the optical principles which have been explained and illustrated in the foregoing pages, to the construction of microscopes. These are distinguished as *simple*, and *compound*; each kind having its peculiar advantages to the Student of Nature. Their essential difference consists in this;—that in the former, the rays of light which enter the eye of the observer proceed directly from the object itself, after having been subject only to a change in their course; whilst in the latter, an enlarged *image* of the object is formed by a lens, which image is viewed by the observer through a simple microscope, as if it were the object itself. The *simple* microscope *may* consist of *one* lens; but (as will be presently shown) it may be formed of *two*, or *even* three; these, however, are so disposed as to produce an action upon the rays of light corresponding to that of a single lens. In the *compound* microscope, on the other hand, not less than two lenses must be employed; one to form the enlarged image of the object, and this, being nearest to it, is called the *object-glass*; whilst the other again magnifies that image, being interposed between it and the eye of the observer, and is hence called the *eye-glass*. A perfect

¹ The mode in which this adjustment is effected, will be more fitly described hereafter (§ 82).

object-glass, as we have seen, must consist of a combination of lenses; and the eye-glass, as we shall presently see (§ 21), is best constructed by placing two lenses in a certain relative position, forming what is termed an *eye-piece*. These two kinds of instrument need to be separately considered in detail.

17. *Simple Microscope*.—In order to gain a clear notion of the mode in which a single lens serves to “magnify” minute objects, it is necessary to revert to the phenomena of ordinary vision. An eye free from any defect has a considerable power of adjusting itself, in such a manner as to gain a distinct view of objects placed at extremely varying distances; but the image formed upon the retina will of course vary in size with the distance of the object; and the amount of detail perceptible in it will follow the same proportion. To ordinary eyes, however, there is a limit within which no distinct image can be formed, on account of the too great divergence of the rays of the different pencils which then enter the eye; since the eye is usually adapted to receive, and to bring to a focus, rays which are parallel or but slightly divergent. This limit is variously stated at from five to ten inches; we are inclined to think from our own observations, that the latter estimate is nearest the truth; that is, although a person with an ordinary vision may see an object much nearer to his eye, he will see little if any more of its details, since what is gained in size will be lost in distinctness. Now the utility of a convex lens interposed between a near object and the eye, consists in its reducing the divergence of the rays forming the several pencils which issue from it; so that they enter the eye in a state of moderate divergence, as if they had issued from an object beyond the nearest limit of distinct vision; and a well-defined picture is consequently formed upon the retina. But not only is the course of the several rays in each pencil altered as regards the rest, by this refracting process, but the course of the pencils themselves is changed, so that they enter the eye under an angle corresponding with that at which they would have arrived from a larger object situated at a greater distance. The picture formed upon the retina, therefore, by any object (Fig. 10),

FIG. 10.




Diagram illustrating the action of the *Simple Microscope*.

corresponds in all respects with one which would have been made by the same object *a b* increased in its dimensions to *A B*, and viewed at the smallest ordinary distance of distinct vision. A “short-sighted” person, however, who can see objects distinctly at a distance of two or three inches, has the same power in his eye alone, by reason of its greater convexity, as that which the person of ordinary vision

gains by the assistance of a convex lens which shall enable him to see at the same distance with equal distinctness. It is evident, therefore, that the magnifying power of a single lens, depending as it does upon the proportion between the distance at which it renders the object visible, and the nearest distance of unaided distinct vision, must be different to different eyes. It is usually estimated, however, by finding how many times the focal length of the lens is contained in ten inches; since, in order to render the rays from the object nearly parallel, it must be placed nearly in the focus of the lens (Fig. 2); and the picture is referred by the mind to an object at the ordinary distance. Thus, if the focal length of a lens be one inch, its magnifying power for each dimension will be ten times, and consequently a hundred superficial; if its focal distance be only one-tenth of an inch, its magnifying power will be a hundred linear, or ten thousand superficial. The use of the convex lens has the further advantage of bringing to the eye a much greater amount of light, than would have entered the pupil from the enlarged object at the ordinary distance, provided its own diameter be greater than that of the pupil; but this can only be the case when its magnifying power is low.

18. It is obviously desirable, especially when lenses of very high magnifying power are being employed, that their aperture should be as large as possible; since the light issuing from a minute object has then to be diffused over a large picture, and will be proportionally diminished in intensity. But the shorter the focus, the less must be the diameter of the sphere of which the lens forms a part; and unless the aperture be proportionally diminished, the spherical and chromatic aberrations will interfere so much with the distinctness of the picture, that the advantages which might be anticipated from the use of such lenses will be almost negated. Nevertheless, the Simple Microscope has been an instrument of extreme value in anatomical research, owing to its freedom from those errors to which the Compound Microscope, as originally constructed, was necessarily subject; the greater certainty of its indications being evident from the fact, that the eye of the observer receives the rays sent forth by the object itself, instead of those which proceed from an image of that object. A detail of the means employed by different individuals, for procuring lenses of extremely short focus, though possessing much interest in itself, would be misplaced here; since recent improvements, as will presently be shown, have superseded the necessity of all these. It may be stated, however, that Leeuwenhoek, De la Torre, and others among the older microscopists, made great use of small globules procured by fusion of threads or particles of glass. The most important suggestion for the improvement of the Simple Microscope composed of a single lens, proceeded some years ago from Dr. Brewster, who proposed to substitute diamond, sapphire, garnet,

and other precious stones of high refractive power, for glass, as the material of single lenses. A lens of much longer radius of curvature might thus be employed, to gain an equal magnifying power; and the aperture would admit of great extension, without a proportional increase in the spherical and chromatic aberrations. This suggestion has been carried into practice with complete success, as regards the performance of lenses executed on this plan; but the difficulties of various kinds in the way of their execution, are such as to render them very expensive; and as they are not superior to the combination now to be described, they have latterly been quite superseded by it. This combination, first proposed by Dr. Wollaston, and known as his *doublet*, consists of two plano-convex lenses, whose focal lengths are in the proportion of one to three, or nearly so, having their convex sides directed towards the eye, and the lens of shortest focal length nearest the object. In Dr. Wollaston's original combination, no perforated diaphragm (or "stop") was interposed; and the distances between the lenses was left to be determined by experiment in each case. A great improvement was subsequently made, however, by the introduction of a "stop" between the lenses, and by the division of the power of the smaller lens between two (especially when a very short focus is required) so as to form a *triplet*, as was first suggested by Mr. Holland.* When combinations of this kind are well constructed, both the spherical and the chromatic aberrations are so much reduced, that the angle of aperture may be considerably enlarged without much sacrifice of distinctness; and hence for all powers above 1-4th inch focus, doublets and triplets are far superior to single lenses. The performance of even the best of these forms of Simple microscope, however, is so far inferior to that of a good Compound microscope as now constructed upon the achromatic principle, that no one who has the command of the latter form of instrument would ever use the *higher* powers of the former. It is for the prosecution of observations, and for the carrying on of dissections, which only require *low* powers, that the Simple microscope is to be preferred; and, consequently, although doublets and triplets afforded the best means of obtaining a high magnifying power, before Achromatic lenses were brought to their present perfection, they are now comparatively little used.

19. Another form of simple magnifier, possessing certain advantages over the ordinary double-convex lens, is that commonly known by the name of the "Coddington" lens. The first idea of it was given by Dr. Wollaston, who proposed to cement together two plano-convex, or hemispherical lenses, by their plane sides, with a stop interposed, the central aperture of which should be equal to 1-5th of the focal length. The great advantage of such a lens is, that the oblique pencils pass, like the centre ones, at right angles with the surface; and that they are consequently but little subject to aberration. The idea was further improved

* "Transactions of the Society of Arts," vol. xlix.

upon by Mr. Coddington, who pointed out that the same end would be much better answered by taking a sphere of glass, and grinding a deep groove in its equatorial part, which should be then filled with opaque matter, so as to limit the central aperture. Such a lens gives a large field of view, admits a considerable amount of light, and is equally good in all directions; but its powers of definition are by no means equal to those of an achromatic lens, or even of a doublet. This form is chiefly useful, therefore, as a hand magnifier, in which neither high power nor perfect definition is required; its peculiar qualities rendering it superior to an ordinary lens of the same power, for the class of objects for which such lenses are applied in this mode. We think it right to state that many of the magnifiers sold as "Coddington" lenses are not really (as we have satisfied ourselves) portions of spheres, but are manufactured out of ordinary double-convex lenses, and will be destitute, therefore, of many of the above advantages. It may be desirable to allude to the magnifier known under the name of the "Stanhope" lens, which somewhat resembles the "Coddington" in appearance, but differs from it essentially in properties. It is nothing more than a double-convex lens, having two surfaces of unequal curvatures, separated from each other by a considerable thickness of glass; the distance of the two surfaces from each other being so adjusted, that when the most convex is turned towards the eye, minute objects placed on the other surface shall be in the focus of the lens. This is an easy mode of applying a rather high magnifying power to scales of butterflies' wings and other similar flat and minute objects, which will readily adhere to the surface of the glass; and it also serves to detect the presence of the larger animalcules, or of crystals in minute drops of fluid, to exhibit the "eels" in paste or vinegar, &c. &c.; but it is almost entirely destitute of value as an instrument of scientific research, and can scarcely be regarded in any higher light than as an ingenious philosophical toy.¹

20. *Compound Microscope*.—In its most simple form, this instrument consists of only two lenses, the "object-glass" and the "eye-glass:" the former, *CD* (Fig. 11), receiving the rays of light direct from the object, *AB*, which is brought into near proximity to it, forms an enlarged and inverted image *A'B'* at a greater distance on the other side; whilst the latter, *LM*, receives the rays which are diverging from this image, as if they proceeded from an object actually occupying its position and enlarged to its dimensions, and these it brings to the eye at *E*, so altering their course as to make that image appear far larger to the eye, precisely as in the case of the simple microscope (§ 16). It is obvious that by the use of the very same lenses, a considerable variety of magnifying power may be obtained, simply by altering their

¹The principal forms of construction of Simple Microscopes, will be described in the next chapter.

position in regard to each other and to the object; for if the eye-glass be carried further from the object-glass, whilst the object is approximated nearer to the latter, the image $A' B'$ will be formed at a greater distance from it, and its dimensions will consequently be augmented. If on the other hand, the eye-glass be brought nearer to the object-glass, whilst the object is removed further from it, the distance of the image will be shortened, and its dimensions proportionably diminished. We shall hereafter see that this mode of varying the magnifying power of compound microscopes may be turned to good account in more than one

FIG. 11.




Diagram of simplest form of Compound Microscope.

FIG. 12.



Diagram of the complete Compound Microscope.


mode (§§ 43, 44); but there are limits to the use which can be advantageously made of it.—The amplification may also be varied by altering the magnifying power of the eye-glass; but here, too, there are limits to the increase; since defects of the object-glass, which are not perceptible when its image is but moderately enlarged, are brought into injurious prominence when the imperfect image is amplified to a much greater extent. In practice, it is generally found much better to vary the power, by employing object-glasses of different foci; an object-glass of *long* focus forming an image, which is not at many times the distance of the object from the other side of the lens, and which, therefore, is not of many times its dimension; whilst an object-glass of *short* focus requires that the object should be so nearly approximated to it, that the distance of the image is a much higher multiple of that of the object, and its dimensions are proportionably larger (§ 8). In whatever mode additional amplification

be obtained, two things must always result from the change: the portion of the surface of the object, of which an image can be formed, must be diminished; and the quantity of light spread over that image must be proportionably lessened.

21. In addition to the two lenses of which the Compound Microscope essentially consists, another (Fig. 12, F F) is usually

introduced between the object-glass and the image formed by it. The ordinary purpose of this lens is to change the course of the rays in such a manner, that the image may be formed of dimensions not too great for the whole of it to come within the range of the eye-piece; and as it thus allows more of the object to be seen at once, it is called the *field-glass*. It is now usually considered, however, as belonging to the ocular end of the instrument,—the *eye-glass* and the *field-glass* being together termed the *Eye-piece*. Various forms of this eye-piece have been proposed by different opticians; and one or another will be preferred, according to the purpose for which it may be required. That which it is most advantageous to employ with Achromatic object-glasses, to the performance of, which it is desired to give the greatest possible effect, is termed the “Huyghenian;” having been employed by Huyghens for his telescopes, although without the knowledge of all the advantages which its best construction renders it capable of affording. It consists of two plano-convex lenses (E E and F F, Fig. 12), with their plane sides towards the eye; these are placed at a distance equal to half the sum of their focal lengths; or, to speak with more precision, at half the sum of the focal length of the eye-glass, and of the distance from the field-glass at which an image of the object-glass would be formed by it. A “stop” or diaphragm, B B, must be placed between the two lenses, in the visual focus of the eye-glass, which is, of course, the position wherein the image of the object will be formed, by the rays brought into convergence by their passage through the field-glass.

By Huyghens, this arrangement was intended merely to diminish the spherical aberration; but it was subsequently shown by Boscovich, that the chromatic dispersion was also in a great part corrected by it. Since the introduction of Achromatic object-glasses for Compound Microscopes, it has been further shown that all error may be avoided by a slight over-correction of these; so that the blue and red rays may be caused to enter the eye in a parallel direction (though not actually coincident), and thus to produce a colorless image. Thus let L M N (Fig. 13), represent the two extreme rays of three pencils, which, without the field-glass, would form a blue image convex to the eye-glass, at B B, and a red one at R R; then, by the intervention of the field-glass, a blue image, concave to the eye-glass, is formed at B' B', and a red one at R' R'. As the focus of the eye-glass is shorter for blue rays than for red rays, by just the difference in the place of these images, their rays, after refraction by it, enter the eye in a parallel direction, and



Section of Huyghenian Eye-piece adapted to over-corrected Achromatic objectives.

produce a picture free from false color. If the object-glass had been rendered perfectly achromatic, the blue rays, after passing through the field-glass, would have been brought to a focus at b , and the red at r ; so that an error would be produced, which would have been increased instead of antagonized by the eye-glass. Another advantage of a well-constructed Huyghenian eye-piece is that the image produced by the meeting of the rays after passing through the field-glass, is by it rendered concave towards the eye-glass, instead of convex, so that every part of it may be in focus at the same time, and the field of view thereby rendered flat.¹

22. Two or more Huyghenian eye-pieces, of different magnifying powers, known as Nos. 1, 2, 3, &c., are usually supplied with a Compound Microscope. The utility of the higher powers will mainly depend upon the excellence of the objectives; for when an achromatic combination of small aperture, which is sufficiently well corrected to perform very tolerably with a low eye-piece, is used with an eye-piece of higher magnifying power (commonly spoken of as a "deeper" one), the image may lose more in brightness and in definition, than is gained by its amplification; whilst the image given by an objective of large angular aperture and very perfect corrections, shall sustain so little loss of light or of definition by "deep eye-piecing," that the increase of magnifying power shall be almost all clear gain. Such an objective, therefore, though of far inferior power in itself, is practically more valuable (as giving a much greater *range* of power with equal efficiency) than a lens of higher power which can only be used effectively with the shallower eye-pieces. Hence the mode in which different achromatic combinations of the same power, whose performance with shallow eye-pieces is nearly the same, are respectively affected by deep eye-pieces, afford a good test of their respective merits; since any defect in the corrections is sure to be brought out by the higher amplification of the image, whilst a deficiency of aperture is manifested by the want of light.

23. An Eye-piece is sometimes furnished with achromatic microscopes, especially for micrometric purposes, which, though composed of only two plano-convex lenses, differs essentially in its construction from the Huyghenian; the field-glass having its convex side upwards, and being so much nearer to the eye-glass, that the image is not formed above it (as at $B B$, Fig. 12), but below it. This eye-piece, which is known as Ramsden's, gives a very distinct view in the central portion of the field; but, as it does not, like the Huyghenian, correct the convexity of the image formed by the object-glass, but rather increases it, the marginal portions of the field of view, when the centre is in focus, are quite indistinct. Hence this eye-piece cannot be recommended for ordinary use; and its chief value to the Microscopist has resulted

¹ Those who desire to gain more information upon this subject than they can from the above notice of it, may be referred to Mr. Varley's investigation of the properties of the Huyghenian eye-piece, in the 51st volume of the "Transactions of the Society of Arts;" and to the article "Microscope," by Mr. Ross, in the "Penny Cyclopædia."

from its adaptation to receive a divided glass micrometer, which may be fitted into the exact plane wherein the image is formed by the object-glass, so that its scale and that image are both magnified together by the lenses interposed between them and the eye. We shall hereafter see, however, that the same end may be so readily attained with the Huyghenian eye-piece (§ 46), that no practical advantage is gained by the use of that of Ramsden. It is affirmed by Mr. Ross, that if the Achromatic principle were applied to the construction of Eye-pieces, the latter is the form with which the greatest perfection would be obtained. That such an adaptation might be productive of valuable results, appears from the success with which Mr. Brooke has employed a triplet objective of one-inch focus, as an eye-piece; the definition obtained by it being very superior to that afforded by the ordinary Huyghenian eye-piece.

24. In the Eye-pieces of compound microscopes of older construction, it was customary to employ a pair of plano or double-convex lenses of longer focus, for the eye-glass, instead of a single plano-convex of shorter focus; the advantage being, that a larger and flatter field could be thereby obtained. A brighter image, a flatter field, and a greater freedom from aberration, than are afforded by any ordinary eye-piece of this kind, may be obtained by the substitution of a combination nearly resembling Herschel's "aplanatic doublet"—namely, a meniscus, having its concave side next the eye, and a double-convex of the form of least aberration,¹ with its flattest side next the object—for the plano-convex eye-glass; and the substitution of a double-convex lens of the form of least aberration, with its flattest side next the object, for the plano-convex field-glass. With such an eye-piece, a field of fourteen inches in diameter (measured at the usual distance of ten inches) may be obtained perfectly flat, and equally distinct and well illuminated over every part. When such an eye-piece, however, is used in conjunction with achromatic objectives, it impairs the definition of their image to such a degree, that their finest qualities are altogether sacrificed. Still there are certain large transparent objects, such as transverse sections of wood, wings of insects, &c., in viewing which a large and flat field is of more importance than perfect definition; since their structure is so coarse, that there is no minute detail to be brought out. Nothing is so effective for the exhibition of these, as an eye-piece of the kind just alluded to, with an objective of about 8 or 4 inches focus; and this may either be a single lens (which, when of such low power, will perform sufficiently well for objects of this class), or a single pair of lenses forming part of a perfect achromatic combination, having its aperture somewhat contracted by a stop.²

¹ The "form of least aberration" is when the radii of the two surfaces are to each other as 1 to 6.

² Some of the lowest French Achromatics answer extremely well for this purpose; and the front pair of the lowest set usually made in this country (that, namely, of 2 inches focus) is sometimes made removable, so that the back pair, which also is very suitable to the class of objects mentioned above, may be employed by itself.

CHAPTER II.

CONSTRUCTION OF THE MICROSCOPE.

25. THE *optical* principles whereon the operation of the Microscope depends, having now been explained, we have next to consider the *mechanical* provisions, whereby they are brought to bear upon the different purposes which the instrument is destined to serve. And first it will be desirable to state those general principles, which have received the sanction of universal experience, in regard to the best arrangement of its constituent parts. Every *complete* Microscope, whether Simple or Compound, must possess, in addition to the lens or combination of lenses which affords its magnifying power, a *stage* whereon the object may securely rest, a *concave mirror* for the illumination of transparent objects from beneath, and a *condensing-lens* for the illumination of opaque objects from above.

I. Now in whatever mode these may be connected with each other, it is essential that *the optical part and the stage should be so disposed, as either to be altogether free from tendency to vibration, or to vibrate together*; since it is obvious that any movement of one, in which the other does not partake, will be augmented to the eye of the observer in proportion to the magnifying power employed. In a badly-constructed instrument, even though placed upon a steady table resting upon the firm floor of a well-built house, when high powers are used, the object is seen to oscillate so rapidly at the slightest tremor, such as that caused by a person walking across the room, or by a carriage rolling by in the street, as to be frequently almost indistinguishable: whereas in a well-constructed microscope, scarcely any perceptible effect will be produced by even greater disturbances.

II. The next requisite is *a capability of accurate adjustment to every variety of focal distance, without movement of the object*. It is now a principle almost universally recognized in the construction of good Microscopes, that the *stage* whereon the object is placed should be a *fixture*; the movement by which the focus is to be adjusted, being effected in the lenses or optical portion. Several reasons concur to establish this principle; of which one of the most important is, that, if the stage be made the movable part,

the adjustment of the illuminating apparatus must be made afresh for every change of magnifying power; whilst if the stage be a fixture, the illumination having been once well adjusted, the object may be examined under a great variety of magnifying powers, without its being changed in any respect. Moreover, if the stage be the movable part, it can never have that firmness given to it which it ought to possess; for it is almost impossible to make a movable stage free from some degree of *spring*, so that, when the hands bear upon it in adjusting the position of an object, it yields in a degree, which, however trifling in itself, becomes unpleasantly apparent with high powers. The mode of effecting the focal adjustment should be such as to allow free range from a minute fraction of an inch to three or four inches, with equal power of obtaining a delicate adjustment at any part. It should also be so accurate, that the optical axis of the instrument should not be in the least altered by movement in a vertical direction; so that, if an object be brought into the centre of the field with a low power, and a higher power be then substituted, it should be found in the centre of *its* field, notwithstanding the great alteration in the focus. In this way much time may often be saved, by employing a low power as a *finder* for an object to be examined by a higher one; and when an object is being viewed by a succession of powers, little or no readjustment of its place on the stage should be required. For the Simple Microscope, in which it is seldom advantageous to use lenses of shorter focus than 1.4th inch (save where doublets are employed, § 18), a *rack-and-pinion* adjustment answers sufficiently well; and this is quite adequate, also, for the focal adjustment of the Compound body, when objectives of low power only are employed. But for any lenses whose focus is less than half an inch, a "fine adjustment," by means of a *screw movement* operating either on the object-glass alone or on the entire body, is of great value; and for the highest powers it is quite indispensable. In some Microscopes, indeed, which are provided with a "fine adjustment," the rack-and-pinion movement is dispensed with: the "coarse adjustment" being given by merely *sliding* the body up and down in the socket which grasps it. But this plan is objectionable, inasmuch as it involves the use of *both* hands in making the "coarse adjustment," for which only *one* should be required; and even then the adjustment cannot be made with nearly the same facility, as by a smooth well-cut rack. The Author's experience, therefore, would lead him to recommend, that if one of these adjustments is to be dispensed with, it should be the "screw" or "fine" adjustment, rather than the "rack" or "coarse," unless the instrument is to be almost exclusively employed for the examination of objects requiring high magnifying powers.¹

¹ In the Microscopes constructed by Mr. Ladd, a *chain-movement* is substituted for the rack-and-pinion; and this has the advantage of being smoother and more sensitive, of being less likely to become unequal by wear, and of being easily tightened if it should

III. Scarcely less important than the preceding requisite, in the case of the Compound Microscope, though it does not add much to the utility of the Simple, is *the capability of being placed in either a vertical or a horizontal position, or at any angle with the horizon*, without deranging the adjustment of its parts to each other, and without placing the eye-piece in such a position as to be inconvenient to the observer. It is certainly a matter of surprise, that Opticians, especially on the Continent, should have so long neglected the very simple means which are at present commonly employed in this country, of giving an inclined position to microscopes; since it is now universally acknowledged, that the vertical position is, of all that can be adopted, *the very worst*. An inclination of about 55° to the horizon will generally be found most convenient for unconstrained observation; and the instrument should be so constructed, as, when thus inclined, to give to the stage such an elevation above the table, that when the hands are employed at it, the arms may rest conveniently upon the table. In this manner a degree of support is attained, which gives such free play to the muscles of the hands, that movements of the greatest nicety may be executed by them; and fatigue of long-continued observation is greatly diminished. Such minutiae may appear too trivial to deserve mention; but no practised microscopist will be slow to acknowledge their value. The stage must of course be provided with some means of supporting the object, when it is itself placed in a position so inclined that the object would slip down unless sustained. There are some objects, however, which can only be seen in a vertical microscope, as they require to be viewed in a position nearly or entirely horizontal; such are dissections in water, urinary deposits, saline solutions undergoing crystallization, &c. For other purposes, again, the microscope should be placed horizontally, as when the camera lucida is used for drawing or measuring. It ought, therefore, to be made capable of every such variety of position.

IV. The last principle on which we shall here dwell, is *simplicity in the construction and adjustment of every part*. Many ingenious mechanical devices have been invented and executed, for the purpose of overcoming difficulties which are in themselves really trivial. A moderate amount of dexterity in the use of the hands is sufficient to render most of these superfluous; and without such dexterity, no one, even with the most complete mechanical facilities, will ever become a good microscopist. Among the conveniences of simplicity, the practised Microscopist will not fail to recognize the saving of time effected by being

"lose time;" whilst its delicacy and smoothness admit of an exact adjustment being made by its means alone, even when high powers are employed. Still, as will be shown hereafter (§ 81), the use of the "fine adjustment" is by no means restricted to this purpose; and it cannot be advantageously dispensed with in a Microscope, which is to be used for any but the most common purposes.

able quickly to set up and put away his instrument. Where a number of parts are to be screwed together before it can be brought into use, interesting objects (as well as time) are not unfrequently lost; and the same cause will often occasion the instrument to be left exposed to the air and dust, to its great detriment, because *time* is required to put it away; so that a slight advantage on the side of simplicity of arrangement, often causes an inferior instrument to be preferred by the working microscopist to a superior one. Yet there is, of course, a limit to this simplification; and no arrangement can be objected to on this score, which gives advantages in the examination of difficult objects or the determination of doubtful questions, such as no simpler means can afford. The meaning of this distinction will become apparent, if it be applied to the cases of the "traversing stage" (§ 37) and the "achromatic condenser" (§ 56). For although the traversing stage may be considered a valuable aid in observation, as facilitating the finding of a minute object, or the examination of the entire surface of a large one, yet it adds nothing to the clearness of our view of either; and its place may in great degree be supplied by the fingers of a good manipulator. On the other hand, the use of the achromatic condenser not only contributes very materially, but is absolutely indispensable, to the formation of a perfect image, in the case of many objects of a difficult class; the want of it cannot be compensated by the most dexterous use of the ordinary appliances; and consequently, although it may fairly be considered superfluous, as regards a large proportion of the purposes to which the Microscope is directed, whether for investigation or for display, yet as regards the particular objects just alluded to, it must be considered as no less necessary a part of the instrument than the achromatic objective itself. Where expense is not an object, the Microscope should doubtless be fitted with *both* these valuable accessories; where, on the other hand, the cost is so limited that only *one* can be afforded, *that* one should be selected which will make the instrument most useful for the purposes to which it is likely to be applied. (See Introduction, pp. 60, 61.)

In the account now to be given, of the principal forms of Microscope readily procurable in this country, it will be the Author's object, not so much to enumerate and describe the various patterns which the several makers of the instrument have produced, as, by selecting from among them those examples which it seems to him most desirable to make known, and by specifying the peculiar advantages which each of these presents, to guide his readers in the choice of the *kind* of Microscope best suited, on the one hand, to the class of investigations they may be desirous of following out, and on the other, to their pecuniary ability. He is anxious, however, that he should not be supposed to mark any preference for the particular instruments

he has selected, over those constructed upon the same general plan by other makers; to have enumerated them all, would obviously be quite incompatible with the plan of his treatise; but he has considered it fair (save in one or two special cases) to give the preference to those makers who have worked out their own plans of construction, and have thus furnished (to say the least) the general designs, which have been adopted with more or less of modification by others.

Simple Microscope.


26. Under this head, the common *hand-magnifier* or pocket-lens first claims our attention; being in reality a Simple Microscope, although not commonly accounted as such. Although this little instrument is in every one's hands, and is indispensable to the Naturalist,—as affording him the means of at once making such *preliminary* examinations as often afford him most important guidance,—yet there are comparatively few who know how to handle it to the best advantage. The chief difficulty lies in the steady fixation of it at the requisite distance from the object, especially when the lens employed is of such short focus, that the slightest want of exactness in this adjustment produces evident indistinctness of the image. By carefully resting the hand which carries the glass, however, against that which carries the object, so that both, whenever they move, shall move together, the observer, after a little practice, will be able to employ even high powers with comparative facility. The lenses most generally serviceable for hand-magnifiers, range in focal length from two inches to half an inch; and a combination of two or three of such in the same handle, with an intervening perforated plate of tortoise-shell (which serves as a diaphragm when they are used together), will be found very useful. When such a magnifying power is desired, as would require a lens of a quarter of an inch focus, it is best obtained by the substitution of a “Coddington” (§ 19) for the ordinary double-convex lens. The handle of the magnifier may be pierced with a hole at the end most distant from the joint by which the lenses are attached to it; and through this may be passed a wire, which, being fitted vertically into a stand or foot, serves for the support of the magnifying lenses in a horizontal position, at any height at which it may be convenient to fix them. Such a little apparatus is a rudimentary form (so to speak) of what is commonly understood as a Simple Microscope; the term being usually applied to those instruments in which the magnifying powers are supported otherwise than in the hand, or, in which, if the whole apparatus be supported by the hand, the lenses have a fixed bearing upon the object (§ 28).

27. *Ross's Simple Microscope.*—This instrument holds an intermediate place between the hand-magnifier and the complete Microscope; being, in fact, nothing less than a lens supported in such a manner, as to be capable of being readily fixed in a

variety of positions suitable for dissecting and for other manipulations. It consists of a circular brass foot, wherein is screwed a short tubular pillar (Fig. 14), which is "sprung" at its upper end, so as to grasp a second tube, also "sprung," by the drawing out of which the pillar may be elongated to about three inches. This carries at its upper end a jointed socket, through which a square bar about $3\frac{1}{2}$ inches long slides rather stiffly; and one end of this bar carries another joint, to which is attached a ring for holding the lenses. By lengthening or shortening the pillar, by varying the angle which the square bar makes with its summit, and by sliding that bar through its socket, almost any position and elevation may be given to the lens, that can be required for the purposes to which it may be most usefully applied; care

being taken in all instances, that the ring which carries the lens should (by means of its joint) be placed horizontally. At A is seen the position which adapts it best for picking out minute shells or for other similar manipulations; the sand or dredgings to be examined being spread upon a piece of black paper, and raised upon a book, a box, or some other support, to such a height, that when the lens is adjusted thereto, the eye may be applied to it continuously without unnecessary fatigue. It will be found advantageous that the foot of the microscope should not stand upon the paper over which the objects are spread, as it is desirable to shake this from time to time, in order to bring a fresh portion of the matters to be examined into view; and, generally speaking, it will be found convenient to place it on the opposite side of the object, rather than on the same side with the ob-

FIG. 14.




Ross's Simple Microscope.

rather than on the same side with the ob-

server. At B is shown the position in which it may be most conveniently set, for the dissection of objects contained in a plate or trough, the sides of which, being higher than the lens, would prevent the use of any magnifier mounted on a horizontal arm. The powers usually supplied with this instrument, are one lens of an inch focus, and a second of either half or a quarter of an inch. By unscrewing the pillar, the whole is made to pack into a small flat case, the extreme portability of which is a great recommendation. Although the uses of this little instrument are greatly limited by its want of stage, mirror, &c., yet for the class of purposes to which it is suited, it has advantages over perhaps every other form that has been devised.

28. *Gairdner's Simple Microscope.*—This little instrument, distinguished like the preceding for its simplicity and portability, is adapted to quite a different class of purposes; namely, the examination of minute transparent objects, especially those contained in fluid, such as Animalcules, Desmidiæ and Diatomacææ, Urinary deposits, &c. It consists (Fig. 15) of a Wollaston's doublet (§ 18), supported upon a handle, with which is also con-

FIG. 15.



Gairdner's Simple Microscope.

connected an elastic slip of brass, carrying a ring which surrounds the projecting centre of the under side of the doublet; this ring is made to approach nearer to, or to recede further from, the doublet, by means of a milled-headed screw which passes through the stem that supports the latter, and bears upon the slip of brass that carries the former; and to the side of it which is furthest from the doublet, a disk of very thin glass is cemented. In using this little instrument, a minute drop of the liquid to be examined is to be placed on the *under* side of the thin glass disk,—that is, on the side *away* from the doublet,—and it is to be covered by another disk, which will be drawn to the fixed disk, and supported in its place by the capillary attraction of the fluid for both. The instrument is then to be so held, that the eye, when applied to the doublet, looks at the light through the film of liquid; and when the focal adjustment is made by means of the milled head, any particles this may contain, of a size to be brought into view by the magnifying power employed, will be distinctly discerned. The instrument is usually constructed with but a single power, adapted to the class of objects for which it is to be employed; thus for the purposes of the botanical or zoological collector, a power of from 70 to 100 diameters is sufficient; whilst for the examination of urinary deposits, a power of 200 or more is desirable. It would not

be difficult so to modify it, however, by making the doublet to screw into a socket, instead of fixing it on the stem, that one power might be substituted for another on the same instrument; and the adjusting screw might then perhaps be dispensed with, since the focal adjustment might probably be made sufficiently well, by turning round the doublet itself in its screwed socket. The object-holder, too, might be so constructed as to receive a greater variety of objects, and even to hold preparations mounted on slips of glass; which would often be a matter of great convenience for class demonstration. All this, however, would add to the complexity and the cost of the instrument; the simplicity and low price of which at present constitute its chief recommendation. Though not suited for the higher purposes of a Microscope (the view of any object afforded by a doublet magnifying 100 or 200 diameters, being far inferior to that presented by only a tolerable achromatic), yet there is a certain class of observations for which it is particularly convenient,—those, namely, which only require a *recognition* of known forms. Thus, the collector of Diatomaceæ, Animalcules, &c., may by its means at once test the general value of the sample he has taken up, and may decide whether to throw it away as worthless, or to reserve it for more minute examination. And the Medical practitioner who is familiar with the aspect of Urinary deposits, may, by this little instrument (which he can carry in his waistcoat-pocket), discriminate on the spot the nature of almost any sediment whose character he may wish to know, without being obliged to have recourse to a more elaborate apparatus.¹

29. *Field's Simple Microscope*.—The general purposes of a simple Microscope are satisfactorily answered by the instrument, which has recently gained the premium awarded by the Council of the Society of Arts, and which is capable of being very effectively used in the examination of most of the objects for which such an instrument is suited. It consists (Fig. 16) of a tubular stem, about five inches high, the lower end of which screws firmly into the lid of the box wherein the instrument is packed when not in use. To the upper end of this stem, the stage is firmly fixed; while the lower end carries a concave mirror. Within the tubular stem is a round pillar, having a rack cut into it, against which a pinion works that is turned by a milled head; and the upper part of this pillar carries a horizontal arm which bears the lenses; so that, by turning the milled head, the arm may be raised or lowered, and the requisite focal adjustment obtained. Three magnifiers are supplied with this instrument; and by using them either separately or in combination (the lens of shortest focus being placed at the bottom, whenever two, or all three, are used together), a considerable range of powers,

¹ This Microscope, the invention of Dr. William Gairdner, of Edinburgh, is made by Mr. Bryson, optician, of that city.

from about five to forty diameters, is obtained. The stage is perforated with a hole at each corner; into any one of which may

FIG. 16.



Field's Simple Microscope.

be fitted a condensing lens for opaque objects (§ 64), or a pair of stage-forceps (§ 66). An aquatic-box for the examination of objects in water (§ 68) is also supplied.¹ This instrument is peculiarly adapted for educational purposes;² being fitted in every particular for the examination of botanical specimens, small insects or parts of insects, water flies, the larger animalcules, and other such objects as young people may readily collect and prepare for themselves; and such as have trained themselves in the application of it

to the study of Nature, are well prepared for the advantageous use of the Compound Microscope. But it also affords to the scientific inquirer all that is essential to the pursuit of such investigations, as are best followed out by the concurrent employment of a Simple and a Compound Microscope, the former being most fitted for the *preparation*, and the latter for the *examination*, of many kinds of objects;³—and it may be easily adapted to the purposes of dissection, by placing it between arm-rests (§ 104), or blocks of wood, or books piled one on another, so as to give a support for the hand on either side, at or near the level of the stage.

30. *Quekett's Dissecting Microscope*.—To the scientific investigator, however, it is generally more convenient to have a larger *stage* than the preceding instrument affords; and in this respect an arrangement devised by Mr. Quekett (Fig. 17) will be found extremely convenient. The stage, which constitutes the principal part of the apparatus, is a plate of brass (bronzed) nearly six inches square, screwed to a piece of mahogany of the same size and about $\frac{3}{8}$ ths of an inch thick; underneath this a folding flap


¹ The price of the instrument, with all these appurtenances, packed in a neat mahogany box, is only *half a guinea*; and the maker, Mr. G. Field, of Birmingham, is bound by his agreement with the Society of Arts to keep it always in stock. See also § 31.

² See Introduction, pp. 60, 61.

³ See Introduction, p. 63.

four inches broad is attached by hinges on each side; and the two flaps are so shaped, that, when closed together, one lies closely upon the other, as shown in Fig. 17, B. These flaps, when opened, and kept asunder by a brass bar, as shown in Fig. 17, A, give a firm support to the stage at a convenient height;

FIG. 17.



Quekett's Dissecting Microscope.

and the bar also carries a socket, into which the stem of the mirror-frame is inserted. At the back of the stage-plate is a round hole, through which a tubular stem slides vertically, that carries at its summit the horizontal arm for the magnifying powers; this stem may or may not be furnished with a rack-and-pinion movement; but the author's experience leads him strongly to recommend that it should be provided with this means of making the focal adjustment; since the sliding action, independently of the greater trouble it always involves, is apt to become uneven and difficult, especially if the surface of the stem should have become roughened by the corrosion of sea-water, or by the action of acids, salines, &c., used as reagents. The same frame which carries the mirror, is also made to carry a lens which serves as a condenser for opaque objects; its stem being then fitted into

a hole in the stage, at one side, or in front of, its central perforation. The instrument is usually furnished with three magnifiers, namely, an inch and half-inch ordinary lenses, and a quarter-inch Coddington (§ 19); and these will be found to be the powers most useful for the purposes to which this instrument is specially adapted. The lenses, mirror, condenser, cross-bar, vertical stem, and milled head, all fit into hollows cut for their reception on the under side of the stage, and are then covered and kept in place by the side flaps; so that, when packed together, and the flaps kept down by an elastic band, as shown in Fig. 17, B, the instrument is extremely portable, furnishing (so to speak) a case for itself. It may be easily made to serve as a Compound microscope, by means of an additional stem and horizontal arm, carrying a light "body." The principal *disadvantages* of this very ingenious and otherwise most convenient arrangement, are that it must be always used with the light *in front* of the observer, or nearly so, since the side-flaps interfere with the access of side-light to the mirror; and that the obstruction of the side-flaps also prevents the hands from having that ready access to the mirror, which is convenient in making its adjustments. These inconveniences, however, are trifling, when compared with the great facilities afforded for scientific investigation by the size and firmness of the stage; and the author can confidently recommend the instrument for all such purposes, from much personal experience of its utility.

Compound Microscope.

The various forms of Compound Microscope may be grouped with tolerable definiteness into two principal classes; one consisting of those instruments, whose size and general plan of construction adapt them only for the ordinary methods of observation; whilst the other includes those which are suited to carry the various accessories, whose use enables the observer not only to work with more facility and certainty, but, in some instances, to gain information respecting the object of his examination, which he could not obtain without them. It is true that some of the most important of these accessories *may* be applied to the smaller and lighter kind of Microscopes; but when it is desired to render the instrument complete by the addition of them, it is far preferable to adopt one of those larger and more substantial patterns, which has been devised with express reference to their most advantageous and most convenient employment. In nearly all the instruments now to be described, the same basis of support is adopted, namely, a triangular "foot," from which arise two uprights; and between these the microscope itself is swung, in such a manner that the weight of its different parts may be as nearly as possible balanced above and below the centres of suspension, in all the ordinary positions of the instrument. This double support was first introduced by Mr. George Jackson, who

substituted two pillars (a form which Messrs. Smith and Beck still retain in their Large Compound Microscope, Fig. 29) for the single pillar connected with the microscope itself by a "cradle-joint" (as in Fig. 20) which was previously in use; but in place of pillars screwed into the tripod base, a pair of flattened uprights, cast in one piece with it, is now generally adopted, with a view both to greater solidity and to facility of construction. Messrs. Powell and Lealand, it will be observed, adopt a tripod support of a different kind (Fig. 28), still, however, carrying out the same fundamental principle, of swinging the microscope itself between two centres. Two different modes of giving support and motion to the "body" will be found to prevail. One consists in its attachment at its base to the transverse "arm," which is borne on the summit of the movable stem, whose rack is acted on by the pinion of the milled head, as in Figs. 18, 27, 28; whilst in the other, the body is supported along a great part of its length by means of a solid "limb," to which is attached the pinion that acts on a rack fixed to the body itself, as in Figs. 21, 22, and 29. The former method has the advantage of enabling the body to be turned aside by the rotation of the transverse arm upon the summit of the stem,—a movement which is often convenient, both as leaving the stage clear for dissection, &c., and as enabling the objectives to be more readily exchanged; but it is subject to the disadvantage, that unless the transverse arm and the body are constructed with great solidity, the absence of support along the length of the latter leaves it subject to vibration, which may become unpleasantly apparent when high powers are used, giving a dancing motion to the objects. With a view of preventing this vibration, Messrs. Powell and Lealand connect the top of the "body" with the back of the transverse arm, by a pair of oblique "stays" (Fig. 28). The second method of support is decidedly superior in steadiness, a perfect freedom from tremor being obtained with less solidity, and therefore with less cumbrousness; the mode in which the rack is applied, moreover, in the microscopes of Messrs. Smith and Beck (most of which are constructed upon this plan) gives to it a peculiar smoothness and easiness of working; but the traversing movement of the body is sacrificed. Although some attach considerable importance to this movement, the author's experience of instruments constructed upon both plans, leads him to give a preference to the second.

31. *Field's Compound Microscope*.—The first of the simpler forms which we shall more particularly describe, is that to which the medal of the Society of Arts has been recently awarded, not as a testimony to the *perfection* of its construction, but as marking the highest degree of excellence among the instruments sent in competition, that seemed consistent with the *cheapness*¹ which

¹ The price of this instrument, complete, with two eye-pieces and two achromatic objectives giving a range of power from about 25 to 200 diameters, condenser on a

was the fundamental requirement (see Introduction, p. 61). The tripod foot (Fig. 18), with its pair of uprights, is of cast iron; and affords a very firm and steady basis of support. The centres of suspension by which the microscope is swung between these up-

Fig. 18.



Field's Compound Microscope.

rights, are attached to the hollow pillar that bears all the other parts. Just above them, when the instrument is in a vertical position, is a milled head on either side, which acts on a rack cut into the stem that rises from the pillar, and carries the body on a transverse arm, thus giving the "coarse" adjustment for focal distance; whilst the "fine" adjustment is given by another milled head (seen edgewise in the figure) in the transverse arm, which turns a screw whose extremity acts upon a lever that produces a slight change in the distance between the object-glass and the object, by elevating or depressing a tube that carries the former,—this tube being

so fitted to the lower end of the body as to slide freely within it, and being pressed downwards by a spring, whilst it is raised upwards by the lever-action just named. The additional advantage is gained by this arrangement (which is the one adopted with some modification by most Microscope makers), namely, that if the object-glass should be carelessly forced down so as to press upon the object, the yielding of the spring-tube prevents any serious injury to the one, and to a certain extent protects the other. The stage, which is firmly attached to the pillar, is furnished on its upper surface with a movable brass ledge, against which the object rests when the stage is inclined in any degree to the horizon; this ledge should slide smoothly and easily from the back to the front of the stage, but should have at the same time sufficient hold upon it to retain its position and to support the object, at whatever point it may be left. At a little distance beneath the stage, there is attached to it a "diaphragm plate," perforated with holes of various sizes for the regulation of the quantity of light admitted to transparent objects (§ 55), and also affording, in one of its positions, a dark background, which is

separate stand, stage-forceps, and live-box, in a mahogany case, is only *three guineas*; and the maker, Mr. G. Field, of Birmingham, is bound by his agreement with the Society of Arts to keep it always in stock, so as to supply any purchaser at once.

useful when opaque objects are being viewed. The stage is perforated at one of its front corners with a hole, into which fits a pair of stage-forceps (§ 66). The mirror, which is concave on one side and plane on the other, is attached, not to the pillar, but to a tube which slides upon it, so that its distance from the under side of the stage may be increased or diminished. The condenser for opaque objects is mounted on a separate stand (§ 64). The simplicity of the construction of this Microscope, and the facility with which all those adjustments may be made that are required for the purposes which it is intended to fulfil, should constitute, with its low price, a great recommendation to those who value a Microscope rather as a means of interesting recreation for themselves, or of cultivating a taste for the study of nature and a habit of correct observation in the young, than as an instrument of scientific research. It is not, of course, to be expected that it should bear comparison, in regard either to the mechanical finish of its workmanship, or to the perfection of its optical effects, with Microscopes of many times its cost; but it is infinitely superior to the best Microscope ever constructed on the old (non-achromatic) plan; and it is greatly to be preferred in its mechanical arrangements to any of the earlier achromatic microscopes, which it at least equals in optical performance.

32. *Highley's Hospital Microscope.*—The scale of this instrument is somewhat larger than that of the preceding, and its workmanship more finished and substantial. The tripod stand, the stage and its fittings, and the mirror, almost exactly resemble those first described; but the body, which is longer, is supported in a different manner. The pillar to which the stage and the mirror are attached, is prolonged upwards, and then forms a kind of "limb," to which is affixed a tube slit down in front; and within this tube the "body" slides up and down, with sufficient freedom to allow of being easily moved, yet with sufficient stiffness to remain firm in any position in which it may be left. In the simple form of this instrument here delineated, the sliding action

FIG. 19.



Highley's Hospital Microscope.

affords the only means of making the "coarse" adjustment (§ 25, II); but a rack-and-pinion movement may be introduced at a trifling additional cost. The "fine" adjustment is made by a milled head in front of the lower end of the body, which acts directly upon a tube sliding within it that carries the objectives. This instrument is particularly adapted, by the roominess of its stage, for the examination of pathological specimens; and, when the body is provided with a rack movement, it forms an unexceptionable microscope for general purposes, and may even be furnished with a movable stage, achromatic condenser, polarizing apparatus, &c.¹

33. *Nachet's Microscope*.—Until a comparatively recent period, all save the most elaborate and expensive forms of Compound Microscope constructed by Continental opticians, were adapted for use in the vertical position only. M. Nachet, however, has now so modified his ordinary pattern, that the instrument may be inclined (like the preceding) at any angle; and he has thus rendered it a very convenient, as well as a cheap and portable Microscope. The basis consists of a somewhat oval foot, with a single pillar rising from a little behind its centre; and at the top of this pillar is a "cradle-joint," which supports the stage and the upright stem that carries the body. The transverse arm, however, is attached, not directly to the summit of this stem, but to a tube which slides over it; and this tube can be raised or lowered by turning the milled head at its summit (which acts upon a screw that enters the stem), whereby a "fine" adjustment is obtained, that acts through the transverse arm upon the body which it carries. The "coarse" adjustment is effected, as in the preceding case, by sliding the body through an outer tube which grasps it; the latter being fixed into the transverse arm. The mode in which the object is supported upon the stage, when this is inclined, is very simple and ingenious, and is in some respects preferable to the sliding-ledge generally used by English makers. Near each side of the stage is seen a somewhat elastic strip or tongue of sheet brass, the front extremity of which is free, but which is attached at its hinder end to a pin that passes through a hole in the stage, in which it works very easily. This pin is prolonged for about $\frac{3}{4}$ ths inch beneath the stage, and then terminates in a broad flat head; and it is surrounded by a slender spiral spring, which, bearing at its two ends against the under side of the stage and the head of the pin, tends to depress the latter, and thus to bring the brass tongue into close apposition with the stage when nothing intervenes, and to bind down anything that may be placed between them. In making use of this little apparatus, it is most convenient to employ both hands, in such a manner that the thumb and forefinger of each shall

¹ The cost of this instrument essentially depends upon the number and magnifying power of the objectives supplied with it; it is usually provided, however, with a 1-inch and $\frac{1}{2}$ -inch; and is then sold (without the rack movement) at £6 10s. This sum, however, does not include either a case or any accessory apparatus.

hold one end of the slip of glass whereon is placed the object under examination, whilst one of the other fingers of each hand is used to push up the head at the end of the pin, so as to lift the tongue from the stage; the slip of glass can then be moved from side to side, or up and down, with the most perfect freedom, and may be firmly secured at any point by ceasing to press upon the heads of the pins, which will then be forced down by the springs, so as to bring the tongues to bear on the slip of glass. When the microscope is used in a vertical position, for the examination of urinary deposits, &c., no means of fixing the object being required, it is convenient to turn the tongues backwards, so as not to occupy any part of the stage. The advantages of this arrangement are the perfect freedom with which the slip of glass can be moved under the objective, either in finding a minute object, or in examining the surface of a larger one; and the facility and exactness with which it is retained at any point, at which it may be desired to fix it. The disadvantages are, the necessity of using both hands to move the object; and the interference of the tongues with the movement of the object from side to side, when it is large enough to require a considerable range; on which last account the plan is unsuited to the use of an aquatic box. The stage is furnished on its under surface with a diaphragm plate, not mounted as a wheel, but sliding in a straight line, which is a less convenient arrangement; and to its lower side is also attached a stem that carries the mirror, the distance of which from the stage is not capable of variation. This instrument is distinguished by its simplicity and cheapness, and by its adaptation to many of the wants of the scientific inquirer.¹ One of its chief disadvantages is the small size (especially the narrowness) of its stage, which cramps the operations of the observer; and hence it will not be found nearly so convenient to the young microscopist, as the equally simple patterns in common use in

FIG. 20.



Nachet's Compound Microscope.

¹ With three objectives and three eye-pieces, giving a range of magnifying powers from about 50 to about 500 diameters, it is sold in Paris for 190 francs.

this country. Those, however, who are carrying on researches upon objects too minute to make this objection felt (such, for example, as urinary deposits), and who need high magnifying powers, without requiring these to possess the greatest attainable perfection, will find this Microscope extremely well suited to their wants. Another instrument constructed by M. Nachet upon the same general plan, but upon a larger scale, is capable of being fitted with Achromatic condenser, Polarizing apparatus, Micrometer eye-piece, Stage movements, &c.; in the arrangement of which accessories, much skilful contrivance is shown.

FIG. 21.



Smith and Beck's Student's Microscope.

The *Binocular Microscope* of the same ingenious Optician will be described further on (§ 40).

34. *Smith and Beck's Student's Microscope*.—Of the patterns yet devised for a microscope of simple construction, which shall yet be capable of answering every essential purpose whether of display or of investigation, that of Messrs. Smith and Beck appears to the Author to be (to say the least) among the best; and he recommends it with the more confidence, since he has for many years employed one of these Microscopes as his own *working* instrument. There is nothing distinctive in the tripod support, or in the mode in which the microscope itself is suspended between the uprights. But the "body" rests for a great part of its length upon a "limb" of solid brass, ploughed into a groove for the reception of the rack which is attached to the body; this groove being of such a form, that the rack is firmly held in it, whilst it slides smoothly through it. The great advantage

of this method of construction over any other in which the rack-and-pinion movement is made to act directly on the body, is that it renders impossible any of that *twist* which tends to throw the object more or less completely out of the field, and secures that exact *centering* which is essential to the optical perfection of the instrument. The upper end of the body is furnished with a "draw tube," by which its length can be increased; and one side

of this is graduated to inches and tenths. The advantages of this arrangement will be explained hereafter (§ 43). The "fine" adjustment is effected by means of a milled head, situated just behind the base of the stem that bears the limb; this acts on a screw, the turning of which (by a contrivance that need not be described in detail) depresses the stem with the limb and body attached to it, so as to bring the objective nearer to the object; whilst if the pressure of the screw be withdrawn, by turning the milled head in the opposite direction, the tubular stem (with the limb and body) is carried upwards by a spiral spring in its interior, thus increasing the distance of the objective from the object. This adjustment is remarkable for its sensitiveness, and for its freedom from any displacing action upon the image. The only other peculiarity that need be noticed in this instrument, is the mode in which the object is borne upon the stage; for, instead of resting against a ledge, it lies upon a kind of fork, which slides in grooves ploughed out of the stage, and which moves with such facility, that the pressure of a single finger upon one of the upright pins at the back of the fork is sufficient to push it in either direction. At the extremity of one of the prongs of this fork, is a "spring clip" for securing the object by a gentle pressure, which is particularly useful when the microscope is placed in a horizontal position for drawing with the camera lucida (§ 49), the stage being then vertical. And at the extremity of the other prong is a hole for the insertion of the pin of the stage-forceps, which thus gains the advantage of the sliding movement of the fork, in addition to its own actions. This instrument can easily be made to receive the addition of an achromatic condenser and of a polarizing apparatus; it *may* also be fitted with a traversing stage, but there is scarcely sufficient room for its working, to render such an addition worth its cost.¹

35. *Smith and Beck's Dissecting Microscope*.—A modification of the preceding pattern has been made for the special purpose of carrying on dissections under the Compound Microscope, without any interference, however, with the use of the instrument for all ordinary purposes. The general plan of the instrument (Fig. 22), as will be at once apparent, is essentially the same as that of the

¹ No working Physiologist or Naturalist can *require*, in the Author's opinion, a better instrument than the above; unless he be directing his attention to some particular class of objects, which need the very highest microscopic refinements for their elucidation. The cost of the instrument, fitted with two eye-pieces, condenser for opaque objects, aquatic box, and stage-forceps (with case) about £7; the cost of the objectives depends upon their magnifying power and upon their angular aperture. Those most serviceable for ordinary purposes are the $1\frac{1}{2}$ inch, $\frac{3}{4}$ inch, $\frac{1}{4}$ inch, and $\frac{1}{2}$ inch, whose respective prices are £3, and 3, 5, and 6 guineas; the first and third, or the second and fourth, of these may be selected in the first instance, and the others added at any time; the addition of the $\frac{1}{4}$ inch (which the unpractised microscopist is scarcely likely to employ to advantage, and which is only useful for a very limited set of purposes) may be postponed until it is really needed and can be effectually employed. More can be seen of most objects by the proper management of such a $\frac{1}{2}$ inch as Messrs. S. and B. now supply, than could have been made out by the $\frac{1}{4}$ of a few years back. These opticians are now constructing a new pattern of Student's Microscope, complete in itself, with two good powers, which will be well adapted to the most important uses of

Student's Microscope; but the stage is much longer from back to front, so as to give more room; and from the back of it rises a strong curved limb for the support of the body, which is made to slide upon it, as in the previous case, by a rack-and-pinion movement. A second milled head is seen above that by which the focal adjustment is made; and this acts by means of a rack upon the draw-tube, which it brings out or shuts in, without the necessity of holding the body with the other hand,—a movement which will be found of very great advantage, when the “erecting

FIG. 22.



eye-piece” (§ 44) is employed for varying the magnifying power. The chief use of this erecting eye-piece, which screws into the lower end of the draw-tube (Fig. 32), in the Dissecting Microscope, is to *erect* the image (as its designation implies), and thus to facilitate the employment of dissecting instruments upon an object under inspection, the selection of minute shells, &c., or other manipulations, which cannot be so conveniently carried on, save after long practice, when the object is inverted. As the “fine” adjustment cannot, in this pattern, be applied to the “limb,” it is attached (if required) to the lower end of the body itself, as in Messrs. Smith and Beck’s larger Microscope (Fig. 29); but for the purposes to which such an instrument as this is usually applied, the fine adjustment is seldom needed, the rack-movement being sufficiently exact and sensitive to furnish all that is needed for low and medium powers.

the Physiologist and Medical Practitioner, without exceeding ten pounds in price. To those, however, who, though obliged to limit their first outlay, contemplate making subsequent additions, the Author would strongly recommend the choice of the instrument described in the text, as one on which such additional expenditure may be more profitably bestowed. Since the above were written, Messrs. Smith and Beck have brought out the “Educational Microscope” there alluded to; and after a careful examination of it, the Author can strongly recommend it as admirably adapted to the purposes for which it is intended. It is fitted with two eyepieces and two objectives, giving a range of powers from 55 to 350 diameters; and may also be furnished with an extra low power for large opaque objects, at a small additional cost. For the additional sum of £5, a Lieberkühn, Parabolic Illuminator, Polarizing Apparatus, Camera Lucida Prism, Aquatic Box, and Zoophyte Trough are supplied; all fitted into the same very portable case, and rendering the instrument extremely complete.

When this addition is made, however, the instrument is adapted to any kind of work to which the preceding can be applied; it can receive the same fittings; and in consequence of the larger dimensions of the stage, a traversing movement may be readily added to it. This Microscope may thus be rendered a very complete instrument; but it will scarcely be so convenient in use, as the instruments which are specially planned for a greater range of adaptations; and the particular advantage it possesses, is for the purpose indicated by its designation.

36. *Warington's Universal Microscope*.—A new set of adaptations for special purposes, called for by new requirements, has been recently devised by Mr. Warington; who, by different combinations of the same very simple materials, has produced an instrument which may be used in four different modes, and which may fairly, therefore, be designated a "universal" microscope. Mr. Warington's original object was to provide an arrangement, whereby the Compound Microscope should be brought to bear upon living objects in an Aquarium, when these might be either in contact with one of the glass sides, or be not far removed from it. This he accomplished by making use of the body of a Student's microscope (§ 34), with the grooved limb in which it slides, and attaching the latter by a strong cradle-joint to a tubular stem, which could be fixed at any height upon the edge of the table that supports the Aquarium, by means of a clamp with a binding screw. Subsequently Mr. W. dispensed with the rack; attaching the cradle-joint at the top of the tubular stem to an outer tube, within which the sliding of the body acts as a "coarse" adjustment; and providing a "fine" adjustment (by an ingenious plan of his own) at the object end of the body itself. To the Author, however, it has seemed far more convenient to retain the rack; and this he has combined with the sliding tube, thus obtaining great facility of adjustment with no perceptible "twist;" and




FIG. 23.

Warington's Universal Microscope, as arranged for viewing objects in an Aquarium.

the arrangement of the apparatus, with this modification, is shown in Fig. 23. If the rack be well cut, there will be no occasion for a "fine" adjustment; since the purposes to which this arrangement is adapted, only require low or moderate powers. When the instrument is set up in the above position, the body may be moved like a swivel from side to side, and it may be inclined downwards at any degree of obliquity; but its most

suitable position will generally be the horizontal, with its axis

FIG. 24.



Warrington's Universal Microscope, as arranged for Dissection in a large trough. (N. B.—By drawing the stem *a* through the clamp, the body may be shifted to such a distance from the wooden base, that the latter need not interfere with the dissecting trough.)

FIG. 25.



Warrington's Universal Microscope, arranged for ordinary use.


directed at right angles to the flat side of the Aquarium. It is obvious that the very same instrument, turned from the horizontal into the vertical position, by attaching the clamp (as in Fig. 24) to the edge of a wooden strutt rising vertically from a horizontal slab, instead of to the edge of a horizontal table, becomes extremely well suited for examining objects which are in course of dissection in a trough too large to be conveniently transferred to the stage of the microscope, for looking over minute shells spread out on a sheet of paper, and for other purposes for which a special form of dissecting microscope has been devised by Messrs. Powell and Lealand. But again, by turning up the \perp shaped support constructed for the last-named purpose, so that it shall rest (as it were) on two legs like the Greek λ , and then clamping the stem that carries the body to its highest edge, the instrument acquires a position very suitable for ordinary microscope work; and nothing is wanted to adapt it to this, save the addition of a stage and a mirror, each of which may be so constructed as to fit into a

brass socket let into the wooden support, thus completing the Microscope in the form represented in Fig. 25. This is not the last of the adaptations of which the instrument is capable; for the wooden support remaining at the same inclination, the body may be brought to the perpendicular, by shifting its stem in the clamp and by altering its angle at the cradle-joint; whilst a horizontal position may be given to the stage, by fitting it into another socket (Fig. 26); in this arrangement,

moreover, the stage acquires an increase of firmness, from the bearing of a plate that projects at right angles from its under surface, upon the inclined face of the wooden support. Thus a dissecting microscope is formed, which has many of the advantages of that of Messrs. Smith and Beck; being subject, however, to the important drawback, that the mirror cannot be so placed as to reflect the light upwards through the axis of the microscope. (A means of remedying this, however, might perhaps be contrived without much difficulty or cost.) On the left side of the slanting support, at a short distance above the stage, is a hole into which may be fitted

either the stem of a condensing lens for opaque objects, or the stem of the stage-forceps; either or both of which may also be fitted into holes in the front corners of the stage. The stage is provided with a sliding ledge for the support of objects in an inclined position; and it might also be furnished, if required, with a diaphragm plate. One of the chief merits of the instrument, however, being lightness and portability, it would not be desirable to encumber it with many accessories. For convenience of packing, the shorter portion of the \perp piece may be connected with the longer by strong pins fitted into sockets, instead of being permanently fixed, so that the two can be readily disconnected and one part laid flat upon the other; and the whole apparatus will then lie within a very small compass. The distinctive peculiarity of this instrument consists in the extreme simplicity of the means by which a variety of useful ends are obtained. It is scarcely one that should be recommended to the beginner; since it is in several respects not so well adapted for *ordinary* work, as the forms already described. But it is a most valuable addition to the Microscopic apparatus of

FIG. 26.



Warington's Universal Microscope, arranged for dissecting on the stage.

the Naturalist; and may be constructed at so trifling an expense, to work with any objectives he already may possess, that a considerable demand may be anticipated for it.¹

37. We now pass to an entirely different class of instruments, —those of which the aim is, not simplicity but perfection; not the production of the best effect with limited means, but the attainment of everything that the Microscope *can* accomplish, without regard to cost or complexity. This object has been certainly carried out by the Opticians of our own country, much more completely than by those of the Continent; and it seems but fair towards the three principal London makers, by whose labors the present admirable results have been attained, that the pattern finally adopted by each should be here delineated and described. Without any invidious preference, the first place may fairly be

FIG. 27.



Ross's Large Compound Microscope.

assigned to the *Large Compound Microscope of Mr. Ross*; not only as being the one which was first brought (in all essential features at least) to its present form, but also because it is that which contains the greatest number of provisions for investigating objects in a variety of different modes. The general plan of Mr. Ross's Microscope will be seen to be essentially the same with that which has been followed by Mr. Field in the simple form of this instrument first described (§ 31), as well as by many other makers; but it is carried out with the greatest attention to solidity of construction, in those parts especially which

are most liable to tremor; and we are informed by Mr. Ross, that every part has been tested by the "inverted pendulum"

¹ This instrument has been made for Mr. Warrington and for the Author by Mr. Salmon, 100 Fenchurch Street: who supplies it, on either plan, without objectives or case; but with condenser and stage-forceps, for 3 guineas.

(an instrument contrived to indicate otherwise insensible vibrations), and either strengthened or reduced as might be found necessary, so as to obtain an *equality of vibration* between the stage and the optical part, which will prevent any perceptible tremor in the image. The "coarse" adjustment is made by the large milled head situated just behind the summit of the uprights, which turns a pinion working into a rack cut on the back of a very strong flattened stem, that carries the transverse arm at its summit; a second milled head (which is here concealed by the stage fittings) is attached to the other end of the axis of the pinion (as in Fig. 18), so as to be worked with the left hand. The "fine" adjustment is effected by the milled head on the transverse arm just behind the base of the "body;" this acts upon the "nose" or tube projecting below the arm, wherein the objectives are screwed. The other milled head seen at the summit of the stem, serves to secure the transverse arm to this, and may be tightened or slackened at pleasure, so as to regulate the traversing movement of the arm; this movement is only allowed to take place in one direction, namely, towards the right side, being checked in the opposite by a "stop," which secures the coincidence of the axis of the body with the centre of the stage and with the axis of the illuminating apparatus beneath it. It is in the movements of the stage, that the greatest contrivance is shown; these are three, namely, a traversing movement from side to side, a traversing movement from before backwards, and a rotatory movement. The traversing movements, which allow the platform carrying the object to be shifted about an inch in each direction, are effected by the two milled heads situated at the right of the stage; and these are placed side by side, in such a position that one may be conveniently acted on by the fore-finger, and the other by the middle finger, the thumb being readily passed from one to the other. The traversing portion of the stage carries the platform whereon the object is laid, which has a ledge at the back for it to rest against; and this platform has a sliding movement of its own, from before backwards, by which the object is first brought near to the axis of the microscope, its perfect adjustment being then obtained by the traversing movement. To this platform, and to the traversing slides which carry it, a rotatory movement is imparted by a milled head, placed underneath the stage on the left hand side; for this milled head turns a pinion which works against the circular rack (seen in the figure) whereby the whole apparatus above is carried round about a third of a revolution, without in the least disturbing the place of the object, or removing it from the field of the microscope. This rotatory movement is useful for two purposes; first, in the examination of very delicate objects by oblique lights, in order that, without disturbing the illuminating apparatus, the effect of the light and shadow may be seen *in every direction*, whereby important additional information is often gained; and,

secondly, in the examination of objects under polarized light, a class of appearances being produced by the rotation of the object between the prisms, which is not developed by the rotation of either of the prisms themselves. Below the stage, and in front of the stem that carries the mirror, is a dovetail sliding bar, which is moved up and down by the milled head shown at its side; this sliding bar carries what is termed by Mr. Ross the "secondary stage" (omitted in the figure for the sake of simplicity), which consists of a cylindrical tube for the reception of the achromatic condenser, the polarizing prism, and other fittings; to this secondary stage, also, a rotatory motion is communicated by the turning of a milled head; and a traversing movement of limited extent is likewise given to it by means of two screws, one on the front and the other on the left hand side of the frame which carries it, in order that its axis may be brought into perfect coincidence with the axis of the "body." The special advantages of this instrument consist in its perfect steadiness, in the admirable finish of its workmanship, and in the variety of movements which may be given both to the object and to the fittings of the secondary stage. Its disadvantages consist in the want of portability, necessarily arising from the substantial mode of its construction; and in the multiplicity of its movable parts, which presents to the beginner an aspect of great complexity. This complexity, however, is much more apparent than real; for each of these parts has an independent action of its own, the nature of which is very soon learned; and the various milled heads are so disposed, that the hand readily (and at last almost instinctively) finds its way from one to the other, so as to make any required adjustment, whilst the eye is steadily directed to the object. To the practised observer, therefore, this multiplication of adjustments is a real saving of time and labor, enabling him to do perfectly and readily what might otherwise require much trouble, besides affording him certain capabilities which he would not otherwise possess at all.

38. *Powell and Lealand's Compound Microscope.*—This instrument, represented in Fig. 28, is far lighter than the preceding in its general "build," without being at all deficient in steadiness; it has not, however, some of those improvements for which Mr. Ross's plan of construction is especially adapted. The three-legged stand gives a firm support to the trunnions that carry the tube to which the stage is attached, and from which a triangular stem is raised, by the rack-and-pinion movement set in action by the double milled head, whereby the "coarse" adjustment of the focus is obtained. The triangular stem carries at its summit the transverse arm, which contains (as in Mr. Ross's Microscope) the lever action of the "fine" adjustment; and this is acted on by the milled head at the back of the arm, whence also pass two oblique stays, which, being attached to the upper part of the body, assist in preventing its

vibration. The stage is provided with a traversing movement in each direction, to the extent of about three-quarters of an inch; this is effected on the plan known as Turrell's, in which

FIG. 28.

the two milled heads are placed on the same axis, instead of side by side, one of them being also repeated on the left hand of the stage, so that the movements may be communicated either by the right hand alone, or by both hands in combination. The platform which carries the object is made to slide, as in the preceding case, on the summit of the traversing apparatus; and it has not only a ledge whereon the object may rest, but also a "spring clip" for securing it whenever the stage may be placed in a vertical position. This platform, moreover, is so connected with the traversing apparatus, that it may be turned round in the direction of its plane; but as



Powell and Lealand's Large Compound Microscope.

this rotation takes place above instead of beneath the traversing apparatus, there is no security that the centre of rotation shall coincide with the axis of the optical portion of the instrument; so that, unless this adjustment have been previously made, the object will be thrown completely out of the field of view when the platform is made to revolve. Hence, although this movement is of great use in facilitating the full examination of an object, by enabling the observer to bring it into the field of view in every variety of position, it does not serve, like the rotatory movement of Mr. Ross's stage, to change the position of the object in regard to the illuminating apparatus, without disturbing the observer's view of it. The condenser for transparent objects, the polarizing apparatus, &c., are here fitted to the under

side of the principal stage itself, instead of to an independent or secondary stage; an arrangement which, though convenient as regards compactness, admits of less variety of adjustment than is afforded by the latter plan. The mirror, instead of being swung loosely upon two centres, is pivoted to one end of a quadrant of brass, of which the other end is pivoted to a strong pin that projects from the sliding tube; a spring being so attached to each of these pivots, as to give to the movements of the mirror that suitable degree of stiffness, which shall prevent it from being disturbed by a passing touch. No instrument can be better adapted than this to all the *ordinary* wants of the Microscopist; there are very few purposes which it cannot be made to answer; and there are many who will consider that its deficiency as to these is counterbalanced (to say the least) by its comparative simplicity and portability, as well as by its lower cost. For the sake, however, of such as may desire the power of obtaining a more oblique illumination, than is permitted by the construction of the stage in the instrument just described, Messrs. P. and L. have recently brought out a new pattern, in which the thickness of the stage is greatly reduced, a sub-stage is provided for the reception of the condenser and other fittings, and the mirror is mounted on a doubly extending arm.

39. *Smith and Beck's Large Microscope.*—The general plan of this instrument (Fig. 29) nearly resembles that of the “dissecting microscope” of the same makers, already noticed (§ 35), so far, at least, as regards the mode of supporting the body, and of effecting the focal adjustments; whilst in the construction of the stage, and in the arrangement of the fittings beneath, it differs from all the microscopes hitherto described. The stage is furnished with the usual traversing movements; but it is distinguished by its *thinness*; and this is of importance in certain cases, as admitting of a more oblique illumination than could otherwise be obtained, and also as allowing the construction of the achromatic condenser (§ 56) to be much simplified. The platform for the object is fitted upon the traversing apparatus, in the same mode as in the microscope last described, and possesses the same kind of rotatory movement. Beneath the stage is a continuation of the gun-metal “limb” which carries the body; and this is ploughed out into a groove for the reception of a sliding-bar, which carries what may be termed the “secondary body,” namely, a short tube (seen beneath the stage), capable of being moved up and down by a milled head, and fitted for the reception of the achromatic condenser, polarizing apparatus, &c. This “secondary body” consequently answers the same purpose as the “secondary stage” of Mr. Ross’s microscope, and its relations to the other parts of the instrument are essentially the same; but it differs in the following particulars:—first, that by being made to work in a groove which is in perfect correspondence with that wherein the principal “body” works (this corre-

spondence being secured by the action of the planing machine that ploughs both grooves), the "secondary" body always has its axis so perfectly continuous with that of the first, that no special adjustment is needed to "centre" the greater part of the illuminating apparatus; and, secondly, that the tube will carry the achromatic condenser at its upper end, the polarising prism at its lower, and the selenite plates between the two, a combination that cannot be made in any other instrument (§ 63). Moreover, as all these fittings are received into a tube of which the exact size and position are assured, the makers of this instrument can supply additional apparatus at any time, with the certainty of its accurate adjustment. This "secondary body," however, has not the rotatory movement possessed by Mr. Ross's "secondary stage;" and to the limited class of purposes, therefore, which that movement is adapted to serve, it cannot be adapted.

The mirror is hung in the usual way between two centres; but the semicircle that carries these, instead of being at once pivoted to the tube which slides upon the cylindrical stem, is attached to an intermediate arm; and by means of this it may be placed in such a position as to reflect light very obliquely upon the object, and thus to bring out a new set of appearances, with which it is very important in certain cases to be acquainted. In regard to weight and complexity, this instrument holds a position intermediate between the two last described. The mode in which the body is supported, appears to the author decidedly preferable to that adopted by the other makers; and though it has the disadvantage of separating the focal adjustments from each other and from the stage motions more widely than is the case in the two preceding instruments, yet the difference is scarcely perceptible in practice. The milled heads acting on the former are both of them in positions in which they are

FIG. 29.



Smith and Beck's Large Compound Microscope.


easily reached by the left hand, when the elbow is resting on the table; whilst the right hand finds the milled heads of the traversing stage and of the secondary body in close proximity to each other. The imperfection of the means of giving rotation to the object, constitutes in this, as in Powell and Lealand's microscope, a point of inferiority to Ross's; the number of cases in which such a movement is important, however, is by no means considerable. On the other hand, the arrangement of the illuminating apparatus in Smith and Beck's Microscope, seems to the author to present some decided advantages over that adopted by either of the other makers; and in point of general excellence of workmanship, this instrument cannot be surpassed.

Without any invidious comparisons, it may be safely said that whoever desires to possess a *first-class* Microscope, cannot do better than select one of the three instruments last described; the excellence of the optical performance of the lenses supplied by their respective makers, being so nearly on a par, that the choice may be decided chiefly by the preference which the taste of the purchaser, or the nature of the researches on which he may be engaged, may lead him to entertain, for one or other of the plans of construction which has now been brought under notice.

40. *Nachet's Binocular Microscope*.—Since that remarkable invention of Prof. Wheatstone, the Stereoscope, has led to a general appreciation of the value of *binocular vision*, in conveying to the mind a notion of the *solid forms* of bodies, various attempts have been made to apply the same principle to the Microscope. To any one who understands the principle of the Stereoscope, a little consideration will make it obvious that this end might be *theoretically* attained, by placing two microscope-bodies at such an angle of inclination, that their respective object-glasses should point to the same object, whilst their eye-pieces should be at the ordinary distance of the right and left eyes from each other; but this *practical* difficulty will obviously and necessarily arise, in bringing the two microscopes into the requisite convergence,—that the axes of the instruments cannot be approximated sufficiently closely at their lower ends, unless the objectives employed should be of a focus so long, that the value of such an instrument would be extremely limited. It was early seen, therefore, that the only feasible method would be to use but a single objective for both bodies; but to bisect the pencils of rays emerging from this lens, so as to cause all those which have issued from the object in such a direction as to pass through either half of it, to be refracted into the body situated on that side; so that the two eyes, applied to the two eye-pieces respectively, shall receive through the two halves of the objective, two magnified images of the object differing from each other in perspective projection, as if the object, actually enlarged to the dimensions of its image, had been viewed by both eyes at once at a moderate distance.

That such a method would produce the Stereoscopic effect, might be anticipated from the result of the very simple experiment of covering the right-hand or the left-hand half of an object-glass of low power, during the examination of any object that lies in oblique perspective; for the two views of it thus obtained, will be found to present just the kind and degree of difference which is observable in stereoscopic pictures. The first attempt to put this plan into execution, seems to have been that of Prof. Riddell, of New Orleans; but the results of his method, as followed by opticians on the European side of the Atlantic, were far from answering the expectations excited by his own description of them. The subject was both theoretically and practically investigated by Mr. Wenham, with much ability (Transactions of the Microscopical Society, new series, Vol. II, p. 1); and a Binocular Microscope on a pattern suggested by him, was constructed by Messrs. Smith and Beck. This, too, was far from satisfactory in its performance, having two capital defects; namely, first, that the view which it gave was often *pseudoscopic*, the projecting portions of the object appearing to be depressed, and *vice versa*; and second, that the two bodies being *united* at a fixed angle of convergence, the distance between their axes could not be conveniently adapted to the varying distances of the eyes of different individuals. The construction adopted by M. Nachet, however, is much more successful. His method is to divide the pencil of

FIG. 30.



Arrangement of Prisms in Nachet's Binocular Microscope.

rays issuing from the objective, by means of a prism (Fig. 30, *p*) whose section is an equilateral triangle; for the rays *a b* on the right side, which enter the flattened surface presented to them, are reflected, by impinging very obliquely against one of the internal faces of the prism, towards the left, emerging again from the prism, as they had entered it, almost at right angles; and in like manner the rays *a' b'* on the left side are reflected towards the right. Each of these pencils is received by a similar prism, which again changes its direction, so as to render it parallel to its original course; and thus the two halves *a b* and *a' b'* of the original pencil are completely separated from each other to any interval that may be required, this interval being determined by the distance between the central and the lateral prisms. In

Fig. 31 is shown the Binocular Microscope constructed by M.

FIG. 31.



Nachet's Binocular Microscope.

Fig. 31 is shown the Binocular Microscope constructed by M. Nachet upon this plan. The arrangement of the base and stage is that commonly employed in French vertical Microscopes; and a stem rises from the back of it, with which the double body is connected by a rack-and-pinion movement that gives the focal adjustment. The apparatus of prisms shown in Fig. 30, is placed between the object-glass and the lower ends of the bodies; and by means of a double-threaded screw acted on by a milled head between the two bodies, they may be separated from, or approximated towards, each other; so that the distance between their axes may be brought to coincide with the distance between the axes of the eyes of the individual observer. The author can confirm by his own experience the statement of M. Nachet, that this instrument is entirely free from that tendency to produce pseudoscopic effects, which is the great drawback in Prof. Riddell's and in Mr. Wenham's arrangements; and it comes so near the theoretical standard of perfection, when used with low powers of moderate aperture, that its performance may be considered highly satisfactory. Its definition, however, when used with higher powers of larger angular aperture, has not yet been rendered sufficiently good, to enable it to afford a satisfactory view of the more difficult class of test-objects; and it may be doubted whether, considering the number of deflections which the rays undergo in their course, such perfect definition is to be anticipated. For although their *general* course on entering and emerging from each prism may be perpendicular to its surfaces, so that they suffer no refraction, many of them will be slightly oblique, and will therefore undergo not only refraction, but also some amount of chromatic dispersion. And it is moreover to be recollected, that when high powers are being employed, and especially such as are of large angular aperture, the smallest departure from exactitude in the focal adjustment gives indistinctness to the image. Now the special object of this instrument being to convey to the mind the notion of the *solid forms* of objects, of which some parts project more than others, it is obvious that the rays proceeding from the projecting parts cannot be so nearly brought to the same focus with those from the receding, as to produce an even tolerably distinct image of both at once. It seems likely to be only with objectives of comparatively low power and small angular aperture, that images suitable for the

production of Stereoscopic effects will be produced ; but for certain classes of objects, this mode of exhibition is most admirably adapted, the solid forms of the *Polycystina* (Chap. X), for example, being brought out by it (especially when they are viewed as opaque, not as transparent objects) with such a reality, as to make them resemble carved ivory balls which the hand feels ready to grasp.

41. The same method of dividing the pencil of rays issuing from the object glass, by a separating prism placed in its course, has been applied by M. Nachet to another purpose,—that of enabling two or more observers to look at the same object at once, which is often a matter not only of considerable convenience, but also of great importance, especially in the demonstration of dissections. The account given by M. Nachet of the construction of this instrument, as adapted for *two* persons, will be found in the "Quarterly Journal of Microscopical Science," Vol. II, p. 72 ; he has subsequently devised another arrangement, by which the form of the separating prism is adapted to divide the pencil into *three* or even *four* parts, each of which may be directed into a different body, so as to give to several observers at one time a nearly identical image of the same object. Of course, the larger the number of secondary pencils into which the primary pencil is thus divided, the smaller will be the share of *light* which each observer will receive ; but this reduction does not interfere with the distinctness of the image, and may be in some degree compensated by a greater intensity of illumination. (See Appendix for a description of American instruments and modifications.)

CHAPTER III.

ACCESSORY APPARATUS.

42. In describing the various pieces of accessory apparatus with which the Microscope may be furnished, it will be convenient in the first place to treat of those which form (when in use) part of the instrument itself, being Appendages either to its Body or to its Stage, or serving for the Illumination of the objects which are under examination; and secondly to notice such as have for their function to facilitate that examination, by enabling the microscopist to bring the Objects conveniently under his inspection.

SECTION 1. APPENDAGES TO THE MICROSCOPE.

43. *Draw-Tube.*—It is advantageous for many purposes, that the Eye-piece should be fitted, not at once into the “body” of the Microscope, but into an intermediate tube; the drawing out of which, by augmenting the distance between the object-glass and the image which it forms in the focus of the eye-glass, still further augments the size of the image in relation to that of the object (§ 20). For although the magnifying power cannot be thus increased with advantage to any considerable extent, yet, if the corrections of the object-glass have been perfectly adjusted, its performance is not seriously impaired by a moderate lengthening of the body; and this may be conveniently had recourse to on many occasions, in which some amplification is desired, intermediate between the powers furnished by any two objectives. Thus if one object-glass give a power of 80 diameters, and another a power of 120, by using the first and drawing the eye-piece, its power may be increased to 100. Again, it is often very useful to make the object fill up the whole, or nearly the whole, of the field of view: thus if an object that is being viewed by transmitted rays, is so far from transparent as to require a strong light to render its details visible, the distinctness of its details is very much interfered with, if, through its not occupying the peripheral part of the field, a glare of light enter the eye around its margin; and the importance of this adjustment is even greater, if opaque objects mounted on black disks are being viewed by

the Lieberkühn (§ 65), since if any light be transmitted to the eye direct from the mirror, in consequence of the disk failing to occupy the centre field, it greatly interferes with the vividness and distinctness of the image of the object. In the use of the Micrometric eye-pieces to be presently described (§§ 45, 46), very great advantage is to be derived from the assistance of the draw-tube; as enabling us to make a precise adjustment between the divisions of the stage micrometer, and those of the eye-piece micrometer; and as admitting the establishment of a more convenient numerical relation between the two, than could be otherwise secured without far more elaborate contrivances. Moreover, if, for the sake of saving room in packing, it be desired to reduce the length of the body, the draw-tube affords a ready means of doing so; since the body may be made to “shut up,” like a telescope, to little more than half its length, without any impairment of the optical performance of the instrument when mounted for use.

44. *Erector*.—It is only, however, in the use of the Erector, that the full value of the draw-tube, and the advantage of giving to it a rack-and-pinion movement of its own (§ 35), come to be fully appreciated. This instrument, first applied to the Compound Microscope by Mr. Lister, consists of a tube about three inches long, having a meniscus at one end and a plano-convex lens at the other (the convex sides being upwards in each case), with a diaphragm nearly half way between them; and this is screwed into the lower end of the draw-tube, as shown in Fig. 32. Its effect is (like the corresponding erector of the Telescope), to antagonize the reversion of the image formed by the object glass, by producing a second reversion, so as to make the image presented to the eye correspond in position with the object. The passage of the rays through two additional lenses, of course occasions a certain loss of light by reflection from their surfaces, besides subjecting them to aberrations whereby the distinctness of the image is somewhat impaired; but this need not be an obstacle to its use for the class of purposes for which it is especially adapted in other respects (§ 35), since these seldom require a very high degree of defining power. By the position given to the Erector, it is made subservient to another purpose, of great utility; namely, the procuring a very extensive *range* of magnifying power, without any change in the objective. For when the draw-tube, with the erector fitted to it, is completely pushed in, the *acting length* of the body (so to speak) is so greatly reduced by the formation of the first image much nearer the objective, that, if a lens of 8-10ths of an inch focus be employed, an object of the diameter of $1\frac{1}{2}$ inch can be taken in,

FIG. 32.



Draw-tube fitted with Erector.

and enlarged to no more than 4 diameters; whilst, on the other hand, when the tube is drawn out to its whole length, the object is enlarged 100 diameters. Of course every intermediate range can be obtained, by drawing out the tube more or less; and the facility with which this can be accomplished, renders such an instrument most useful in various kinds of research, especially those in which it is important, after *finding* an object with a lower power, to *examine* it under a higher amplification; since this may be done, without either a change of objectives, or a transfer of the object to another microscope fitted with a different power. It is when the draw-tube is thus made subservient to the use of the Erector, that the value of its rack-and-pinion adjustment is most felt; for by giving motion to the milled head which acts upon this (Fig. 22) with one hand, whilst the other hand is kept upon the milled head which moves the whole body (it being necessary to shorten the distance between the object and the objective, in proportion as the distance of the image from the objective is increased), the observer—after a little practice in the working together of the two adjustments—may almost instantaneously alter his power to any amount of amplification which he may find the object to require, without ever losing a tolerably distinct view of it. This can scarcely be accomplished without the rack movement; since, if both hands be required to make the alteration of the draw-tube, the readjustment of the focus must be effected subsequently.

45. *Micrometer*.—Although some have applied their micrometric apparatus to the stage of the microscope, yet it is to the Eye-piece that it may be most advantageously adapted.¹ The *cobweb micrometer*, invented by Ramsden for Telescopes, is probably, when well constructed, the most perfect instrument that the Microscopist can employ. It is made by stretching across the field of a “positive” eye-piece (§ 23) two very delicate parallel wires or cobwebs, one of which can be separated from the other by the action of a fine-threaded screw, the head of which is divided at its edge into a convenient number of parts, which successively pass by an index as the milled head is turned. A portion of the field of view on one side is cut off at right angles to the cobweb threads, by a scale formed of a thin plate of brass having notches at its edge, whose distance corresponds to that of the threads of the screw, every fifth notch being made deeper than the rest for the sake of ready enumeration. The object being brought into such a position that one of its edges seems to touch the stationary thread, the other thread is moved by the micrometer screw, until it appear to lie in contact with the other edge of the object; the number of entire divisions on the scale

¹ The Stage-micrometer constructed by Fraunhofer is employed by many continental Microscopists; but it is subject to this disadvantage,—that any error in its performance is augmented by the *whole* magnifying power employed; whilst a like error in the Eye-piece-micrometer is increased by the magnifying power of the eye-piece alone.


shows how many complete turns of the screw must have been made in thus separating the threads; while the number to which the index points on the milled head, shows what fraction of a turn may have been made in addition. It is usual, by employing a screw of 100 threads to the inch, to give to each division of the scale the value of 1-100th of an inch, and to divide the milled head into 100 parts; but the *absolute* value of the divisions is of little consequence, since their micrometric value depends upon the objective with which the instrument may be employed. This must be determined by means of a ruled slip of glass laid upon the stage; and as the distance of the divisions even in the best ruled slip is by no means uniform,¹ it is advisable to take an average of several measurements, both upon different slips, and upon different parts of the same slip. Here the draw-tube will be of essential use, in enabling the microscopist to bring the value of the divisions of his Micrometer to *even numbers*. Thus, suppose that with a 1-4th-inch object-glass, the tube being pushed in, a separation of the lines by one entire turn and 37-100ths of another were needed to take in the space between two lines on the ruled slip, whose actual distance is 1-1000th of an inch; then it is obvious that 137 divisions on the milled head are equivalent with that power to a dimension of 1-1000th of an inch, or the value of each division is 1-137,000th of an inch. But as this is an awkward number for calculation, the magnifying power may be readily increased by means of the draw-tube, until the space of 1-1000th of an inch shall be represented by a separation of the cobweb threads to the extent of 150 divisions; thus giving to each division the much more convenient value of 1-150,000th of an inch. The Microscopist who applies himself to researches requiring micrometric measurement, should determine the value of his Micrometer with each of the objectives he is likely to use for the purpose; and should keep a table of these determinations, recording in each case the extent to which the tube has been drawn out, as marked by the graduated scale of inches which it should possess. The accuracy with which measurements may be made with this instrument, is not really quite so minute as it appears to be; for it is found practically that when the milled head is so graduated, that, by moving it through a single division, the cobweb threads are separated or approximated by no more than 1-10,000th of an inch, it needs to be moved through *four* divisions, for any change in the position of the threads to be made sensible to the eye. Consequently, if three entire turns, or 300 divisions, were found to separate the threads so far as to coincide with a distance of 1-1000th of an inch on the ruled glass under a 1-8th of an inch

¹ Of the degree of this inequality, some idea may be formed from the statement of Hannover, that the value of the different divisions of a glass ruled by Chevalier to 1-100th of a millimetre, varied between the extreme ratios of 31-36, the mean of all being 34.


objective, although each division of the milled head, will thus represent 1-300,000th of an inch, yet the smallest measurable space will be four times that amount, or 1-75,000th of an inch. With the 1-12th inch objective, the smallest measurable space may be about 1-100,000th of an inch.

46. The expensiveness of the cobweb-micrometer being an important obstacle to its general use, a simpler method is more commonly adopted, which consists in the insertion of a transparent scale into the focus of the eye-piece, on which the image of the object is seen to be projected. By Mr. Ross, who first devised this method, the "positive" eye-piece was employed, and a glass plate ruled in squares was attached beneath its field-glass, at such a distance that it and the image of the object should be in focus together; and the value of these squares having been determined with each of the objectives, in the manner already described, the size of the object was estimated by the proportion of the square that might be occupied by its image. While the use of the positive eye-piece, however, renders the definition of the ruled lines peculiarly distinct, it impairs the definition of the object; and the "negative" or common Huyghenian eye-piece is now generally preferred. The arrangement devised by Mr. G. Jackson allows the divided glass to be introduced into the ordinary eye-piece (thus dispensing with the necessity for one specially adapted for micrometry), and greatly increases the facility and accuracy with which the eye-piece scale may be used. This scale is ruled like that of an ordinary measure (*i. e.* with every tenth line *long*, and every fifth line half its length), on a slip of glass, which is so fitted into a brass frame (Fig. 33, B), as to have a slight motion

FIG. 33.



Mr. Jackson's Eye-piece Micrometer.



towards either end; one of its extremities is pressed upon by a small fine milled-headed screw which works through the frame, and the other by a spring (concealed in the figure) which antagonizes the screw. The scale thus mounted is introduced through a pair of slits in the eye-piece tube, immediately above the diaphragm (Fig. 33, A), so as to occupy the centre of the field; and it is brought accurately into focus by unscrewing the glass nearest to

the eye, until the lines of the scale are clearly seen. The value

of the divisions of this scale must be determined, as in the former instance, by means of a ruled stage-micrometer, for each objective employed in micrometry (the drawing out of the eye-piece tube enabling the proportions to be adjusted to even and convenient numbers); and this having been accomplished, the scale is brought to bear upon the object to be measured, by moving the latter as nearly as possible into the centre of the field, and then rotating the eye-piece in such a manner, that the scale may lie across that diameter which it is desired to measure. The pushing-screw at the extremity of the scale being then turned, until one edge of the object is in exact contact with one of the long lines, the number of divisions which its diameter occupies is at once read off by directing the attention to the other edge,—the operation, as Mr. Quekett justly remarks, being nothing more than laying a rule across the body to be measured. This method of measurement may be made quite exact enough for all ordinary purposes, provided, in the first place, that the eye-piece scale be divided with a fair degree of accuracy; and secondly, that the value of its divisions be ascertained (as in the case of the cobweb-micrometer) by *several* comparisons with the scale laid upon the stage. Thus if, by a mean of numerous observations, we establish the value of each division of the eye-piece scale to be 1-12,500th of an inch, then, if the image of an object be found to measure $3\frac{1}{2}$ of those divisions, its real diameter will be $3\frac{1}{2} \times \frac{1}{12500}$ or 1-3571st of an inch.¹ Now as, with an objective of 1-12th inch focus, the value of the divisions of the eye-piece scale may be reduced to 1-25,000th of an inch, and as the eye can estimate a fourth part of one of the divisions with tolerable accuracy, it follows that a magnitude of as little as 1-100,000th of an inch can be measured with a near approach to exactness, and that this instrument cannot be fairly considered as ranking much below the cobweb-micrometer in minute accuracy. At any rate, it is sufficiently precise (when due care is employed) for all ordinary purposes; and it has the great advantage of cheapness and simplicity. Whatever method be adopted, if the measurement be made in the Eye-piece, and not on the stage, it will be necessary to make allowance for the adjustment of the object-glass to the thickness of the glass that covers the object, since its magnifying power is considerably affected by the separation of the front pair of lenses from those behind it (§ 83). It will be found convenient to compensate for this alteration, by altering the draw-tube in such a manner as to neutralize the effect produced by the adjustment of the objective; thus giving one uniform value to the divisions of the eye-piece scale, what-

¹ The calculation of the dimensions is most simplified by the adoption of a decimal scale; the value of each division being made, by the use of the draw-tube adjustment, to correspond to some aliquot part of a ten-thousandth or a hundred-thousandth of an inch, and the dimensions of the object being then found by simple multiplication:—Thus (to take the above example) the value of each division in the decimal scale is .00008, and the diameter of the object is .00028.

ever may be the thickness of the covering-glass : the amount of the alteration required for each degree must of course be determined by a series of measurements with the stage-micrometer, and should be recorded on the table of the micrometric values of the several objectives.


47. *Goniometer*.—When the Microscope is employed in researches on minute crystals, a means of measuring their angles is provided by the adaptation of a goniometer to the eye-piece. The simplest form (contrived by Schmidt and made by Ross) which answers sufficiently well for all ordinary purposes, essentially consists merely of a “positive” eye-piece, with a single cobweb-thread stretched across it diametrically in a circular frame capable of rotation ; the edges of this frame are graduated in degrees, and a vernier is also attached to the index, whereby fractional parts of degrees may be read off. By rotating the frame carrying the thread, so that it shall lie successively in the directions of the two sides of the crystal, the angle which they form is at once measured by the difference of the degrees to which the index points on the two occasions. For the cobweb-thread, a glass plate, ruled with parallel lines at about the 1-50th of an inch asunder, may be advantageously substituted ; since it is not then necessary to bring the crystal into such a position as to lie along the diametrical thread, but its angle may be measured by means of any one of the lines to which it happens to be nearest. If a higher degree of precision be required than this instrument is fitted to afford, the *Double-refracting Goniometer*, invented by Dr. Leeson, may be substituted ; for a description of which (too long to be introduced here) the reader is referred to Dr. L.’s account in the “Proceedings of the Chemical Society,” Part xxxiii, and to Mr. Quekett’s “Practical Treatise on the Microscope.” (See Appendix for a description of a Micrometer and Goniometer, by Prof. J. L. Smith.)

48. *Indicator*.—When the Microscope is used for the purpose of demonstrating to others such objects as may not be at once distinguished by the uninitiated eye, it is very useful to introduce into the eye-piece, just over the diaphragm, a small steel hand pointing to nearly the centre of the field ; to whose extremity the particular portion of the image which the observer is intended to look at, is to be brought by moving the object. The hand may be so attached, as to be readily turned back when not required ; leaving the field of the eye-piece quite free. This little contrivance, which was devised by Mr. J. Quekett, is appropriately termed by him the *indicator*.

49. *Camera Lucida*.—Various contrivances may be adapted to the eye-piece, in order to enable the observer to see the image projected upon a surface whereon he may trace its outlines. The one most generally employed is the *Camera Lucida prism* contrived by Dr. Wollaston for the general purposes of delineation ; this being fitted on the front of the eye-piece, in place of the

“cap” by which it is usually surmounted. The Microscope being placed in a horizontal position, as shown in Fig. 34, the rays which pass through the eye-piece into the prism, sustain such a total reflection

FIG. 34.



Microscope arranged with Camera Lucida for Drawing or Micrometry.

from its oblique surface, that they come to its upper horizontal surface at right angles to their previous direction; and the eye being so placed over the edge of this surface, that it receives these rays from the prism through part of the pupil, whilst it looks beyond the prism, down to a white-paper surface on the table, with the other half, it sees the image so strongly and clearly projected upon that surface, that the only difficulty in tracing it arises from a certain incapacity which seems to exist in some individuals, for seeing the image and the tracing-point at the same time. This difficulty (which is common to all instruments devised for this purpose) is lessened by the interposition of a slightly convex lens in the position shown in the figure, between the eye and the paper, in order that the rays from the paper and tracing-point may diverge at the same angle as those which are received from the prism; and it may be generally got over altogether, by experimentally modifying the relative degrees of light received from the object and from the paper. If the image be too bright, the paper, and the tracing-point, and the outline it has made, are scarcely seen; and either less light may be allowed to come from the object, or more light (as by a taper held near) may be thrown on the paper and tracing-point. Sometimes, on the other hand, measures of the contrary kind must be taken. Instead of the prism, some microscopists prefer a *speculum* of polished steel, of smaller size than the ordinary pupil of the eye, fixed at an angle of 45° in front of the eye-piece; and this answers exactly the same purpose as the preceding, since the rays from the eye-piece are reflected vertically upwards to the central part of the pupil placed above the mirror, whilst, as the eye also receives rays from the paper and tracer, in the same direction, through the peripheral portion of the pupil, the image formed by the microscope is visually projected downwards, as in the preceding case. This disk, the in-

vention of the celebrated anatomist Soemmering, is preferred by some microscopic delineators to the camera lucida. The fact is, however (as the author can testify from his own experience), that there is a sort of "knack" in the use of each instrument, which is commonly acquired by practice alone; and that a person habituated to the use of either of them, does not at first work well with another. A different plan is preferred by some microscopists, which consists in the substitution of a *plate* of neutral-tint or darkened glass for the oblique mirror; the eye receiving at the same time the rays of the microscopic image, which are obliquely reflected to it from the surface of the glass, and those of the paper, tracing-point, &c., which come to it through the glass. It is so extremely useful to the microscopist, to be able to take outlines with one or other of these instruments, that every one would do well to practise the art. Although some persons at once acquire the power of seeing the image and the tracing-point with equal distinctness, the case is more frequently otherwise; and hence no one should allow himself to be baffled by the failure of his first attempt. It will sometimes happen, especially when the prism is employed, that the want of power to see the pencil is due to the faulty position of the eye, too large a part of it being over the prism itself. When once a good position has been obtained, the eye should be held there as steadily as possible, until the tracing shall have been completed. It is essential to keep in view, that the proportion between the size of the tracing and that of the object, is affected by the height of the eye above the paper; and hence that if the microscope be placed upon a support of different thickness, or the eye-piece be elevated or depressed by a slight inclination given to the body, the scale will be altered. This it is of course peculiarly important to bear in mind, when a series of tracings is being made of any set of objects which it is intended to delineate on a uniform scale; or when the camera lucida (or any similar arrangement) is employed for the purpose of *Micrometry*. All that is requisite to turn it to this account, is an accurately divided stage-micrometer, which, being placed in the position of the object, enables the observer to see its lines projected upon the surface upon which he has drawn his outline; for if the divisions be marked upon the paper, the average of several be taken, and the paper be then divided by parallel lines at the distance thus ascertained (the spaces being subdivided by intermediate lines, if desirable), a very accurate scale is furnished, by which the dimensions of any objects drawn in outline under the same power may be minutely determined. Thus if the divisions of a stage-micrometer, the real value of each of which is 1-200th of an inch, should be projected with such a magnifying power, as to be at the distance of an inch from one another on the paper, it is obvious that an ordinary inch-scale applied to the measurement of an outline, would give its dimensions in two-hundredths of an inch, whilst each fifth of that scale

would be the equivalent of a thousandth of an inch. When a sufficient magnifying power is used, and the scale thus made is minutely divided, great accuracy may be obtained. It has been by the use of this method, that Mr. Gulliver has made his admirable series of measurements of the diameters of the Blood-corpuscles of different animals.

50. *Object-Glass Holder*.—In Microscopes of the old construction, whose objectives were single lenses, these were not unfrequently mounted near the periphery of a circular disk pivoted to the lower end of the body, in such a manner that any desired power might at once be brought into use by merely rotating the disk. Since the introduction of achromatic object-glasses, this method has been until recently abandoned; every "power" being separately connected with the extremity of the body, so as not to admit of any substitution, save by screwing off one objective and screwing on another. The old method, however, has been partially reverted to by Mr. C. Brooke; who has contrived a *holder* into which two objectives may be screwed, and which, being attached to the "nose" of the body, enables either of them to be brought into position, by simply turning the arm on its pivot. This is an extremely convenient arrangement, and might easily be carried further if desired; since, by having a tri-radiate or quadri-radiate arm, three or four powers might be thus brought into use successively, with as much facility as two. The principal objection to the general use of such an appendage, lies in the nicety of workmanship that is required to obtain that exact "centering," which is needed to bring the axis of the objective into precise continuity with that of the body; and unless this be attained, the performance of the instrument is greatly impaired. In microscopes of the old construction, the *other* imperfections were so great, that none but an excessive deficiency in this respect would attract attention. The convenience of such an instantaneous change in the power of the objective is very great; since it is continually desirable to obtain a general view of an object with a low power, and to examine the parts of it in detail under a higher amplification, with as little expenditure of time and trouble as possible.

51. *Object-Marker*.—All Microscopists occasionally, and some continually, feel the need of a ready means of *finding*, upon a glass slide, the particular object or portion of an object, which they desire to bring into view; and various contrivances have been suggested for the purpose. Where different magnifying powers can be readily substituted one for another, as by the use of the Erector (§ 44) or of the Object-glass holder (§ 50), no special means are required; since, when the object has been found by a low power, and brought into the centre of the field, it is rightly placed for examination by any other objective. Even this slight trouble, however, may be saved by the adoption of more special methods; among the simplest of which is *marking* the position of

the object on the surface of the thin glass which covers it. The readiest mode of doing this, when the object is large enough to be distinguished by the naked eye, is to make a small ring round it with a fine camel-hair pencil dipped in Indian ink; but when the object is not thus visible, the slide must be laid in position on the stage, the object "found" in the microscope, the condenser adjusted to give a bright and defined circle of light, and then, the microscope-body being withdrawn, the black ring is to be marked around the illuminated spot. The same end, however, may be more precisely as well as more neatly accomplished, by attaching an *object-marker* to the objective itself. That of Mr. Tomes consists simply of an ivory cap, fitting over the 1.4th-inch objective, having its extremity narrowed down (like that of the objective itself), but perforated in the centre, so as to form a minute ring; the object having been "found" and brought into the centre of the field, the cap is placed upon the objective, the ring is blackened with Indian ink, and then, being carefully brought by the focal adjustment into contact with the surface of the glass, it stamps on this a minute circle enclosing the object. A more elaborate contrivance of a similar kind, for marking a circle round the object by a diamond point, attached to a cap fitting on the objective, has been recently described by Mr. Bridgman ("Quarterly Microscopical Journal," vol. iii, p. 237); this has the advantage of admitting a variation in the size of the circle, and also of substituting a delicate line for the broad ring which may partly obscure some neighboring object; but, on the other hand, the very delicacy of the diamond marks prevents them from being readily distinguished, and some kinds of glass are so apt to "star" when marked with a diamond point, that cracks or splinters may extend from the circle over the object it is intended to indicate. The most unobjectionable and satisfactory mode of "finding" an object, however, is, in the Author's opinion, that which is afforded by a graduation of the movable parts of the stage, in the manner to be presently described (§ 53).

52. *Lever Stage*.—The general arrangement of the Traversing Stage, now usually adapted to all high class Microscopes, has been already explained (§§ 37–39); and though the details are differently constructed by the several makers, yet the general principle is, that a lateral or *horizontal* movement is given to the object-platform by one milled head, and a front to back or *vertical* movement (the microscope being supposed to be placed in an inclined position) by another. The stage may be so constructed, however, that motion shall be given to the object-platform by means of a *lever* acting upon it in any required direction; this being accomplished by making the object-platform slide laterally on an intermediate plate, and by making the latter slide vertically upon the fixed stage-plate which forms the basis of the whole; each pair of plates being connected by dovetailed slides and grooves. Thus the object-platform may be readily made to

traverse, not merely horizontally or vertically, but, by the simultaneous sliding of both plates, in any intermediate direction. This is especially convenient in following the movements of Animalcules, &c., for which purpose this lever-stage is to be preferred to the ordinary form: its use being attended with this particular facility, that, as the motion of the hand is reversed by the lever, so that the object moves in the opposite direction, and as the motion of the object is again reversed to the eye by the microscope, the image moves in the same direction as the hand; and thus, with a little practice, even the most rapid swimmer may be kept within the field by the dexterous management of the lever. For general purposes, however, the ordinary traversing stage will be found most convenient.

53. *Object-Finder*.—Either kind of movable stage admits of a simple addition, which very much facilitates the “finding” of minute objects mounted in slides, that are not distinguishable by the naked eye; such, for example, as the particular forms that present themselves in Diatomaceous deposits. This “finder” consists of two graduated scales, one of them *vertical*, attached to the fixed stage-plate, and the other *horizontal*, attached to an arm carried by the intermediate plate; the first of these scales enables the observer to “set” the vertically sliding plate to any determinate position in relation to the fixed plate, while the second gives him the like power of setting the horizontally sliding plate by the intermediate. In order to make use of these scales, it is of course necessary that the sliding and rotating platform on which the object immediately rests, should be always brought into one constant position upon the traversing plates beneath; this is accomplished by means of a pair of *stops*, against which it should be brought to bear. So, again, this sliding plate or object-platform should itself be furnished with a “stop” for the glass slide to abut against, so as to secure this being always laid in the same position. These stops may be made removable, so as not to interfere with the ordinary working of the stage. Now supposing an observer to be examining a newly-mounted slide, containing any objects which he is likely to wish to find on some future occasion; he first lays the slide on the object-platform, with its lower edge resting on the ledge, and its end abutting against the lateral stop, and brings the object-platform itself into its fixed place against the stops; then if, on sweeping through the slide, he meet with any particular form worthy of note, he reads off its position upon the two scales, and records it in any convenient mode. The scale may be divided to 50ths of an inch, and each of these spaces may be again halved by the eye; the record may perhaps be best made thus,—*Triceratium favus* $\frac{2}{184}$;—the upper number always referring to the upper scale, which is the horizontal, and the lower to the vertical. Now whenever the Microscopist may wish again to bring this object under examination, he has merely to lay the slide in the same position on

the platform, to bring the platform itself into its fixed place on the traversing plate below, and then to adjust the traversing plates themselves by their respective scales. Even a non-movable stage may have a similar pair of scales adapted to it; the vertical scale being so placed, as to mark the position into which the object-platform is brought by sliding it up or down; and the horizontal scale being marked upon the object-platform itself, so as to allow the observer to note the precise position of the end of the glass slide. Thus let it be supposed that, by shifting the slide from side to side, and by moving the object-platform up or down, a certain object has been brought into the field; if the place of the object-platform and of the slide be then noted by the vertical and the horizontal scales, the object may be found at any future time without difficulty, by readjusting the slide and the object-platform to the same numbers.¹ The numbers referring to each object may either be marked upon the slides themselves, like the names of the objects, or may be recorded with these in a separate list, referring to the slides by figures alone. The general adoption of such a plan, though involving a little more labor at first, would prove in the end to be a great saving both of time and trouble.

54. *Magnetic Stage*.—If a stage be unprovided with a traversing movement of any kind, there is no means of allowing the object to be moved in all directions with smoothness and facility, and yet of holding it in any position in which the Microscopist desires to retain it, more convenient and more ready of application, than is furnished by magnetic attraction. A magnetic stage was originally proposed by Mr. King of Bristol; but seems to have been first brought into efficient practical action by Mr. G. Busk. His plan consists in attaching two semicircular magnets to the under side of the stage, so as nearly to surround its aperture, and in inserting, for the conveyance of the magnetic force to the upper side, four soft iron pegs, which slightly project above its surface; over these an object-bearer of soft iron, with its under surface ground smooth and true, will slide so readily, as to admit of very easy and precise adjustment of the

¹ The first of the above plans, to the utility and accuracy of which the Author can bear strong testimony, was suggested by Mr. Okeden in the "Quart. Microsc. Journal," vol. iii, p. 166. The second had been previously suggested by Mr. E. G. Wright in the same Journal, vol. i, p. 302; the descriptions of both are made clear by figures. Other "finders" are described and figured by Mr. J. Tyrrel, Mr. T. E. Amyot, and Mr. Bridgman, at pp. 234 and 302-304 of the last-named volume. It appears to the Author that Mr. Okeden's plan might be adopted with very little trouble or expense in every Microscope possessed of a stage movement, and Mr. Wright's in every Microscope with a fixed stage but movable object-carrier; and that it would be very desirable for every microscope that may be made hereafter, to be furnished with such scales. If the different makers could agree upon some common system of graduation, in the same way as Microscopists have adopted 3 inches by 1 as the standard dimension of object-slides, much trouble would be saved to observers at a distance from one another, who might wish to examine each others' objects; for the numerical reference attached to each object would then enable it to be found by every observer, whose stage should be graduated upon the same method.

place of the object by a steady and practised hand. Another arrangement, which has some advantages over the preceding, has been proposed by Mr. J. B. Spencer; this consists in surrounding the aperture of the stage by a ring of soft iron, the surface of which projects very slightly above the brass plate, the magnets (cut out of sheet-steel) being attached to the under side of the object-bearer. This method is perhaps more readily applicable than Mr. Busk's to the stage of *any* microscope, and will probably interfere less with its other fittings.¹

55. *Diaphragm-Plate*.—No microscope stage should ever be without a diaphragm-plate fitted to its under surface, for the sake of restricting the amount of light reflected from the mirror, and of limiting the angle at which its rays impinge on the object (see Figs. 18 and 21). This plate should always be at least half an inch below the object, since it is otherwise comparatively inoperative; and thus, whilst it may be fixed immediately beneath a movable stage whose thickness serves to remove it sufficiently far, it should be fixed on the end of a short tube forming a sort of *well* on the under side of the stage, when this consists of but a single fixed plate. The diaphragm-plate should be perforated with holes of several different sizes, in the largest of which it is convenient to fit a ground-glass (this, by means of a screw-socket, may be made removable at pleasure), the use of which is to diffuse a soft and equable light over the field, when large transparent objects (such as sections of wood) are under examination; between the smallest and the largest aperture, there should be an unperforated space, to serve as a dark background for opaque objects. The diaphragm-plate itself, the "well" of the stage, in fact every part through which light passes to the object from beneath, should be blackened, in order to avoid the interference that would be occasioned by irregularly reflected rays. The edge of the diaphragm-plate should be notched at certain intervals, and a spring catch fitted so as to drop into the notches, in order that each aperture may be brought into its proper central position. This simple arrangement, in combination with the mirror (which should be concave on one side and plane on the other) and side-condenser (§ 64), affords to the Microscopist all the means of illuminating his objects, whether transparent or opaque, which are ordinarily requisite: to bring out the highest powers of the instrument, however, more refined methods of illumination are required; and a far greater variety of treatment is needed in the case of many objects, the determination of whose true characters is a matter of difficulty, even under every advantage which can be derived from assistance of this kind.

56. *Achromatic Condenser*.—In almost every case in which an objective of 1-4th inch or any shorter focus is employed, its per-

¹ For a more detailed description, with illustrative figures, of Mr. Busk's Magnetic Stage, see "Quarterly Microsc. Journal," vol. ii, p. 280; and for Mr. Spencer's, vol. iii, p. 173.

formance is greatly improved by the interposition of an achromatic combination between the mirror and the object, in such a manner that the rays reflected from the former shall be brought to a focus in the spot to which the objective is directed. This may be accomplished sufficiently well for ordinary purposes, by adapting a French triple combination of about 1-4th inch focus, to the end of a tube $1\frac{1}{2}$ inch long, which shall slide within another tube fitted to the opening in the stage, by the bayonet catch or any similar connection that gives attachment to the diaphragm-plate. If this be correctly centred in the first instance, and the workmanship of the microscope be good, no more expensive arrangement will be required, by such at least as may be satisfied with that degree of perfection, which suffices for the clear discernment of all but the most difficult objects. The sliding movement of the tube, especially if it be accomplished by a lever-action (as suggested by Mr. Quekett), is quite sufficient for the adjustment of the focus; and the removal of the outer lens adapts it for use with objectives below 1-4th inch, to whose performance it often affords important assistance. In the most perfect arrangement of the Achromatic Condenser, however, such as is now adapted to all first-class instruments made in this country, the achromatic combination is one specially adapted to the purpose; and is so mounted as to insure the greatest accuracy of its adjustments. By Mr. Ross it is supported by what he terms the "secondary stage" (§ 37); and by Messrs. Smith and Beck it is carried upon the summit of the "cylindrical fitting" which answers the same purpose (§ 39); whilst by Messrs. Powell and Lealand, it is attached by a bayonet-catch to the under side of the fixed stage-plate (§ 38). In either case it is provided with a pair of milled-headed screws (Fig. 36), which give it a slight degree of horizontal motion in transverse directions, for the purpose of procuring an accurate centring; and where, as in Messrs. Powell and Lealand's instrument, the focal adjustment is not given by the movement of the carriage which bears it, a rack and pinion is attached for this purpose to the tube of the condenser itself. In order that the Achromatic Condenser should be made to afford the greatest possible variety of modifications of the illuminating pencil, it requires to be furnished with a diaphragm-plate (as first suggested by Mr. Gillett) immediately behind its lenses; and this should be pierced with holes of such a form and size, as to be adapted to cut off in various degrees, not merely the peripheral, but also the central parts of the illuminating pencil. The former of these purposes is of course accomplished, by merely narrowing the aperture which limits the passage of the rays through the central part of the lens; the latter, on the other hand, requires an aperture as large as that of the lens, having its central part more or less completely occupied by a solid disk, which may so nearly fill the circle, as to leave but a mere ring through which the light may pass. Such

apertures are shown in the diaphragm-plates in Figs. 35 and 36. The Condenser thus completed is constructed on three different plans by the three principal makers, in accordance with the different arrangements of their respective stages. By Mr. Ross, who originally carried Mr. Gillett's plans into operation, the diaphragm-plate has the shape of a short frustrum of a cone (Fig. 35), so attached to the condenser, that the portion of the plate which passes through it shall cut it transversely; each aperture is indicated by a number on the dial; and a spring-catch is so arranged, as to mark when any one of the apertures is in its right place, and to show its number. The thinness of the stage in Messrs. Smith and Beck's microscope, allows the diaphragm-plate to be made upon the ordinary plan (Fig. 36), since it can be brought sufficiently near to the lenses of the condenser, without coming into too close contiguity with the stage; and this is obviously the simplest and most convenient arrangement. By Messrs. Powell and Lealand, again,—their stage being too thick to allow of the diaphragm-plate being placed beneath it, without removing that plate from its proper position behind the lenses of the condenser,—the diaphragm-plate is made so small that it can be received into the interior of the stage (Fig. 37), but is rotated by a milled head beneath; and the edge of this is marked by numbers, each signifying a particular aperture, and thus marking by its position *which* aperture is in use. As, however, the smallness of the diaphragm-plate

so limits the number of apertures, that the desirable variety could not be afforded by it alone, a second plate is made to rotate immediately beneath it upon the same axis (like the hour and minute hands of a watch), by means of a second milled head, numbered at its edge like the first; and the apertures in the diaphragm-plate being simple circles, the centres of these are covered by stops of different sizes, supplied by the second or "stop"-plate; by which very ingenious arrangement, a great variety of combinations may be obtained, all of them indicated by the numbering on the two milled heads.


57. *Reflecting Prisms*.—Every mirror composed of glass silvered at the back, gives, as is well known, a double reflection; namely, a principal image from the metallic surface, and a secondary im-

FIG. 35.



Ross's Achromatic Condenser.

FIG. 36.



Smith and Beck's Achromatic Condenser.

age from the surface of the glass in front of it. This secondary image, it has been thought, interferes with the perfect performance of the achromatic condenser; and hence, for obtaining the

FIG. 37.



Powell and Lealand's Achromatic Condenser.


most satisfactory definition, some Microscopists prefer to direct the axis of the microscope to the source of light (the mirror being turned aside); whilst others, feeling the inconvenience of the position thus required, have recourse to a *prism* which shall give the required reflection with only a single image. The prism usually employed (having been originally applied to this purpose by M. Dujardin) has plane surfaces, and acts, therefore, as the equivalent of a plane mirror. A reflecting prism has been devised, however, by Mr. Abraham (optician of Liverpool), which is intended by him to take the place both of mirror and achromatic condenser, though its action (as it seems to the author) must rather be that of the ordinary concave mirror; this has one of its surfaces hollowed out to receive one side of a double-convex lens, the other side of which acts as the emergent surface of the prism, causing the rays as they pass through it to converge; and the prism itself being composed of flint-glass, whilst the lens is of crown, no chromatic dispersion of the rays is produced, though the spherical aberration is not corrected.

58. *White-Cloud Illuminators*.—It being universally admitted that the light of a bright white cloud is the best of all kinds of illumination for nearly every kind of Microscopic inquiry, various attempts have been made to obtain such light, from the direct rays either of the sun or of a lamp, by what may be called an artificial cloud. Some have replaced the plane mirror by a surface of pounded glass or of carbonate of soda, or (more commonly) by a disk of plaster of Paris, the latter being decidedly the preferable method; but a sufficiently bright light is not thus obtained, unless a condenser be employed to intensify the illumination of the mirror. Such a condenser may be most conveni-

ently attached by a jointed arm to the frame which carries the disk, according to the method of Messrs. Powell and Lealand, shown in Fig. 38; the frame itself being made to fit upon the mirror, and to turn with it in every direction. Another very simple, and for many purposes very efficient mode of obtaining a white-cloud illumination (invented by Mr. Handford) consists in coating the back of a concave plate of glass, like that employed in the ordinary concave mirror, with white zinc paint, instead of silvering it; and then mounting this in a frame, which may be fitted (like the plaster of Paris disk just described) over the ordinary mirror. A concave surface of plaster of Paris, moreover, might easily be obtained by casting it when fluid upon the convex surface of such a plate. When a concavity is thus given to the white surface, its performance with low powers is much improved; but with high powers, a special condensation of the light must be adopted, and the arrangement above described seems the simplest that could be devised. It is open, however, to certain objections, which become apparent when very high powers are used and difficult objects are under examination; and to obtain the most perfect white-cloud illumination possible, is the object of the apparatus devised by Mr. Gillett. This consists of a small camphine lamp, placed nearly in the focus of a parabolic speculum, which reflects the rays either at once upon a disk of roughened enamel or upon a second (hyperbolic) speculum which reflects them upon such a disk. A very pure and concentrated light is thus obtained; and as the forms of the incident pencils are broken up by the roughened surface, that surface takes the place of the lamp, as the source from which the rays primarily issue. The advantage of this illumination is specially felt, in the examination of objects of the most difficult class under the highest powers.

59. *Oblique Illuminators.*—It is frequently desirable to obtain a means of illuminating transparent objects with rays of more obliquity than can be reflected to them from the mirror, even when this is thrown as much as its mounting will permit out of the axis of the Microscope (§ 39), or than can be transmitted by the ordinary achromatic condenser, even when all but its marginal aperture is stopped out. Such oblique light may be used in two entirely different modes. The rays, although very far out of the axis of the microscope, may still not make too great an angle with it to fall beyond the aperture of the objective; and thus, entering its peripheral portion after their passage through the object, they will form the image in the ordinary way. The advantage of such oblique illumination, arises from its power of bringing out markings which cannot be seen when only direct rays are employed; and when the rays come only from one side,

FIG. 38.



White-Cloud Illuminator.


so as to throw a strong shadow, and either the stage or the illuminator is made to rotate, so that the light shall fall upon the object *successively* in every azimuth, information may often be gained respecting the nature of these markings, which can be acquired in no other mode. But the direction given to the rays may be so oblique, that they shall not enter the object-glass at all; in this case, they serve to illuminate the object itself, which shines by the light whose passage it has interrupted; and as the observer then receives no other light than that which radiates from it, the object (provided it be of a nature to stop enough light) is seen *bright* upon a *dark field*. Each of these methods has its advantages for particular classes of objects; and it is advisable, in all doubtful cases, to have recourse to every variety of oblique illumination that shall present the object under a different aspect. Almost every Microscopist who has especially devoted his attention to the more difficult *lined* or *dotted* objects, has devised his own particular arrangement for oblique illumination, and feels confident of its superiority to others. To give a full description of all, would be quite unsuitable to our present object; those, therefore, will be specially noticed, which have already acquired *general* approval; whilst such as have only been recommended by *individuals*, will be simply referred to. As they have little in common, save their purpose, it seems scarcely possible to classify them according to any other character, than that afforded by the *direction* which they give to the oblique rays; some of them bringing these to bear on the object from *one side* alone, and others from *all sides*.

60. One of the earliest methods devised for obtaining oblique light, was the *eccentric prism* of M. Nacet; which, occupying the place of the achromatic condenser, and like it receiving its light from the mirror, has its surfaces so arranged, as to throw a converging pencil of rays on the under side of the object, whose axis is at an angle of about 40° with the axis of the microscope. One great convenience of this instrument lies in the power of giving revolution to the prism, by simply turning it in its socket, so as to direct the oblique rays upon the object from every side successively, without moving the stage. Its principal disadvantages consist in the limitation of its aperture (producing a deficiency of light), in the want of correction for its chromatic aberration, and in the absence of any power of varying the obliquity of the illuminating pencil.¹ All these disadvantages seem to be remedied by the plan of oblique illumination recently proposed by Mr. Sollitt, of Hull, which consists in the employment of an Achromatic condenser of very long focus and large aperture, mounted in such a manner as to enable its axis to be inclined to that of the microscope through a wide angular range; a con-

¹ A full description of M. Nacet's prism, and a mathematical investigation of its properties, by Mr. G. Shadbolt, will be found in the "Transactions of the Microscopical Society" (1st series), vol. iii, p. 74, *et seq.*

denser of this description he states to be suitable also for all ordinary purposes. ("Quart. Microsc. Journ.," vol. iii, p. 87.) Such an instrument, when its axis does not form a very large angle with that of the microscope, may receive its light from the plane mirror, especially if this be so mounted as to be capable of being turned considerably out of the visual axis; but when its position is too oblique for the light to be thus supplied to it, recourse must be had to rays either proceeding direct from their source (such as a lamp or a bright cloud), or directed at the requisite angle by a reflector placed in a suitable position. For this latter purpose, a rectangular prism (§ 57), mounted on a separate stand, will be found very convenient. By many observers, a combination of the *reflecting* and *refracting* powers of a prism is preferred, which causes the rays to be at once reflected by a plane surface, and concentrated by lenticular surfaces; so that the prism answers the purpose of mirror and condenser at the same time. Such a prism was first constructed by Amici; and it may be either mounted on a separate base, or attached to some part of the microscope-stand. The mounting adopted by Messrs. Smith and Beck, and shown in Fig. 39, is a very simple and convenient

FIG. 39.



Amici's Prism for oblique illumination.

one; this consists in attaching the frame of the prism to a sliding bar, which works in dovetail grooves on the top of a cap that may be set on the cylindrical fitting beneath the stage; the slide serves for the regulation of the distance of the prism from the axis of the microscope, and consequently of the obliquity of the illumination; whilst its distance beneath the stage is adjusted by the rack-movement of the cylindrical fitting. In this manner, an illuminating pencil of almost any degree of obliquity may be readily obtained; but there is no provision for the correction of its aberrations. Such a provision is afforded by the achromatic prism of Mr. Abraham (§ 57), which may be mounted in the manner just described. And the same object is attained by an arrangement devised by Mr. Grubb, a Dublin optician, of which Dr. Robinson, of Armagh, speaks very highly; the prism having its aberrations corrected for a lamp placed at a given distance in the plane of the stage; and being mounted in such a manner as to be capable of travelling (like Mr. Sollitt's condenser) through an angular range of as much as 120° ("Quart. Microsc. Journ.," vol. iii, p. 166). In all of these methods, the obliquity of the illumination is practically limited by the construction of the stage, and especially by the relation which its thickness bears to the diameter of its lower aperture. The thinner the stage, and the larger its lower aperture, the more oblique will be the rays which *may* be transmitted through it; and in admitting an extreme obliquity of illumination, the

thin stage recently introduced into some of the best Microscopes (§§ 38, 39) possesses a great advantage over all whose thickness is greater. On the other hand, it is when the rays are most oblique, that the greatest advantage is gained by making them fall upon the object from every side in succession; and where this cannot be accomplished (as in the case of Nacet's prism) by the rotation of the illuminating apparatus, the rotatory movement must be given to the object. It is obvious that, for this purpose, a revolving stage which keeps the object constantly in the field (§ 37), is decidedly preferable to one which does not possess such a movement; but the means have not yet been found of obtaining this advantage without some sacrifice of the other.

61. Whenever the rays are directed with such obliquity, as not to be received into the object-glass at all, but are sufficiently retained by the object, to render it (so to speak) self-luminous, we have what is known as the *black ground illumination*; to which the attention of Microscopists generally was first drawn by the Rev. J. B. Reade, in the year 1838, although it had been practised some time before, not only by the Author but by several other observers. For low powers whose angular aperture is small, and for such objects as do not require any more special provision, a sufficiently good "black ground" illumination may be obtained by means of the concave mirror alone, especially when it is so mounted as to be capable of a more than ordinary degree of obliquity. In this manner it is often possible, not merely to bring into view features of structure that might not otherwise be distinguishable, but to see bodies of extreme transparency (such, for instance, as very minute Animalcules) that are not visible when the field is flooded (so to speak) by direct light; these presenting the beautiful spectacle of phosphorescent points rapidly sailing through a dark ocean. Where the mirror cannot be placed in a position oblique enough to give this effect, a black ground illumination sufficiently good for many purposes may be obtained by Mr. Reade's original method; which consisted in dispensing with the mirror altogether, and in placing the lamp and ordinary condensing-lens (§ 64) in such a position beneath and to one side of the stage, as to throw upon the under side of the object a pencil of rays too oblique to enter the object-glass after passing through it. Another very simple mode, which answers sufficiently well for low powers and for the larger objects which these are fitted to view, consists in the substitution, for the achromatic condenser, of a plano-convex lens of great convexity, forming a large segment of its sphere, with a central stop to cut off the direct rays; for the rays passing through the marginal portion of this *Spotted Lens*, being strongly refracted by its high curvature, are made to converge at an angle too wide for their entrance into an objective of moderate aperture, and thus the field is left dark; whilst all the light stopped by the object serves (as it were) to give it a luminosity of its own. Neither of the

foregoing plans, however, will answer well for objectives of high power, having such large angles of aperture that the light must fall *very* obliquely to pass beyond them altogether. Thus if the pencil formed by the "spotted lens" have an angle of 60° , its rays will enter a 1.4th-inch objective of 70° , and the field will not be darkened. For obtaining a greater degree of obliquity, Mr. Wenham has contrived a *Parabolic Speculum*,¹ having its apex cut off, so that the object might be placed in the focus, to which all rays parallel to its axis are reflected; and the direct rays being checked by a stop placed behind it, the object is illuminated only by those which are reflected to it from all sides of the interior of the parabola at a very oblique angle. As the thickness of the glass slide on which the object is mounted, was found by Mr. W. to produce a very sensible aberration in the rays converging towards it, he interposed a meniscus lens, having such a curvature as to produce a counteracting aberration of an opposite kind. The circular opening at the bottom of the wide tube (Fig. 40) that carries the speculum, may be fitted with a diaphragm, adapted to cover any portion of it that may be desired; and by giving rotation to this diaphragm, rays of great obliquity may be made to fall upon the object from every azimuth in succession (§ 60). A like purpose was aimed at in the *Annular Condenser* of Mr. Shadbolt,² which consists of a ring of glass, whose surface was so shaped as to present a prismatic section; the inclination of the outer side being such as to produce a total reflection of the rays impinging on it, and to direct these through the inner side of the ring, so as to fall at a very oblique angle upon the object, from every azimuth of the circle. A combination of both methods is adopted in the *Parabolic Illuminator* (Fig. 40), now supplied by Messrs. Smith and Beck; for this consists of a paraboloid of glass, resembling a cast of the interior of Mr. Wenham's parabolic speculum, but reflecting the rays which fall upon the outer surface of the glass, like Mr. Shadbolt's annular prism. It has the advantage of being more easily constructed than the parabolic speculum, and is little, if at all, inferior to it in performance; but it requires that an appropriate "stop" should be adapted to it, for each objective with which it is to be used: whilst in Mr. Wenham's speculum, the requisite adaptation for the angular aperture of the objective is made by altering the position of the stop by means of the central stem; the effect of which alteration is to cut off a larger and larger proportion of the least oblique rays, the more nearly the stop is approximated to the ob-

FIG. 40.



Parabolic Illuminator.

¹ "Transactions of the Microscopical Society" (1st Series), vol. iii, p. 85.

² Op. cit. p. 132.


ject; and thus to illuminate it more and more exclusively by those which meet at the widest angle. In using either of these illuminators, the rays which are made to fall upon them should be parallel, consequently the *plane* mirror should always be employed; and when, instead of the parallel rays of daylight, we are obliged to use the diverging rays of a lamp, these should be rendered as parallel as possible, previously to their reflection from the mirror, by the interposition of the "bull's eye" condenser (§ 64) so adjusted as to produce this effect.

62. For the exhibition of those classes of objects which are suitable for "black ground" illumination, and which are better seen by light sent into them from every azimuth, than they are by a pencil, however bright, incident in one direction only, no more simple, convenient, and efficient means could probably be found, than that which is afforded by the "spotted lens" for low powers, and by the "parabolic illuminator" for powers as high as 1-4th or 1-5th of an inch focus;—the use of the latter with the highest powers, being rendered disadvantageous by the great reduction in the amount of light, occasioned by the necessity for cutting off of all the rays reflected from the paraboloid, which fall upon the object within the limits of their angle of aperture. One of the great advantages of this kind of illumination consists in this: that, as the light *radiates from* each part of the object as its proper source, instead of merely *passing through* it from a more remote source, its different parts are seen much more in their normal relations to one another, and it acquires far more of the aspect of solidity. The rationale of this is easily made apparent by holding up a glass vessel with a figured surface between one eye and a lamp or a window, so that it is seen by transmitted light alone; for the figures of its two surfaces are then so blended together to the eye, that unless their form and distribution be previously known, it can scarcely be said with certainty which markings belong to either. If, on the other hand, an opaque body be so placed behind the vessel, that no rays are transmitted directly through it, whilst it receives adequate illumination from the circumambient light, its form is clearly discerned, and the two surfaces are differentiated without the least difficulty.

63. *Polarizing Apparatus.*—In order to examine transparent objects by polarized light, it is necessary to employ some means of *polarizing* the rays before they pass through the object, and to apply to them, in some part of their course between the object and the eye, an *analyzing* medium. These two requirements may be provided for in different modes. The *polarizer* may be either a bundle of plates of thin glass, used in place of the mirror, and polarizing the rays by reflection; or it may be a "single image" or "Nicol" prism of Iceland Spar, which is so constructed as to transmit only one of the two rays into which a beam of ordinary light is made to divaricate on passing through this substance; or it may be a plate of Tourmaline, or one of

the artificial tourmalines composed of the disulphate of iodine and quinine, now known by the designation of "Herapathite," after the name of their discoverer. Of these methods, the "Nicol" prism is the one generally preferred; the objection to the reflecting polarizer being, that it cannot be made to rotate; the tourmaline being undesirable, on account of the color which it imparts when sufficiently thick to produce an effective polarization; and the crystals of Herapathite being seldom obtained perfect, of sufficient size to afford a good illumination. The polarizing prism is usually fitted into a tube (Fig. 41, A, *a*) with a large milled head (*c*) at the bottom, by which it is made to rotate in a collar (*b*) that is attached to the microscope; this collar may be fitted to the under side of the stage-plate, or, where a secondary stage is provided, it may be attached to this; in the microscope of Messrs. Smith and Beck, it screws into the lower part (*b*) of a tube (Fig. 41, B) that slides into the "cylindrical fitting" beneath the stage (Fig. 29). The analyzer, which

FIG. 41.



Fitting of Polarizing Prism in Smith and Beck's Microscope.

FIG. 42.




Fitting of Analyzing Prism upon the Eye-piece.

may be either a "Nicol" prism, a Tourmaline, or a crystal of Herapathite, is usually placed either in the interior of the microscope, or between the eye-piece and the eye. If it be a prism, it is mounted in a tube, which may either be screwed into the lower end of the body in the situation of the erector (Fig. 32), or may be fitted over the eye-piece in place of its ordinary cap (Fig. 42); in the former situation it has the advantage of not limiting the field, but it stops a considerable proportion of the light; in the latter, it detracts much less from the brightness of the image, but cuts off a good deal of the margin of the field. A plate of Tourmaline or Herapathite, if obtainable of sufficient size and freedom from color, has a decided advantage above the Nicol prism, as an analyzer, in being free from both these inconveniences; and it may be set in a cap which fits over the ordinary cap of the eye-piece. For bringing out certain effects of color by the use of Polarized light (Chap. XX), it is desirable to interpose a plate of Selenite beneath the polarizer and

the object; and it is advantageous that this should be made to revolve. A very convenient mode of effecting this, is to mount the selenite plate in a revolving collar, which fits into the upper end (*a*) of the tube (Fig. 41, B) that receives the polarizing prism. In order to obtain the greatest variety of coloration with different objects, films of selenite of different thickness should be employed; and this may be accomplished by substituting one for another in the revolving collar. A still greater variety may be obtained by mounting three films, which separately give three different colors, in a frame resembling that in which hand-magnifiers are usually mounted, so that they may be used singly or in double or triple combinations; as many as thirteen different tints may thus be obtained; but the advantage of revolution is sacrificed. When the construction of the microscope does not readily admit of the connection of the selenite plate with the polarizing prism, it is convenient to make use of a plate of brass (Fig. 43) somewhat larger than the glass

FIG. 43.



Selenite Object-Carrier.

slides in which objects are ordinarily mounted, with a ledge near one edge for the slide to rest against, and a large circular aperture into which a glass is fitted, having a film of selenite cemented to it; this "selenite stage" or object-carrier being laid upon the stage of the microscope, and the slide containing the object being placed upon it, the effect of the selenite is obtained, as in the previous arrangement; and by an ingenious modification contrived by Dr. Leeson, the ring into which the selenite plate is fitted being made movable, one plate may be substituted for another, whilst rotation may be given to the ring by means of a tangent-screw fitted into the brass plate. Such a "selenite stage" answers every purpose that can be required; but as there is no provision for using two or three plates in combination, it is necessary to have a distinct selenite plate for every modification of colors that may be desired.


A very beautiful effect may be obtained with certain kinds of semi-opaque objects, by illuminating them by means of a "spotted lens" (§ 61), with a polarizer of Herapathite placed at such a distance above it as to receive the converging hollow pencil near its termination in the object, and an analyzer of the usual description,—a combination devised by Mr. Furze;¹ for the solidity which this mode of oblique illumination imparts to certain objects, is remarkably heightened by the play of colors afforded by the polarization of the light. When the polarizing apparatus is being employed with any but the lowest powers, it is very advan-

¹ "Transactions of the Microscopical Society" (2d series), vol. iii, p. 63.

tageous to use the achromatic condenser in combination with it; this combination, which cannot be made in ordinary microscopes, is provided for in that of Messrs. Smith and Beck, by the "cylindrical fitting" so often referred to, which can receive the polarizing prism at its lower end, and the achromatic condenser at its upper, whilst the selenite plate or plates may be interposed between them.¹

64. *Illuminators for Opaque Objects.*—All objects through which sufficient light cannot be *transmitted* to enable them to be viewed in the modes already described, require to be illuminated by rays, which, being thrown *upon* the surface under examination, shall be *reflected* from it into the microscope; and this mode of viewing them may often be advantageously adopted in regard to semi-transparent or even transparent objects, for the sake of the diverse aspects it affords. Among the various methods devised for this purpose, the one most generally adopted consists in the use of a *condensing lens*, either attached to the microscope, or mounted upon a separate stand, by which the rays proceeding from a lamp or from a bright sky are made to converge upon the object. For the efficient illumination of large opaque objects, such as injected preparations, it is desirable to employ a "bull's-eye" condenser (which is a plano-convex lens of short focus, two or three inches in diameter), mounted upon a separate stand, in such a manner as to allow of being placed in a great variety of positions. The mounting shown in Fig. 44 is perhaps one of the best that can be adopted: the frame which carries the lens is borne at the bottom upon a swivel-joint, which allows it to be turned in any azimuth; whilst it may be inclined at any angle to the horizon, by the revolution of

FIG. 44.




Bull's-Eye Condenser.

¹ For an account of the nature and properties of Polarized Light, which would be out of place in the present treatise, see the chapters on that subject in Dr. Golding Bird's "Manual of Natural Philosophy," Dr. Pereira's "Lectures on Polarized Light," New Ed., edited by Prof. Baden Powell, or any modern treatise on Optics.

the horizontal tube to which it is attached, around the other horizontal tube which projects from the stem; by the sliding of one of these tubes within the other, again, the horizontal arm may be lengthened or shortened; the lens may be secured in any position (as its weight is apt to drag it down when it is inclined, unless the tubes be made to work, the one into the other, more stiffly than is convenient) by means of a tightening collar milled at its edges; and finally the horizontal arm is attached to a spring socket, which slides up and down upon a vertical stem. The optical effect of such a lens differs according to the side of it turned towards the light, and the condition of the rays which fall upon it. The position of *least* spherical aberration, is when its *convex* side is turned towards *parallel* or towards the *least diverging* rays; consequently, when used by daylight, its *plane* side should be turned towards the *object*; and the same position should be given to it, when it is used for procuring converging rays from a lamp, the lamp being placed four or five times farther off on one side, than the object is on the other. But it may also be employed for the purpose of reducing the diverging rays of the lamp to parallelism, for use either with the parabolic illuminator (§ 61), or with the side-reflector to be presently described; and the *plane* side is then to be turned towards the lamp, which must be placed at such a distance from the condenser, that the rays which have passed through

FIG. 45.



Ordinary Condensing Lens.

the latter shall form a luminous circle equal to it in size, at whatever distance from the lens the screen may be held. Even where the large "bull's-eye" condenser is provided, it is well to have a smaller condensing lens in addition; and this, which is usually a double-convex lens, may either be mounted on a separate base (Fig. 45), or may be attached to some part of the microscope. (In Messrs. Smith and Beck's large microscope, Fig. 29, two sockets with binding-screws, one for the condensing lens, the other for the side-reflector, are seen in the "limb.") This condensing lens is sufficient by itself for most ordinary purposes; and it may


also be used to obtain a greater concentration of the rays already brought into convergence by the bull's-eye (§ 93).

65. The illumination of opaque objects may be effected by *reflection*, as well as by *refraction*; and a very advantageous means

of using the light of a lamp for this purpose, is afforded by the *Side-Reflector* contrived by Mr. Ross. This is a highly polished concave speculum (Fig. 46), which can be placed above and to one side of the object; and which is so mounted as to be capable of being placed in every kind of position, according to the place

of the lamp, and the degree of obliquity of the illumination required. The squared stem, with which the speculum is connected by several intermediate joints, may be fitted to a socket, either in the stage or in some part of the microscope-stand, like that of the smaller condensing lens. The light reflected by the speculum upon the object, may be either that which falls on it direct from the lamp, or may come to it through the intervention of the bull's-eye, arranged so as to throw parallel rays upon the speculum (§ 64). The prisms already described as in use for the illumination of transparent objects by the reflection of light from beneath, may also be employed, by an inversion of their position, for the illumination of opaque objects from above. In Continental Microscopes, the prism is frequently attached to the lower end of the body; but this is an undesirable mode of supporting it, since the illumination is disturbed by every alteration in the distance between the body and the object. This seems to be provided against by the mounting of the prism in Mr. Grubb's microscope (§ 60), which allows it to be used at any angle either above or below the stage. A mode of illuminating opaque objects by a small concave speculum reflecting the light directly down upon it, was formerly much in use, but is now comparatively seldom employed. This concave speculum, termed a "Lieberkühn" from the celebrated Microscopist who invented it, is made to fit upon the end of the objective, having a perforation in the centre for the passage of the rays from the object to the lens; and it receives its light from the mirror beneath, the object being so mounted as only to stop out the central portion of the rays that are reflected upwards. The curvature of the speculum is so adapted to the focus of the object-glass, that, when the latter is duly adjusted, the rays reflected up to it from the mirror shall be made to converge strongly upon the part of the object that is in focus; consequently, unless (as is sometimes done) the speculum should be mounted on a tube sliding over the "nose" of

FIG. 46.



Side-Reflector.


the microscope, and capable of being adjusted to the different distances required by the several objectives, a separate speculum is required for every object-glass. The disadvantages of this mode of illumination are chiefly these:—first, that by sending the light down upon the object almost perpendicularly, there is scarcely any shadow, so that the inequalities of its surface, and any minute markings which it may present, are but faintly or not at all seen; second, that the size of the object must be so limited by that of the speculum, as to allow the rays to pass to its marginal portion; and third, that a special mode of mounting is required, to allow the light to be reflected from the mirror around the margin of the object. The first objection may be in some degree removed, by turning the mirror considerably out of the axis, so as to reflect its light obliquely upon the Lieberkühn, which will then send it down obliquely upon the object; the illumination, however, will not even then be so good as that which is afforded by the side-reflector. The mounting of opaque objects in wooden slides (Chapter V), which affords in many cases the most convenient means of preserving them, completely prevents the employment of the Lieberkühn in the examination of them; and they must either be set, for this purpose, upon disks which afford them no protection, or in glass cells with a blackened background. The cases wherein the Lieberkühn is most useful, are those in which it is desired to examine small opaque objects, such as can be held in the stage-forceps (§ 66), or laid upon a slip of glass, with lenses of half inch focus or less; since a stronger light can be thus concentrated upon them, than can be easily obtained by side-illumination. In every such case, a black background must be provided, of such a size as to fill the field, so that no light shall come to the eye direct from the mirror, and yet not large enough to create any unnecessary obstruction to the passage of the rays from the mirror to the speculum. With each Lieberkühn is commonly provided a blackened stop of appropriate size, having a well-like cavity, and mounted upon a pin which fits into a support connected with the under side of the stage; but though the “dark well” serves to throw out a few objects with peculiar force, yet, for all ordinary purposes, a spot made with black paper or black sealing-wax-varnish upon a slip of glass will answer the required purpose very effectually, the slip being simply laid upon the stage beneath the object.

SECTION 2. APPARATUS FOR THE PRESENTATION OF OBJECTS.

66. *Stage-Forceps*.—Every Microscope should be furnished with a pair of Stage-forceps (Fig. 47) for holding minute objects beneath the object-glass. They are mounted by means of a joint upon a pin, which fits into a hole either in the corner of the stage itself, or in the object-platform; the object is inserted by pressing the pin that projects from one of the blades, whereby it is sepa-

rated from the other; and the blades close again, so as to retain the object when the pressure is withdrawn. By sliding the wire stem which bears the forceps through its socket, and by moving that socket vertically upon its joint, and the joint horizontally

FIG. 47.



Stage-Forceps.


upon the pin, the object may be brought into the field precisely in the position required; and it may be turned round and round, so that all sides of it may be examined by simply giving a twisting movement to the wire stem. The other extremity of the stem often bears a small brass box filled with cork, and perforated with holes in its side; this affords a secure hold to common pins, to which disks of card, &c., may be attached, whereon objects are mounted for being viewed with the Lieberkühn. This method of mounting was formerly much in vogue, but has been less employed of late, since the Lieberkühn has fallen into comparative disuse.

67. *Glass Stage-Plate.*—Every Microscope should be furnished with a piece of plate-glass, about 4 in. by $1\frac{1}{2}$ in., to one margin of which a narrow strip of glass is cemented, so as to form a ledge. This is extremely useful, both for laying objects upon (the ledge preventing them from sliding down when the microscope is inclined), and for preserving the stage from injury by spilling of sea-water or other saline or corrosive liquids, when such are in use. Such a plate not only serves for the examination of transparent, but also of opaque objects; the dark background being furnished by the diaphragm-plate (§ 55), and the condensing-lens being so placed as to throw a side-light upon them. A small addition may be conveniently made to the glass stage-plate, which adapts it for use as a *Growing-Slide*. A circular aperture, of about the diameter of a test-tube, is made near one end of the plate (the length of which, for this purpose, had better be not less than 5 inches), and in this is to be fitted a little cup, formed of the end of a test-tube, about three-quarters of an inch deep, in such a manner that its rim shall project a little above the surface of the plate. The cup may be closed by an ordinary cork, or (to avoid the danger of splitting it) by a disk of glass cemented to a ring of cork which shall embrace the exterior of the tube; but a small aperture must be left, by grinding a notch in the rim of the cup, sufficient to admit the passage of two or three threads of lamp-cotton. The manner in which the "growing-slide" is used, is this:—Supposing we wish to follow the changes undergone by some minute Alga or Infusorium, which

we have just detected in a drop of liquid under examination upon an ordinary slip of glass (and covered with thin glass),—we transfer this slip to the “growing-slide,” fill the cup with distilled water mixed with a small proportion of the water in which the organism was found, and then so arrange the threads (previously moistened with distilled water), that they shall pass from the cup to the edge of the liquid in which the object is contained. Thus, as the water evaporates from beneath the thin glass, the threads will afford a continuous supply; and the threads will not become dry, until the whole of the liquid has been absorbed by them and has been dissipated by evaporation. Fresh supplies may, of course, be introduced into the cup from time to time, as may be needed, so as to prevent any loss of liquid from beneath the thin glass; and in this manner, the most important requisite for the continued growth of aquatic organisms,—a constant supply of liquid, without an exclusion of air,—may be secured.¹

68. *Aquatic Box or Animalcule Cage*.—This, also, is an appendage with which every Microscope should be provided, so varied and so constant is its utility. It consists of a short piece of wide brass tube, fixed perpendicularly at one end into a flat plate of brass (Fig. 48) which is perforated by an aperture equal in

FIG. 48.



Aquatic Box or Animalcule Cage, as seen in perspective at A, and in section at B.


diameter to that of the tube, and having its opposite extremity closed by a disk of glass (B b); over this fits a cover, formed of a piece of tube just large enough to slide rather stiffly upon that which forms the box, closed at the top by another disk of glass (B a). The cover being taken off, a drop of the liquid to be examined, or any thin object which can be most advantageously looked at in fluid, is placed upon the lower plate; the cover is then slipped over it, and is pressed down until the drop of liquid be spread out, or the object be flattened, to the degree most convenient for observation. If the glass disk which forms the lid be cemented or burnished into the brass ring which carries it, a small hole should be left for the escape of air or superfluous fluid; and this hole may be closed up with a morsel of wax, if

¹ See the “Micrographic Dictionary,” by Dr. Griffith and Mr. Henfrey, Introduction, p. xx.

it be desired to prevent the included fluid from evaporating. But as it is desirable that this glass should be thin enough to allow a 1-4th-inch objective to be employed for the examination of Animalcules, &c., and as such thin glass is extremely apt to be broken, it is a much better plan to furnish the brass cover with a screw-cap, which holds the brass disk with sufficient firmness, but permits it to be readily replaced when broken; and as the looseness of this fitting gives ample space for the escape of air or fluid around the margin of the disk, no special aperture is needed. It is always desirable, if possible, to prevent the liquid from spreading to the edge of the disk; since any objects it may contain are very apt, in such a case, to be lost under the opaque ring of the cover; this is to be avoided by limiting the quantity of the liquid introduced, by laying it upon the centre of the lower plate, and by pressing down the cover with great caution, so as to flatten the drop equally on all sides, stopping short when it is spreading too close to the margin. With a little practice, this object may in general be successfully attained; but if so much superfluous liquid should have been introduced, that it has flooded the circumference of the inclosed space, and exuded around the edge of the disk, it is better to wipe the whole perfectly dry, and then to introduce a fresh drop, taking more care to limit its quantity and to restrain it within convenient bounds. If the box be well constructed, and the glass disks be flat, they will come into such close contact, that objects of extreme thinness may be compressed between them; hence not only may such small animals as Water-fleas (*Entomostraca*) be restrained from the active movements which preclude any careful observation of their structure,—and this without any permanent injury being inflicted upon them,—but much smaller creatures, such as Wheel-Animalcules (*Rotifera*), or *Bryozoa*, may be flattened out, so as to display their internal organization more clearly, and even the larger *Infusoria* may be treated in like manner. The working Microscopist will find it of great advantage to possess several of these aquatic boxes, of different sizes; and one or two of them may have the glass cover of stronger glass than the rest, and firmly fixed in its rim, so that, if the cover be made to slide equably on the box, the instrument (in hands accustomed to careful manipulation) may be made to answer the purpose of a *compressorium* (§ 70).

69. *Zoophyte Trough*.—For the examination of living aquatic objects, too large to be conveniently received into the Aquatic Box, the Zoophyte trough contrived by Mr. Lister may be employed with great advantage. This consists of a trough of the shape represented in Fig. 48, formed of plates and slips of plate-glass cemented together by marine glue; of a loose vertical plate of glass, just so much smaller than the front or back of the inside of the trough, as to be able to move freely between its sides; and of a horizontal slip of glass, whose length equals that of the

inside bottom of the trough, but whose breadth is inferior by the



Zoophyte Trough.

thickness of the plate just mentioned. The trough being filled with water (fresh or salt, as the case may be), the horizontal slip is laid at the bottom, and the vertical plate is placed in contact with the front of the trough, its lower margin being received into the space left at the front edge of the horizontal slip, which serves to hold it there, acting as a kind of hinge; a small ivory wedge is then inserted between the *front* glass of the trough

and the upper part of the vertical plate, which it serves to press backwards; but this pressure is kept in check by a little spring of bent whalebone, which is placed between the vertical plate and the *back* glass of the trough. By moving the ivory wedge up or down, the amount of space left between the upper part of the vertical plate and the front glass of the trough can be precisely regulated; and as their lower margins are always in close apposition, it is evident that the one will incline to the other, with a constant diminution of the distance between them, from above downwards. Hence a Zoophyte, or any similar body, dropped into this space, will descend until it rests against the two surfaces of glass, and will remain there in a situation extremely convenient for observation; and the regulating-wedge, by increasing or diminishing the space, serves to determine the level to which the object shall fall. Of these troughs, again, it is convenient for the working Microscopist to be furnished with several, of different sizes; and in one of them *Chara* or *Nitella* may be kept growing in a state very convenient for observation. A similar trough may be provided for this last purpose, however, by dispensing with the vertical plate and horizontal slip altogether, and approximating the front and back plates so that only a very narrow space is contained between them; in this case it is convenient to let the upper lip of the back plate project considerably beyond that of the front plate, as objects may then be much more readily inserted between them; and the back plate may also be conveniently made to project beyond the sides of the trough, as would be useful, too, in the case of larger troughs. If it be wished to *grow Chara*, &c., in a thin trough of this kind, the trough, whenever it is not under observation, should be immersed in a tumbler or jar of water, since the plant will not flourish in a very limited supply.

70. *Compressorium*.—The purpose of this instrument is to apply a graduated pressure to objects, whose structure can only be made out when they are thinned by extension. For such as will bear tolerably rough treatment, a well-constructed Aquatic Box may be made to answer the purpose of a compressor; but

there is a very large class, whose organization is so delicate as to be confused or altogether destroyed by the slightest excess of pressure; and for the examination of such, an instrument in which the degree of compression can be regulated with precision, is almost indispensable. Various plans of construction have been proposed; but none among them appears to the Author to present so many advantages as the one represented in Fig. 50, the general plan of which was originally devised by Schiek of Berlin, but the details of which have been modified by M. de Quatrefages, who has constantly employed this instrument in his elaborate and most successful researches on the organization of the Marine Worms. It consists of a plate of brass between 3 and 4

inches long, and from $1\frac{1}{4}$ to $1\frac{1}{2}$ inch broad, having a central aperture of from $\frac{1}{2}$ to $\frac{3}{4}$ of an inch. This central aperture is covered on its upper side by a disk of thin glass,

which may be cemented to the brass plate by Canada balsam; and the under side of it is bevelled away, so that the thickness of the edge shall not interfere with the approach of the objective to its margin, when that side is made the uppermost. Near one extremity of the plate is a strong vertical pin, that gives support to a horizontal bar which turns on it as on a swivel; through the end of this bar that projects beyond the plate, there passes a screw with a milled head; and at the other end is jointed a second bar, against one end of which the screw bears, whilst the other carries a frame holding a second disk of thin glass. This frame is a small circular plate of glass, having an aperture equal in size to that of the large plate; to its under side, which is flat, a disk of thin glass is cemented by Canada balsam, while its upper side is bevelled off as it approaches the opening, for the purpose just now specified; and by being swung between pivots in a semicircle of brass, which is itself pivoted to the movable arm, it is made capable of a limited movement in any direction. The upper disk with the apparatus which supports it, having been completely turned aside round the swivel-joint, the object to be compressed is laid upon the lower disk; the upper disk is then turned back so as to lie precisely over it, and by the action of the milled-head screw, is gradually approximated to the lower, to which the pivot movements of its frame allow it to take up a parallel position, whatever may be the inclination of the bar. As it is frequently of great importance to be able to look at either side of the object under compression, the principal plate is provided with two pins at the extremity farthest from the milled head, which, being exactly equal in length to the swivel-pin, afford, with it, a support to the instrument, when it

FIG. 50.



Compressorium.

is so turned that the side represented as undermost in the figure, shall be uppermost; and it is in order that high powers may be used in this case as in the other, that the disk which then covers the object is made of thin glass, instead of being (as in the original form of the instrument) a piece of thick glass plate. That a thin glass disk is more liable to fracture under pressure, than a thick one, is no serious objection to its use for this purpose; since the lower one is not more likely to break than the upper one; and either may be replaced with extreme facility, by simply warming the part of the instrument to which it is attached, so as to loosen the cement that holds it. And the advantage of being able to view an object under a high power, *from either side*, will be most fully appreciated by every one who has been much engaged in the class of observations, which this instrument is specially adapted to facilitate.

FIG. 51.



Fishing Tubes.


If this Compressorium be made of sufficient size to admit an ordinary glass slide between the vertical pins, an object may be subjected to compression, and afterwards removed for examination out of the compressor, without transferring it from one glass to another, which is frequently an advantage. In this case, it will be convenient that the thin glass disk should be "countersunk" into the upper surface of the principal plate, so as to form one level with it.

71. *Dipping Tubes*.—In every operation in which small quantities of liquid, or small objects contained in liquid, have to be dealt with by the Microscopist, he will find it a very great convenience to be provided with a set of tubes of the forms represented in Fig. 51, but of somewhat larger dimensions. These were formerly designated as "fishing tubes;" the purpose for which they were originally devised having been the fishing out of Water-Fleas, aquatic Insect-Larvæ, the larger Animalcules, or other living objects distinguishable either by the unaided eye or by the assistance of a magnifying glass, from the vessels that may contain them. But they are equally applicable, of course, to the selection of minute Plants; and they may be turned to many other no less useful purposes, some of which will be specified hereafter. When it is desired to secure an object which can be seen either with the eye alone or with a magnifying glass, one of these tubes is passed down into the liquid, its upper orifice having been previously closed by the forefinger, until its lower orifice is close above the object; the finger being then removed, the liquid suddenly rises into the tube, probably carrying

the object up with it; and, if this is seen to be the case, by putting the finger again on the top of the tube, its contents remain in it when the tube is lifted out, and may be deposited on a slip of glass or on the lower disk of the aquatic box, or, if too copious for either receptacle, may be discharged into a watch-glass. In thus fishing for any but the minutest objects, it will be generally found convenient to employ the open-mouthed tube *c*; and when its contents have been discharged, if they include but a single object of the desiderated kind, this may be taken up by one of the finer tubes, *A*, *B*, or, if more convenient, the whole superfluous fluid may be sucked up by the mouth, and the object left with no more than is suitable; or, if there be many of the desired objects in the fluid first selected, these may be taken up from it, one by one, by either of the finer tubes.

72. *Forceps*.—Another instrument so indispensable to the Microscopist as to be commonly considered an appendage to the Microscope, is the Forceps for taking up minute objects; many forms of this have been devised, of which one of the most convenient is represented in Fig. 52, of something less than the

FIG. 52.



Forceps.

actual size. As the forceps, in marine researches, have continually to be plunged into sea-water, it is better that they should be made of brass or of German silver, than of steel, since the latter rusts far more readily; and as they are not intended (like dissecting forceps) to take a firm grasp of the object, but merely to hold it, they may be made very light, and their spring part slender. As it is essential, however, to their utility, that their points should meet accurately, it is well that one of the blades should be furnished with a guide-pin, passing through a hole in the other.

The foregoing constitute, it is believed, all the most important pieces of Apparatus, which can be considered in the light of Accessories to the Microscope. Those which have been contrived to afford facilities for the preparation and mounting of Objects, will be described in a future chapter (Chap. V). It may be thought that some notice ought to be taken of the *Frog Plate* and *Fish Pan*, with the former of which many Microscopes are supplied, whilst the latter has scarcely yet gone altogether out of use. But the Author having been accustomed to gain all the advantages of these, by methods far more simple, whilst at least equally efficacious, does not consider them as presenting any advantages which render it desirable to expend time or space in giving a detailed account of them; and he will explain the methods alluded to, under the appropriate head.

CHAPTER IV.

MANAGEMENT OF THE MICROSCOPE.

73. *Support.*—The Table on which the Microscope is placed, when in use, should be one whose size enables it also to receive the various appurtenances which the observer finds it convenient to have within his reach, and whose steadiness is such as to allow of his arms being rested upon it without any yielding; it should, moreover, be so framed, as to be as free as possible from any tendency to transmit the vibrations of the building or floor whereon it stands.¹ The manner in which the Microscope itself is constructed, however, will have a great influence on the effect of any such disturbing cause; since, if the whole instrument move together, scarcely any tremulousness will be produced in the image, by vibrations which cause it to “dance” most unpleasantly, if the body and stage of the Microscope oscillate independently of each other. Hence, in choosing a Microscope, it should always be subjected to this test, and should be unhesitatingly rejected if the result be unfavorable. It is of course to be borne in mind, that any vibration, either of the object or of the optical apparatus, in which the other does not partake, will be much more apparent when high magnifying powers are used, than when the object is amplified in a much less degree, the motion of the object being magnified in precisely the same ratio with the object itself; hence if, when the microscope is thus tested with high powers, it is found to be free from fault, its steadiness with low powers may be assumed; but, on the other hand, a Microscope which may give an image free from perceptible tremor when the lowest powers only are employed, may be quite unfit for use with the highest.

74. *Light.*—Whatever may be the purposes to which the Microscope is applied, it is a matter of the first importance to secure a pure and adequate illumination. There is scarcely any class

¹ The working Microscopist will find it a matter of great convenience to have a Table specially set apart for this purpose; furnished with drawers in which are contained the various accessories he may require for the preparation and mounting of objects. If the Microscope be one which is not very readily taken out from and put back into its case, it is very convenient to cover it with a large bell-glass; which may be so suspended from the ceiling, by a cord carrying a counterpoise at its other end, as to be raised or lowered with the least possible trouble, and to be entirely out of the way when the Microscope is in use. Similar but smaller bell-glasses are also useful for the protection of objects, which are in course of being examined or prepared, and which it is desirable to seclude from dust.

of objects, for the examination of which *good daylight* is not to be preferred to any other kind of light; but *good lamplight* is preferable to bad daylight. When daylight is employed, the Microscope should be placed near a window, whose aspect should be (as nearly as may be convenient) *opposite* to the side on which the sun is shining; for the light of the sun reflected from a bright cloud, is that which the experienced Microscopist will almost always prefer, the rays proceeding from a cloudless blue sky being by no means so well fitted for his purpose, and the dull lurid reflection of a dark cloud being the worst of all. The *direct* rays of the sun are far too powerful to be used with advantage, unless its intensity be moderated, either by reflection from a plaster-of-paris or some other "white-cloud" mirror (§ 58), or by passage through some imperfectly transparent medium. The *moderator* contrived by Mr. Rainey for lamp or gas-light (§ 75), has been found to answer equally well for direct sunlight; the glare and heating power of which it so effectually subdues, as to destroy all tendency to injure the most delicate object, or to confuse the observer's view of it; whilst an illumination is obtained by its means, whose intensity renders it superior for certain purposes to anything else. The young Microscopist is earnestly recommended to make as much use of *daylight* as possible; not only because, in a large number of cases, the view of the object which it affords is more satisfactory than that which can be obtained by any kind of lamp-light, but also because it is much less "trying" to the eyes. So great, indeed, is the difference between the two in this respect, that there are many who find themselves unable to carry on their observations for any length of time by lamp-light, although they experience neither fatigue nor strain from many hours' continuous work by daylight.

75. When recourse is had to Artificial light, it is of great importance, not only that it should be of good quality, but that the arrangement for furnishing it should be suitable to the special wants of the Microscopist. Thus, although a wax or composition candle affords a very pure light, yet its use is attended with two inconveniences, which render its use very undesirable when any better light can be obtained;—namely, the constant flickering of the flame, which is not sufficiently prevented by surrounding it with a chimney; and the continual alteration in its level, which is occasioned by the consumption of the candle. The most useful light for ordinary use, is that furnished by the steady and constant flame of the lamp, fed either with oil, camphine, or gas; the wick or burner should be cylindrical or "argand;" it should be capable of adjustment to any height above the table; and a movable shade should be provided, by which the light may be prevented from coming direct to the observer's eyes, or from diffusing itself too widely through the room. These requisites are supplied by the lamp commonly known as the "University"

or "reading" lamp, which has a circular foot with a vertical stem, on which the oil-reservoir (carrying with it the burner) and the shade, can be fixed at any convenient height. French and German lamps on the same general construction, but having the reservoir contrived on the "bird-fountain" principle, are also to be obtained, being largely imported for the use of watch-makers; these have the advantage of burning out all their oil, which is not the case with the ordinary "reading"-lamp, as it does not burn well except when full or nearly so; but they are usually destitute of a shade, which, however, can be easily added. Lamps of either kind are sometimes constructed on the "solar" principle, which increases the purity and intensity of the light, but at the same time not only diminishes the diameter of the flame, but also produces an inconvenient transverse "break" near its lower part. The best kind of light which an oil-lamp can furnish, is that yielded by the "Moderator" lamps which have of late come into such general use; but they have this important drawback, that they contain in themselves no adjustment for varying the elevation of the burner, and that their construction is such as to give no facilities for any arrangement of this kind. The same objection applies to the Camphine-lamps in ordinary use; but a small camphine-lamp has been constructed for the special use of Microscopists, which is capable of being placed on an adjustable stand, so that its flame may be raised or lowered to any desired level. The light of this lamp is whiter and more intense than that of any other, and it may be used with advantage for certain very delicate observations (§ 58); but for the ordinary purposes of the Microscopist it is not so convenient, the surface of flame from which the light can be received by the mirror or condenser, being limited by the peculiar construction which the combustion of camphine requires. To every one who has a supply of gas at command, the use of it for his microscope-lamp (by means of a flexible tube) strongly recommends itself, on account of its extreme convenience, and its freedom from any kind of trouble. The lamp should be constructed on the general plan already described, the burner being made to slide up and down on a stem rising perpendicularly from a foot, which also carries a shade; and the burner should be one which affords a bright and steady cylindrical flame, either "Leslie's" or the "cone"-burner being probably the best. Even the best light supplied by a gas-lamp, however, is inferior in quality to that of a good oil-lamp; and is more injurious and unpleasant to the eye. Hence the interposition of some kind of artificial medium, adapted to keep back the yellow rays, whose predominance in the lamp-flame is the chief source of its injurious action, is especially required when gas-light is used. This may be partly effected, by the simple expedient of using a chimney of bluish glass, known as "Leblond's;" but, in addition, it is advantageous to cause the light to pass through a screen of bluish-black or

neutral-tint glass; and it will then be nearly purified as to quality, though much reduced in intensity.¹ Mr. Rainey, who has paid great attention to the best means of obtaining a good illumination by artificial light, recommends, as the best *moderator*, one piece of dark blue glass, free from any tint of red, another of very pale blue with a slight shade of green, and two of thick white plate-glass, all cemented together with Canada balsam; this, as already stated, may be used with sun-light, as well as with lamp-light. The Microscopist who wishes to render the artificial light which he may be in the habit of using, as pure as possible, will do well to compare with daylight (as suggested by Dr. Griffith, who seems to have been the first to employ tinted glass with this object); furnishing himself with several pieces of glass of different shades, substituting one for another, and altering their distances from the lamp,² until he has succeeded in so tempering its rays, that the field of his Microscope, or the object under view, is not more colored when illuminated by the artificial light reflected from the mirror, than it is when the mirror is so turned as to reflect good light from a white cloud.

76. *Position of the Light.*—When the Microscope is used by daylight it will usually be found most convenient to place it in such a manner, that the light shall be at the left hand of the observer. It is most important that no light should enter his eye, save that which comes to it through the Microscope; and the access of direct light can scarcely be avoided, when he sits with his face to the light. Of the two sides, it is more convenient to have the light on the *left*; first, because it is not interfered with by the right hand, when this is employed in giving the requisite direction to the mirror, or in adjusting the illuminating apparatus; and secondly, because, as most persons employ the right eye rather than the left, the projection of the nose serves to cut off those lateral rays, which, when the light comes from the right side, glance between the eye and the eye-piece. In order to prevent, still more completely, the access of false light, it is desirable, if it be otherwise convenient, that when daylight is employed, its source should be a little behind the observer: but as it will then, by falling upon the stage, interfere with the view of any object which is imperfectly transparent (§ 87), it may be necessary to keep it from doing so, by the interposition of a screen. When Artificial light is employed, the same general precautions should be taken. The lamp should always be placed on the left side,

¹ A gas-lamp provided with these and other appurtenances for regulating the illumination, and also with a water-bath and mounting-plate, is supplied by Mr. S. Highley, Fleet Street.

² The nearer the colored glass is approximated to the flame, the less modification will it produce in its rays; since *their* intensity varies in different parts of their course, inversely with the square of their distance from the illuminating centre, whilst *its* influence is a constant quality. Hence a pale-blue glass placed near the mirror, or between the mirror and the stage, has more effect than a chimney of much deeper blue immediately surrounding the flame.

unless the use of the mirror be dispensed with, or some special reason exist for placing it otherwise. If the object under examination be transparent, the lamp should be placed at a distance from the eye about midway between that of the stage and that of the mirror; if on the other hand, the object be opaque, it should be at a distance about midway between the eye and the stage; so that its light may fall, in the one case upon the mirror, in the other case upon the stage, at an angle of about 45° with the axis of the microscope. The passage of direct rays from the flame to the eye, should be guarded against by the interposition of the lamp-shade; and no more light should be diffused through the apartment, than is absolutely necessary for other purposes. If observations of a very delicate nature are being made, it is desirable, alike by daylight and by lamp-light, to exclude all lateral rays from the eye, as completely as possible; and this may be readily accomplished, by means of a *shade* attached to the eye-piece of the microscope. Such a shade may be made most simply of an oblong piece of card-board, having a circular hole cut in it, by which it may fit upon the eye-piece or the upper part of the body; its two ends should be turned up, so as to cut off all lateral light; its upper side should also be turned up, so as to cut off the light from the front; and a notch should be cut in its lower edge, in the proper position to receive the nose. It may be either painted black, or may be covered with black cloth or velvet.

77. *Care of the Eyes.*—Although most Microscopists acquire a habit of employing only *one* eye (generally the right), yet it will be decidedly advantageous to the beginner, that he should learn to use either eye indifferently; since by employing and resting each alternately, he may work much longer, without incurring unpleasant or injurious fatigue, than when he always employs the same. Whether or not he do this, he will find it of great importance to acquire the habit of *keeping open the unemployed eye*. This, to such as are unaccustomed to it, seems at first very embarrassing, on account of the interference with the microscopic image, which is occasioned by the picture of surrounding objects, formed upon the retina of the second eye; but the habit of restricting the attention to that impression only which is received through the microscopic eye, may generally be soon acquired; and when it has once been formed, all difficulty ceases. Those who find it unusually difficult to acquire this habit, may do well to learn it in the first instance with the assistance of the shade just described; the employment of which will permit the second eye to be kept open without any confusion. The advantage of the practice, in diminishing the fatigue of long-continued observation, is such, that no pains are ill bestowed by the Microscopist, which are devoted to early habituation to it. There can be no doubt that the habitual use of the Microscope for many hours together, especially by lamp-light, and with high magnifying

powers, has a great tendency to injure the sight. Every Microscopist who thus occupies himself, therefore, will do well, as he values his eyes, not merely to adopt the various precautionary measures already specified, but rigorously to observe the simple rule of *not continuing to observe, any longer than he can do so without fatigue.*

78. *Care of the Microscope.*—Before the Microscope is brought into use, the cleanliness and dryness of its glasses ought to be ascertained. If dust or moisture should have settled on the Mirror, this can be readily wiped off. If any spots should show themselves on the field of view, when it is illuminated by the mirror, these are probably due to particles adherent to one of the lenses of the Eye-piece; and this may be determined by turning the eye-piece round, which will cause the spots also to rotate, if their source lies in it. It may very probably be sufficient to wipe the upper surface of the eye-glass (by removing its cap), and the lower surface of the field-glass; but if, after this has been done, the spots should still present themselves, it will be necessary to unscrew the lenses from their sockets, and to wipe their inner surfaces; taking care to screw them firmly into their places again, and not to confuse the lenses of different eye-pieces. Sometimes the eye-glass is obscured by dust of extreme fineness, which may be carried off by a smart puff of breath; the vapor which then remains upon the surface being readily dissipated, by rapidly moving the glass backwards and forwards a few times through the air. And it is always desirable to try this plan in the first instance; since, however soft the substance with which the glasses are wiped, their polish is impaired in the end by the too frequent performance of the process. The best material for wiping glass, is a piece of soft wash-leather, from which the dust it generally contains has been well beaten out. If the Object-glasses be carefully handled, and be kept in their boxes when not in use, they will not be likely to require cleansing. One of their chief dangers, however, to which they are liable in the hands of an inexperienced Microscopist, arises from the neglect of precaution in using them with fluids; which, when allowed to come in contact with the surface of the outer glass, should be wiped off as soon as possible. In screwing and unscrewing them, great care should be taken to keep the glasses at a distance from the surface of the hands; since they are liable not only to be soiled by actual contact, but to be dimmed by the vaporous exhalation from skin which they do not touch. This dimness will be best dissipated, by moving the glass quickly through the air. It will sometimes be found, on holding an object-glass to the light, that particles either of ordinary dust, or more often of the black coating of the interior of the microscope, have settled upon the surface of its back lens; these are best removed by a clean and dry camel-hair pencil. If any cloudiness or dust should still present itself in an object-glass, after its front and back surfaces

have been carefully cleansed, it should be sent to the maker (if it be of English manufacture) to be taken to pieces, as the amateur will seldom succeed in doing this without injury to the work; the foreign combinations, however, being usually put together in a simpler manner, may be readily unscrewed, cleansed, and screwed together again. Not unfrequently an objective is rendered dim by the cracking of the cement by which the lenses are united, or by the insinuation of moisture between them; this last defect occasionally arises from a fault in the quality of the glass, which is technically said to "sweat." In neither of these cases has the Microscopist any resource, save in an Optician experienced in this kind of work; since his own attempts to remedy the defect are pretty sure to be attended with more injury than benefit.

79. *General Arrangement of the Microscope for Use.*—The inclined position of the instrument, already so frequently referred to, is that in which Observation by it can be so much more advantageously carried on than it can be in any other, that this should always be had recourse to, unless particular circumstances render it unsuitable. The precise inclination that may prove to be most convenient, will depend upon the "build" of the Microscope, upon the height of the observer's seat as compared with that of the table on which the instrument rests, and lastly, upon the tallness of the individual; and it must be determined in each case by his own experience of what suits him best,—that which he finds *most comfortable*, being that in which he will be able not only to work the longest, but to see most distinctly. The selection of the object-glasses and eye-pieces to be employed, must be entirely determined by the character of the object. Large objects presenting no minute structural features, should always be examined in the first instance by the *lowest* powers, whereby a general view of their nature is obtained; and since, with lenses of comparatively long focus and small angle of aperture, the precision of the focal adjustment is not of so much consequence as it is with the higher powers, not only those parts can be seen which are exactly in focus, but those also can be tolerably well distinguished, which are not precisely in that plane, but are a little nearer or more remote. When the general aspect of an object has been sufficiently examined through low powers, its details may be scrutinized under a higher amplification; and this will be required in the first instance, if the object be so minute, that little or nothing can be made out respecting it, save when a very enlarged image is formed. The power needed in each particular case, can only be learned by experience; that which is most suitable for the several classes of objects hereafter to be described, will be specified under each head. In the general examination of the larger class of objects, the range of power that is afforded by the "erector" in combination with the "draw-tube" (§ 44), will be found very useful; whilst for the ready exchange

of a low power for a high one, great convenience is afforded by Mr. Brooke's object-glass holder (§ 50).

80. When the Microscopist wishes to augment his magnifying power, he has a choice between the employment of an Objective of shorter focus, and the use of a deeper Eye-piece. If he possess a complete series of objectives, he will generally find it best to substitute one of these for another, without changing the eye-piece for a deeper one; but if his "powers" be separated by wide intervals, he will be able to break the abruptness of the increase in amplification which they produce, by using each objective first with the shallower, and then with the deeper eye piece. Thus if a Microscope be only provided with two objectives, of $\frac{1}{2}$ inch and $\frac{1}{4}$ inch focus respectively, and with two eye-pieces, one nearly double the power of the other (as is the case with Messrs. Smith and Beck's new Educational Microscope, p. 103, *note*), such a range as the following may be obtained,—55, 100, 200, 350 diameters; or, with two objectives of somewhat shorter focus, and with deeper eye-pieces (as is the case with an instrument in the Author's possession, constructed by Kellner of Wetzlar, whose Microscopes have acquired for themselves a deservedly high reputation),—88, 176, 350, 700 diameters. The use of the "draw-tube" (§ 43) enables the Microscopist still further to vary the magnifying power of his instrument, and thus to obtain almost any exact number of diameters he may desire, within the limits to which he is restricted by the focal length of his objectives. The advantage to be derived, however, either from "deep eye-piecing," or from the use of the draw-tube, will mainly depend upon the quality of the object-glass. For if it be imperfectly corrected, its errors are so much exaggerated, that more is lost in definition than is gained in amplification; whilst, if its aperture be small, the loss of light is an equally serious drawback. On the other hand, a combination of perfect construction and wide angle of aperture, will sustain this treatment with so little impairment in the perfection of its image, that a magnifying power may be obtained by its use, such as, with an inferior instrument, can only be derived from an objective of much shorter focus combined with a shallow eye-piece.¹ In making any such comparisons, it must be constantly borne in mind that the real question is, *what can be seen?* It is always desirable for the purposes of research, to employ the *lowest* power with which the details of structure can be clearly made out; since, the lower the power, the less is the liability to error from false appearances, and the better can the mutual relations of the different parts of the object be appreciated. Hence in testing the optical quality of a Microscope, the question should always

¹ The 4-10ths object-glass of Messrs. Smith and Beck was specially distinguished by the Jurors of the Great Exhibition, as affording, by the use of deep eye-pieces and the draw tube, a power fully equivalent in the resolution of difficult tests, to that which, a few years previously, could only have been given by an objective of $\frac{1}{8}$ th inch.

be,—not, what is its *greatest* magnifying power,—but, what is the *least* magnifying power under which it will show objects of a given degree of difficulty.

81. In making the Focal Adjustment, when low powers are used, it will scarcely be necessary to employ any but the *coarse movement*; provided that the rack be well cut, the pinion work in it smoothly and easily, without either “spring,” “loss of time,” or “twist,” and the milled head be large enough to give the requisite leverage. All these are requisites which should be found in every well-constructed instrument; and its possession of them should be tested, like its freedom from vibration, by the use of high powers, since a really good coarse adjustment should enable the observer to “focus” an objective of 1-8th inch with precision. What is meant by “spring” is the alteration which may often be observed to take place on the withdrawal of the hand; the object which has been brought precisely into focus, and which so remains as long as the milled head is between the fingers, becoming indistinct when the milled head is let go. The source of this fault may lie either in the rack-movement itself, or in the general framing of the instrument, which is so weak as to allow of displacement by the mere weight or pressure of the hand; should the latter be the case, the “spring” may be in great degree prevented, by carefully abstaining from *bearing on* the milled head, which should be simply *rotated* between the fingers. By “loss of time” is meant the want of sufficient readiness in the action of the pinion upon the rack, so that the milled head may be moved a little in either direction, without affecting the body; thus occasioning a great diminution in the sensitiveness of the adjustment. This fault may sometimes be detected in Microscopes of the best original construction, which have gradually worked loose, from the constancy with which they have been in employment; and it may often be corrected by tightening the screws that bring the pinion to bear against the rack. And by “twist” it is intended to express that apparent movement of the object across the field, which results from a real displacement of the axis of the body to one side or the other, owing to a want of correct fitting in the working parts. As this last fault depends entirely upon bad original workmanship, there is no remedy for it; but it is one which most seriously interferes with the convenient use of the instrument, however excellent may be its optical performance. In the use of the coarse adjustment with an objective of short focus, extreme care is necessary to avoid bringing it down upon the object, to the injury of one or both; for although the spring with which the tube for the reception of the object-glass is furnished, whenever the fine adjustment is immediately applied to this (§ 31), takes off the violence of the crushing action, yet such an action, even when thus moderated, can scarcely fail to damage or disturb the object, and *may* do great mischief to the lenses. Where no such spring-

tube is furnished, the fine adjustment being otherwise provided for, or being not supplied at all, still greater care is of course required. It is here, perhaps, well to notice, for the guidance of the young Microscopist, that the *actual* distance between the object-glass and the object, when a distinct image is formed, is always considerably less than the *nominal* focal length of the object-glass; thus, the distance of the 1 inch or 2-3 inch object-glass may be little more than half an inch; that of the 4-10 inch may be but little more than one-tenth of an inch; that of a 1-4 or a 1-5 inch may scarcely exceed one-twentieth; that of a 1-8 inch may not be one-fortieth; and that of a 1-12 or a 1-16 inch may be so close as not to admit the intervention of a piece of glass no more than one-hundredth of an inch in thickness. The reason of this is, that the focal length of an Achromatic objective is estimated by that of the Single lens with which it agrees in the size of the image it forms, and therefore in magnifying power (*e. g.*, it is said to be of 1 inch focus, when its power is equivalent to that of a single lens, which brings parallel rays to a point at an inch distance); whilst from its being composed of a combination of lenses, the point from which that focal *distance* has really to be measured, is not at the surface of its front lens, but at some distance behind it, in the interior of the combination. One more precaution it may be well to specify;—namely, that either in changing one object for another, or in substituting one objective for another, save when powers of such focal length are employed as to remove all likelihood of injury, the “body” should be turned to one side, where the construction of the Microscope admits of this displacement, or (where it does not) should have its distance from the stage increased by the “coarse movement.” This precaution is absolutely necessary, when objectives of short focus are in use, to avoid injury either to the lenses or to the object; and when it is habitually practised with regard to these, it becomes so much like an “acquired instinct,” as to be almost invariably practised in other cases.

82. In obtaining an exact Focal Adjustment with object-glasses of less than half an inch focus, it will be generally found convenient to employ the *fine movement*; and as recourse will frequently be had to its assistance for other purposes also, it is very important that it should be well constructed and in good working order. The points to be particularly looked to in testing it, are for the most part the same with those already noticed in relation to the coarse movement. It should work smoothly and equably, producing that *graduated* alteration of the distance of the object-glass from the object, which it is its special duty to effect, without any jerking or irregularity. It should be so sensitive, that any movement of the milled head should at once make its action apparent, by an alteration in the distinctness of the image, when high powers are employed, without any “loss of

time."¹ And its action should not give rise to any twisting or displacing movement of the image, which ought not to be in the least degree disturbed by any number of rotations of the milled head, still less, by a rotation through only a few degrees. One great use of the "fine adjustment" consists in bringing into view different *strata* of the object, and this in such a gradual manner that their connection with one another shall be made apparent. Whether an opaque or a transparent object be under examination, only that part can be perfectly discerned under any power, which is exactly in focus; and when high powers of large aperture are employed, this is the only part that can be seen at all. A minute alteration of the focus often causes so entirely different a set of appearances to be presented, that, if this alteration be made abruptly, their relation to the preceding can scarcely be even guessed at; and the gradual transition from the one to the other, which the fine adjustment alone affords, is therefore necessary to the correct interpretation of either. To take a very simple case:—The transparent body of a certain animal being traversed by vessels lying in different planes, one set of these vessels is brought into view by one adjustment, another set by "focussing" to a different plane; and the connection of the two sets of vessels, which may be the point of most importance in the whole anatomy of the animal, may be entirely overlooked, for want of a fine adjustment, the graduated action of which shall enable one to be traced continuously into the other. What is true even of low and medium powers, is of course true to a still greater degree of high powers; for although the "coarse movement" may enable the observer to bring any stratum of the object into accurate focus, it is impossible for him by its means to secure that *transitional* "focussing," which is often so much more instructive than an exact adjustment at any one point. A clearer idea of the nature of a doubtful structure is, in fact, often derived from what is caught sight of *in the act* of changing the focus, than by the most attentive study and comparison of the different views obtained by any number of separate "focussings." The experienced Microscopist, therefore, when examining an object of almost any description, constantly keeps his finger upon the milled head of the "fine movement," and watches the effect produced by its revolution upon every feature which he distinguishes; never leaving off, until he be satisfied that he has scrutinized not only the entire *surface*, but the entire *thickness* of the object. It will often happen, that, where different structural features present themselves on different planes, it will be difficult or even impossible to determine which of them is the nearer and


¹ It will sometimes happen that the "fine movement" will seem not to act, merely because it has been so habitually worked in one direction rather than the other, that its screw has been turned too far. In that case, nothing more is required for its restoration to good working order, than turning the screw in the other direction, until it shall have reached about the middle of its range of action.

which the more remote (it being the special result of the ordinary mode of viewing objects by transmitted light, that such differences are obliterated), unless, by the use of the "fine movement," it be ascertained, when they are successively brought into focus, whether the object-glass has been moved *towards* or *away from* the object. Even this, however, will not always succeed in certain of the most difficult cases, in which the difference of level is so slight as to be almost inappreciable;—as, for instance, in the case of the markings on the siliceous *loricæ* of the Diatomaceæ (Fig. 80).

83. When objectives of short focus and of wide angular aperture are being employed, something more is necessary than exact focal adjustment; this being the *Adjustment of the Object-glass* itself, which is required to neutralize the disturbing effect of the glass cover upon the course of the rays proceeding from the object (§ 15). For this adjustment, it will be recollected, a power of altering the distance between the front pair and the remainder of the combination is required; and this power is obtained in the following manner. The front pair of lenses is fixed into a tube (Fig. 53, A), which slides over an interior tube (B) by which

the other two pairs are held; and it is drawn up or down by means of a collar (c), which works in a furrow cut in the inner tube, and upon a screw-thread cut in the outer, so that its revolution in the plane to which it is fixed by the one tube, gives a vertical movement to the other. In one part of the outer tube, an oblong slit is made, as seen at D, into which projects a small tongue, screwed on the inner tube; at the side of the former two

horizontal lines are engraved, one pointing to the word "uncovered," the other to the word "covered;" whilst the latter is crossed by a horizontal mark, which is brought to coincide with either of the two lines by the rotation of the screw-collar, which moves the outer tube up or down. When the mark has been made to point to the line "uncovered," it indicates that the distance of the lenses of the object-glass is such, as to make it suitable for viewing an object without any interference from thin glass; when, on the other hand, the mark has been brought, by the revolution of the screw-collar, into coincidence with the line "covered," it indicates that the front lens has been brought into



Section of an Adjusting Object-Glass.

such proximity with the other two, as to produce a "positive aberration" in the objective, fitted to neutralize the "negative aberration" produced by the interposition of a glass cover of a certain thickness. It is evident, however, that unless the particular thickness of glass for which this degree of alteration is suited, be always employed for this purpose, the correction cannot be exact; and means must be taken for adapting it to every grade of thickness, which may be likely to present itself in the glass covers. Unless this correction be made with the greatest precision, the enlargement of the angle of aperture, to which our Opticians have of late applied themselves with such remarkable success, becomes worse than useless; being a source of diminished instead of increased distinctness in the details of the object, which are far better seen with an objective of greatly inferior aperture, possessing no special adjustment for the thickness of the glass. The following general rule is given by Mr. Wenham, for securing the most efficient performance of an object-glass with any ordinary object:—"Select any dark speck or opaque portion of the object, and bring the outline into perfect focus; then lay the finger on the milled head of the fine motion, and move it briskly backwards and forwards in both directions from the first position. Observe the expansion of the dark outline of the object, both when within, and when without, the focus. If the greater expansion, or coma, is when the object is *without* the focus, or furthest from the objective, the lenses must be placed further asunder, or towards the mark 'uncovered.' If the greater coma is when the object is *within* the focus, or nearest to the objective, the lenses must be brought closer together, or towards the mark 'covered.' When the object-glass is in proper adjustment, the expansion of the outline is exactly the same both within and without the focus." A different indication, however, is afforded by such "test-objects" as present (like the Podura-scale and the Diatomaceæ) a set of distinct dots or other markings. For "if the dots have a tendency to run into lines when the object is placed *without* the focus, the glasses must be brought closer together; on the contrary, if the lines appear when the object is *within* the focal point, the object must be further separated.¹ When the angle of aperture is very wide, the difference in the aspect of any severe test under different adjustments becomes at once evident; markings which are very distinct when the correction has been exactly made, disappearing almost instantaneously when the screw-collar is turned a little way round."²

¹ See "Quart. Journ. of Microsc. Science," vol. ii, p. 138.

² Mr. Wenham remarks (*loc. cit.*), not without justice, upon the difficulty of making this adjustment, even in the Objectives of our best Opticians; and he states that he has himself succeeded much better, by making the outer tube the fixture, and by making the tube that carries the other pairs slide within this; the motion being given by the action of an inclined slit in the revolving collar, upon a pin that passes through a longitudinal slit in the outer tube, to be attached to the inner. The whole range of adjust-

84. Although the *most perfect* correction required for each particular object (which depends, not merely upon the thickness of its glass cover, but upon that of the fluid or balsam in which it may be mounted) can only be found by experimental trial, yet for all ordinary purposes, the following simple method, first devised by Mr. Powell, will suffice. The object-glass, adjusted to "uncovered," is to be "focussed" to the object; its screw-collar is next to be turned, until the surface of the glass cover comes into focus, as may be perceived by the spots or striæ by which it may be marked; the object is then to be again brought into focus by the "fine movement." The edge of the screw-collar being now usually graduated, the particular adjustment which any object may have been found to require, and of which a record has been kept, may be made again without any difficulty. By Messrs. Smith and Beck, however, who first introduced this graduation, a further use is made of it. By experiments such as those described in the last paragraph, the correct adjustment is first found for any particular object, and the number of divisions observed, through which the screw-collar must be moved in order to bring it back to 0° , the position suitable for an uncovered object. The thickness of the glass cover must then be measured by means of the "fine movement:" this is done by bringing into exact focus, first the object itself, and then the surface of the glass cover, and by observing the number of divisions through which the milled head (which is itself graduated) has passed in making this change. A definite ratio between that thickness of glass, and the correction required in that particular objective, is thus established; and this serves as the guide to the requisite correction for any other thickness, which has been determined in like manner by the "fine movement." Thus, supposing a particular thickness of glass to be measured by 12 divisions of the milled head of the fine movement, and the most perfect performance of the object-glass to be obtained by moving the screw-collar through 8 divisions, then a thickness of glass measured by 9 divisions of the milled head, would require the screw-collar to be adjusted to 6 divisions in order to obtain the best effect. The ratio between the two sets of divisions is by no means the same for different combinations; and it ought to be determined for each objective by its maker, who will generally be the best judge of the best "points" of his lenses; but when this ratio has been once ascertained, the adjustment for any thickness of glass with which the object may happen to be covered, is readily made by the Microscopist himself. Although this method appears somewhat more complex than that of Mr. Powell, yet it is more perfect; and when the ratio between the two sets of divisions has been once determined, the adjustment does not really involve

ment is thus performed within a third part of a revolution, with scarcely any friction, and with such an immediate transition from good to bad definition, that the best point is made readily apparent.


more trouble. Another use is made of this adjustment by Messrs. Smith and Beck; namely, to correct the performance of the objectives, which is disturbed by the increase of distance between the objective and the eye-piece, that is occasioned by the use of the draw-tube (§ 43). Accordingly, they mark a scale of inches on the draw-tube (which is useful for many other purposes), and direct that for every inch the body is lengthened, the screw-collar of the objective shall be moved through a certain number of divisions.

85. *Arrangement for Transparent Objects.*—If the object be already “mounted” in a slide, nothing more is necessary, in order to bring it into the right position for viewing it, than to lay the slide upon the object-platform of the stage, and to support it in such a position (by means of the sliding ledge or other contrivance) that the part to be viewed is, as nearly as can be guessed, in the centre of the aperture of the stage, and therefore in a line with the axis of the body. If the object be not “mounted,” and be of such a kind that it is best seen dry, it may be simply laid upon the glass stage-plate (§ 67), the ledge of which will prevent it from slipping off when the microscope is inclined, and a plate of thin glass may be laid over it for its protection, if its delicacy should seem to render this desirable. If, again, it be disposed to curl up, so that a slight pressure is needed to flatten or extend it, recourse may be had to the use of the aquatic box (§ 68), or of the compressorium (§ 70), no liquid, however, being introduced between the surfaces of glass. In a very large proportion of cases, however, either the objects to be examined are already floating in fluid, or it is preferable to examine them in fluid, on account of the greater distinctness with which they may be seen; if such objects be minute, and the quantity of liquid be small, the drop is simply to be laid on a slip of glass, and covered with a plate of thin glass; if the object or the quantity of liquid be larger, it will be better to place it in the aquatic box; whilst, if the objects have dimensions which render even this inconvenient, the zoophyte trough (§ 69) will afford the best medium for its examination. If it be wished to have recourse to *compression*, for the expansion or flattening of the object, this may be made upon the ordinary slide, by pressing down the thin glass cover with a pointed stick; and this method, which allows the pressure to be applied where it may chance to be most required, will generally be found preferable for delicate portions of tissue which are easily spread out, and which, in fact, require little other compression than is afforded by the weight of the glass cover, and by the capillary attraction which draws it into proximity with the slide beneath. A firmer and more enduring pressure may be exerted by the dexterous management of a well-constructed aquatic box; and this method is peculiarly valuable for confining the movements of minute animals, so as to keep them at rest under

the field of the microscope, without killing them. It is where a firm but graduated pressure is required, for the flattening out of the bodies of thin semi-transparent animals, without the necessity of removing them from the field of the microscope, that the compressorium is most useful. Wherever the first and simplest of the above methods can be had recourse to, it is the preferable one; since the object, when on a glass slide, can be subjected to the Achromatic Condenser, Polariscope, Oblique Illumination, &c., with far more convenience than when removed to a plane above the stage, as it must be when the aquatic box is used. Whether the object be submitted to examination on a slip of glass, or in the aquatic box or compressorium, it must be first brought approximately into position, and supported there, just as if it were in a mounted slide. The precise mode of effecting this will differ, according to the particular plan of the instrument employed; thus in some, it is only the ledge itself that slides along the stage; in others, it is a carriage of some kind, whereon the object-slide rests; in others, again, it is the entire platform itself that moves upon a fixed plate beneath.

86. Having guided his object, as nearly as he can do by the unassisted eye, into its proper place, the Microscopist then brings his light (whether natural or artificial) to bear upon it, by turning the mirror in such a direction as to reflect upon its under surface the rays which are received by itself from the sky or the lamp. The *concave* mirror is that which should always be first employed, the *plane* being reserved for special purposes; and it should bring the rays to convergence in or near the plane in which the object lies (Fig. 54). The distance at which it should be ordinarily set beneath the stage, is that at which it brings parallel rays to a focus; but this distance should be capable of elongation, by the lengthening of the stem to which the mirror is attached; since the rays diverging from a lamp at a short distance, are not so soon brought to a focus. The correct focal adjustment of the mirror may be judged of, by its formation of images of window-bars, chimneys, &c., upon any semi-transparent medium placed in the plane of the object. It is only, however, when

FIG. 54.



Arrangement of Microscope for Transparent Objects.

small objects are being viewed under high magnifying powers, that such a concentration of the light reflected by the mirror is either necessary or desirable; for with large objects, seen under low powers, the field would not in this mode be equably illuminated. The diffusion of the light over a larger area may be secured, either by shifting the mirror so much above or so much below its previous position, that the pencil will fall upon the object whilst still converging or after it has met and diverged; or, on the other hand, by the interposition of a plate of ground glass in the course of the converging pencil,—this last method, which is peculiarly appropriate to lamp-light, being very easily had recourse to, if the diaphragm plate, as formerly recommended (§ 55), have had its largest aperture filled with such a diffused medium. The eye being now applied to the Eye-piece, and the body being “focussed,” the object is to be brought into the exact position required, by the use of the traversing movement, if the stage be provided with it; if not, by the use of the two hands, one moving the object-slide from side to side, the other pushing the ledge, fork, or holder that carries it, either forwards or backwards, as may be required. It is always to be remembered, in making such adjustments by the direct use of the hands, that, owing to the inverting action of the microscope, the motion to be given to the object, whether lateral or vertical, must be precisely opposed to that which its image *seems* to require, save when the Erector (§ 44) is employed. When the object has been thus brought fully into view, the Mirror may require a more accurate adjustment. What should be aimed at, is the diffusion of a clear and equable light over the entire field; and the observer should not be satisfied, until he has attained this object. If the field should be darker on the one side than on the other, the mirror should be slightly turned in such a direction as to throw more light upon that side; perhaps in so doing, the light may be withdrawn from some part previously illuminated; and it may thus be found that the pencil is not large enough to light up the entire field. This may be owing to one of three causes: either the cone of rays may be received by the object too near to its focal apex, the remedy for which lies in an alteration in the distance of the mirror from the stage; or, from the very oblique position of the mirror, the cone is too much narrowed across one of its diameters, and the remedy must be sought in a change in the position either of the microscope or of the lamp, so that the face of the mirror may not be turned so much away from the axis of vision; or, again, from the centre of the mirror being out of the optical axis of the instrument, the illuminating cone is projected obliquely, an error which can be rectified without the least difficulty. If the cone of rays should come to a focus in the object, the field is not unlikely to be crossed by the images of window-bars or chimneys, or the form of the lamp-flame may be

distinguished upon it; the former must be got rid of by a slight change in the inclination of the mirror; and if the latter cannot be dissipated in the same way, the lamp should be brought a little nearer.

87. The equable illumination of the entire field having been thus obtained, the *quantity* of light to be admitted should be regulated by the Diaphragm-plate (§ 55). This must depend very much upon the nature of the object, and upon the intensity of the light. Generally speaking, the more transparent the object, the less light does it need for its most perfect display; and its most delicate markings are frequently only made visible, when the major part of the cone of rays has been cut off. Thus the movement of the *cilia*,—those minute vibratile filaments, with which almost every Animal is provided in some part of its organism, and which many of the humbler Plants also possess,—can only be discerned in many instances, when the light is admitted through the smallest aperture. On the other hand, the less transparent objects usually require the stronger illumination which is afforded by a wider cone of rays; and there are some (such as semi-transparent sections of fossil teeth) which, even when viewed with low powers, are better seen with the intenser light afforded by the Achromatic Condenser. In every case in which the object presents any considerable obstruction to the passage of the rays through it, great care should be taken to protect it entirely from *incident* light; since this extremity weakens the effect of that which is received into the microscope by transmission. It is by daylight that this interference is most likely to occur: since, if the precautions already given (§ 76) respecting the use of lamp-light be observed, no great amount of light *can* fall upon the upper surface of the object. The observer will be warned that such an effect is being produced, by perceiving that there is a want, not only of brightness, but of clearness, in the image; the field being veiled, as it were, by a kind of thin vapor; and he may at once satisfy himself of the cause, by interposing his hand between the stage and the source of light, when the immediate increase of brilliancy and of distinctness will reveal to him the occasion of the previous deficiency in both. Nothing more is necessary for its permanent avoidance, than the interposition of an opaque screen (blackened on the side towards the stage) between the window and the object; care being of course taken, that the screen does not interfere with the passage of light to the mirror. Such a screen may be easily shaped and adapted either to be carried by the stage itself, or by the stand for the condenser; but it is seldom employed by Microscopists, as it interferes with access to the left side of the stage; and the interposition of the hand, so often as it may be needed, is more frequently had recourse to in preference, as the more convenient expedient. The young Microscopist who may be examining transparent objects by daylight, is recommended never to omit

ascertaining, whether the view which he may obtain of them, is in any degree thus marred by incident light.

88. Although the illumination afforded by the mirror alone is quite adequate for a very large proportion of the purposes for which the Microscope may be profitably employed (nothing else having been used by many of those who have made most valuable contributions to Science by means of this instrument), yet, when high magnifying powers are employed, and sometimes even when but a very moderate amplification is needed, great advantage is gained from the use of the *Achromatic Condenser*. The various modes in which this may be constructed, and may be fitted to the Microscope, have been already described (§ 56); we have now to speak of the manner of using it. The lenses with which the Condenser is provided should be made to separate from each other, in such a manner that two or three distinct powers should be afforded; the complete combination should be used with objectives of 1-5th inch focus or less; the front lens should be removed with objectives of from half to a quarter of an inch focus; and the second lens may be removed, so that the back lens will be alone employed, when it is desired to use the condenser with objectives of less than half an inch focus. It is of the greatest importance that the Condenser should be accurately adjusted, both as to the coincidence of its optical axis with that of the Microscope itself, and as to its focal distance from the object. The *centring* may be most readily accomplished, by so adjusting the distance of the condenser from the stage (by the rack-and-pinion action, or the sliding movement, with which it is always provided), that a sharp circle of light shall be thrown on any semi-transparent medium laid upon it; then, on this being viewed through the Microscope with an objective of sufficiently low power to take in the whole of it, if this circle be not found to be concentric with the field of view, the axis of the condenser must be altered by means of the milled-head tangent screws with which it is provided. The *focal adjustment*, on the other hand, must be made under the objective which is to be employed in the examination of the object, by turning the mirror in such a manner as to throw upon the visual image of the object (previously brought into the focus of the Microscope) an image of a chimney or window-bar, if daylight be employed, or of the top, bottom, or edge of the lamp-flame, if lamp-light be in use; such a vertical movement should be given to the condenser, as may render the view of this as distinct as possible; and the direction of the mirror should then be sufficiently changed to displace these images, and to substitute for them the clearest light that can be obtained. It will generally be found, however, that although such an exact focussing gives the most perfect results by daylight, yet that by lamp-light the best illumination is obtained, when the condenser is removed to a somewhat greater distance from the object, than that at which it gives

a distinct image of the lamp. In every case, indeed, in which it is desired to ascertain the effect of *variety* in the method of illumination, the effects of alterations in the distance of the condenser from the object should be tried; as it will often happen that delicate markings become visible when the condenser is a little *out of focus*, which cannot be distinguished when it is precisely *in focus*. The diaphragm-plate with which all the best forms of Achromatic Condenser are now furnished, enables the observer not only to vary the angle of his illuminating pencils through a range of from 20° to 80° , but also to stop off the central portion of the pencil, so as to allow only its most oblique rays to pass; and the contrast presented by the aspect of many objects, according as the size and form of the aperture in the diaphragm-plate limits the rays transmitted by the condenser to those of the central or those of the peripheral portion of the pencil, is often so marked, as to show beyond question the great importance of this mode of varying the illumination. When the condenser is employed, the *plane* Mirror may often be substituted with advantage for the concave; the chief effect of this exchange being to diminish the quantity of light, without altering the angle of the illuminating pencil. It must be borne in mind, in making such an alteration, that the plane mirror reflects parallel or (if from a lamp) diverging rays, instead of the converging rays reflected by the concave mirror; so that the focus of the condenser is likely to require readjustment. For objects of great delicacy and transparency, the "white-cloud" illumination (§ 58) may be had recourse to with advantage; or, if it be desired that the illuminating pencil should be free from the error imparted by the double reflection of the mirror, the mirror may be turned aside, and in its stead the lamp (if the observation be made by artificial light) may be placed in the axis of the microscope; or the mirror may be replaced by "Dujardin's prism" (§ 57), which will be equally available either by daylight or by lamp-light.

89. Should it be desired, however, to try the effect of very oblique light upon an object, the Achromatic Condenser must be removed (unless, as in Mr. Sollitt's arrangement, § 130, it be so constructed as to be capable of inclination to the axis of the Microscope), and other means must be employed. The simplest method, where the mirror is mounted on an "arm" (Fig. 29), is to turn it to one side so as to reflect the rays at a considerable angle; and where this cannot be done, nearly the same effect is produced by placing the lamp in the direction from which it is desired that the oblique rays should proceed, and interposing an ordinary condensing lens between it and the object. Or, if the Microscopist be provided with the means of mounting a "Dujardin's prism" on a separate stand, he may place it in such a position as to reflect light from any point required: and he may concentrate that light by an ordinary condenser. The possession

of Amici's prism, however (which serves both as mirror and condenser, § 60), will save the necessity of any other provision of this kind. It is when objects are thus illuminated by oblique light, and when their markings are of such a kind as to be *best* or to be *only* shown by light falling upon them in one particular direction, that we derive the greatest advantage from the power of giving a rotatory movement either to the object or to the illuminating apparatus. Thus suppose that an object be

marked by longitudinal striæ, too faint to be seen by ordinary direct light; the oblique light most fitted to bring them into view, will be that proceeding in either of the directions c or d; that which falls upon it in the directions A and B, tending to obscure the striæ rather than to disclose them. But, moreover,

if the striæ should be due to furrows or prominences which have one side inclined and the other side abrupt, they will not be brought into view indifferently by light from c or from d, but will be shown best by that which makes the strongest shadow; hence if there be a projecting ridge, with an abrupt side looking towards c, it will be best seen by light from d; whilst if there be a furrow with a steep bank on the side of c, it will be by light from that side that it will be best displayed. But it is not at all unfrequent for the longitudinal striæ to be crossed by others; and these transverse striæ will usually be best seen by the light that is least favorable for the longitudinal; so that, in order to bring them into distinct view, either the illuminating pencil or the object must be moved a quarter round. The revolving action with which the stage of Mr. Ross's Microscope is provided (§ 37), enables this movement to be given to the object without any displacement of its image, which, of course, executes, to the eye of the observer, a rotation in the opposite direction. In other microscopes, however, it is difficult to give a rotation to the object, by causing the object-platform to turn upon its axis, without throwing the object out of the field (§ 38); though this *may* be accomplished, by such an adjustment of the traversing movement, as shall bring the centre of the tube on which that platform turns round, into the visual axis of the microscope—or, if this adjustment cannot be conveniently made in the first instance, by keeping the right hand constantly in action upon the milled heads of the stage movement, whilst the left hand rotates the object-platform, so as, by means of the former, to correct the displacement of the object occasioned by the latter. It may be sufficient, however, to examine the object in several different positions, so that the appearances it presents in each may be compared, without thus watching the transition from one to the other.

90. There are many kinds of transparent objects, especially such as either consist of thin plates, disks, or spicules of siliceous or calcareous matter, or contain such bodies, which are peculiarly

well seen under the *black-ground* illumination (§§ 61, 62); for not only does the brilliant luminosity which they then present, contrasting remarkably well with the dark ground behind them, show their forms to extraordinary advantage; but this mode of illumination imparts to them an appearance of solidity, which they do not exhibit by ordinary transmitted light (§ 62); and it also frequently brings out surface-markings, which are not otherwise distinguishable. Hence, when any object is under examination, that can be supposed to be a good subject for this method, the trial of it should never be omitted. For the low powers, the use of the "spotted lens" will be found sufficiently satisfactory; for the higher, the paraboloid should be employed (§ 61). Similar general remarks may be made, respecting the examination of objects by *polarized* light. Some of the most striking effects of this kind of illumination, are produced upon bodies whose particles have a crystalline aggregation; and hence it may often be employed with great advantage to bring such bodies into view, when they would not otherwise be distinguished; thus, for example, the *raphides* of Plants are much more clearly made out by its means, in the midst of the vegetable tissues, than they can be by any other. But the peculiar effects of polarized light are also exerted upon a great number of other organized substances, both Animal and Vegetable; and it often reveals differences in the arrangement or in the relative density of their component particles, the existence of which would not otherwise have been suspected; hence, the Microscopist will do well to have recourse to it, whenever he may have the least suspicion that its use can give him an additional power of discrimination.

91. *Arrangement for Opaque Objects.*—Although a large proportion of the objects best suited for Microscopic examination are either in themselves sufficiently transparent to admit of being viewed by light *transmitted* through them, or may be made so by appropriate means, and although that method (where it *can* be adopted) is generally the one best fitted for the elucidation of the details of their structure, yet there are many objects of the most interesting character, the opacity of which entirely forbids the use of this method, and of which, therefore, the *surfaces* only can be viewed, by means of the *incident* rays which they *reflect*. These are, for the most part, objects of comparatively large dimensions, for which a low magnifying power suffices; and it is specially important, in the examination of such objects, not to use a lens of shorter focus than is absolutely necessary for discerning the details of the structure; since, the longer the focus of the objective employed, the less is the indistinctness produced by inequalities of the surface, and the larger, too, may be its aperture, so as to admit a greater quantity of light, to the great improvement of the brightness of the image. It is surprising how little attention has been given by Opticians to the construction of objectives suitable for this purpose. In their zeal for the im-


provement of the higher powers of the Microscope, they have thought comparatively little of the lower; and in Continental Microscopes, it is rare to meet with an objective which will give even a tolerable view of a large opaque object. The Author, indeed, well remembers the time, when it was not thought worth while, even by English Opticians, to construct Achromatic object-glasses of less than an inch focus; and the production of objectives of $1\frac{1}{2}$ inch and 2 inch focus has been chiefly called for, in consequence of their value in displaying anatomical preparations in which the bloodvessels have been injected with coloring matter. The view which is afforded of large opaque objects, however, by a Compound Microscope, furnished with even an imperfectly corrected Achromatic object-glass, giving a magnifying power of 20 or 25 diameters, is so greatly preferable to that which is given by any Simple Microscope, that no instrument that is intended for general research should be unfurnished with such a power. It is especially required in Microscopes that are to be used for Educational purposes; since it is most important that the young should be trained in a knowledge of the wonders and beauties of the familiar objects around them; and an objective of low power and wide aperture, adapted to the examination of a large surface at once, affords a means of displaying these, such as can be afforded in no other way, save by the use of the Erector and draw-tube (§ 44). A microscope furnished with these appendages, need not be supplied with an objective of longer focus than 1 inch or 8-10ths inch; but the Author would strongly recommend to such as do not possess them, that they should give to a "dividing" $1\frac{1}{2}$ inch or 2 inch (in which the front lens is removable, and is replaced by a perforated cap that limits the aperture of the back lens, which is then employed by itself, having a focus of about 3 inches) a preference over such as do not thus supply the extremely low power which he recommends.¹

92. The mode of bringing opaque objects under view, will differ according to their "mounting," and to the manner in which it is desired to illuminate them. If the object be mounted in a "slide" of glass or wood, upon a large opaque surface, the slide must be laid on the stage in the usual manner, and the object brought as nearly as possible into position by the eye alone (§ 84). If it be not so mounted, it may be simply laid upon the glass stage-plate, resting against its ledge; and the diaphragm-plate must then be so turned, as to afford it a black background.

¹ A single pair (flint and crown) of about 2 inch focus, was constructed at the Author's request, some years since, by Messrs. Smith and Beck, for the special purpose of exhibiting injected preparations, and other opaque objects; and its performance has been so satisfactory to him, that he was induced to urge upon the Microscopic Committee appointed by the Society of Arts, that the Educational Microscope for which they invited competition (§ 31), should be furnished with such a power. This recommendation having been adopted, the instrument selected has been specially fitted for the class of objects above alluded to.

For all ordinary purposes, a plano or double-convex lens, of about $1\frac{1}{2}$ inch diameter, and 2 inches focus, either mounted upon a separate stand (as in Fig. 45), or so attached by a jointed support to the Microscope itself as to admit of being placed in any required position, will answer extremely well as a Condenser. If Daylight be employed, the microscope should be so placed that the strongest light may fall obliquely upon the stage, and preferably from the left hand side; there will then be no difficulty in so disposing this condenser, as to afford an illumination sufficient for almost any kind of object, provided the quality of

FIG. 55.



Arrangement of Microscope for Opaque Objects.

the light itself be good. Direct sunlight cannot be here employed, without the production of an injurious glare, and the risk of burning the object; but the sunlight reflected from a bright cloud is the best light possible. The condenser should always be placed at right angles to the direction of the illuminating rays, and at a distance from the object which will be determined by the size of the surface to be illuminated and by the kind of light required. If the magnifying power employed be high, and the field of view be consequently limited, it will be desirable so to adjust the lens, as to bring the cone of rays to a point upon the part of the object under examination; and this adjustment can only be rightly made whilst the object is kept in view under the microscope, the condenser being moved in various modes, until that position has been found for it in which it gives the best light. If, on the other hand, the power be low, and it be desired to spread the light equably over a large field, the condenser should be placed either within or beyond its focal distance; and here, too, the best position will be ascertained by trial. It will often be desirable, also, to vary both the obliquity of the light, and the direction in which it falls upon the object; the aspect of which is greatly affected by the manner in which the shadows are projected upon its surface, and in which the lights are reflected from the various points of it. There are many objects, indeed, distinguished by their striking appearance when the light falls upon them on one side, which are entirely destitute both of brilliancy of color and of sharpness of outline,

when illuminated from the opposite side. Hence it is always desirable to try the effect of changing the position of the object; which, if it be "mounted," may be first shifted by merely reversing the place of the two ends of the slide, and then, if this be not satisfactory, may be more completely as well as more gradually altered, by making the object-platform itself revolve, where the stage is fitted with such a movement: if, however, the object be not mounted, but be simply resting on the stage-plate, it may be readily shifted by hand. With regard to the obliquity of the illuminating rays, it is well to remark, that if the object be "mounted" under a glass cover, and the incident rays fall at too great an angle with the perpendicular, a large proportion of them will be reflected, and the brilliancy of the object will be greatly impaired.

93. The same general arrangement must be made, when Artificial light is used for the illumination of opaque objects; the lamp being placed in such a position in regard to the stage, that its rays may fall in the direction indicated in Fig. 55; and these rays being collected and concentrated by the condenser, as already directed. As the rays proceeding from a lamp within a short distance are already diverging, they will not be brought by the condenser to such speedy convergence, as are the parallel rays of daylight; and it must, therefore, be further removed from the object, to produce the same effect. By modifying the distance of the condenser from the lamp and from the object respectively, the cone of rays may be brought nearly to a focus, or it may be spread almost equally over a large surface, as may be desired. In the illumination of opaque objects, the inferiority of artificial to solar light is not so perceptible as in the case of transparent objects; and the former has the advantage of being more easily concentrated to the precise degree, and of being more readily made to fall in the precise direction, that may be found most advantageous. Moreover, the contrast of light and shadow will be more strongly marked, when no light falls upon the object except that proceeding from the lamp used for its illumination, than it can be when the shadows are partially lightened by the rays which fall upon the object from every quarter, as must be the case if it be viewed by daylight. If the ordinary condensing lens do not afford a sufficient illumination, the large "bull's-eye" condenser (§ 64) may be employed; its convex side being turned towards the lamp, when it is desired to bring its rays into the most complete convergence. And, if a still more concentrated light be required for the illumination of a small object under a high power, the small condenser may be so placed as to receive the cone where it is reduced to its own size; since, by its means, the rays may be brought to a more exact convergence than they can be by the bull's-eye alone. In this manner, very minute bodies may be viewed as opaque objects under a tolerably high magnifying power; provided that the

brasswork of the extremities of the objectives be so bevelled off, as to allow the illuminating cone to have access to the object.¹ No method of illuminating large opaque objects by lamp-light is more effective, than the reflection of light from a concave speculum placed near the side of the object (§ 65); this not only affords a brilliant light, which may be equably spread over as large a surface as may be required, but may, by the mode in which it is jointed to its supports, be made to throw its rays upon the object at a great variety of angles, without the necessity of moving the lamp, whereby the direction in which the best illumination can be gained, is readily ascertained. If a more intense light and a greater concentration be required, than the speculum will afford by reflecting the diverging rays of the lamp, these may be rendered parallel or slightly convergent by the interposition of the bull's-eye condenser, which, for such a purpose, must have its *plane* side turned towards the lamp. This speculum cannot be so advantageously used by daylight, the ordinary condensing lens being then decidedly preferable.

94. If the object which it is desired to examine be of small size, and of a shape and character that render it unsuitable to be laid upon the glass stage-plate, or to be turned over so as to bring each side in turn into the most advantageous position,—as is the case, for example, with the capsules of Mosses, the mouths of which cannot be conveniently brought into view in this mode,—it may be grasped in the stage-forceps (§ 66), which afford great facility for this kind of manipulation; or, if it be too minute or delicate to be thus held, it may be taken up upon the head of a small pin, by moistening this with saliva or with a little thin gum-water; and the pin may then be either held in the stage-forceps, or may be run into the cork at its opposite extremity. By careful manipulation, every part of such an object may be brought under view successively, and may be exposed to every variety of illumination. It is in viewing objects supported in this mode, that the utility of the Lieberkühn (§ 65) is chiefly felt; for, as the stage-forceps needs to be shifted into different positions, so that the object is sometimes raised above and sometimes depressed below the level of the stage, in order to present it under a different aspect, the side illumination, whatever be its source, needs to be newly adjusted with each change in the position of the object; whilst the Lieberkühn adjusts itself, so to speak, when the object is brought into focus. If the mirror be so mounted that it can be turned considerably out of the axis of the microscope, and the aperture of the stage be sufficiently large, a light of considerable obliquity may be reflected from the Lie-

¹ Since the introduction of the Parabolic illuminator, the occasions on which advantageous recourse can be had to the examination of minute objects with high powers by incident light, have become much less numerous; since these objects are for the most part sufficiently transparent to admit of being illuminated by that instrument; and when they are so, the view of them which it affords is generally much superior to any that can be gained by the method of illumination described above.

berkühn ; thus enabling it to afford a kind of illumination, which, as already remarked, is usually much more valuable than that produced by the nearly perpendicular rays sent down by it on the object, when the mirror is placed in the axis. Whenever the Lieberkühn is employed, care must be taken that the direct light from the mirror be entirely stopped out by the interposition of a "dark well" or of a black disk, of such a size as to *fill* the field given by the particular objective employed, but not to pass much beyond it. An ingenious combination of a *hemispherical* Lieberkühn with the Paraboloid (§ 61) has been devised by Mr. Wenham, for the illumination of minute opaque objects by very oblique rays,¹ and Mr. C. Brooke has attached a small *plane* speculum to objectives of 1-8th and 1-12th inch focus (which cannot be otherwise advantageously employed with that illuminator), in such a manner that its surface is level with, or very little below, that of the outer lens, so as to reflect downwards upon the object those extreme pencils of rays which pass by the aperture of the object-glass. In either case, an oblique illumination from one side only may be obtained, by shutting off either half of the lower aperture of the paraboloid. These contrivances for the examination of minute objects with high powers by incident light, have scarcely yet received the attention they deserve.

95. *Errors of Interpretation.*—The correctness of the conclusions which the Microscopist will draw, regarding the nature of any object, from the visual appearances which it presents to him, when examined in the various modes now specified, will necessarily depend in great degree upon his previous experience in microscopic observation, and upon his knowledge of the class of bodies to which the particular specimen may belong. Not only are observations of *any* kind liable, as already remarked (Introduction, pp. 39-41), to certain fallacies arising out of the previous notions which the observer may entertain, in regard to the constitution of the objects or the nature of the actions to which his attention is directed; but even the most practised observer is apt to take no note of such phenomena as his mind is not prepared to appreciate. Thus, for example, it cannot be doubted that many Physiologists must have seen those appearances in thin slices of Cartilage, which are now interpreted as denoting its *cellular* organization, without in the least degree suspecting their real import, which Schwann was the first to deduce from the study of the development of that tissue; it was not known before his time, "what cells mean" in Animal organization; and the retinal pictures which now suggest the idea of them to the mind of even the tyro in the study of Histology (p. 54), passed almost entirely unnoticed by keen-sighted and intelligent Microscopists previously to 1839. Errors and imperfections of this kind can only be corrected, it is obvious, by general advance in scientific knowledge; but the history of them affords a useful

¹ "Quart. Journ. of Microsc. Science," vol. ii, p. 155.

warning against hasty conclusions drawn from a too cursory examination. If the history of almost *any* scientific investigation were fully made known, it would generally appear, that the stability and completeness of the conclusions finally arrived at, had only been attained after many modifications, or even entire alterations, of doctrine. And it is, therefore, of such great importance to the correctness of our conclusions, as to be almost essential, that they should not be finally formed and announced, until they have been tested in every conceivable mode. It is due to Science, that it should be burdened with as few false facts and false doctrines as possible. It is due to other truth-seekers, that they should not be misled, to the great waste of *their* time and pains, by *our* errors. And it is due to ourselves, that we should not commit our reputation to the chance of impairment, by the premature formation and publication of conclusions, which may be at once reversed by other observers better informed than ourselves, or may be proved to be fallacious at some future time, perhaps even by our own more extended and careful researches. *The suspension of the adjudgment, whenever there seems room for doubt*, is a lesson inculcated by all those Philosophers who have gained the highest repute for practical wisdom; and it is one which the Microscopist cannot too soon learn, or too constantly practise.

96. Besides these general warnings, however, certain special cautions should be given to the young Microscopist, with regard to errors into which he is liable to be led, by the misinterpretation of appearances peculiar to objects thus viewed, even when the very best instruments are employed. Thus the sharpness of the outline of any transparent object is impaired by a change in the course of the rays that merely *pass by* it, which is termed *Inflection* or *Diffraction*. If any opaque object be held in the course of a cone of rays diverging from a focus, the shadow which it will form upon a screen held to receive it, will not possess a well-defined *edge*, but will have as its boundary a shaded *band*, gradually increasing in brightness from the part of the screen on which the shadow is most intense, to that on which the illumination is most complete. If the light be homogeneous in its quality, the shaded band will possess no colors of its own; but if the light be decomposable, like the ordinary solar beam, the band will exhibit prismatic fringes.¹ It is obvious that such a diffraction must exist in the rays transmitted through the substance, as well as along the edges, of transparent objects; and that it must interfere with the perfect distinctness, not merely of their outlines, but of their images, the various markings of which are *shadows* of portions that afford obstacles, more or

¹ This phenomena is explained on the Undulatory Theory of light, by the disturbance which takes place in the onward propagation of waves, when subsidiary centres of undulation are developed by the impact of the principal undulations on obstacles in their course; the chromatic dispersion being due to the inequality in the lengths of the undulations proper to the severally colored rays.

less complete, to the perfectly free transmission of the rays. There are many objects of great delicacy, in which the "diffraction-band" is liable to be mistaken for the indication of an actual substance; on the other hand, the presence of an actual substance of extreme transparency, may sometimes be doubted or denied, through its being erroneously attributed to the "diffraction-band."¹ No rules can be given for the avoidance of such errors, since they can only be escaped by the discriminative power which education and habit confer. The practised Microscopist, indeed, almost instinctively makes the requisite allowance for diffraction; and seldom finds himself embarrassed by it, in the interpretation of the visual appearances which he obtains through a good instrument. Besides this unavoidable result of the inflection of the rays of light, there is a peculiar phenomenon attendant upon oblique illumination at certain angles in one direction; which consists in the production of a double image, or a kind of overlying shadow, sometimes presenting markings equally distinct with those of the object itself. This image, which is not unlike the secondary spectrum formed by reflection from the outer surface of a silvered-glass mirror, has been called the "diffracting spectrum;" but its origin does not really lie in the diffraction of the luminous rays, since on the one hand it cannot be explained according to the laws of diffraction, and on the other it may be traced to an entirely different cause. An object thus illuminated is seen by two different sets of rays; those, namely, of *transmitted* light, which pass through it obliquely from the source of the illumination to the opposite side of the object-glass; and those of *radiated* light, which, being intercepted by the object, are given off from it again in all directions. (The latter alone are the rays whereby the images are formed in any kind of "black-ground" illumination, §§ 61, 62.) Two different images will be formed, when the illuminating pencil is very oblique, and the angular aperture of the object-glass is wide; one of them by the light transmitted to one extreme of its aperture, the other by the light radiated to its general surface; and one or the other of these images may be stopped out, by covering that portion of the lens which receives, or that which does not receive, the transmitted pencil. This "diffracting-spectrum" may be produced at pleasure, in an object illuminated by direct light and seen with a large aperture, by holding a needle or a horse-hair before the front lens, so as to split the aperture into two parts.

97. Errors of interpretation arising from the imperfection of the Focal adjustment, are not at all uncommon amongst young

¹ Thus the account given by Prof. Sharpey and the Author, of the structure of Muscular Fibre (Chap. XVIII), has been called in question by observers who had not seen their preparations, on the ground that the "diffraction-band" had not been allowed for. To whatever the appearance in question (Fig. 326) may be due, there cannot be the slightest question that it does *not* arise from diffraction.

Microscopists. With lenses of high power, and especially with those of large angular aperture, it very seldom happens that all the parts of an object, however small and flat it may be, can be in focus together; and hence the focal adjustment being exactly made for one part, everything that is not in exact focus is not only more or less indistinct, but is often wrongly represented. The indistinctness of outline will sometimes present the appearance of a pellucid border, which, like the diffraction-band, may be mistaken for actual substance. But the most common error is that which is produced by the reversal of the lights and shadows, resulting from the refractive powers of the object itself; thus, the bi-concavity of the blood-disks of Human (and other Mammalian) blood, occasions their centres to appear *dark*, when in the focus of the Microscope, through the dispersion of the light which it occasions; but when they are brought a little within the focus, by a slight approximation of the object-glass, the centres appear brighter than the peripheral parts of the disks (Fig. 315). The same reversal presents itself in the case of the markings of the Diatomaceæ; for these, when the surface is exactly in focus, are seen as light hexagonal spaces, separated by dark partitions; and yet, when the surface is slightly beyond the focus, the hexagonal areas are dark, and the intervening partitions light (Fig. 80). The best means of avoiding errors of interpretation arising from this source, lies in the employment of the *lowest* powers with which the particular structures can be distinguished; since, if the different parts of the surface and margin of the object can be simultaneously brought so nearly into focus that a distinct view may be gained of all of them at once, no false appearances will be produced, and everything will be seen in its *real* aspect.

98. A very important and very frequent source of error, which sometimes operates even on experienced Microscopists, lies in the refractive influence exerted by certain peculiarities in the form or constitution of objects, upon the rays of light transmitted through them; this influence being of a nature to give rise to appearances in the image, which suggest to the observer an idea of their cause that may be altogether different from the reality. A very characteristic illustration of the fallacy resulting from external configuration, is furnished by the notion which long prevailed among Microscopic observers, and which still lingers in the public mind, of the *tubular* structure of the Human hair. This notion has no other foundation, than the existence of a bright band down the axis of the hair, which is due to the convergence of the rays of light occasioned by the *convexity* of its surface, and which is equally shown by any other transparent cylinder; and it is unmistakably disproved by the appearances presented by thin transverse sections of Hair, which show that it is not only filled up to its centre with a medullary substance, but that its centre is sometimes even darker than the surrounding

part (Fig. 311). Of the fallacy which may sometimes arise from diversities in the refractive power of the internal parts of an object, we have an equally "pregnant instance" in the misinterpretation of the nature of the *lacunæ* and *canaliculi* of Bone (Fig. 300), which were long supposed to be solid corpuscles with radiating filaments of peculiar opacity, instead of being, as is now universally admitted, minute chambers with diverging passages, excavated in the solid osseous substance. For just as the convexity of its surfaces will cause a transparent cylinder to show a bright axial band, so will the concavity of the internal surfaces of the cavities or tubes hollowed out in the midst of highly refracting substances, occasion a divergence of the rays passing through them, and consequently render them so dark that they are easily mistaken for opaque solids. That such is the case with the so-called "bone-corpuscles," is shown by the effects of the infiltration of Canada balsam through the osseous substance; for when this fills up the excavations,—being nearly of the same refractive power with the bone itself, and being also quite transparent, and (in thin laminæ) quite colorless,—it obliterates them altogether. So, again, if a person who is unaccustomed to the use of the microscope should chance to have his attention directed to a preparation mounted in liquid or in balsam, that might chance to contain *air-bubbles*, he will be almost certain to be so much more strongly impressed by the appearance of these, than by that of the object, that his first remark will be upon the number of strange-looking black rings which he sees, and his first inquiry will be in regard to their meaning.

99. No experienced Microscopist could now be led astray by such obvious fallacies as those alluded to; but it is necessary to dwell upon them, as warnings to those who have still to go through the same education. The best method of learning to appreciate the class of appearances in question, is the comparison of the aspect of globules of Oil in water, with that of globules of Water in oil, or of bubbles of Air in water or Canada-balsam. This comparison may be very readily made by shaking up some oil with water to which a little gum has been added, so as to form an emulsion; or by simply placing a drop of oil of turpentine and a drop of water together on a slip of glass, laying a thin glass cover upon them, and then moving the cover several times backwards and forwards upon the slide.¹ Now when such a mixture is examined with a sufficiently high magnifying power, all the globules present nearly the same appearance, namely, dark margins with bright centres; but when the test of alteration of the focus is applied to them, the difference is at once revealed; for whilst the globules of Oil surrounded by water become *darker* as the object-glass is *depressed*, and *lighter* as it is *raised*, those of

¹ If this latter mode be adopted, it is preferable, as suggested by the authors of the "Micrographic Dictionary" (Introduction, p. xxxii), to color the oil of turpentine with alkanet, or some similar substance, for its more ready distinction.

Water surrounded by oil become *more luminous* as the object-glass is *depressed*, and *darker* as it is *raised*. The reason of this lies in the fact, that the high refracting power of the oil causes each of its globules to act like a double-convex lens of very short focus; and as this will bring the rays which pass through it into convergence *above* the globule (*i. e.* between the globule and the objective), its brightest image is given, when the object-glass is removed somewhat further from it than the exact focal distance of the object. On the other hand, the globule of water in oil, or the minute bubble of air in water or balsam, acts, in virtue of its inferior refractive power, like a double-concave lens; and as the rays of this diverge from a virtual focus *below* the globule (*i. e.* between the globule and the mirror), the spot of greatest luminosity will be found, by causing the object-glass to approach within the proper focus. Now in the "protoplasm" of the cells of the lower Plants, and in the "sarcode" of the lower animals, oil-particles and *vacuoles* (or void spaces) are often interspersed; and present, at first sight, so very striking a resemblance, that the inexperienced observer may well be pardoned for mistaking the "vacuoles" for larger globules of a material more refractive than the gelatinous substance around them. But the difference in the effects of alterations of focus on the two sets of appearances, at once serves to make evident the difference of their causes; and this, moreover, is made obvious by the effect of oblique light, which will cause the strongest shadow to exhibit itself on opposite sides, in the two cases respectively. It will be obvious that minute elevations and depressions of the *surface* of the object will exert an influence upon the course of the rays which it transmits, very similar to that which proceeds from the presence of globular spaces, filled with transparent substances of greater or less refracting power, in its interior; and that the discrimination between the two may be made by the same means. For if the dots appear more luminous as the object-glass is raised, and darker as it is depressed, they may be interpreted as being due to *convexities* upon the surface; but if the contrary is the case, they may be referred to concavities.

100. Among the sources of fallacy by which the young Microscopist is liable to be misled, one of the most curious is the *Molecular Movement* which is exhibited by the particles of nearly all bodies that are sufficiently finely divided, when suspended in water or other fluids. This movement was first observed in the fine granular particles, which exist in great abundance in the contents of the pollen-grains of plants (sometimes termed the *fovilla*), and which are set free by crushing these grains; and it was imagined that they indicated the possession of some special vital endowment of these particles, analogous to that of the spermatozoa of animals. In the year 1827, however, it was announced by Dr. Robert Brown, that numerous other substances, organic and inorganic, when reduced to a state of equally minute

division, exhibit a like movement, so that it cannot be regarded as indicative of any endowment peculiar to the fovilla-granules; and subsequent researches have shown, that there is no known exception to the rule, that such motion takes place in the particles of *all* substances, though some require to be more finely divided than others, before they will exhibit it. Nothing is better adapted to show it, than a minute portion of gamboge, indigo, or carmine, rubbed up with water; for the particles of these substances, which are not dissolved, but only suspended, are of sufficiently large size to be easily distinguished with a magnifying power of 250 diameters, and are seen to be in perpetual locomotion. Their movement is chiefly of an oscillatory kind; but they also rotate backwards and forwards upon their axes, and they gradually change their places in the field of view. It may be observed that the movement of the smallest particles is the most energetic, and that the largest are quite motionless, whilst those of intermediate size move, but with comparative inertness. The movement is not due (as some have imagined) to evaporation of the liquid; for it continues, without the least abatement of energy, in a drop of aqueous fluid that is completely surrounded by oil, and is therefore cut off from all possibility of evaporization; and it has been known to continue for many years, in a small quantity of fluid enclosed between two glasses in an air-tight case. It is, however, greatly accelerated, and rendered more energetic, by Heat; and this seems to show that it is due, either directly to some caloric changes continually taking place in the fluid, or to some obscure chemical action between the solid particles and the fluid, which is indirectly promoted by heat. It is curious that the closer the conformity between the specific gravity of the solid particles and that of the liquid, the less minute need be that reduction in their size which is a necessary condition of their movement; and it is from this that the substances just named are so favorable for the exhibition of it. On the other hand, the particles of metals, which are from seven to twelve times as heavy as water, require to be reduced to a minuteness many times greater than that of the particles of carmine or gamboge, before they become subject to this curious action. In any case in which the motions of very minute particles, of whatever kind, are in question, it is necessary to make allowance for this "molecular movement;" and the young Microscopist will therefore do well to familiarize himself with its ordinary characters, by the careful observation of it in such cases as those just named, and in any others in which he may meet with it.

101. *Comparative Values of Object-Glasses; Test Objects.*—In estimating the comparative values of different object-glasses, regard must always be had to the *purpose* for which each is designed; since it is impossible to construct a combination, which shall be equally serviceable for *every* requirement. It is commonly assumed, that an Objective which will show certain *test-*

objects, must be very superior, for everything else, to a glass which will not show these; but this is known to every practical Microscopist to be a great mistake,—the very qualities which enable it to resolve the more difficult “tests” being incompatible with those which make it most useful in all the ordinary purposes of scientific investigation. Four distinct attributes have to be specially considered, in judging of the character of an object-glass, viz.: (1) its *defining power*, or power of giving a clear and distinct image of all well-marked features of an object, especially of its boundaries; (2) its *penetrating power*, or power of enabling the observer to *look into* the structure of objects;¹ (3) its *resolving power*, by which it enables closely-approximated marking to be distinguished; and (4) the *flatness of the field* which it gives.

I. The “Defining power” of an objective mainly depends upon the *perfection of its correctness*, both for Spherical and for Chromatic aberration (§§ 9, 15); and it is an attribute essential to the satisfactory performance of *any* objective, whatever be its other qualities. Good definition may be more easily obtained with lenses of *small* or *moderate*, than with lenses of *large* angular aperture; and in the aim to extend the aperture, the perfection of the definition is not unfrequently impaired. An experienced Microscopist will judge of the defining power of a lens by the quality of the image which it gives of almost any object with which he may be familiar; but there are certain “tests,” to be presently described, which are particularly appropriate for the determination of it. Any imperfection in defining power is exaggerated, as already pointed out (§§ 22, 80), by the use of “deep” eye-pieces; so that, in determining the value of an objective, it is by no means sufficient to estimate its performance under a low eye-piece, an image which appears tolerably clear when moderately magnified, being often found exceedingly deficient in sharpness when more highly amplified. The use of the draw-tube (§ 43) affords an additional means of testing the defining power; but this cannot be fairly had recourse to, unless an alteration be made in the adjustment for the thickness of the glass that covers the object (§ 84), in proportion to the lengthening of the body, and the nearer approximation of the object to the objective which this involves.

II. The “Penetrating power” of an object-glass (good definition being of course presupposed) mainly depends upon the

¹ The Author is aware that he is here employing the term “Penetration” in a sense very different from that which it was intended to convey by Dr. Goring, who first applied it to designate a certain quality of Microscopic objectives. But he considers that what was termed “penetration” by Dr. Goring may be far more appropriately designated as *resolving power*; this term having been long in use to express the parallel attribute of Telescopes, as regards the separation of the diffused luminosity of Nebulæ into distinct points of light. The term Penetration, having been thus set free, may well be applied (as above) in what seems its *natural* meaning; and the Author (who has long been in the habit of employing it in this sense) may refer to the Report of the Jury of the “Great Exhibition” of 1851, as giving an authoritative sanction to the above use of it

degree of distinctness with which parts of the object that are *a little out of focus* can be discerned; and this will be found to vary greatly in different objectives, being, within certain limits, in an *inverse* proportion to the extent of the angle of aperture. This is very easily understood on optical principles. The *central* rays of any pencil undergo the *least* refraction or change in their course; the *peripheral* rays, the *most*. The greater the change, the greater is the difference between the amounts of refraction respectively undergone by rays coming off from points at slightly different distances; and the greater, when the focal adjustment is correct for *one* of these points, will be the indistinctness of the image of *the other*. Hence an objective of comparatively limited aperture may enable the observer to gain a view of *the whole* of an object, the several parts of whose structure lie at different distances from it, sufficiently good to afford an adequate idea of the relation of those parts to each other; whilst if the same object be looked at with an objective of very wide angle of aperture, which only enables what is *precisely* in focus to be seen at all, each part can only be *separately* discerned, and the mutual relations of the whole cannot be brought into view. The want of this "penetrating power" is a serious drawback in the performance of many objectives, which are distinguished by the possession of other admirable qualities. The possession of a high measure of it is so essential, in the Author's opinion, to the satisfactory performance of those objectives which are to be employed for the *general purposes of scientific investigation*, that he cannot consider its deficiency to be compensated by the possession of any degree of the resolving power, whose use is comparatively limited.

III. The "Resolving power," by which very minute markings, —whether lines, striæ, or dots,—are discerned and clearly separated from each other, may be said to stand in direct relation (a perfect definition being presupposed) to the extent of its angle of aperture, and consequently to the obliquity of the rays which it can receive from the several points of the surface of the object. This is not so much the case, where the markings depend upon the interposition of opaque or semi-opaque particles in the midst of a transparent substance, so that the lights and shadows of the image represent the absolute degrees of greater or less transparency in its several parts; as it is where, the whole substance being equally transparent, the markings are due to the refracting influence which inequalities of the surface exert upon the course of the rays that pass through it. It may be readily perceived, on a little reflection, that the information given about such inequalities by rays of light transmitted axially through the object, must be very inferior to that which can be gained from rays of light transmitted obliquely; and thus it happens that, as already explained, many such markings are seen by oblique illumination (as, for instance, by the use of the central stop in

the condenser, § 56), which could not be seen, under the same object-glass, by light transmitted more nearly in the axis of the microscope. When an object, however, is seen by *transmitted* light, no degree of obliquity in the illuminating rays can be useful, which exceeds that at which the object-glass can receive them: but the illumination of objects which are seen by *radiated* light (§ 62), depends upon these very rays; and thus it is that the "black-ground" illumination by the paraboloid or by any other effective contrivance (§ 61), will often bring surface-markings into view, which cannot be seen by transmitted light. An object-glass of very wide aperture, however, will receive, even with ordinary illumination, so many rays of great obliquity, that the same kind of effect will be produced, as by oblique illumination with an objective of smaller aperture; but when, with such an objective, oblique illumination is used, a greater resolving power is obtained, than any combination of smaller angular aperture *can* possess. In comparing the resolving power of different object-glasses, it is obviously essential to a correct judgment, that the illumination should be the same; for it will often happen that an observer who knows the "points" of his own instrument, will "bring out" tests, which another, with object-glasses of much greater capability, does not resolve, simply for want of proper management. Moreover, it must be borne in mind that great resolving power may exist, even though the definition may be far from exact; since the former depends more upon angle of aperture, than upon the perfection of the corrections: and yet there cannot be the slightest question, that, of two objectives of the same focal length, one perfectly corrected up to a moderate angle of aperture, the other with the wider aperture but less perfectly corrected, the former will be the one most suitable to the *general* purposes of the Microscopist.

IV. The "Flatness of the field" afforded by the object-glass, is a condition of great importance to the advantageous use of the Microscope; since the real extent of the field of view practically depends upon it. Many objectives are so constructed, that, even with a perfectly flat object, the foci of the central and of the peripheral parts of the field are so different, that when the adjustment is made for one, the other is entirely indistinct. Hence, when the central portion is being looked at, no more information is gained respecting the peripheral, than if it had been altogether "stopped out." With a really good object-glass, not only should the image be distinct even to the margin of the field, but the marginal portion should be as free from chromatic fringes or from indistinctness of outline, as the central portion. In many microscopes of inferior construction, the imperfection of the objectives in this respect, is masked by the contraction of the aperture of the diaphragm in the eye-piece (§ 21), which limits the dimensions of the field; and the performance of one objective within this limit may scarcely be distinguishable from

that of another, although, if the two were compared under an eye-piece of larger aperture, their difference of excellence would be at once made apparent, by the perfect correctness of one to the margin of the field, and by the entire failure of the other in every part save its centre. In estimating the relative merits of two lenses, therefore, as regards this condition, the comparison should of course be made under the same Eye-piece.

V. It may be safely affirmed, that the *most perfect Object-glass* is that which combines all the preceding attributes, in the highest degree in which they are compatible one with another. But, as has just been shown, two of the most important, namely—penetrating power and resolving power,—stand in such opposite relations to the angular aperture, that the highest degree of which each is in itself capable, can only be attained by some sacrifice of the other; and, therefore, of two objectives which are respectively characterized by the predominance of these opposite qualities, one or the other will be preferred by the Microscopist, according to the particular class of researches which he may be carrying on; just as a man who is about to purchase a horse, will be guided in his choice by the kind of work for which he destines the animal. Hence it shows, in the Author's estimation, just as limited an appreciation of the practical applications of the instrument, to estimate the merits of an object-glass by its capability of showing certain lined or dotted "tests," without any reference to its penetrating or defining power; as it would be if a man should estimate the merits of a horse *merely* by the number of seconds within which he could run a mile, or by the number of pounds he could draw; without any reference, in the first case, either to the weight he could carry, or to the length of time during which he could maintain his speed; and in the second case, either to the rate of his draught, or to his power of continuing the exertion. The greatest capacity for *speed* alone, the power of sustaining it not being required, and burden being reduced almost to nothing, is that which is sought in the Racer; the greatest *power of steady draught*, the rate of movement being of comparative little importance, is that which is most valued in the cart-horse; but for the ordinary carriage-horse or roadster, the highest merit lies in such a *combination* of speed and power with endurance, as cannot coexist with the greatest perfection of either the two first. The Author feels it the more important that he should express himself clearly and strongly on this subject, as there is a great tendency at present, both among amateur Microscopists and among Opticians, to look at the attainment of that "resolving power" which is given by angular aperture, as the one thing needful; those other attributes which are of far more importance in almost every kind of scientific investigation, being comparatively little thought of; and he therefore ventures here to repeat the remarks he made upon this subject in his recent Presidential Address to the Microscopical Society, of the

correctness of which he has been since assured, by the approval of many of those who have most successfully employed the Microscope in Physiological investigations. "The superiority in resolving power possessed by object-glasses of large angular aperture, is obtained at the expense of other advantages. For even granting that there is no sacrifice of that most important element *defining* power (which can only be secured, with a very wide angle, by the utmost perfection in all the corrections), yet the adequate performance of such a lens can only be secured by the greatest exactness in the adjustments. Only that portion of the object which is *precisely* in focus, can be seen with an approach to distinctness, everything that is in the least degree out of it being imbedded (so to speak) in a thick fog; it is requisite, too, that the adjustment for the thickness of the glass that covers the object, should exactly neutralize the effect of its refraction; and the arrangement of the mirror and condenser must be such as to give to the object the best possible illumination. If there be any failure in these conditions, the performance of a lens of very wide angular aperture is *very much inferior* to that of a lens of moderate aperture; and except in very experienced hands, this is likely to be generally the case. Now to the working Microscopist, unless he be studying the particular classes of objects which expressly require this condition, it is a source of great inconvenience and loss of time to be obliged to be continually making these adjustments; and a lens, which, when adjusted for a thickness of glass of 1-100", will perform without much sensible deterioration with a thickness either of 1-80" or of 1-120", is practically the best for all ordinary purposes. Moreover, a lens of moderate aperture has this very great advantage, that the parts of the object which are less perfectly in focus, can be much better seen; and therefore that the relation of that which is most distinctly discerned, to all the rest of the object, is rendered far more apparent. Let me remind you, further, that almost all the great achievements of Microscopic research have been made by the instrumentality of such objectives as I am recommending. There can be no question about the large proportion of the results which Continental microscopists may claim, in nearly all departments of minute anatomical, physiological, botanical, or zoological investigation, since the introduction of this invaluable auxiliary; and it is well known that the great majority of their instruments are of extremely simple construction, and that their objectives are generally of very moderate angular aperture. Moreover, if we look at the date of some of the principal contributions which this country has furnished to the common stock, such as the 'Odontography' of Professor Owen, the 'Researches into the Structure of Shell' carried out by Mr. Bowerbank and myself, the 'Physiological Anatomy' of Messrs. Todd and Bowman, the first volume of the 'Histological Catalogue,' by Professor Quekett, and the 'British Desmi-

deæ' of Mr. Ralfs, we find sure reason to conclude that these researches *must* have been made with the instrumentality of lenses, which would in the present day be regarded as of very limited capacity. I hope that, in these remarks, I shall not be understood as in any way desirous to damp the zeal of those, who are applying themselves to the perfectionizing of achromatic objectives. I regard it as a fortunate thing for the progress of science, that there are individuals whose tastes lead them to the adoption of this pursuit; who stimulate our instrument-makers to go on from one range to another, until they have conquered the difficulties which previously baffled them; and then apply themselves to find out some new tests, which shall offer a fresh difficulty to be overcome. But it is not the *only*, nor can I regard it as the *chief* work of the Microscope, to resolve the markings upon the Diatomacææ, or tests of the like difficulty; and although I *should* consider this as the highest object of ambition to our makers, if the performances of such lenses with test-objects were any fair measure of their general utility, yet as I think that I have demonstrated that the very conditions of their construction render them inferior in this respect for the purposes of ordinary microscopic research, I would much rather hold out the reward of high appreciation (*we* have no other to give) to him who should produce the *best working microscope*, adapted to all ordinary requirements, *at the lowest cost*. It does not seem to me an unapt simile, to compare the devotees of large angular apertures to the gentleman of the 'turf.' It is, I believe, generally admitted, that the breeding of a class of horses distinguished by speed and 'blood,' which is kept up by the devotion of a certain class of our countrymen to the noble sport of racing, is an advantage to almost every breed of horses throughout the country; tending, as it does, to develop and maintain a high standard in these particulars. But no one would ever think of using a race-horse for a roadster or a carriage-horse; knowing well that the very qualities which most distinguish him as a racer, are incompatible with his suitability for ordinary work. And so I think that the 'breeders' of first-class Microscopes (if I may so designate them) are doing great service, by showing to what a pitch of perfection certain kinds of excellence may be carried, and by thus improving the standard of ordinary instruments; notwithstanding that, for nearly all working purposes, the latter may be practically superior."

102. *Test Objects*.—It is usual to judge of the optical perfection of a Microscope, by its capacity for exhibiting certain objects, which are regarded as *tests* of the merits of its object-glasses; these tests being of various degrees of difficulty; and that being accounted the best instrument, which shows the most difficult of such tests. Now it must be borne in mind, that only *two* out of the four qualities which have been just enumerated,—namely, *defining* power, and *resolving* power,—can be estimated by any

of these tests; and the greater number of them, being objects whose surface is marked by lines, striæ, or dots, are tests of *resolving* power, and thus of angular aperture only. Hence, as already shown, an objective may show very difficult *test-objects*, and yet may be very unfit for ordinary use. Moreover, these *test-objects* are only suitable to object-glasses of very short focus and high magnifying power; whereas the greater part of the real *work* of the Microscope is done with objectives of comparatively low power; and the enlargement of the angular aperture, which enables even these to resolve (under deep eye-pieces) many objects which were formerly considered adequate tests for higher powers, is by no means an unmixed good. In estimating the value of an object-glass, it should always be considered for what purpose it is intended; and its merits should be judged of according to the degree in which it fulfils that purpose. We shall therefore consider, what are the attributes proper to the several "powers" of object-glasses,—*low*, *medium*, and *high*;—and what are the objects by its mode of exhibiting which, it may be fairly judged.

I. By object-glasses of *low* power, we may understand any whose focal length is *greater than half an inch*. The "powers" usually made in this country are of 2 in. or $1\frac{1}{2}$ in. focus (these being sometimes made to divide, so as to leave a power of about 3 in. focus), 1 in., and 8-10ths or 2-3ds in.; and they give a range of amplification of from 12 to 60 diameters with the shallower eye-piece, and of from 18 to 90 diameters with the deeper. These are the objectives which are most used in the examination of opaque objects, and of transparent objects of large size and of comparatively coarse texture; and the qualities most desirable in them, are a sufficiently large aperture to give a *bright* image, combined with such accurate definition as to give a *clear* image, with penetrating power sufficient to prevent any moderate inequalities of surface from seriously interfering with the distinctness of the entire picture, and with perfect flatness of the image when the object itself is flat. For the 2 in. or $1\frac{1}{2}$ in. objective, no ground of judgment is better, than the manner in which it shows such an "injected" preparation as the interior of a Frog's lung (Fig. 331) or a portion of the villous coat of the Monkey's intestine (Fig. 328); for the aperture ought to be sufficient to give a bright image of such objects, by ordinary daylight, without the use of a condensing-lens; the border of every vessel should be clearly defined, without any thickness or blackness of edge; every part of such an object that comes within the field, should be capable of being made out when the focal adjustment is adapted for any other part; whilst, by making that adjustment a medium one, the whole should be seen without any marked indistinctness. If the aperture be too small, the image will be dark; if it be too large, details are brought into view (such as the separateness of the particles of the vermilion injection) which

it is of no advantage to see, whilst, through the sacrifice of penetration, those parts of the object which are brought exactly into focus being seen with over-minuteness, the remainder are enveloped in a thick fog, through which even their general contour can scarcely be seen to loom; and if the corrections be imperfectly made, no line or edge will be seen with perfect sharpness. For *defining* power, the Author has found the pollen-grains of the Hollyhock, or any other flower of the *Mallow* kind (Fig. 189), viewed as an opaque object, a very good test; the minute spines with which it is beset, being but dimly seen with any save a *good* object-glass of these long foci, and being really well exhibited only by adding such power to the eye-piece, as will exaggerate any want of definition on the part of an inferior lens. For flatness of field, no test is better than a section of Wood (Fig. 165) or a large Echinus-spine (Fig. 237), under an eye-piece that will give a field of the diameter of from 9 to 12 inches. Such objects ought to be very well shown by the divided lens of 2 in. or 3 in. focus; but, as its corrections are rendered imperfect by the removal of the front pair, its defining power is necessarily impaired, and cannot be made even tolerable, save by such a curtailment of the aperture as detracts from the brightness of the image. The general performance of object-glasses of 1 in. and 8-10ths in. focus, may be partly judged of by the manner in which they show such injections as those of the Gill of the Eel (Fig. 330) or of the Bird's Lung (Fig. 332), which require a higher magnifying power for their resolution than those previously named; still better, perhaps, by the mode in which they exhibit a portion of the wing of some Lepidopterous insect, having well-marked scales; the same qualities should here be looked for, as in the case of the lowest powers; and a want of either of them is to be distinguished in a similar manner. The increase of angular aperture which these lenses may advantageously receive, should render them capable of resolving all the easier "test" scales of Lepidoptera, such as those of the *Morpho menelaus* (Fig. 279), in which, with the deeper eye-piece, they should show the transverse as well as the longitudinal markings. The tongue of the common Fly (Fig. 287) is one of the best transparent objects for enabling a practised eye to estimate the general performance of object-glasses of these powers; since it is only under a really good lens, that all the details of its structure can be clearly made out; and an objective which shows *this* well, may be trusted to for any other object of its kind. For flatness of field, sections of small Echinus-spines are very good tests. The exactness of the corrections in lenses of these foci, may be judged of by the examination of objects which are almost sure to exhibit color, if the correction be otherwise than perfect; this is the case, for example, with the glandulæ of Coniferous wood (Fig. 161), the centres of which ought to be clearly defined under such objectives, and ought to be quite free from color; and also with the

tracheæ of Insects (Fig. 291), the spires of which ought to be distinctly separated from each other, without any appearance of intervening chromatic fringes.

II. We may consider as object-glasses of *medium* power, those which range from half to one-fifth of an inch focus; whose magnifying power is from about 100 to 250 diameters under the shallower eye-piece, and from about 150 to 375 diameters with the deeper. These cannot be advantageously employed in the examination of opaque objects, save of such as are of unusual minuteness; but their great value lies in the information they enable us to obtain, regarding the details of organized structures and of living actions, by the examination of properly prepared *transparent* objects by transmitted light. It is to these lenses, that the remarks already made respecting angular aperture (§ 101) especially apply; since it is in them that the greatest difference exists, between the ordinary requirements of the scientific investigator, and the special needs of those who devote themselves to the particular classes of objects for which the greatest resolving power is required. A moderate amount of such power is essential to the value of every objective within the above-named range of foci; thus, even a good half-inch should enable the markings of the larger scales of the *Polyommatus argus* (azure-blue butterfly) to be distinguished, these being of the same kind with those of the Menelaus, but more delicate, and should clearly separate the dots of the small or "battledoor" scales (Fig. 280) of the same insect, which, if unresolved, are seen as coarse longitudinal lines; a good 4-10ths in. should resolve the larger scales of the *Podura* (Fig. 281) without difficulty; and a good 1-4th or 1-5th in. should bring out the markings on the smaller scales of the *Podura*, and should resolve the markings on the *Pleurosigma hippocampus* into longitudinal and transverse lines. Even the 4-10ths (a power for which Messrs. Smith and Beck have attained a deserved celebrity) *may* be made with an angle of aperture sufficiently wide to resolve the objects named as fair tests for the powers above it; and so the 1-4th inch *may*, by the enlargement of its angular aperture to 120° (which has been accomplished by Mr. Ross) be made to exhibit the more difficult Diatomacæ. But it will be found that, in such object-glasses, the difficulty of making the most advantageous use of them, and the loss of penetrating power which necessarily attends the excessive extension of their angular aperture, are most serious drawbacks to their practical utility in the hands of the Anatomical or Physiological investigator; for whose purposes, such a resolving power as will show the easier tests first enumerated, combined with perfect definition, with a fair amount of penetrating power, and with flatness of field, constitute the best combination. For defining power, very good tests are found in the complex hairs of many animals, such as the Indian *Bat* (Fig. 310, c), and the *Dermestes* (Fig. 282, b). And for that combination of the seven-

ral attributes which the Author thinks most important, he has found no test more valuable and positive, as regards objectives of from 4-10ths to 1-5th inch focus, than Mr. Lealand's preparations of Muscular fibre (Fig. 326). In every case, the objective should be tested with the deeper, as well as with the shallower eye-piece; and the effect of this will be a fair test of its merits. Where markings are indistinguishable under a certain objective, merely because of their minuteness or their too close approximation, they may be enlarged or separated by a deeper eye-piece, provided that the objective be well corrected. But if, in such a case, the image be darkened or blurred, so as to be rather deteriorated than improved, it may be concluded that the objective is of inferior quality, having either an insufficient angular aperture, or being imperfectly corrected, or both.

III. All object-glasses of less than 1-5th of an inch focus, may be classed as *high* powers; the focal lengths to which they are ordinarily constructed are 1-6th, 1-8th, 1-12th, and 1-16th of an inch respectively; and the magnifying powers they are fitted to afford, range from about 320 to 850 diameters with the shallower eye-piece, and from 480 to 1300 diameters with the deeper. By the use of still deeper eye-pieces, a power of 2000 or more may be easily obtained; but nothing seems to be really gained by such high amplification. Moreover, as the 1-12th inch object-glass may have its angular aperture extended to the utmost limits compatible with the reception of rays from any object, it does not seem that anything can be gained by a reduction of the focal distance to the 1-16th inch; and the latter being a more difficult combination, as well to construct as to use, both Opticians and Microscopists have of late years found it advantageous to limit themselves to the 1-12th, which gives an amplification of about 650 diameters with the shallower eye-piece, and of about 1000 with the deeper. The use of this class of objectives is much more restricted than that of the preceding. They are not employed for the ordinary purposes of scientific investigation; and their value chiefly lies in the power which they afford, of tracing out certain points of minute structure, which the objectives of medium power may only doubtfully indicate, and of exhibiting certain classes of very difficult striated or dotted objects, which they cannot resolve. Hence it is obvious that, with regard to object-glasses of *this* class, "resolving power" (coupled with "defining power") is the highest requisite, "penetration" and "flatness of field" being of secondary account; and that the value of an objective may *here* be fairly estimated by its angular aperture, provided that its aberrations be exactly corrected. Of angular aperture and definition, very good tests are afforded by the lines artificially ruled by M. Nobert, and by the more "difficult" species of Diatomaceæ. What is known as "Nobert's Test" is a plate of glass, on a small space of which, not exceeding a fiftieth of an inch in breadth, are ruled ten or more series of lines, form-

ing as many separate bands of equal breadth; in each of these bands, the lines are ruled at a certain known distance; and the distances are so adjusted in the successive bands, as to form a regularly diminishing series, and thus to present a succession of tests of progressively increasing difficulty. The distances of the lines differ on different plates; all the bands in some series being resolvable under a good objective of 1-4th inch focus, whilst the closest bands in others defy the resolving power of a 1-12th inch objective of large aperture. Thus a "test-plate" whose *widest* lines are at a distance from each other of 1-1000th of a Paris line, or of 1-11,200th of an English inch, and whose *closest* lines are at 1-5000th of a line, or 1-56,000th of an inch, from each other, will serve as a very fair test for the angular aperture and defining power of object-glasses below 1-4th inch focus; the superiority of each in these particulars, being judged of by the number of bands which it will resolve into well-defined lines, and by the sharpness and clearness of these lines; while the performance of a 1-4th inch objective may be accounted very satisfactory, if it will enable them all to be clearly distinguished. But if the first of the bands should have an interval of only 1-4000th of a Paris line, or 1-45,000th of an English inch, between its lines, and the last should have its lines approximated to 1-10,000th of a Paris line, or 1-112,000th of an English inch, then only a few of the easier bands will be resolved by the 1-4th inch, a few more by the 1-8th inch, and even the 1-12th inch will probably not enable any band to be distinctly resolved, whose lines are closer than 1-7000th of a Paris line, or 1-79,000th of an English inch. At present, therefore, the existence of separate lines of a narrower interval than this, is a matter of *faith* rather than of *sight*; but there can be no reasonable doubt that the lines do exist; and the resolution of them would evince the extraordinary superiority of any objective, or of any system of illumination, which should enable them to be distinguished. The mathematical certainty with which the degree of approximation of these lines may be ascertained, and the gradation of the series which they present, gives to M. Nobert's test-plate a very high value for the determination of the relative merits of different objectives, of that class, at least, in which angular aperture and definition are of the first importance; whilst it also serves to test the degree in which these capabilities are possessed by object-glasses of medium power, in which other attributes also have to be considered. The value of the minuter *Diatomaceæ*, as furnishing, in their surface-markings, admirable test-objects for the highest powers of the Microscope, was first made known by Messrs. Harrison and Sollitt, of Hull, in 1841; and it cannot be questioned that this discovery has largely contributed to the success of the endeavors which have since been so effectually made, to perfect this class of objectives, and to find out new methods of using them to the best advantage. The nature of these markings will be described

hereafter; and it will be sufficient in this place to give a table of the average distances of the lineation of different species,¹ which will serve to indicate their respective degrees of difficulty as "tests." The greater part of those which are now in use for this purpose, are comprehended in the genus *Pleurosigma* of Prof. W. Smith, which includes those *Naviculæ* whose "frustules" are distinguished by their sigmoid (S-like) curvature (§ 184).

	Lines in 1-1000th of an inch.
1. <i>Pleurosigma littorale</i> ,	24
2. <i>Pleurosigma Hippocampus</i> ,	30 long., 40 trans.
3. <i>Pleurosigma strigile</i> ,	36
4. <i>Pleurosigma strigosum</i> ,	44
5. <i>Pleurosigma elongatum</i> ,	48
6. <i>Pleurosigma angulatum</i> ,	52
7. <i>Pleurosigma Spenceri</i> ,	55 long., 50 trans.
8. <i>Pleurosigma fasciola</i> ,	64
9. <i>Pleurosigma obscurum</i> ,	75
10. <i>Pleurosigma macrum</i> ,	85
11. <i>Nitzschia sigmoidea</i> ,	85
12. <i>Navicula rhomboides</i> ,	85

The first seven of the foregoing may be resolved, with judicious management, by good 1-4th or 1-5th in. objectives; the remainder require the 1-8th or 1-12th in., for the satisfactory exhibition of their markings. Several very difficult tests of this description have been furnished by Prof. Bailey of West Point (U. S.), among them the very beautiful *Grammatophora subtilissima* and the *Hyalodiscus subtilis*; the latter, being of discoid form, and having markings which radiate in all directions, very much like those of an engine-turned watch, is a useful test for observers who have not facilities for obtaining oblique light in any direction; since, whatever may be the azimuth from which the oblique pencil may proceed, some portion of the disk will always

¹ This table is taken from Prof. W. Smith's admirable Monograph on the Diatomaceæ; and it includes most of the species usually employed as tests. These should always be mounted between two pieces of thin glass, according to the method hereafter to be described (§ 122), in order to avoid, as much as possible, the production of aberrations in the illuminating pencil. The number of lineations must be considered as an average, the extremes sometimes varying to a considerable amount on either side. A much higher estimate is given by Messrs. Harrison and Sollitt in the "Quart. Journ. of Microsc. Science," vol. ii, p. 62; the *Pleurosigma fasciola* being reckoned by them to contain 90 lines in 1-1000th of an inch, the *Nitzschia sigmoidea* 100 lines, and a species cited as *Navicula arcus* (which can scarcely be the one so named by Ehrenberg, and termed by Prof. W. Smith *Eunotia arcus*) no less than 130. The last they speak of as "so extremely difficult, that, in order even to catch a glimpse of its delicate markings, the observer must be in possession of glasses of a very large angle of aperture and the finest definition, have the most careful management of oblique light, and in addition be possessed of a large share of patience." The Author cannot but believe that there is some error in these measurements; since, as the well-defined lines upon Nobert's test-plate have not yet been resolved, when they have approximated more closely than the highest numbers mentioned in Prof. W. Smith's table, it can scarcely be imagined possible that the delicate markings of a *Navicula* should even be "glimpsed," if they be as much closer than those of the species previously accounted most difficult, as those of the latter are than those of the easiest.

be in the best possible position in regard to the light, whereas, in the case of other finely-lined tests, it is only when the most favorable position has been attained, perhaps after tedious and troublesome trials, that the markings are displayed.¹

103. *Determination of Magnifying Power.*—The last subject to be here adverted to, is the mode of estimating the magnifying power of Microscopes, or, in other words, the number of times that any object is magnified. This will of course depend upon a comparison of the *real* size of the object, with the *apparent* size of the image; but our estimate of the latter will depend upon the distance at which we assume it to be seen, since, if it be projected at different distances from the eye, it will present very different dimensions. Opticians generally, however, have agreed to consider *ten inches* as the standard of comparison; and when, therefore, an object is said to be magnified 100 diameters, it is meant that its visual image, projected at 10 inches from the eye (as when thrown down by the Camera Lucida, § 49) upon a surface at that distance beneath, has 100 times the actual dimensions of the object. The measurement of the magnifying power of Simple or Compound Microscopes by this standard is attended with no difficulty. All that is required is a stage-micrometer accurately divided to a small fraction of an inch (the 1-100th will answer very well for low powers, the 1-1000th for high), and a common foot-rule divided to tenths of an inch. The micrometer being adjusted to the focus of the objective, the rule is held parallel with it, at the distance of ten inches from the eye. If the second eye be then opened, whilst the other is looking at the object, the circle of light included within the field of view, and the object itself, will be seen faintly projected upon the rule; and it will be very easy to mark upon the latter the apparent distances of the divisions on the micrometer, and thence to ascertain the magnifying power. Thus, supposing each of the divisions of 1-100th of an inch to correspond with $1\frac{1}{2}$ inch upon the rule, the linear magnifying power is 150 diameters; if it correspond with half an inch, the magnifying power would be 50 diameters. If, again, each of the divisions of the 1-1000th inch micrometer correspond to 6-10ths of an inch upon the rule, the magnifying power is 600 diameters; and if it correspond to $1\frac{2}{3}$ inch, the magnifying power is 1200 diameters. In this mode of measurement, the estimate of *parts* of tenths on the rule can only be made by guess; but greater accuracy may be obtained by projecting the micrometer-scale with the Camera Lucida at the distance of ten inches from the eye, marking the intervals on paper, taking an average of those, and repeating this with the compasses ten times along the inch-scale. Thus, if the space given by one of the divisions of the 1-1000th-inch micrometer,

¹ See Prof. Bailey's interesting memoirs in Vols. II and VII of the "Smithsonian Contributions to Knowledge."

repeated ten times along the rule, gave 6 inches and $2\frac{1}{2}$ tenths, the value of each division would be $\cdot 625$ of an inch, and the magnifying power 625. The *superficial* magnifying power is of course estimated by *squaring* the linear; but this is a mode of statement never adopted by scientific observers, although often employed to excite popular admiration, or to attract customers, by those whose interest is concerned in doing so.¹

¹ An ingenious method has been devised by Prof. Harting, of Utrecht, for determining "the utmost limits of penetrating and separating power possessed by a Microscope," by using as test-objects the very reduced images of various bodies formed by air-bubbles in gum-mucilage. The mode of obtaining and employing these images for the above purpose, will be found in the "Quarterly Journal of Microscopical Science," vol. i, p. 292.

CHAPTER V.

PREPARATION, MOUNTING, AND COLLECTION OF OBJECTS.

UNDER this head it is intended to give such *general* directions respecting the preparation, mounting, and collection of Objects, as will supersede the necessity of frequent repetition when each particular class is described ; and also to enumerate the materials and appliances, which will be required or found advantageous.

SECTION 1. PREPARATION OF OBJECTS.

104. *Microscopic Dissection.*—The separation of the different parts of an Animal or Vegetable structure by dissection, so as to prepare any portion for being minutely examined under the Microscope should be accomplished, so far as may be found practicable, with the naked eye ; but the best mode of doing this, will depend in great degree upon the size and character of the object. Generally speaking, it will be found advantageous to carry on the dissection under water, with which alcohol should be mingled where the substance has been long immersed in spirit. The size and depth of the vessel should be proportioned to the dimensions of the object to be dissected ; since, for the ready access of the hands and dissecting instruments, it is convenient that the object should neither be far from its walls, nor lie under any great depth of water. Where there is no occasion that the bottom of the vessel should be transparent, no kind of dissecting-trough is more convenient, than that which every one may readily make for himself, of any dimensions he may desire, by taking a piece of sheet gutta percha of adequate size and stoutness, warming it sufficiently to render it flexible, and then turning up its four sides, drawing out each corner into a sort of spout, which serves to pour away its contents when it needs emptying. The dark color of this substance enables it to furnish a background, which assists the observer in distinguishing delicate membranes, fibres, &c., especially when magnifying lenses are employed ; and it is hard enough, without being too hard, to allow of pins being fixed into it, both for securing the object, and for keeping apart such portions as it is useful to put on the stretch. When glass or earthenware troughs are employed, a piece of sheet-cork, loaded with lead, must be provided, to an-

swer the same purposes. In carrying on dissections in such a trough, it is frequently desirable to concentrate additional light upon the part which is being operated on, by means of the smaller condensing lens (Fig. 45); and when magnifying power is wanted, it may be supplied either by a single lens, mounted after the manner of Ross's Simple Microscope (Fig. 14, B), or by a Compound body mounted as in one of Mr. Warington's arrangements (Fig. 24). Portions of the body under dissection, being floated off when detached, may be conveniently taken up from the trough by placing a slip of glass beneath them (which is often the only mode in which delicate membranes can be satisfactorily spread out); and may be then placed under the microscope for minute examination, being first covered with thin glass, beneath the edges of which is to be introduced a little of the liquid wherein the dissection is being carried on. Where the body under dissection is so transparent, that more advantage is gained by transmitting light through it, than by looking at it as an opaque object, the trough should have a glass bottom; and for this purpose, unless the body be of unusual size, some of the glass "cells" to be hereafter described (§§ 136, 137) will usually answer very well. The finest dissections may often be best made upon ordinary slips of glass; care being taken to keep the object sufficiently surrounded by fluid. For work of this kind, no simple instrument is more generally serviceable than Mr. Quekett's Dissecting Microscope (Fig. 17); but if higher magnifying powers be needed than this will conveniently afford, recourse may be had to Smith and Beck's Dissecting Microscope (Fig. 29), which for this purpose should always be furnished with the Erector (Fig. 32). A particular arrangement of the light, devised many years since by the Author, will enable an expert dissector to prosecute his work with the naked eye, to an extent for which a lens would otherwise be required. This consists in giving to the object the same kind of black-ground illumination, as is now in common use for a very different purpose; and nothing more is necessary to afford it, than to attach to the under side of the stage a sort of "well," composed of a tube blackened in its interior, about $1\frac{1}{2}$ inch long, of the same diameter as the opening of the stage-plate, into the lower extremity of which a diaphragm or a ground-glass may be fitted, for the purpose of diminishing or of softening the light. The slide being laid upon the stage, and the mirror being so turned as to illuminate the object, the eye is to be so placed (the arm carrying the magnifiers being turned to one side) that the object is seen against the dark background afforded by the side of the well. In this manner, fibres of extreme minuteness, or other particles of extraordinary delicacy, can be clearly distinguished, such as could otherwise be scarcely discerned at all without the assistance of a magnifier. And the further the dissection can be carried in this mode, the less difficulty will be found in completing it, when the simple or

compound Microscope is brought to bear upon it. Whenever a dissection is being made upon the stage of a microscope, it is desirable that support should be provided for the hands on either side. This may be given by books or blocks of wood piled up to the requisite height; but in place of *flat* "rests," it is much more convenient to provide a pair of *inclined planes*, sloping away from the stage at an angle of about 30° below the horizon. These may be either solid blocks of wood, or (which is much less cumbrous) they may be made of two boards hinged together, one giving the inclined plane, which rests at one end upon the table, while the other, standing vertically, affords the requisite elevation to the extremity which abuts against the stage.

105. The instruments used in Microscopic dissection, are for the most part of the same kind as those which are needed in ordinary minute Anatomical research, such as scalpels, scissors, forceps, &c.; the fine instruments used in operations upon the eye, however, will commonly be found most suitable. A pair of delicate scissors curved to one side, is extremely convenient for cutting open tubular parts; these should have their points blunted; but other scissors should have fine points. A pair of fine-pointed scissors (Fig. 56), one leg of which is fixed in a handle, and the other kept

FIG. 56.



Spring-Scissors.

apart from it by a spring, so as to close by the pressure of the finger and to open of itself, will be found (if the blades be well sharpened on a hone) much superior to any kind of knives, for cutting through delicate tissues with as little disturbance of them as possible: Swammerdam is said to have made great use of this instrument in his elaborate insect-dissections. Another cutting instrument much used by some dissectors, may be designated as a miniature of the shears used in shearing sheep, or as a cutting-forceps; the blades of such an instrument may be prevented from springing too far asunder, by means of a regulating-screw (as in the "microtome" of M. Strauss-Durckheim) or by some other kind of check; and the cutting action, being executed by the opposed pressure of the finger and thumb, may be performed with great precision. A pair of small straight forceps, with fine points, and another pair of curved forceps, will be found useful in addition to the ordinary dissecting-forceps. Of all the instruments contrived for delicate dissections, however, none are more serviceable than those which the Microscopist may make for himself out of ordinary needles. These should be fixed in light wooden handles¹ (the cedar sticks used for

¹ Special *needle-holders* (like miniature port-crayons) have been made for this purpose; and although they afford the facility of lengthening or shortening the acting point of the needle at will, and also of carrying a reserve store of needles at the other end, yet the Author would decidedly recommend the use of the wooden handles, of which a large stock may be obtained for a trifle.

camel-hair pencils, or the handles of steel-pen-holders, will answer extremely well), in such a manner that their points should not project far,¹ since they will otherwise have too much "spring:" much may be done by their mere *tearing* action; but if it be desired to use them as *cutting* instruments, all that is necessary is to give them an edge upon a hone. It will sometimes be desirable to give a finer point to such needles, than they originally possess; this also may be done upon a hone. A needle with its point bent to a right angle, or nearly so, is often useful; and this may be shaped by simply heating the point in a lamp or candle, giving to it the required turn with a pair of pliers, and then hardening the point again by reheating it and plunging it into cold water or tallow.

106. *Cutting Sections of Soft Substances.*—Most important information respecting the structure of many substances, both Animal and Vegetable, may be obtained by cutting sections of

FIG. 57.



Curved Scissors for cutting Thin Sections.

them, thin enough to be viewed as transparent objects. Where the substances are soft, no other instrument is necessary for this purpose, than a sharp knife, which may be best made with a thin two-edged blade like that of a lancet; considerable practice is needed, however, to make effectual use of it; and some individuals acquire a degree of dexterity, which others never succeed in attaining. In cutting sections of Animal tissues, which, owing to the quantity of water they contain, do not present a sufficiently firm resistance, it is often desirable to half-dry these, by exposing small pieces freely to the air, with the aid of a gentle warmth if required; when this desiccating process has been carried sufficiently far, thinner sections can be cut, than could possibly have been made in the original state of the tissue; and the texture, after a short maceration in water, almost entirely recovers its pristine characters. There are certain tissues, however, which will not bear to be thus treated, and of which it is sufficient to examine an extremely minute portion; and for making sections of these, such a pair of scissors as is represented in Fig. 57 will often be found very useful; since, owing to the curvature of the blades,² the two extremities of a

¹ The following is the mode in which the Author has found it convenient to mount his needles for this and other purposes:—The needle being held firmly in a pair of pliers grasped by the right hand, its point may be forced into the end of a cedar or other stick held in the left, until it has entered to the depth of half an inch or more; the needle is then cut off to the desired length (the eye end being thus got rid of); and being then drawn out, the truncated end is forced into the hole previously made by the point, until it cannot be made to penetrate further, when it will be found to be very securely fixed. The end of the handle which embraces it, may then be bevelled away round its point of insertion.

² It is difficult to convey by a drawing the idea of the real curvature of this instrument, the blades of which, when it is held in front view, curve—not to either side—

section taken from a flat surface will generally be found to thin away, although the middle of it may be too thick to exhibit any structure. Where only a moderate degree of thinness is required, either in consequence of the transparency of the tissue, or because it is not desired to exhibit its minutest details, the two-bladed knife contrived by Prof. Valentin (Fig. 58) may be employed with advantage. The blades are attached to each other at their lower end by a screw, in such a manner that their "spring" tends to keep them apart; and their distance is regulated by pushing the little rivet backwards or forwards in the

FIG. 58.



Valentin's Knife.


slit through which it works. The knife should be dipped in water before using, or, still better, the section should be made under water, as the instrument works much better when wet; after use, it should be carefully washed and dried, a piece of soft leather being passed between the blades. If any water have found its way into the part through which the rivet works, the movable blade should be detached by taking out its screw, and each blade should be cleaned separately.¹

107. *Cutting Sections of Harder Substances.*—There is a large class of substances, both Animal and Vegetable, which are too hard to admit of sections being made in the manner just described, but of which extremely thin slices can be made by a sharp cutting instrument, if only they be properly held and supported, more especially when the thickness of the section can be regulated by a mechanical contrivance; such are, in particular, the Stems and Roots of Plants, and the Horns, Hoofs, Cartilages, and similarly firm structures of Animals. Various costly machines have been devised for this purpose, some of them characterized by great ingenuity of contrivance and beauty of workmanship; but every purpose to which these are adapted, will be found to be answered by a very simple and unexpensive little instrument, which may either be held in the hand, or (which is preferable) may be firmly attached by means of a T-shaped piece of wood (as in Fig. 59), to the end of a table or work-bench.

but towards the observer; these scissors being, as the French instrument-makers say, *courbés sur le plat*. As an example of the utility of such an instrument to the Microscopist, the Author may cite the curious demonstration given a few years since, by Dr. Aug. Waller, of the structure of the gustative papillæ, by snipping off the papillæ from the living human tongue, which may be done with no more pain than the prick of a pin would occasion.

¹ An improved form of this instrument is constructed by Mr. Matthews of Portugal Street; the blades being made with a convex instead of a straight edge, their distance from each other being regulated by a milled-head screw, and their separation for cleaning being more easily accomplished.

This instrument essentially consists of an upright hollow cylinder



Section Instrument.

of brass, with a kind of piston which is pushed from below upwards by a fine-threaded screw, turned by a large milled head; at the upper end, the cylinder terminates in a brass table, which is made to present a perfectly flat surface. At one side is seen a small milled head, which acts upon a "binding-screw," whose extremity projects into the cavity of the cylinder, and serves to compress and steady anything that it holds. A cylindrical stem of wood, a piece of horn, whalebone, cartilage, &c., is to be fitted to the

interior of the cylinder, so as to project a little above its top, and is to be steadied by the "binding-screw;" it is then to be cut to a level by means of a sharp knife or razor, laid flat upon the table. The milled head is next to be moved through such a portion of a turn, as may very slightly elevate the substance to be cut, so as to make it project in an almost insensible degree above the table; and this projecting part is to be sliced off with a knife, previously dipped in water. The best knife for this purpose is a razor, ground flat (instead of concave) on one side, but having still a concave surface on the other; the flat side is to be laid downwards upon the table; and the motion given to the edge should be a combination of *drawing* and *pressing*. (It will be generally found that better sections are made, by working the knife *from* the operator, than *towards* him.) When one slice has been thus taken off, it should be removed from the blade by dipping it into water, or by the use of a camel-hair brush; the milled head should be again advanced, and another section taken; and so on. Different substances will be found both to *bear* and to *require* different degrees of thickness; and the amount that suits each can only be found by trial. It is advantageous to have the large milled head graduated, and furnished with a fixed index; so that this amount having been once determined, the screw shall be so turned as to always produce the exact elevation required. Where the substance of which it is desired to obtain sections by this instrument, is of too small a size or of too soft a texture to be held firmly in the manner just described, it may be placed between the two vertical halves of a cork of suitable size to be pressed into the cylinder; and the cork, with the object it grasps, is then to be sliced in the manner already described, the small section of the latter being carefully taken off the knife, or floated away from it, on each occasion, to prevent it from being

lost among the lamellæ of cork which are removed at the same time. The special methods of preparation which are required in the case of the various substances, of which sections may be conveniently cut by this instrument, will be described under their several heads.

108. *Grinding and Polishing of Sections.*—Substances which are too hard to be sliced with a cutting instrument in the manner last described,—such as bones, teeth, shells, corals, fossils of all kinds, and even some recent vegetable tissues,—can only be reduced to the requisite thinness for Microscopical examination, by grinding down thick sections, until they become so thin as to be transparent. The general method of making such preparations will be here described;¹ but those special details of management which particular substances may require, will be given when these substances are respectively described. The first thing to be done, will usually be to procure a *section* of the substance, as thin as it can be safely cut. Most substances not siliceous may be divided by the fine saws used by artisans for cutting brass; but there are some bodies (such as the enamel of teeth, and porcellanous shells), which, though merely calcareous, have their mineral particles arranged in such a peculiar state of aggregation, as to make it very difficult and tedious to divide them in this mode; and it is much the quicker operation to *split* them with a disk of soft iron (resembling that used by the lapidary) charged at its edge with diamond-dust, which may be driven in an ordinary lathe. Where waste of material is of no account, a very expeditious method of obtaining pieces fit to grind down, is to detach them from the mass with a strong pair of “cutting-pincers,” or, if it be of small dimensions, with “cutting-pliers;” and a flat surface must then be given to it, either by holding it to the side of an ordinary grindstone, or by rubbing it on a plate of lead (cast or planed to a perfect level) charged with emery, or by a strong toothed file, the former being the most suitable for the *hardest* substances, the latter for the *toughest*. There are certain substances, especially calcareous fossils of wood, bone, and teeth, in which the greatest care is required in the performance of these preliminary operations, on account of their extreme friability; the vibration produced by the working of the saw or the file, or by grinding on a rough surface, being sufficient to disintegrate even a thick mass, so that it falls to pieces under the hand; such specimens, therefore, it is requisite to treat with great caution, dividing them by the smooth action of the wheel, and then rubbing them down upon nothing rougher than a very fine “grit.” Where (as often happens) such specimens are sufficiently porous to admit of the penetration of Canada balsam, it will be desirable, after soaking them in tur-

¹ The following directions do not apply to *Siliceous* substances; as sections of these can only be prepared by those who possess a regular Lapidary's apparatus, and who have been specially instructed in the use of it.

pentine for a while, to lay some liquid balsam upon the parts through which the section is to pass, and then to place the specimen before the fire or in an oven for some little time, so as first to cause the balsam to run in, and then to harden it; by this means the specimen will be rendered much more fit for the processes it has afterwards to undergo. It not unfrequently happens, that the small size, awkward shape, or extreme hardness of the body, occasions a difficulty in holding it either for cutting or grinding; in such a case, it is much better to attach it to the glass in the first instance, by any side that happens to be flattest; and then to rub it down by means of the "hold" of the glass upon it, until the projecting portion has been brought to a plane, and has been prepared for permanent attachment to the glass. This is the method which is generally most convenient to pursue with regard to small bodies; and there are many which can scarcely be treated in any other way, than by attaching a number of them to the glass at once, in such a manner as to make them mutually support one another.¹

109. The mode in which the operation is then to be proceeded with, depends upon whether the section is to be ultimately set up in Canada balsam (§ 125), or is to be mounted dry (§ 122), or in fluid (§ 132). In the former case, the following is the plan to be pursued. The flattened surface is to be polished, by rubbing it with water on a "Water-of-Ayr" stone, on a hone or "Turkey" stone, or on a new stone recently introduced under the name of the "Arkansas" stone; the first of the three is the best for all ordinary purposes; but the two latter, being much harder, may be employed for substances which resist it.² When this has been sufficiently accomplished, the section is to be attached with Canada balsam to a slip of thick, well-annealed glass; and, as the success of the final result will often depend upon the completeness of its adhesion to this, the means of most effectually securing that adhesion will now be described in detail. Some Canada balsam, previously rendered somewhat stiff by the evaporation of part of its turpentine, is to be melted on the glass slip, so as to form a thick drop, covering a space somewhat larger than the area of the section; and it should then be set aside to cool, during which process, the bubbles that may have formed in it will usually burst. When cold, its hardness should be tested, which is best done by the edge of the thumb-nail; for it should be with difficulty indented by its pressure, and yet should not be

¹ Thus, in making horizontal and vertical sections of Foraminifera, as it would be impossible to cut them through, they must be laid close together in a bed of hardened Canada balsam on a slip of glass, in such positions, that, when rubbed down, the plane of section shall traverse them in the desired directions; and one flat surface having been thus obtained for each, this must be turned downwards, and the other side ground away.

² As the flatness of the polished surface is a matter of the first importance, that of the stones themselves should be tested from time to time; and whenever they are found to have been rubbed down on any one part more than on another, they should be flattened on a paving-stone with fine sand, or on the lead-plate with emery.

so resinous as to be brittle. If it be too soft, as indicated by its too ready yielding to the thumb nail, it should be boiled a little more; if too hard, which will be shown by its chipping, it should be re-melted and diluted with more fluid balsam, and then set aside to cool as before. When it is found to be of the right consistence, the section should be laid upon its surface, with the polished side downwards; the slip of glass is next to be gradually warmed until the balsam is softened, special care being taken to avoid the formation of bubbles; and the section is then to be gently pressed down upon the liquefied balsam, the pressure being at first applied rather on one side than over its whole area, so as to drive the superfluous balsam in a sort of wave towards the other side, and an equable pressure being finally made over the whole. If this be carefully done, even a very large section may be attached to glass, without the intervention of any air-bubbles; if, however, they should present themselves, and they cannot be expelled by increasing the pressure over the part beneath which they are, or by slightly shifting the section from side to side, it is better to take the section entirely off, to melt a little fresh balsam upon the glass, and then to lay the section upon it as before.

110. When the section has been thus secured to the glass, and the attached part thoroughly saturated (if it be porous) with hard Canada balsam, it may be readily reduced in thickness, either by grinding or filing as before, or, if the thickness be excessive, by taking off the chief part of it at once by the slitting-wheel. So soon, however, as it approaches the thinness of a piece of ordinary card, it should be rubbed down with water on one of the smooth stones previously named, the glass slip being held beneath the fingers with its face downwards, and the pressure being applied with such equality, that the thickness of the section shall be (as nearly as can be discerned) equal over its entire surface. As soon as it begins to be translucent, it should be placed under the Microscope (particular regard being had to the precaution specified in § 86), and note taken of any inequality; and then, when it is again laid upon the stone, such inequality may be brought down, by making special pressure with the fore-finger upon the part of the slide above it. When the thinness of the section is such as to cause the water to spread around it between the glass and the stone, an excess of thickness on either side may often be detected, by noticing the smaller distance to which the liquid extends. In proportion as the substance attached to the glass is ground away, the superfluous balsam which may have exuded around it will be brought into contact with the stone; and this should be removed with a knife, care being taken, however, that a margin be still left round the edge of the section. As the section approaches the degree of thinness which is most suitable for the display of its organization, great care must be taken that the grinding process be not carried too far; and fre-

quent recourse should be had to the Microscope, which it is convenient to have always at hand, when work of this kind is being carried on. There are many substances whose intimate structure can only be displayed in its highest perfection, when a very little more reduction would destroy the section altogether; and every Microscopist who has occupied himself in making such preparations, can tell of the number which he has sacrificed in order to attain this perfection. Hence if the amount of material be limited, it is a good rule to stop short as soon as a *good* section has been made, and to lay it aside—"letting well alone"—whilst the attempt is being made to procure a *better* one; if this should fail, another attempt may be made, and so on, until either success has been attained, or the whole of the material has been consumed,—the *first* section, however, still remaining: whereas, if the first, like every successive section, be sacrificed in the attempt to obtain perfection, no trace will be left to "show what has been." In judging of the appearance of sections in this stage under the Microscope, it is to be remembered that its transparency will subsequently be considerably increased by mounting in Canada balsam (§ 125); this is particularly the case with fossils, to which a deep hue has been given by the infiltration of some coloring matter; and with any substances whose particles have a molecular aggregation, that is rather amorphous than crystalline. When a sufficient thinness has been attained, the section may generally be "mounted" in Canada balsam; and the mode in which this must be managed, will be detailed hereafter (§ 129).

111. As there are certain substances, however, the view of whose structure is impaired by mounting in Canada balsam, and which should therefore be mounted either dry or in fluid, a different method of procedure must be adopted with them. If tolerably thin sections of them can be cut in the first instance, or if they are of a size and shape to be held in the hand whilst they are being roughly ground down, there will be no occasion to attach them to glass at all; it is frequently convenient to do this at first, however, for the purpose of obtaining a "hold" upon the specimen; but the surface which has been thus attached, must afterwards be completely rubbed away, in order to bring into view a stratum which the Canada balsam shall not have penetrated. As none but substances possessing considerable toughness, such as bones and teeth, can be treated in this manner, and as these are the substances which are most quickly reduced by a coarse file, and are least liable to be injured by its action, it will be generally found possible to bring the sections to a considerable thinness, by laying them upon a piece of cork or soft wood held in a vice, and operating upon them first with a coarser and then with a finer file. When this cannot safely be carried further, the section must be rubbed down upon that one of the fine stones already mentioned (§ 109), which is found best to suit it; as long as the section is tolerably thick, the finger may be used to press and move it: but as soon as the finger

itself begins to come into contact with the stone, it must be guarded by a flat slice of cork, or by a piece of gutta percha, a little larger than the object. Under either of these, the section may be rubbed down until it has been reduced to the requisite degree of tenuity; but even the most careful working, on the finest-grained stone, will leave its surface covered with scratches, which not only detract from its appearance, but prevent the details of its internal structure from being as readily made out, as they can be in a polished section. This polish may be imparted, by rubbing the section with putty powder (peroxide of tin) and water, upon a leather strap, made by covering the surface of a board with buff-leather, having three or four thicknesses of cloth, flannel, or soft leather beneath it; this operation must be performed on both sides of the section, until all the marks of the scratches left by the stone shall have been rubbed out; when the specimen will be fit for mounting, after having been carefully cleansed from any adhering particles of putty powder.

112. *Chemical Actions.*—One important part of the preparation of Microscopic objects, is often effected by the use of Chemical Reagents. These may be employed, either for the sake of *removing* substances of which it is desired to get rid, in order to bring something else into view; or for the sake of *detecting* the presence of particular substances in the object under examination. Thus, in order to obtain the animal basis of Shell, Bone, Tooth, &c., it is necessary to dissolve away the calcareous portion of these tissues by the use of acids; a mixture of nitric and muriatic acids is preferable; and this should be added, little by little, to a considerable bulk of water, until a disengagement of gas be perceived to commence from the surface of the specimen. Care should always be taken not to hurry the process by adding too much acid, since, when the animal membrane is of very delicate consistence, it is liable to be dissolved; and in some cases it is better to allow the action to go on for many weeks, adding only a drop or two of acid at a time. When siliceous particles are to be removed (such as those which form the *loricae* of the Diatomaceæ), for the sake of leaving the organic membrane in a state adapted to separate examination, hydrofluoric acid must be employed as the menstruum. It is sometimes necessary, on the other hand, to get rid of the organic matter, for the sake of obtaining the mineral particles in a separate state, as in the case of the spicules of Sponges, Gorgoniæ, &c., this may be done either by incineration, or (which is generally preferable) by boiling or macerating for a long time in a solution of caustic potash. In separating from Guano, again, the siliceous skeletons of Diatomaceæ, &c., which it may contain, muriatic and nitric acids are largely used, to dissolve away every part of the mass on which they will act; the microscopic organisms for which search is made, being contained in a few grains of sediment which are left when a pound of pure guano is thus treated.

113. In applying Chemical Reagents to Microscopic objects for the purpose of *testing*, it is necessary to use great care not to add too much at once; and it is better that the test-bottle itself should afford the means of regulating the quantity, than that an additional rod or tube should be required. Two modes have been devised for this purpose. One consists in drawing the neck of the test-bottle to a capillary orifice, and covering it with a cap which fits around it; and the fluid is caused to flow from this, drop by drop, by the warmth of the hand applied to the bottle, which causes an expansion of the air it may contain.¹ When these bottles are emptied, they must be refilled by expelling the air by heat, and placing the capillary orifice under the surface of the fluid to be introduced, which will then be forced in as the bottle cools; this process may need to be repeated two or three times (care being taken that the heat applied be not so great as to crack the bottle); but it is better not to fill the bottle more than half-full, in order that air enough may be left for the warmth of the hand to act upon. The other arrangement for applying minute quantities of test-liquids, consists in the elongation of the stopper, which is drawn to a fusiform point, so as to serve as the test-rod for its own bottle.² This enables either a mere trace, or several ordinary drops, of the reagent to be applied at once; for the elongated stopper will take up a considerable quantity, a larger or smaller proportion of which (as desired), may be left behind, by bringing the lower part of the stopper into contact with the inside of the neck of the bottle, as it is being withdrawn. Whichever plan is made use of, great care should be taken to avoid carrying away from the slide to which the test-liquid is applied, any loose particles which may be upon it, and which may be thus transferred to some other object, to the great perplexity of the Microscopist. It is better, indeed, not to deposit the drop of test-liquid on the slide in immediate contact with the substance to which it is to be applied; but to bring the two into contact after the test-bottle has been withdrawn.

114. The following are the Test-Liquids most frequently needed.

1. Solution of *Iodine* in water (1 gr. of iodine, 3 grs. of iodide of potassium, 1 oz. of distilled water) turns starch blue, and cellulose brown; it also gives an intense brown to albuminous substances.

2. Dilute *Sulphuric Acid* (one of acid to two or three parts of water) gives to cellulose that has been previously dyed with iodine, a blue or purple hue; also, when mixed with a solution of sugar, it gives a rose-red hue, more or less deep, with nitrogenous substances and with bile (Pettenkofer's test).

¹ A set of 12 test-bottles on this plan, packed in a box, is supplied by Mr. Highley, of Fleet Street.

² Bottles of this pattern, which was devised by Dr. Griffith, are sold by Mr. Ferguson of Giltspur Street.

3. Solution of chloride of zinc, iodine, and iodide of potassium, made in the following way:—Zinc is dissolved in hydrochloric acid, and the solution is permitted to evaporate, in contact with metallic zinc, until it attains the thickness of a syrup; this syrup is then saturated with iodide of potassium, and iodine is last added. This solution (which is known as Schultz's test) serves, like the preceding, to detect the presence of cellulose, and has the advantage over sulphuric acid of being less destructive to the tissues. Each will sometimes succeed where the other fails; consequently, in doubtful cases, both should be employed.

4. Concentrated *Nitric Acid* gives to albuminous substances an intense yellow; when diluted with about two or three parts of water, it is very useful in separating the elementary parts of many Animal and Vegetable tissues, when these are boiled in it.

5. *Acetic Acid* (diluted with from three to five parts of water) is a most useful test-liquid to the Animal Histologist, from its power of dissolving, or at least of reducing to a state of such transparency that they can no longer be distinguished, certain membranes, fibres, &c.; whilst others are brought strongly into view.

6. *Acid Nitrate of Mercury* (Millon's test) colors albuminous substances red.

7. Solution of caustic *Potash* or *Soda* (the latter being generally preferable) has a remarkable solvent effect upon many organic substances, both Animal and Vegetable.

8. *Alcohol* dissolves resinous substances and many vegetable coloring matters, and renders most vegetable preparations more transparent; on the other hand, by its coagulating action, it renders many animal tissues (as nerve-fibres) more opaque, and thus brings them into greater distinctness.

9. *Ether* dissolves not only resins, but oils and fats.

SECTION 2. MOUNTING OF OBJECTS.

115. The Microscopist not merely desires to prepare objects for examination, but, where possible, to preserve them in such a manner that they may be inspected at any future time. This may be so effectually accomplished in regard to many substances, that they undergo no kind of change, however long they may be retained; and even delicate structures whose composition renders them peculiarly liable to decay, may often be kept, by complete seclusion from the air, and by immersion in a preservative fluid, in a state so nearly resembling that in which they were at first prepared, that they will continue, during an indefinite length of time, to exhibit their original characters with scarcely any deterioration. The method of "mounting" objects to be thus preserved, will differ, of course, both according to their respective natures, and also to the mode in which they are to be viewed,

whether as *transparent* or as *opaque* objects. Thus they may be set up dry, or in Canada balsam, or in some preservative liquid; they may need to be simply covered with thin glass, or they may require to be surrounded by a "cell;" if they are to be viewed by transmitted light, they must always have glass below them; but if they are to be seen by the light reflected from their surfaces, they may often be preferably mounted on wood, card, or some other material which itself affords a black background. In almost all cases in which transparent objects are to be mounted, use will have to be made of the slips of glass technically called *slides* or *sliders*, and of *covers* of thin glass, and it will therefore be desirable to treat of these in the first instance.

116. *Glass Slides*.—The kind of glass usually employed for mounting objects, is that which is known as "flatted crown;" and it is now almost invariably cut, by the common consent of Microscopists in this country, into slips measuring 3 in. by 1 in.; for objects too large to be mounted on these, the size of 3 in. by 1½ in. may be adopted. Such slips may be purchased, accurately cut to size, and ground at the edges, for so little more than the cost of the glass, that few persons to whom time is an object would trouble themselves to prepare them; it being only when glass slides of some unusual dimensions are required, or when it is desired to construct "built up cells" (§§ 136, 137), that a facility of cutting glass with a glazier's diamond becomes useful. The glass slide prepared for use, should be free from veins, air bubbles, or other flaws, at least in the central part on which the object is placed; and any whose defects render them unsuitable for ordinary purposes, should be selected and laid aside for uses to which the working Microscopist will find no difficulty in putting them. As the slips vary considerably in thickness, it will be advantageous to separate the *thick* from the *thin*, and both from those of *medium* substance: the last may be employed for mounting ordinary objects; the second for mounting delicate objects to be viewed by the high powers, with which the achromatic condenser is to be used, so as to avoid any unnecessary refraction of the illuminating pencil by the thickness of the plate which it has to traverse beneath the object; whilst the first should be set aside for the attachment of objects which are to be ground down, and for which, therefore, a stronger mounting than usual is desirable. Where very hard substances have to be thus operated on, it is advantageous to attach them in the first instance to pieces of very thick plate-glass; only transferring them to the ordinary slides, when they have been reduced to nearly the requisite thinness (§ 129).


117. *Thin Glass*.—The older Microscopists were obliged to employ thin laminae of *talc*, for covering objects to be viewed with lenses of short foci; but this material, which was in many respects objectionable, is now entirely superseded by the thin glass manufactured for this express purpose by Messrs. Chance

of Birmingham, which may be obtained of various degrees of thickness, from 1-20th to 1-250th of an inch. This glass, being unannealed, is very hard and brittle; and much care and some dexterity are required in cutting it. This should be done with the *writing* diamond; and it is advantageous to lay the thin glass upon a piece of wetted plate-glass, as its tendency to crack and "star" is thereby diminished. For cutting *square* or other *rectangular* covers, nothing but a flat rule is required. For cutting *rounds* or *ovals*, on the other hand, it is necessary to have "guides" of some kind. The simplest, which are as effective as any, consist of pieces of flat brass plate, perforated with holes of the various sizes desired; or curtain-rings, with a piece of wire soldered on either side: these being held firmly down on the thin glass with two fingers of the left hand, the writing-diamond is carried round the inner margin of the aperture with the right; care being taken that, in so doing, the diamond be made to revolve on its own axis, which is needful both that it may mark the glass, and also that the beginning and the end of the cut may join.¹ Where a number of such "rounds" are being cut at once, it saves much trouble, as well as risk of loss by breakage in separating them, to cut the glass first into strips, whose breadth shall equal the diameter of the rounds. But it is very convenient to use up for this purpose any odd pieces of glass, whose shape may render them unsuitable for being cut into "squares" without much waste. The pieces of thin glass thus prepared for use, should be sorted, not only according to size and shape, but also according to thickness. The thinnest glass is of course most difficult to handle safely, and is most liable to fracture from accidents of various kinds; and hence it should only be employed for the purpose for which it is absolutely needed,—namely, the mounting of objects which are to be viewed by the highest powers. The thickest pieces, again, may be most advantageously employed as covers for large cells in which objects are mounted in fluid (§§ 136, 137), to be viewed by the low powers, whose performance is not sensibly affected by the aberration thus produced. And the pieces of medium thinness will be found those most serviceable for all ordinary purposes; neither being, on the one hand, difficult to handle; nor, on the other, interfering with the clearness of the image formed by medium powers of moderate aperture, even when no special adjustment is made for the aberration they produce (§ 101).

¹ A very elegant little instrument, for the purpose of cutting thin glass rounds, contrived by Mr. Shadbolt, and another, of a more substantial character, invented by Mr. Darker, will be found described in Mr. Quekett's "Practical Treatise." These instruments, however, are rather adapted for the use of those who have occasion to prepare such rounds in large quantities, than for the ordinary working Microscopist, who will find the method above described answer his requirements sufficiently well. Indeed it is in some respects superior; since a firm pressure made by the ring or plate on the glass round, tends to prevent the crack from spreading into it. To every one to whom the saving of time is a greater object than the expenditure of a few shillings, it is strongly recommended that these "rounds" should be purchased ready cut; as they may be obtained of any required size and thinness, at a very moderate cost.

118. The exact thickness of any piece of glass may be determined without difficulty, by placing it edgeways on the stage of the microscope (holding it in the stage-forceps), and measuring its edge by the eye-piece micrometer (§ 46). A much more ready means is afforded, however, by the *Lever of Contact* (Fig. 60) devised by Mr. Ross for this express purpose. This instrument

FIG. 60.



Lever of Contact.

consists of a small horizontal table of brass, mounted upon a stand, and having at one end an arc graduated into 20 divisions, each of which represents 1-1000th of an inch, so that the entire arc measures 1-50th of an inch; at the other end is a pivot, on which moves a long and delicate lever of steel, whose extremity points to the graduated arc, whilst it has very near its pivot a sort of projecting tooth, which bears at * against a vertical plate of steel that is screwed to the horizontal table. The piece of thin glass to be measured, being inserted between the vertical plate and the projecting tooth of the lever, its thickness in thousandths of an inch is given by the number on the graduated arc, to which the extremity of the lever points. Thus, if the number be 8, the thickness of the glass is $\cdot 008$, or 1-125th of an inch. A very elegant little instrument, which is used by watchmakers for measuring the thickness of thin plates and wires, may be obtained at a much less cost from the dealers in Swiss tools; this answers the purpose equally well; but the "value" of its scale must be determined by gauging the thickness of a piece of glass, or the diameter of a fine wire, and comparing the number of divisions which it indicates, with the micrometrical measurement of the same body obtained by the microscope. When the glass covers have been sorted according to their thickness, it will be found convenient to employ those of one particular thickness for each particular class of objects; since, when one object is being examined after another, no readjustment of the objective will then be required for each. This will be found a great saving of time and trouble, when high powers are in use. It is undesirable to employ glass covers of greater thickness than 1-140th ($\cdot 007$) of an inch, with any object-glass whose aperture exceeds 75° ; and for object-glasses of 120° and upwards, the glass cover should not exceed 1-250th ($\cdot 004$) of an inch.

119. On account of the extreme brittleness of the thin glass, it is desirable to keep the pieces, when cut and sorted, in some

fine soft powder, such as starch. Before using it, however, the Microscopist should be careful to clean it thoroughly; not merely for the sake of removing foulnesses which would interfere with the view of the object, but also for the sake of getting rid of adherent starch-grains, the presence of which might lead to wrong conclusions, and also of freeing the surface from that slight greasiness, which, by preventing it from being readily wetted by water, frequently occasions great inconvenience in the mounting of objects in fluid. The thicker pieces may be washed and wiped without much danger of fracture, if due care be employed; but the thinner require much precaution; and in cleansing these, the simple method devised by Mr. Spencer will be found very useful. This consists in the use of a pair of round flat disks, about $1\frac{1}{2}$ inch in diameter, made of wood or metal covered with chamois leather, and furnished with handles; for when a piece even of the thinnest glass is laid upon one of these, it may be rubbed clean with the other, and any amount of pressure may be used, without the least risk of breaking it. Previously to doing this, however, it will be advantageous to soak the pieces for a time in strong sulphuric acid, and then to wash them in two or three waters; if greasiness, however, be their chief fault, they should be soaked in a strong infusion of nut-galls; with which it will be also advantageous to cleanse the surface of glass slides that are to be used for mounting objects in liquid.

120. *Varnishes and Cements*.—There are three very distinct purposes, for which cements that possess the power of holding firmly to glass, and of resisting, not merely water, but other preservative liquids, are required by the Microscopist; these being (1) the attachment of the glass covers to the slides or cells containing the object, (2) the formation of thin cells of cement only, and (3) the attachment of the glass plate or tube-cells to the slides. The two former of these purposes are answered by liquid cements or *varnishes*, which may be applied without heat; the last requires a *solid cement* of greater tenacity, which can only be used in the melted state. The Varnishes used for mounting objects in liquid, should always be such as contain *no mixture of solid particles*. This is a principle on which the Author, from an experience of many years, is disposed to lay great stress; having often made trial, at the recommendation of friends, of varnishes which were said to have been greatly improved by thickening with litharge or lamp-black; and having always found that, although they might stand well for a few weeks or months, they became porous after a greater lapse of time, allowing the evaporation of the liquid and the admission of air. He has himself found none more durable than that known as japanner's *Gold-size*, which may be obtained at almost every color-shop; for although this, when newly made, is apt to be somewhat too thin, so as to tend to *run in* beneath the glass cover, it may be easily

prevented by employing *very liquid* Gold-size, and by using it in extremely small quantity in the first instance; since whenever the glass cover lies perfectly flat on its bed, and the fluid beneath extends to the edges, the thin layer of this varnish dries very quickly, without any tendency to run in. When this has completely set, a second layer should be applied; and a layer of *Asphalte* over the whole will add to its security, and improve the appearance of the mounting. The danger of running in appears to the Author to be the greatest, when, in consequence of the use of old and viscid gold-size, the layer is too thick, and is long in drying. His experience leads him to distrust *Asphalte* when used alone, as being liable to admit air after a lengthened period of drying. He has recently learned from Mr. Tomes, that he finds saturated solution of arsenious acid to be a very good medium for mounting delicate preparations of animal structures. If it be allowed to become so thick, however, as not to be easily worked by the brush, it is quite unfit for use. There are few preservative liquids with which gold-size may not be employed; since it is not acted on by any aqueous solution, and resists moderately diluted spirit; oil of turpentine being its only true solvent.¹ Many Microscopists prefer the solution of shell-lac in naphtha, which is sold under the name of *Liquid Glue*; this dries more quickly than gold-size, but is more brittle when completely hardened, and does not, in the Author's opinion, adhere so firmly and enduringly to glass; and it is, moreover, more easily acted on by diluted alcohol than the preceding. Of late, a solution of *Asphalte* in drying-oil or turpentine, sometimes known under the name of "Brunswick-black," has come much into use. It is extremely easy and pleasant to work with, and dries quickly; but it is brittle when dry, and is disposed to crack, not merely when subject to any "jar," but also (after some time) spontaneously. This evil may be corrected, according to Mr. Brooke, by adding to it a little solution of Caoutchouc in mineral naphtha. Oil of turpentine is the solvent for this varnish, as for gold-size; and brushes which have been used with either, may be cleansed by that menstruum; those which have been used with liquid glue, may be cleansed with naphtha. For mounting objects *dry* (§ 122), or for giving a finished appearance to mountings which have been made by one or other of the foregoing cements, varnishes may be used, which, from containing coloring particles, or from being acted on by the preservative liquids employed, could not be safely laid on in the first instance. Among the most convenient of this kind, are varnishes made by dissolving red or black or any other colored *Sealing-wax* in strong alcohol; these are more to be recommended for their appearance, however, than for their

¹ The Author has preparations mounted with gold-size as much as twelve years ago, which have remained perfectly free from leakage; the precaution having been taken, to lay on a thin coat of varnish every two or three years.

tenacity, being very apt to lose their hold upon the glass after a time; and the Author, having suffered much injury to his preparations from trusting to them, would recommend that, even in mounting objects dry, some other cement should be first used, by which the glass cover should be attached to the slide, the sealing-wax varnish being only laid on as a finish. If a black varnish be desired for such a purpose, it may be readily made by mixing gold-size with a small quantity of lamp-black; this dries quickly, and is free from brittleness; but, for the reason already mentioned, it should not be used in the first instance to mount objects in fluid, although it may be laid as a finish over gold-size or asphalte. For making cement-cells (§ 134), either asphalte, gold-size, or liquid glue may be employed, the first being on the whole preferable; the varnish termed *Black Japan* also makes very good and durable cells, if the glasses to which it has been applied be exposed to the heat of an oven, not raised so high as to cause them to "blister."

121. Although Canada balsam has been sometimes used as a cement, and has the advantage of being worked with extreme convenience, yet it is so apt to crack when hardened by time, that a slight "jar" will cause the cell to spring away from the glass to which it has been attached. Hence, if employed at all for fixing cells to glass slides, its use should be limited to those plate-cells which afford a large surface of attachment (§ 136), or to those very thin tube-cells (§ 135), which cannot be so conveniently attached with marine glue, and of which the cover may be secured to the slide by spreading the ring of gold-size round the margin of the cell itself (§ 138). Care should be taken, in applying the Canada balsam, that it be sufficiently hardened by heat, but that it be not so heated as to become brittle (§ 109): the general method of using it for this purpose, is the same as that which must be practised in the case of marine glue. The superfluous balsam left after pressing down the cell, is to be removed, first by scraping with a heated knife, and then with a rag dipped in oil of turpentine, after which it is desirable to give the glass surface a final cleansing with alcohol. For all kinds of cells (§§ 135, 137) except those just mentioned, the proper cement is *Marine Glue*, which is a mixture of shell-lac, caoutchouc, and naphtha, now extensively employed; being distinguished by its extraordinary tenacity, and by its power of resisting solvents of almost every kind. Different qualities of this substance are made for the several purposes to which it is applied; that which is most suitable to the wants of the Microscopist, is known in commerce as GK 4. As this cement can only be applied hot, and as it is a great saving of trouble to attach a considerable number of cells at the same time, a *Mounting-Plate* should be provided, which will furnish the requisite heat to several slides at once. Such a surface may be afforded by the top of a stove; but it is better to have one which can be used at all seasons, and the heat

of which can be precisely regulated at pleasure. A very simple apparatus much used for this purpose, consists of a small table of brass or iron plate, about 6 inches long and 2 broad, with legs about 4 inches high, either screwed into its four corners, or so jointed to them as to fold down; this is set over a small spirit-lamp, the flame of which is regulated to give the heat required. The Author has found it much preferable, however, to lay the plate on one of the rings of a small "retort-stand" (used in Chemical operations), which admits of being shifted to any height that may be desired, so that the heat applied may be precisely graduated; or, if a gas lamp be employed for the ordinary purposes of illumination, its stem may be fitted with a sliding-ring, which will carry either a hot plate or a water-bath.¹ It is more convenient, however, to have two such plates, laid on two rings; one being allowed to cool with the slides upon it, whilst the other is being heated. The glass slides and cells which are to be attached to each other, must first be heated on the mounting-plate; and some small cuttings of marine-glue are then to be placed, either upon that surface of the cell which is to be attached, or upon that portion of the slide on which it is to lie, the former being perhaps preferable. When they begin to melt, they may be worked over the surface of attachment by means of a needle-point; and in this manner, the melted glue may be uniformly spread, care being taken to pick out any of the small gritty particles which this cement sometimes contains. When the surface of attachment is thus completely covered with liquefied glue, the cell is to be taken up with a pair of forceps, turned over and deposited in its proper place on the slide; and it is then to be firmly pressed down with a stick (such as the handle of the needle) or with a piece of flat wood, so as to squeeze out any superfluous glue from beneath. If any air-bubbles should be seen between the cell and the slide, these should if possible be got rid of by pressure, or by slightly moving the cell from side to side; but if their presence results, as is sometimes the case, from deficiency of cement at that point, the cell must be lifted off again, and more glue applied at the required spot. Sometimes, in spite of care, the glue becomes hardened and blackened by overheating; and as, in this case, it will not stick well to the glass, it is preferable not to attempt to proceed, but to lift off the cell from the slide, to let it cool, and then to repeat the process. When the cementing has been satisfactorily accomplished, the slides should be allowed to cool gradually, in order to secure the firm adhesion of the glue; and this is readily accomplished, in the first instance, by pushing each, as it is finished, towards one of the extremities of the plate, which is of course cooler than the centre. If two plates are in use, the heated plate may then be readily moved away upon the ring which supports

¹ Both these fittings are adapted to the Gas lamp supplied for the use of Microscopists by Mr. S. Highley (§ 75).

it, the other being brought down in its place; and as the heated plate will be some little time in cooling, the firm attachment of the cells will be secured. If, on the other hand, there be only a single plate, and the operator desire to proceed at once in mounting more cells, the slides already completed should be carefully removed from it, and laid upon a *wooden* surface, the slow conduction of which will prevent them from cooling too fast. Before they are quite cold, the superfluous glue should be scraped from the glass with a small chisel or awl; and the surface should then be carefully cleansed with a solution of potash, which may be rubbed upon it with a piece of rag covering a stick shaped like a chisel. The cells should next be washed with a hard brush and soap and water, and may be finally cleansed by rubbing with a little weak spirit and a soft cloth. In cases in which *appearance* is not of much consequence, and especially in those in which the cell is to be used for mounting large opaque objects, it is decidedly preferable not to scrape off the glue too closely round the edges of attachment, as the "hold" is much firmer, and the probability of the penetration of air or fluid much less, if the immediate margin of glue be left, both outside and inside the cell.

122. *Mounting Objects Dry.*—There are certain objects, which, even when they are to be viewed by *transmitted* light, are more advantageously seen when simply laid on glass, than when they are immersed either in fluid or in balsam. This is the case especially with sections of bones and teeth, much of whose internal structure is obliterated by the penetration of fluid; and also with the scales of Lepidopterous and other Insects, whose minute surface-markings are far more distinct when thus examined, than when treated in any other way.¹ For preserving such objects, it is of course desirable that they should be protected by a cover; and this must be so attached to the glass slide, as to keep the object in place, besides being itself secured. For this purpose sealing-wax varnish is often used, but is unsuitable on account of its brittleness when dry; gold size mixed with lamp-black is much to be preferred, and, if carefully laid on, will not tend to run in between the cover and the slide. If the object have any tendency to curl up, or to keep off the cover from the slide by its own "spring," it will be useful, while applying the varnish, to make use of pressure, such as that afforded by the little implement represented in Fig. 62. This pressure should

¹ It is affirmed by two high authorities on all that relates both to the theoretical and practical action of Object-glasses of large aperture (namely, Prof. Robinson and Mr. Wenham), that the effect of mounting delicate test-objects in Canada balsam is practically to reduce the aperture, since no rays can diverge after passing through a stratum of this substance, at a greater angle than 85° or 90°. Hence they recommend that the "difficult" *Diatomaceæ* should *not* be mounted in balsam, if they are to be viewed by objectives of 120° or 130°. Their position is disputed, however, by Prof. Bailey (U. S.), who affirms that balsam-mounted specimens are preferable as test-objects. Those who are interested in this question, will do well to consult the papers of these gentlemen, in the 2d and 3d volumes of the "Quart. Journ. of Microsc. Science."


not be remitted, until the varnish is dry enough to hold down the cover by itself. For mounting delicate objects, the *thinner* slides should be selected (§ 116); and for very difficult test-objects, it is advantageous to employ thin glass below as well as above the specimens, for the sake of diminishing the aberration which the illuminating pencil sustains in its passage to the object, and for allowing the achromatic condenser to approach the object as closely as possible. For this purpose, the simplest method is to take a slip of wood, of the ordinary size of the glass slide (3 in. by 1 in.), with a central aperture of from 3 to 5-8ths of an inch; to cover this aperture with a "square" or "round" of thin glass of sufficient size to project considerably beyond it; to lay the object upon this glass, and to protect it with a cover of rather smaller size, which should be fastened down all round by varnish, to prevent the entrance of moisture; and finally to secure both glasses to the wooden slide, by gumming down over them a piece of paper, with a perforation of the same size as that of the slide itself.

123. For dry-mounting *opaque* objects, the method adopted must vary with the mode in which the object is to be illuminated; since, if a side-condenser or reflector is to be employed, the whole slide may be opaque; whereas if the Lieberkühn is preferred, the object should be placed on a disk of appropriate size (§ 65), supported in such a mode as to admit the light all around it. For the former purpose, the Author has devised the following simple method, which he has found to afford peculiar conveniences. Let there be provided a cedar slide of the kind just described, a piece of card of the same dimensions, and a piece of *dead-black* paper, rather larger than the aperture of the slide, if a dark mounting be desired, which is preferable for most objects; this piece of paper is to be gummed to the middle of the card, and then, some stiff gum having been previously spread over one side of the slide (care being taken that there is no superfluity of it immediately around the aperture), this is to be laid down upon the card, and subjected to pressure.¹ An extremely neat "cell" will thus be formed for the reception of the object, the depth of which will be determined by the thickness of the slide, and the diameter by the size of the perforation; and it will be found convenient to use slides of various thicknesses and having apertures of different sizes. The cell should always be deep enough for its wall to rise above the object; but, on the other hand, if it be too deep, its wall will interfere with the oblique incidence of the light upon any object that may be near its periphery. The object, if flat or small, may be attached by ordinary gum-mucilage; if, however, it be large, and the part of it to be attached have an irregular surface, it is desirable to afford a "bed" to this by gum

¹ It will be found a very convenient plan, to prepare a large number of such slides at once; and this may be done in a marvellously short time, if the slips of card have been previously cut to the exact size in a bookbinder's press.

thickened with starch. The complete protection thus given to the object, is the great recommendation of this method; since, when objects are simply fastened on black paper gummed on a slip of glass, their protection from its surface renders them constantly liable to accidents; as many know, to their cost, who have used that mode of mounting. But this is by no means its only convenience. It is far cheaper than mounting objects in glass cells, which is the only other mode of affording them protection, save the use of pill-boxes in the manner to be presently described. It allows the slides, not only to range in the ordinary cabinets, but also to be laid one against another, and to be packed closely in cases or secured by elastic bands; and this last plan is extremely convenient, not merely for the saving of space, but also for preserving the objects from dust. Should any more special protection be required, a thin glass cover may be laid over the top of the cell, and secured there by gummed paper; but this will, of course, occasion a slight projection, which will expose the glass cover to the risk of fracture when the slide is pressed against others; and the Author's experience leads him to conclude, that the mode of packing just described affords a security from dust, that is scarcely less effectual than a thin glass cover. Further, the card on the under surface affords a great convenience for writing on the slide the name and other particulars of the object. If the object be so large as to project above the surface, even when the thickest slides are used which it is convenient to employ, an additional protection may be afforded, by gluing a couple of strips of wood of adequate thickness along the edges of the slide;¹ or by gumming a second slide to the face of the first, taking care that its aperture be large enough to prevent obstruction to oblique light. Very delicate, flat objects, on the other hand, even when to be viewed by incident light, should be mounted on glass and protected by a cover, in the same manner as transparent objects; a dark background being furnished to them, either by the "dark well" (§ 65) or by the closed diaphragm (§ 55).

124. Objects which are to be viewed by the Lieberkühn should be mounted either on flat disks of card or cork, or, if protection be desired, in cups of proper depth, resembling very shallow pill-boxes, which may be made with or without covers,² of any size

¹ A very convenient kind of slide, for mounting large opaque objects, is made by Messrs. Carpenter and Westley, 24 Regent Street, by ploughing out a groove in a strip of wood, so that its section presents this form . The object may be mounted upon the bottom of the groove; or upon a disk of card fitted into an aperture in the slide, and so held there by a brass ring, that it may be taken out and held between the stage-forceps, so as to be viewed with the Lieberkühn. These slides are commonly made five inches long, and are fitted for the reception of four objects; for the sake of uniformity, however, as well as for preventing them from overbalancing on the stage, it might be convenient that they should be made three inches long, and that each should only receive one or (at most) two objects.

² Round boxes with glass covers are now coming into very extensive use for the preservation of Natural History specimens of various kinds, in such a manner that the con-

desired, and lined with black paper, by any pill-box maker. The disks or pill-box cells may be attached to glass slides by gum, so as to range in an ordinary object-cabinet; and this mode will be generally adopted by such as lay stress upon uniformity, and prefer the easiest methods of exhibiting their objects. As there are many opaque objects, however, which can only be well judged of when different sides are presented to the Microscope, there is a great advantage in mounting these in such a manner, as to admit of their being turned at various angles; and this may be done by attaching the disk with sealing-wax or some other cement, to a pin, which may be either held between the blades of the stage-forceps, or passed into the cork box at its other extremity (§ 66). If the Microscopist should be pursuing the study of any class of objects which renders it desirable to mount a large number in this mode, the most convenient plan is to glue two pieces of cardboard to the two sides of a piece of rather thick chamois leather; one of the surfaces of this sandwich-like board should be covered with dead-black paper; and disks of any desired diameter may then be cut out with a punch, and mounted upon a pin by simply passing it through the stratum of leather. The pill-box mounting will be less advantageous for this purpose, since, if the object be completely buried in the cell, there will be less power of seeing it on any but its upper surface; a pin may be secured to the bottom of such a box, however, by gumming over it a piece of stout paper. Disks may easily be punched, also, out of sheet gutta percha of any convenient thickness; or short cylinders may be made, by cutting up the thick cords made for lathe-bands; these are all readily penetrated by a heated pin, and no further trouble is necessary to attach them to it. Protective wells may also be made by cutting off short pieces from a gutta percha tube, and attaching these to the disks by a gentle heat. For the reception of boxes or disks thus mounted on pins, a drawer should be provided with a thick cork bottom, into which the pin is to be inserted far enough to prevent risk of displacement.

125. *Mounting Objects in Canada Balsam.*—This method of mounting is suitable to a very large proportion of those objects, which are to be viewed by transmitted light, and whose texture is not affected by the loss of the aqueous fluid they may contain; and it has many advantages over the mounting of the like objects dry. For, in the first place, as it fills up the little inequalities of their surface, even where it does not actually penetrate their substance, it increases their transparency by doing away with irregular refractions of the light in its way through them,

tents of each box, whilst protected and kept together, are at the same time presented to the eye. The Author has found the smallest and shallowest boxes of this kind that can be made, especially when lined with black paper, extremely useful for keeping Foraminifera and other Microscopic organisms in quantities; and also for mounting larger specimens for the Microscope as above described.


and gives them the aspect of perfect smoothness ; this is well seen in the case of sections of *Shell*, &c., which, when thus mounted, do not require a high polish (§ 110). But, secondly, where the structure, although itself hard, is penetrated by internal vacuities, the balsam, by filling these, prevents that obscuration resulting from the interposition of air-spaces, and from additional internal surfaces of reflection, by which the transmitted rays are distorted, and a large proportion of them lost: this is well seen in the case of the *Foraminifera*, and of sections of the "test" and "spines" of *Echinida*, whose intimate structure can be far better made out, when they are thus mounted, than when mounted dry, although their substance is (for the most part at least) itself so dense, that the balsam cannot be imagined to penetrate it; and likewise with dry *Vegetable* preparations, which are perhaps also affected in the manner to be next described. Thirdly, there are very many structures of great interest to the Microscopist, whose appearance is extraordinarily improved by this method of mounting, in consequence of a specific effect which the balsam has in combining (so to speak) with their component elements, so as to render them far more transparent than before: this effect is seen in the case of all dry preparations of *Insect* structure, especially of such as consist of their hard external tegument or of parts derived from this; also in the various *horny* tissues (hairs, hoof, horn, &c.) of the higher animals; and likewise in many organized substances, both recent and fossil, which are penetrated by calcareous matter in an *amorphous* condition. Besides these advantages, the mounting of objects in Canada balsam affords one of the easiest methods of fixing and preserving them; and consequently, it is almost always had recourse to, in the case of such transparent objects as do not need to be preserved in fluid; save where, in virtue of the action just described, it impairs the distinctness of surface-markings, or obliterates internal cavities or canals, which constitute the most important features of the object.

126. Canada Balsam, being nothing else than a very pure *Turpentine*, is a natural combination of resin with the essential oil of turpentine. In its fresh state, it is a viscid liquid, easily poured out, but capable of being drawn into fine threads; and this is the condition in which the Microscopist will find it most desirable to use it for the mounting of objects generally. The balsam may be conveniently kept in a glass bottle or jar with a wide mouth, being taken up as required by a small glass rod drawn to a blunt point, such as is used by Chemists as a "stirrer;" and if, instead of a cork or stopper, this bottle should be provided with a tall hollow "cap," the glass rod may always stand in the balsam, with its upper end projecting into the cap. In taking out the balsam, care should be taken not to drop it prematurely from the rod; and not to let it come into contact with the interior of the neck or with the mouth of the jar. Both

these mischances may be avoided, by not attempting to take up on the rod more than it will properly carry; and by holding it in a horizontal position, after drawing it out from the bottle, until the slip on which it is to deposit the balsam, is just beneath its point. The Author has himself been in the habit of employing, instead of a hollow cap to his balsam jar, a disk of light wood, simply lying on its mouth; the centre of this disk is perforated with a hole, just large enough to allow the glass rod to fit it rather stiffly; and the rod is to be passed into the jar, only so far as to dip by its point into the balsam, being pushed further down as the level of the balsam is lowered. In this manner, a small quantity of balsam may be taken up with far less risk of *messing* it, than when the rod dips into it for two or three inches; if due care be taken not to allow the balsam to come into contact with the lip of the jar, the cover never sticks to it, but is readily lifted off upon the rod; and although it might seem an incumbrance in the use of the rod, yet it is not practically found to be so. If the balsam be kept too long, it becomes, through the loss of part of its volatile oil, too stiff for convenient use, and may be thinned by mixing it at a gentle heat with pure oil of turpentine; this mixture, however, does not produce that thorough incorporation of the constituents which exists in the fresh balsam; and it is consequently preferable to use in other ways the balsam which has become somewhat too stiff, and to have recourse to a fresh supply of liquid balsam for mounting purposes. When Canada balsam is to be employed as a *cement*, as for attaching sections, &c., to glass slides (§ 109), it should be in a much stiffer condition; since, if it be dropped on the slide in too liquid a state, it will probably spread much wider, and will lie in a thinner stratum than is desirable. This hardening process may be carried to any extent that may be desired, by exposing the balsam in an uncorked jar (the mouth of which however, should be covered with paper, for the sake of keeping off dust) to a continual gentle heat, such as that of a water-bath.

127. In mounting objects in Canada balsam, it is convenient to be provided with certain simple instruments, the use of which will save much time and trouble. For the heat required, a

FIG. 61.




Slider Forceps.

Spirit lamp is by far the best source; both as admitting of easy regulation, and as being perfectly free from smoke. Where a number of objects are being mounted on the same occasion, it will be found convenient to employ either a water-bath covered

with a flat plate of metal, or a similar metal plate supported at such a distance above the lamp flame (§ 121), as not to become more heated than it would be through a water-bath. For holding the slide, either whilst it is being heated over the flame, or whilst it is being subsequently cooled, the Wooden Forceps (Fig. 61) contrived for this purpose by Mr. Page, will be found extremely convenient; this, by its elasticity, affords a secure grasp to a slide of any ordinary thickness, the wooden blades being separated by pressure upon the brass studs; and the lower stud, with the bent piece of brass at the junction of the blades, affords a level support to the forceps, which thus, while resting upon the table, keeps the heated glass from contact with its surface. Besides a pair of fine-pointed Metal forceps for holding the object to be mounted, there should be another of a commoner kind for taking up the glass cover, the former being liable to be soiled with balsam. A pair of stout needles mounted in wooden handles (§ 105), will be found indispensable, both for manipulating the object, and for breaking or removing air-bubbles; and if these handles be cut to a flat surface at the other extremity, they will serve also to press down the glass covers, for which purpose a pointed stick also is useful. For holding down these covers, whilst the balsam is cooling, if the elasticity of the objects should tend to make them spring up, nothing is more simple or convenient than a compressor made by a slight alteration of the "American clothes peg," which is now in general use in this country for a variety of purposes; all that is necessary being to rub down the opposed surfaces of the "clip" with a flat file, so that they shall be parallel to each other when an ordinary slide with its cover is interposed between them (Fig. 62). Great care should be taken to keep these implements free from soils of balsam; since the slides and glass covers are certain to receive them. The readiest mode of cleansing the needles (their "temper" being a matter of no consequence for these purposes) is to heat them red hot in the lamp, so as to burn off the balsam; and then carefully to wipe them. The forceps, both of wood and of metal, should be cleansed with oil of turpentine or with rectified spirit.

128. Much of the success of mounting objects in this mode will depend upon their previous preparation. They should have been previously well cleansed with water, from which they should be transferred into proof-spirit, as this will dry out much more readily than water. If they have any greasiness of surface or of substance, this should be removed by maceration in ether or in oil of turpentine; and maceration in turpentine is also very useful in preparing the way for the penetration of the balsam into

FIG. 62.



Spring Press.

substances which are unusually opaque. A long-continued maceration in turpentine, moreover, assists in freeing the specimen from air-bubbles; as it gradually creeps into spaces which are otherwise unoccupied, and, when the object is transferred to the balsam, the free miscibility of the two substances causes its place to be partly taken by the latter. In mounting an ordinary object, a sufficient quantity of liquid balsam should be laid in the centre of the slide; this should be warmed, but not boiled; and any air-bubbles which may make their appearance, should either be caused to burst by touching them with the needle-point, or should be drawn to one side. The object, if it can be held in the fine-pointed forceps, should then be plunged into the drop of balsam; and, if it be not completely covered, a little more balsam should be applied over it, care being taken as before to prevent over-heating, and to get rid of the bubbles as they rise. If, on the other hand, the object be in a finely divided condition (such, for example, as a collection of fossil Infusoria, of Spongespicules, &c.), it is better to place it first in the position it is to occupy on the slide, then to place near it a drop of balsam, which is to be warmed and freed from bubbles as before; the slide is then to be so inclined, as to cause the balsam to run towards the object, through and over which it will probably diffuse itself without any air-bubbles; but if any should arise, they may be broken by a heated needle-point. If the object contain numerous large air-spaces with free openings, and be one whose texture is not injured by heat, as is the case with Foraminifera, the air may often be got rid of by boiling it in the balsam; for the heat, causing the air to expand, drives out a large proportion of it; this will be replaced, if it be allowed partly to cool, by the entrance of balsam; and then, by a second heating, the balsam being boiled within the cavities, its vapor expels the remaining air, and, on the condensation of the vapor, the liquid balsam runs in and takes its place. For this method to succeed, however, it is essential that the balsam be prevented from becoming hard through boiling, by the addition of fresh liquid balsam from time to time; and it will often be found that, should vacuities remain which boiling does not remove, these contract or altogether disappear if the slide be kept for a few days at a gentle heat, the semi-fluid balsam being gradually forced into their place by the pressure of the surrounding air. There are many textures, however, which are extremely injured by a very slight excess of heat, having a tendency to curl up and to become stiff and brittle; and objects containing these are at once spoiled by boiling them in balsam. In such cases, it is much better to have recourse to the assistance of the air-pump; for by placing the slide, with the object immersed in very liquid balsam, upon a tin or copper vessel filled with hot water, under the receiver, and then exhausting this, the air-bubbles will be drawn forth, and, on the readmission of the air, the balsam is forced by its

pressure into the place which they occupied. Some objects, however, retain the air with such tenacity, as to require the repetition of the exhausting process two or three times: and in this case it is preferable to use camphine or oil of turpentine instead of balsam, on account of its greater fluidity, and to warm even this to a temperature of about 100°. There are certain cases, on the other hand, in which it is desirable to retain, instead of expelling, the air contained within the cavities of the object. Thus, if minute insects (such as Fleas) be displayed as transparent objects to show the ramifications of the tracheæ, or if it be wished that a section of Tooth or Bone should be so mounted in balsam as to exhibit its canaliculi, the previous maceration in oil of turpentine should be never employed, and the balsam employed should be some which has been previously hardened; this being melted without the use of more heat than is necessary, the object should be surrounded by it, and the cover put on as quickly as possible; and the slide should then be laid upon a surface of stone or metal, the good conducting power of which causes the balsam to cool rapidly, and thus diminishes its tendency to penetrate into the substance of the object.

129. When the object is already attached to the glass slide, the mounting in Canada balsam is usually a matter of very little difficulty. If it be a soft tissue which has been spread out and allowed to dry upon the glass for the purpose of securing it in its place, all that is necessary in the first instance is to dry it thoroughly, to shave or scrape it with a sharp knife if it should seem too thick, and to moisten its surface with oil of turpentine if it should not readily "take" the balsam. The slide is then very gently warmed, a sufficient quantity of balsam is spread over the surface of the specimen, care being taken that it "takes" it in every point, and the glass cover is put on. If the preparation cover a large area, great care should be taken in letting down the cover *gradually* from one side, so as to drive a wave of balsam before it which shall sweep away air-bubbles; raising it a little, and introducing a small quantity of fresh balsam, if any vacuity present itself as it descends. The preferable mode, however, of mounting thin sections of hard bodies, will depend in great degree upon the size of the section and the tenacity of its substance. Where its area is great and its texture brittle, its removal from the glass on which it has been ground down, to another slip, cannot be accomplished, even by the most dexterous management, without considerable risk of breaking it; and although, by the friction of the glass upon the stone, the edges of this will probably have been scratched or roughened, yet this is a dis-sight about which the scientific Microscopist will care but little, as it only affects the saleable value of such objects. Nothing more will in this case be necessary, than to lay some liquid Canada balsam on the surface of the section, to warm it gently, and then to place on it a thin glass cover of suitable dimensions, gently

pressing this down wherever the balsam happens to be thickest, and endeavoring to drive all air-bubbles before a wave of liquid, until they are entirely expelled, or at any rate are driven beyond the margin of the section. If this operation be not at once successful,—either a few large air-bubbles, or a great number of smaller ones, which cannot be got rid of by gentle pressure, being visible between the surface of the section and the covering-glass,—it is better at once to remove the cover by gentle warmth, and to repeat the operation with an additional supply of balsam, rather than to attempt to drive out the bubbles by any manipulation. Whatever treatment be adopted, special care should always be taken not to apply so much heat as to *melt* the hard balsam *beneath* the section,¹ or to *boil* the thin balsam *above*; in the former case, the loosening of the section from the glass is very apt to be followed by the detachment of some portions of it from the rest, whilst the glass cover is being pressed down; and in the latter, the production of bubbles very seriously embarrasses the operation. If the heat should unfortunately be carried so far as to *boil* the cement *beneath* the section, there will be little chance, if its area be large, of getting rid of the bubbles thus produced, without removing it altogether from the glass to which it was attached, or, at any rate, without pushing it along the glass, in such a way as to slide it away from the bubbles; in that case, the part towards which it is moved should always be well supplied with balsam, and the bubbles that remain should be drawn away or broken with the needle-point; after which, the section being slid back to its original position, it is probable that no bubbles may be found beneath it. In cases, however, in which the *appearance* of the preparations is an object of much consideration, and in which the tenacity of the substance and the small size of the section prevent much risk of its breaking in the transfer, it may be loosened from the glass to which it was first attached, either by heat, or by soaking in ether. The former being the simplest and readiest method, is the one most commonly practised; the only difficulty lies in lifting off the specimen without breaking it; and this may best be done, by means of a camel-hair brush dipped in oil of turpentine. The glass to which the section is to be transferred, should have a large spot of liquid balsam laid in the proper place; the object is to be laid on this, and its upper surface covered with the like balsam; and then, the thin glass cover being placed upon it, this is to be gently pressed down in the manner already described. If ether be had recourse to, the slide should be placed in a wide-mouthed bottle of that liquid, which should then be corked or stopped; and after a time, the section will be found to be lying detached in it, whence it may be taken up either by the forceps

¹ It will often be found convenient to turn the slide with the face downwards, and to apply a gentle heat directly to the thin glass *cover*, and to the balsam *above* the object, instead of heating this through the glass *slide* and the balsam *beneath* it.

or camel-hair brush. Such a transfer will often be found advantageous, before the final completion of the reducing process; for it will occasionally happen that we find something in the structure of the specimen, which will be best displayed by rubbing it down afresh on the side first attached to the glass; and, when a number of small sections are being made at once (which it is often very convenient to do, not only in the case already mentioned, § 108, but in many others), it not only saves time, but insures the accurate flattening of the surface in grinding, to fix several upon the same slip, and to work them down together, until the requisite thinness has been nearly attained, when they must be transferred to separate slips, and finished one by one. In either case, the reattachment must of course be made like the original attachment, with balsam which has been first hardened (§ 109).

130. When the Balsam employed in mounting has remained in the liquid condition here recommended, the glass cover will not be secure from displacement until the balsam has become harder. This change it will require a long time to undergo, unless the aid of a gentle continuous warmth be afforded. Nothing is more suitable for this purpose, than the warmth of a chimney-piece immediately above the fire-place; as it is quite sufficient to produce the effect in the course of a few days, whilst there is no danger of its becoming excessive; but in default of this convenience, an oven carefully regulated, or (still better) a water-bath, may be employed. Whether either of these means be adopted, or the slides be put aside for the balsam to be hardened by time, they should always be laid in the *horizontal* position, that their covers may not be caused by gravitation to slip down from their places. It may be better, before submitting the slides to this hardening process, to scrape from their surface any superfluous balsam that does not immediately surround the glass cover; but the knife should never be carried so near to the edge of this, as to run any risk of displacing it; and it is much better to defer the final cleaning of the slide, until the attachment of the cover has become firm. The remaining balsam may then be scraped away with a knife or small chisel, the implement being warmed if the balsam be very stiff; the slide should then be rubbed with a rag dipped in oil of turpentine, until every perceptible soil of balsam is removed, especial care being taken to cleanse the surface and edges of the glass cover; and as this will itself leave a certain resinous film, it is better to give the slide a final cleansing with Alcohol. If its surface should have been considerably smeared with balsam, it is very convenient, after scraping away all that can be removed in that manner, to scrub it with a tooth-brush or nail-brush, first letting fall on it a few drops of turpentine or spirit of wine; and there is less risk of displacing the glass cover in this mode, than in rubbing it in any other way. The menstrua which serve thus to

cleanse the slides, of course answer equally well for cleansing the hands. The most ready solvent for balsam is Ether; but the ordinary use of this being interdicted by its costliness, and by the quickness with which it is dissipated by evaporation, Alcohol or Oil of Turpentine may be used in its stead.

131. *Preservative Fluids.*—Objects which would lose their characters in drying, can of course only be preserved in anything like their original condition, by mounting in fluid; and the choice of the fluid to be employed in each case will depend upon the character of the object, and the purpose aimed at in its preservation. As specific directions will be given hereafter in regard to most of the principal classes of Microscopic preparations, little more will be required in this place, than an enumeration of the preservative fluids, with a notice of their respective qualities. For very minute and delicate Vegetable objects, especially those belonging to the orders Desmidiaceæ and Diatomaceæ, nothing seems to produce less alteration in the disposition of the endochrome, or serves better to preserve their color, than *Distilled Water*; provided that, by the complete exclusion of air, the vital processes and decomposing changes can be alike suspended. This method of mounting, however, is liable to the objection that confervoid growths sometimes make their appearance in the preparation; and it is preferable to make some addition to the water, which shall be unfavorable to their development. Saturation with *camphor*, or shaking up with a few drops of *creasote*, will sometimes answer; but if the preservation of color be not an object, nothing is better than the addition of about a tenth part of *alcohol*; and where the loss of color would be objectionable, the solution of a grain of bay-salt and a grain of alum in an ounce of water will give it the requisite qualities. For larger preparations of Algæ, &c., what is called *Thwaites's Fluid* may be employed; this is prepared by adding to one part of rectified spirit as many drops of *creasote* as will saturate it, and then gradually mixing up with it in a mortar some prepared chalk with 16 parts of water; an equal quantity of water saturated with camphor is then to be added, and the mixture, after standing for a few days, is to be carefully filtered. A liquid of this kind also serves well for the preservation of many animal preparations; but it is said by Dr. Beale to become turbid when thus employed in large quantity; and he recommends the following modification. Mix 3 drachms of *creasote* with 6 ounces of wood naphtha, and add in a mortar as much prepared chalk as may be necessary to form a smooth thick paste; water must be gradually added to the extent of 64 ounces, a few lumps of camphor thrown in, and the mixture allowed to stand for two or three weeks in a lightly covered vessel, with occasional stirring; after which it should be filtered, and preserved in well-stopped bottles. Of late years, *Glycerine* has been much employed as a preservative fluid; it allows the colors of vegetable substances to be

retained, but, unless much diluted, it alters the disposition of the endochrome; and confervoid growths are apt to make their appearance in it. The best proportion seems to be one part of glycerine to two parts of camphor-water. The preparation known as *Deane's Gelatine* is one of the most convenient media for preserving the larger forms of Confervæ and other Microscopic Algæ, as well as sections of such as are still more bulky. This is prepared by soaking 1 oz. of gelatine in 4 oz. of water until the gelatine is quite soft, and then adding 5 oz. of honey previously raised to boiling heat in another vessel; the whole is then to be made boiling hot, and when it has somewhat cooled, but is still perfectly fluid, 6 drops of creasote and $\frac{1}{2}$ oz. of spirit of wine, previously mixed together, are to be added, and the whole is to be filtered through fine flannel. This composition, when cold, forms a very stiff jelly; but it becomes perfectly fluid on the application of a very slight warmth, and may then be used like any other preservative liquid, care being taken, however, that the slide and the glass cover are themselves gently warmed before it comes into contact with them. A mixture of gelatine and glycerine has been employed for the same purpose. Some Vegetable Anatomists make great use of a solution of *Chloride of Calcium* (muriate of lime) in three parts of water; its chief advantage being, that owing to its deliquescent property, it does not dry up, even when the cell is not perfectly closed in. It has, however, the disadvantages of not preserving colors, and of altering the disposition of the cell-contents by its endosmotic power. For the preservation of microscopic preparations of *Animal* tissues, a mixture of one part of Alcohol and five of water will generally answer very well, save in regard to the removal of their colors. Where the preservation of these is aimed at, the best medium will usually be *Goadby's Solution*, which is made by dissolving 4 oz. of bay salt, 2 oz. of alum, and 4 grains of corrosive sublimate, in 4 pints of boiling water; this should be carefully filtered before it is used; and for all delicate preparations it may be diluted with an equal bulk, or even with twice its bulk of water. This solution must not be used where any calcareous texture, such as shell or bone, forms part of the preparation; and one of Mr. Goadby's other solutions (8 oz. of bay salt and 2 grs. of corrosive sublimate, to a quart of water,—or, in cases where the coagulating action of corrosive sublimate on albuminous matters would be an objection, the substitution of 20 grains of arsenious acid) may be used in its stead; or Thwaites's fluid, or Dr. Beale's modification of it, or Deane's Gelatine may be tried. It is often quite impossible to predicate beforehand what preservative fluid will answer best for a particular kind of preparation; and it is consequently desirable, where there is no lack of material, always to mount the same object in two or three different ways, marking on each slide the method

employed, and comparing the specimens from time to time, so as to judge how each is affected.

132. *Of Mounting Objects in Fluid.*—As a general rule, it is desirable that objects which are to be mounted in fluid, should be soaked in the particular fluid to be employed, for some little time before mounting; since, if this precaution be not taken, air-bubbles are very apt to present themselves. It is sometimes necessary, in order to secure the displacement of air contained in the specimen, to employ the air-pump in the mode already directed (§ 128); but it will sometimes be found sufficient to immerse the specimen for a few minutes in alcohol (provided that this does not do any detriment to its tissues), which will often penetrate where water will not make its way; and when the spirit has driven out the air, the specimen may be removed back to water, which will gradually displace the spirit. When Deane's Gelatine is used, however, all that can be done, will be to drain the object of its superfluous water before applying the liquefied medium; but as air-bubbles are extremely apt to arise, they must be removed by means of the air-pump, the gelatine being kept in a liquid state by the use of a vessel of hot water, as in the case of Canada balsam. In dealing with the small quantities of fluid required in mounting microscopic objects, it is essential for the operator to be provided with the means of transferring very small quantities from the vessel containing it, to the slide, as well as of taking up from the slide what may be lying superfluous upon it. The straight and curved-pointed "dipping-tubes" (Fig. 51, A, B) may be made to answer this purpose; but it is much better that tubes for this purpose be furnished with a bulb, like that of the Chemist's "pipette," and that their orifices be drawn to a fine point. The fluid is drawn into the tubes by suction, and expelled by the pressure of the breath; the curved-pointed tube will generally be the best for introducing fluid beneath the glass cover, and the straight-pointed for simply filling cells or for taking up superfluous fluid. The Author has of late found very great convenience in the use of a small glass syringe, the orifice of which is slightly curved and drawn to a fine capillary point; for as the syringe works independently of the mouth, its orifice may be applied in any way that may be found convenient; and when the mouth is freed from the efforts of suction and ejection, the eyes can be better employed in watching the operation. Besides the pipettes and the syringe, some blotting-paper, of the most bibulous kind that can be procured, will be found very useful.

133. There are certain objects of extreme thinness, which require no other provision for mounting them in fluid, than an ordinary glass slide, a thin glass cover, and some gold-size or asphalte (§ 120). The object having been laid in its place, and a drop of the fluid laid upon it (care being taken that no air-space remains beneath the under side of the object and the sur-

face of the slide), the glass cover is then to be laid upon it, one side being first brought into contact with the slide, and the other being gradually lowered, in such a manner that the air shall be all displaced before the fluid. If any air-bubbles remain in the central part of the space between the cover and the slide, the former must be raised again, and more fluid should be introduced; but if the bubbles be near the edge, a slight pressure on that part of the cover will often suffice to expel them, or the cover may be a little shifted so as to bring them to its edge. There are some objects, however, whose parts are liable to be displaced by the slightest shifting of this kind; and it is more easy to avoid making air-bubbles, by watching the extension of the fluid as the cover is lowered, and by introducing an additional supply when and where it may be needed, than it is to get rid of them afterwards without injury to the object. When this end has been satisfactorily accomplished, all that is needed is first to remove all superfluous fluid from the surface of the slide, and from *around* the edge of the cover, with a piece of blotting-paper, taking care not to draw away any of the fluid from *beneath* the cover, or (if any have been removed accidentally) to replace what may be deficient; and then to make a circle of asphalte or gold-size around the cover, taking care that it "wets" its edges, and advances a little way upon its upper surface. When this first coat is dry, another should be applied, particular care being taken that the cement shall fill the angular furrow at the margin of the cover. In laying on the second coat, it will be convenient, if the cover be round, to make use of the whirling-table (Fig. 63); and if the slide be so carefully laid upon it, that the glass cover is exactly concentric with its axis, the whirling-table may be used even for the first application of the varnish; a slight error in this respect, however, may occasion the displacement of the cover. By far the greater number of preparations which are to be preserved in liquid, however, should be mounted in a "Cell" of some kind, which forms a *well* of suitable depth, wherein the preservative liquid may be retained. This is *absolutely necessary* in the case of all objects, whose thickness is such as to prevent the glass cover from coming into close approximation with the slide; and it is *desirable*, wherever that approximation is not such as to cause the cover to be drawn to the glass slide by capillary attraction, or wherever the cover is *sensibly* kept apart from the slide by the thickness of any portion of the object. Hence it is only in the case of objects of the most extreme tenuity, that the "cell" can be advantageously dispensed with;—the danger of *not* employing it, in many cases in which there is no difficulty in mounting the object without it, being that after a time the cement is apt to run in beneath the cover, which process is pretty sure to continue, when it may have once commenced.

134. *Cement-Cells*.—When the Cells are required for mounting very thin objects, they may be advantageously made of varnish

only, by the use of the ingenious little instrument (Fig. 63) contrived by Mr. Shadbolt. This consists of a small slab of ma-

FIG 63.



Shadbolt's Turn-table for making Cement-Cells.

hogany, into one end of which is fixed a pivot, whereon a circular turn-table of brass, about three inches in diameter, is made to rotate easily, a rapid motion being given to it by the application of the fore-finger to the

milled head seen beneath. The glass slide being laid upon the turn-table, in such a manner that its two edges shall be equidistant from the centre (a guide to which is afforded by a circle of an inch in diameter, traced upon the brass), and being held by the springs with which it is furnished, a camel-hair pencil dipped in the varnish to be used (Brunswick black or Asphalte is the best) is held in the right hand, so that its point comes into contact with the glass, a little within the guiding circle just named. The turn-table being then put into rotation with the left hand, a ring of varnish of suitable breadth is made upon the glass; and if the slide be set aside in a horizontal position, this ring will be found, when dry, to have lost the little inequalities it may have at first presented, and to present a very level surface. If a greater thickness be desired than a single application will conveniently make, a second layer may be laid on after the first is dry. It is convenient to prepare a number of these cells at once, since, when "the hand is in," they will be made more dexterously than when the operation is performed only once; and it will be advantageous to subject them to the warmth of a slightly heated oven, whereby the flattening of their surface will be more completely assured. The Microscopist will find it a matter of great convenience to have a stock of these cells always by him, ready prepared for use.

135. *Thin Glass Cells*.—For the reception of objects too thick for varnish cells, but not thicker than ordinary thin glass, it is advantageous to construct cells of glass; and these may be made in one of two ways, either by grinding down the cross sections of glass tubes (§ 137) until they have been reduced to the desired thinness, or by perforating a plate of thin glass with an aperture of the desired size; and then cementing the ring or the plate to the glass slide with marine glue. The former plan is liable to the objection, that in reducing the glass rings to the desired thinness, they are extremely liable to crack or break, and that their attainable forms are limited. The latter will generally answer very well, if care be taken in the selection of a *flat* piece of thin glass; and the perforation, if due precaution be employed, may be made of any size or form that may be desired. For making *round* cells, the perforated pieces that sometimes re-

main entire after the cutting of disks (§ 117) may be employed, the disks often falling out of themselves when the glass is laid aside for a few days; and thus the same piece of thin glass may afford a plate, which, when cemented to a glass slide forms a cell, and a disk suitable as the cover to a cell of somewhat smaller size. There is great danger, however, of the cracking of the surrounding glass, especially when the disk is of large size; and it will generally be found a saving of trouble, to employ the method recommended by Dr. L. Beale. This consists in attaching a piece of thin glass to one of the glass rings of which the deeper cells are made (§ 137), of any form that may be desired, by means of marine glue, first laid upon the latter, and melted upon the hot plate; when the glue is quite cold, the point of a round or semicircular file is sharply thrust through the centre of the thin glass, which is carefully filed to the size of the interior of the ring; and the ring being then heated a second time on the hot plate, the thin glass plate may be readily detached from it, and at once cemented upon the glass slide. The success of this simple process depends upon the very firm and intimate adhesion of the thin glass to the ring, which prevents any crack from running into the part of the thin glass that is attached to it, however roughly the file may be used. By having many of the rings on the hot plate at once, and operating with them in turn, a great number of cells can be made in a short time; and such large thin cells may be made in this mode, as could scarcely be fabricated (on account of the extreme brittleness of this glass) by any other. A *press*, consisting of two plates of brass screwed together, holding the thin glass between them, has been devised by Mr. C. Brooke for the same purpose; but the foregoing method has the advantage, not only of requiring no special apparatus, but also of enabling the form and size of the perforation to be readily varied. After the thin glass has been cemented to the slide, it is desirable to roughen its upper surface, by rubbing it upon a leaden or pewter plate (§ 108) with fine emery; since the gold-size or other varnish adheres much more firmly to a "ground" than to a polished surface. Although the thin glass cell requires much more trouble in its preparation than the cement cell, yet it is decidedly to be preferred for any very choice objects; since, if any air should find admission, it is more readily detected; and the remounting of the object may be accomplished in the same cell, with very little disturbance of its position.

136. *Plate Glass and Shallow Cells.*—For mounting objects of somewhat greater thickness than can be included within thin glass cells, *shallow* cells may be made by drilling apertures of the desired size in pieces of plate-glass of the requisite thickness, and by attaching these with marine glue to glass slides (Fig. 64). Such holes may be made not merely circular (A) but oval (C); and a very elongated perforation may be made, by drilling two

holes at the required distance and then connecting them by cutting out the intermediate space (B). These operations, however, can scarcely be performed by any but regular glass-cutters, and, being troublesome, are expensive; hence the plate-glass cells have been generally superseded, either by *tube-cells* (§ 137) or by

built-up cells. Although the former may be reduced to any degree of shallowness that may be desired, and are made of most of the sizes and forms that can be ordinarily needed, yet for extra sizes or peculiar forms, shallow cells may be easily built up after the following very simple and effective method. A piece of plate glass, of a thickness that shall give the desired depth to the cell, is to be cut to the dimensions of its outside wall; and a strip is then to be cut off with the diamond from each of its edges, of such breadth as shall leave the interior piece equal in its dimensions to the cavity of the cell, that is desired. This piece being rejected, the four strips are then to be cemented upon the glass slide in their original position, so that the diamond cuts shall fit together with the most exact precision; and the upper surface is then to be ground flat with emery upon the pewter plate, and left rough as before. This plan answers admirably for constructing such large shallow cells as are required for the mounting of Zoophytes and similar objects.




Plate Glass Cells.


Having had occasion, during the last few months, to mount a large number of objects in shallow cells, the Author has adopted the recommendation of a friend, to make use of cells which are sunk by grinding out a concave in the thickness of a glass plate. These, until recently, were costly; but they are now made in large quantities, and their price has been so much reduced, that they can be obtained more cheaply than any other kind. For objects whose shape adapts them to the form and depth of the concavity, these cells will be found peculiarly advantageous; since they do not hold air-bubbles so tenaciously as do those with perpendicular walls; and there is no cemented plate or ring to be loosened from its attachment, either by a sudden jar or by the lapse of time. For transparent objects, however, they are less suitable (unless manufactured with more care than is usually given to them) than they are for opaque; since the concave bottom is seldom so highly polished, as to be free from scratches and rough-

nesses, which greatly interfere with the appearance of the picture. Cells of this kind may be obtained, from Messrs. Jackson, Oxford Street, either of round or oval form, and not only ground out of slides of the usual size (3 in. by 1) and thickness, but also hollowed in pieces of plate-glass of larger dimensions.

137. *Deep and Built-up Cells.*—The *deep* cells which are required for mounting Injections and other microscopic preparations of considerable size and thickness, *may* be made by drilling through a piece of thick plate glass (Fig. 64, D); but for the reason already given, the drilled cells are now seldom used, their place having been taken, either by tube-cells, or by the deep built-up cells to be presently described. The *tube-cells* are made by cutting transverse sections of thick-walled glass tubes of the required size, grinding the surfaces of these rings to the desired thinness, and then cementing them to the glass slides with marine glue. Not only may *round* cells (Fig. 65, A, B), of any diameter and any depth that the Microscopist can possibly require, be made by this


simple method, but oval, square-shaped, or oblong cells (C, D) are now made, of the forms and sizes that he is most likely to want, by flattening the round glass-tube whilst hot, or by blowing it within a mould. The facility with which such cells may be made, and the security they afford, have caused the deep cells *built up* of separate strips of glass (Fig. 66) to be comparatively little employed, except in cases where some very unusual size or shape (A) may be required, or where it is necessary that not merely the top and bottom, but also the sides of the object, should

be clearly seen (B). The perfect construction of these requires a nicety of workmanship which few amateurs possess, and the expenditure of more time than Microscopists generally have to spare; and as it is consequently preferable to obtain them ready made, directions for making them need not here be given. A new plan of making deep cells, however, has been lately introduced by Dr. L. Beale; which, though it does not give them side walls possessing the same flatness with those of the built-up cells, adapts them to serve most of the purposes for which these are required, and makes them more secure against leakage; whilst it has the advantage of being so easy and simple, that



Tube-Cells, Round and Quadrangular.

any one may put it into practice. A long strip of plate glass is to be taken, whose breadth is equal to the desired depth of the cell, and whose length must be



equal to the sum of that of all its sides. This strip is to be carefully bent to a right angle in the blow-pipe flame, at three points previously indicated by marks so placed as to show where the angles should fall; and the two ends, which will thus be brought into contact at right angles, are to be fused together. Thus a large square well, slightly rounded at the

angles, will be formed; and this, being very brittle, should be allowed to cool very gradually, or, still better, should be annealed in an oven. It must then be ground quite true on its upper and lower edges, either on the lead plate with emery, or on a flat stone with fine sand; and it may then be cemented to a glass slide in the usual way.

138. *Mounting Objects in Cells.*—In mounting an object in a cell, the first attention will of course be given to the cleanness of the interior of the cell, and of the glass cover which is to be placed on it; this having been secured, the cell is to be filled with fluid by the pipette or syringe; and any minute air-bubbles which may be seen adhering to its bottom or sides, must be removed by the needle: the object, previously soaked in fluid resembling that with which the cell is filled, is then to be placed in the cell, and should be carefully examined for air-bubbles on all sides, and also by looking up from below. When every precaution has been taken to free it from these troublesome intruders, the cover may be placed on the cell, one side being first brought down upon its edge and then the other; and if the cell have been previously brimming over with fluid (as it ought to be) it is not likely that any air-space will remain. If, however, any bubbles should present themselves beneath the cover, the slide should be inclined, so as to cause them to rise towards the highest part of its circumference, and the cover slipped away from that part, so as to admit of the introduction of a little additional fluid by the pipette or syringe; and when this has taken the place of the air-bubbles, the cover may be slipped back into its place.¹ All superfluous fluid is then to be taken

¹ Mr. Quekett and some other practised manipulators recommend that the edges of the cell and that of the disk of glass be smeared with the gold-size or other varnish employed, before the cell is filled with fluid; but the Author has found this practice objectionable, for two reasons.—first, because it prevents the cover from being slipped to one side (which is often desirable), without its being soiled by the varnish,—and second, because when the edge of the cell has been thus made to “take” the varnish, that which is afterwards applied for the closure of the cell is more likely to run in, than if the whole of the surface covered by the glass is moistened with water.

up with blotting-paper; and particular care should be taken thoroughly to dry the surface of the cell and the edge of the cover, since the varnish will not hold to them if they be in the least damp with water. Care must also be taken, however, that the fluid be not drawn away from between the cover and the edge of the cell on which it rests; since any deficiency here is sure to be filled by varnish, the running in of which is particularly objectionable. These minutiae having been attended to, the closure of the cell may be at once effected, by carrying a thin layer of gold-size or asphalte around and upon the edge of the glass cover, taking care that it touches every point of it, and fills the angular channel which is left around its margin. If the wall of the cell be very thin, it will be advantageous to include it in the ring of varnish; so that this shall hold down the cover, not only on the cell, but on the slide beneath. The Author has found it advantageous, however, to delay closing the cell for some little time after the superfluous fluid has been drawn off; for as soon as evaporation beneath the edges of the cover begins to diminish the quantity of fluid in the cell, air-bubbles often begin to make their appearance, which were previously hidden in the recesses of the object; and in the course of half an hour, a considerable number are often collected. The cover should then be slipped aside, fresh fluid be introduced, the air-bubbles removed, and the cover put on again; and this operation should be repeated, until it fails to draw forth any more air-bubbles. It will, of course, be observed, that, if the evaporation of fluid should proceed far, air-bubbles will *enter* beneath the cover; but these will show themselves on the *surface* of the fluid; whereas those which arise from the object itself, are found in the deeper parts of the cell. Much time may be saved, however, and the freedom of the preparation from air-bubbles may be most effectually secured, by placing the cell, after it has been filled in the first instance, in the vacuum of an air-pump; and if several objects are being mounted at once, they may all be subjected to the exhausting process at the same time. The application of the varnish should be repeated after the lapse of a few hours, and may be again renewed with advantage several times in the course of a week or two; care being taken that each layer covers the edges, as well as the whole surface, of that which preceded it.

189. The presence of air-bubbles, in any preparation mounted in fluid, is to be particularly avoided, not merely on account of its interference with the view of the object, but also because, when air-spaces, however small, once exist, they are almost certain to increase, until at last they take the place of the entire fluid, and the object remains dry. Even with the most experienced manipulators, however, this misfortune not unfrequently occurs; being sometimes due to the obstinate entanglement of air-bubbles in the object, when it was originally mounted;

and sometimes to the perviousness of some part of the cement, which has allowed a portion of the contained fluid to escape, and air to find admission. In either case, so soon as an air-bubble is seen in such a preparation, the attempt should be made to prevent its increase, by laying on an additional coat of varnish; but if this should not be successful, the cover should be taken off, and the specimen remounted, so soon as the fluid has escaped to such a degree as to leave any considerable portion of it uncovered.

140. *Importance of Cleanliness.*—The success of the result of any of the foregoing operations is greatly detracted from, if, in consequence of the adhesion of foreign substances to the glasses whereon the objects are mounted, or to the implements used in the manipulations, any extraneous particles are brought into view with the object itself. Some such will occasionally present themselves, even under careful management; especially fibres of silk, wool, cotton, or linen, from the handkerchiefs, &c., with which the glass slides may have been wiped, and grains of starch, which often remain obstinately adherent to the thin glass covers kept in it. But a careless and uncleanly manipulator will allow his objects to contract many other impurities than these; and especially to be contaminated by particles of dust floating through the air, the access of which may be readily prevented by proper precautions. It is desirable to have at hand a well-closed cupboard furnished with shelves, or a cabinet of well-fitted drawers, or a number of bell-glasses upon a flat table, for the purpose of securing our glasses, objects, &c., from this contamination, in the intervals of the work of preparation; and the more readily accessible these receptacles are, the more use will the Microscopist be likely to make of them. Great care ought, of course, to be taken, that the liquids employed for mounting should be freed, by effectual filtration, from all floating particles; and both these and the Canada balsam should be kept in well-closed bottles.

141. *Labelling and Preserving of Objects.*—Whenever the mounting of an object has been completed, its name ought to be at once marked on it, and the slide should be put away in its appropriate place. Some inscribe the name on the glass itself, with a writing diamond; whilst others prefer to gum a label¹ on the slide; and others, again, cover one or both surfaces of the slide with colored paper, and attach the label to it. In the case of objects mounted dry or in balsam, the latter method has the advantage of rendering the glass cover more secure from displacement by a slight blow or “jar,” when the varnish or balsam may have become brittle by the lapse of years. Instead, however, of attaching the white label on which the name of the object is written, outside the colored paper with which the slide is

¹ Very neat gummed labels, of various sizes and patterns suitable to the wants of the Microscopist, are sold by the “Drapers’ Stationers” in the City.

covered, it is better to attach the label to the glass, and to punch a hole out of the colored paper, sufficiently large to show the name, in the part corresponding to it; in this manner the label is prevented from falling off, which it frequently does when attached to the glass without protection, or to the outside of the paper cover. When objects are mounted in fluid, either with or without cells, paper coverings to the slides had better be dispensed with; and besides the name of the object, it is desirable to inscribe on the glass that of the fluid in which it is mounted. For the preservation of objects, the pasteboard boxes now made at a very reasonable cost, with wooden racks, to contain 6, 12, or 24 slides, will be found extremely useful. In these, however, the slides must always stand upon their edges; a position which, besides interfering with that ready view of them which is required for the immediate selection of any particular specimen, is unfavorable to the continued soundness of preparations mounted in fluid. Although such boxes are most useful, indeed almost indispensable, to the Microscopist, for holding slides which he desires (for whatever purpose) to keep for a while constantly at hand, yet his regularly classified series is much more conveniently stored in a Cabinet containing numerous very shallow drawers, in which they lie flat and exposed to view. Such cabinets are now prepared for sale under the direction of our principal Opticians, with all the improvements that experience has suggested. In order to prevent the warping of the thin wood of which the bottoms of the drawers are usually made, whereby their sliding action is obstructed, it has been found advantageous to substitute strained canvas or papier mache. Again, in order to antagonize the disposition of the slides to slip one over another in the opening or shutting of the drawers, it has been found preferable to arrange them in such a manner, that they lie with their ends (instead of their long sides) towards the front of the drawer, and to interpose a cross-strip of wood, lying parallel to the front of the drawer, between each row. It is very convenient, moreover, for the front of the drawer to be furnished with a little tablet of porcelain, on which the name of the group of objects it may contain can be written in pencil, so as to be readily rubbed out; or a small frame may be attached to it, into which a slip of card may be inserted for the same purpose.

SECTION 3. COLLECTION OF OBJECTS.

142. A large proportion of the objects with which the Microscopist is concerned, are derived from the minute parts of those larger organisms, whether Vegetable or Animal, the collection of which does not require any other methods than those pursued by the ordinary Naturalist. With regard to such, therefore, no special directions are required. But there are several most interesting and important groups, both of Plants and Animals,

which are themselves, on account of their minuteness, essentially microscopic; and the collection of these requires peculiar methods and implements, which are, however, very simple,—the chief element of success lying in the knowledge *where* to look, and *what* to look for. In the present place, *general* directions only will be given; the particular details relating to the several groups, being reserved for the account to be hereafter given of each.

143. All the Microscopic organisms in question, being aquatic, must be sought for in pools, ditches, streams, or other collections of water; through which some of them freely move, whilst others attach themselves to the stems and leaves of aquatic plants, or even to pieces of stick or decaying leaves, &c., that may be floating on the surface or submerged beneath it, while others, again, are to be sought for in the muddy sediments at the bottom. Of those which have the power of free motion, some keep near the surface, whilst others swim in the deeper waters; but the situation of many depends entirely upon the light, since they rise to the surface in sunshine, and subside again afterwards. The Collector will therefore require a means of obtaining samples of water at different depths, and of drawing to himself portions of the larger bodies to which the microscopic organisms may be attached. For these purposes, nothing is so convenient as a rod about five feet long, which may be divided into two pieces jointed together; and the farther extremity of this rod should be pierced with a hole, passing for some distance into its length.¹ Into this hole, as a socket, may be fitted either of the three implements which the Collector may happen to require. If he desires to take up samples of the water, he will need a wide-mouthed bottle, containing about 2 oz. This may be attached to the extremity of the rod, by simply passing round its neck a strap of thin whale-bone or sheet gutta percha, the two ends of which are to be brought together and inserted into the socket, in which they may be secured by a plug of soft wood or cork. The bottle being held sideways with its mouth partly below the water, the surface may be skimmed; or, if it be desired to bring up a sample of the liquid from below, or to draw into the bottle any bodies that may be loosely attached to the submerged plants, the bottle is to be plunged into the water with its mouth downwards, carried into the situation in which it is desired that it should be filled, and then suddenly turned with its mouth upwards. If, again, the organisms which it may be desired to collect, are of sufficient size to be strained out of the water by a piece of fine muslin, a ring-net should be fitted into the socket of the rod. This may be made by sewing the muslin bag to a ring of stout wire, furnished with a projecting stem which may be inserted by means of a cork into the socket of the rod. But it is more convenient

¹ Cheap fishing-rods are now sold at the toy shops, which answer this purpose extremely well, the last or slenderest joint being laid on one side; its socket in the last joint but one, being well adapted to receive the fittings above described.

that the muslin should be made removable; and this may be provided for (as suggested in the "Micrographic Dictionary," Introduction, p. xxiv) by the substitution of a wooden ring, grooved on its outside, for the wire ring; the muslin being strained upon it by a ring of vulcanized India rubber, which lies in the groove, and which may be readily slipped off and on, so as to allow a fresh piece of muslin to be put in the place of that which has been last used. For bringing up portions of larger Plants, either for the sake of examining their own structure, or for obtaining the growths which may be parasitic upon them, a cutting-hook, shaped somewhat like a sickle, may be fitted into the socket of the rod.

144. The Collector should also be furnished with a number of bottles, into which he may transfer the samples thus obtained. These it will be convenient to have of two kinds; one set wide-mouthed, and capable of being closely corked, for minute Plants; the other set with narrower mouths, having short pieces of tube passed through the corks, for the purpose of containing Animalcules without depriving them of air. The former kind, however, may be safely employed for Animalcules, if they be not above two-thirds filled (so as to leave an adequate air-space), and be not kept long closed. Such bottles should be fitted into cases, in which several may be carried at once without risk of breakage.¹ Whilst engaged in the search for Microscopic objects, it is desirable for the collector to possess a means of at once recognizing the forms which he may gather, where this is possible, in order that he may decide whether the "gathering" is, or is not, worth preserving; for this purpose either a powerful "Coddington" or "Stanhope" lens (§ 19), or a Gairdner's Simple Microscope (§ 28), will be found most useful, according to the class of objects of which the collector is in search. The first will answer very well for Zoophytes and the larger Diatomaceæ; but the second or third will be needed for Desmidiaceæ, the smaller Diatomaceæ, and Animalcules.

¹ The bottles in which smelling-salts are now commonly sold, having the corks fitted into disks of turned wood, are very convenient, both in size and shape, for the purposes of the Microscopist; cases containing 3, 4, 6, or 8 such bottles, are made by Mr. Ferguson, of Giltspur Street. The wide-mouthed bottles with screw caps, made by the York Glass Company, are also extremely convenient.

CHAPTER VI.

MICROSCOPIC FORMS OF VEGETABLE LIFE.—PROTOPHYTES.

145. IN commencing our survey of these wonders and beauties of Life and Organization, which are revealed to us by the assistance of the Microscope, it seems on every account the most appropriate to turn our attention in the first instance to the Vegetable Kingdom; and to begin with those humblest members of that kingdom, whose form and structure, and whose very existence, in many cases, are only known to us through its use. For those who desire to make themselves familiar with microscopic appearances, and to acquire dexterity in microscopic manipulation, cannot do better than educate themselves by the study of those comparatively simple forms of organization, which the Vegetable fabric presents; since a facility in minute dissection and in microscopic analysis may be thus acquired, which will save much expenditure of time and labor, that might be unprofitably applied, without such apprenticeship, to the attempt to unravel the complexities of Animal organization. But further, the scientific Histologist (p. 49) looks to the careful study of the structure of the simplest forms of Vegetation, as furnishing the key (so to speak) that opens the right entrance to the study of the elementary Organization, not merely of the higher Plants, but of the highest Animals. And in like manner, the scientific Physiologist looks to the complete knowledge of their life history, as furnishing the surest basis for those general notions of the nature of Vital Action, which the advance of science has shown to be really well founded, only when they prove equally applicable to both kingdoms. But further, a peculiar interest attaches itself at the present time, to everything which throws light upon the debated question of the boundary between the two kingdoms; a question which is not less keenly debated among Naturalists, than that of many a disputed frontier has been between adjacent Nations. For many parts of this border country have been taken and retaken several times; their inhabitants (so to speak) having first been considered, on account of their general appearance, to belong to the Vegetable Kingdom,—then, in consequence of some movements being observed in them, being claimed by the Zoologists,—then, on the ground of their

evidently plant-like mode of growth, being transferred back to the Botanical side,—then, owing to the supposed detection of some new feature in their structure or physiology, being again claimed as members of the Animal Kingdom,—and lastly, on the discovery of a fallacy in these arguments, being once more laid hold of by the Botanist, with whom, for the most part, they now remain. For the attention which has been given of late years to the study of the humblest forms of Vegetation, has led to the knowledge of so many phenomena, among what must be *undoubtedly* regarded as Plants, which would formerly have been considered unquestionable marks of animality, that the discovery of the like phenomena among the doubtful beings in question, so far from being any evidence of *their* animality, really affords a probability of the opposite kind.

146. In the present state of Science, it would be very difficult, and is perhaps impossible, to lay down any definite line of demarcation between the two kingdoms; since there is no single character by which the Animal or Vegetable nature of any organism can be tested. Probably the one which is most generally applicable, among those lowest organisms which most closely approximate to one another, is—not, as formerly supposed, the presence or absence of spontaneous motion, but the dependence of the being for nutriment upon *organic compounds* already formed, which it takes (in some way or other) into the *interior* of its body, or its possession of the power of obtaining its own alimentary matter by absorption, from the *inorganic elements* on its *exterior*. The former is the characteristic of the *Animal Kingdom* as a whole; the latter is the attribute of the *Vegetable*; and although certain apparently exceptional cases *may* exist, yet these do not seem to occur among the group in which such a means of distinction is most useful to us. For we shall find that those *Protozoa*, or simplest Animals, which seem to be composed of nothing else than a mass of living jelly (Chaps. IX, X), are supported as exclusively, either upon other *Protozoa*, or upon *Protophyta*, which are humble Plants of equal simplicity, as the highest Animals are upon the flesh of other animals, or upon the products of the Vegetable Kingdom; whilst these *Protophytes*, in common with the highest Plants, draw *their* nourishment from water, carbonic acid, and ammonia, and are distinguished by their power of liberating oxygen, through the decomposition of carbonic acid, under the influence of sunlight. And we shall, moreover, find, that even such *Protozoa* as have neither stomach nor mouth, receive their alimentary matter direct into the very substance of their bodies, in which it undergoes a kind of digestion; whilst the *Protophyta* absorb through their external surface only, and take in no solid particles of any description. With regard to motion, which was formerly considered the distinctive attribute of animality, we now know, not merely that many *Protophytes* (perhaps all, at some period or

other of their lives) possess a power of spontaneous movement, but also that the instruments of motion, when these can be discovered, are of the very same character in the Plant as in the Animal; being little hair-like filaments termed *cilia* (from the Latin *cilium*, an eyelash), by whose rhythmical vibration the body of which they form part is propelled in definite directions. The peculiar contractility of these cilia cannot be accounted for in either case, any better than in the other; all we can say is, that it seems to depend upon the continued vital activity of the living substance of which these filaments are prolongations; and that this contractile substance has a composition essentially the same in the Plant as in the Animal.

147. The plan of organization throughout the Vegetable kingdom presents this remarkable feature of uniformity,—that the fabric of the highest and most complicated Plants, consists of nothing else than an aggregation of the bodies termed *cells*, every one of which, amongst the lowest and simplest forms of Vegetation, may maintain an independent existence, and may multiply itself most indefinitely, so as to form vast assemblages of similar bodies. And the essential difference between the plans of structure in the two cases lies in this,—that the cells produced by the self-multiplication of the primordial cell of the Protophyte, are all mere repetitions of it and of one another, each living *by* and *for* itself,—whilst those produced by the like self-multiplication of the primordial cell in the Oak or Palm, not only remain in mutual connection, but undergo a progressive “differentiation,” a fabric being thereby developed, which is composed of a number of distinct organs (stem, leaves, roots, flowers, &c.), each of them characterized by specialities not merely of external form but of intimate structure (the ordinary type of the cell undergoing various modifications, to be described in their proper place, Chap. VIII), and performing actions peculiar to itself, which contribute to the life of the Plant *as a whole*. Hence, as was first definitely stated by Schleiden (see Introduction, p. 43), it is in the *life-history of the individual cell*, that we find the true basis of the study of Vegetable Life in general. And we shall now inquire, therefore, what information on this point we derive from Microscopic research. In its most completely developed form, the Vegetable Cell may be considered as a closed membranous bag or vesicle, containing a fluid cell-sap; and thus we have to consider separately the *cell-wall* and the *cell-contents*. The “cell-wall” is composed of two layers, of very different composition and properties. The *inner* of these, which has received the name of *primordial utricle*, appears to be the one first formed, and most essential to the existence of the cell; it is extremely thin and delicate, so that it escapes attention so long as it remains in contact with the external layer; and it is only brought into view when separated from this, either by developmental changes (Fig. 107), or by the influence of reagents which

cause it to contract by drawing forth part of its contents (Fig. 175). Its composition is indicated, by the effects of reagents, to be *albuminous*; that is, it agrees with the formative substance of the Animal tissues, not only in the proportions of oxygen, hydrogen, carbon, and nitrogen which it contains, but also in the nature of the compound formed by the union of these elements. The external layer, on the other hand, though commonly regarded as the proper "cell-wall," is generated on the surface of the primordial utricle, after the latter has completely inclosed the cavity and its contents, so that it takes no essential part in the formation of the cell. It is usually thick and strong in comparison with the other, and may often be shown to consist of several layers. In its chemical nature it is altogether dissimilar to the primordial utricle; for it is essentially composed of *cellulose*, a substance containing no nitrogen, and nearly identical with starch. The relative offices of these two membranes are very different; for whilst there are many indications that the primordial utricle continues to participate actively in the vital operations of the cell, it seems certain that the cellulose-wall takes no concern in them, but is only their product; its function being simply protective. The contents of the vegetable-cell, being usually more or less deeply colored, have received the collective designation of *endochrome* (or internal coloring substance); and they essentially consist of a layer of colorless "protoplasm" (or organizable fluid, containing albuminous matter in combination with dextrine or starch-gum) in immediate contact with the primordial utricle, within which is the more watery cell-sap, particles of chlorophyll or coloring substance being diffused through both, or through the former only.

148. But although these component parts may be made out without any difficulty in a large proportion of Vegetable cells, yet they cannot be distinguished in some of those humble organisms, which are nearest to the border ground between the two kingdoms. For in them we find the "cell-wall" very imperfectly differentiated from the "cell-contents;" the former not having by any means the firmness of a perfect membrane, and the latter not possessing the liquidity which elsewhere characterizes them. And in some instances, the cell appears to be represented only by a mass of endochrome, so viscid as to retain its external form without any liminary membrane, though the superficial layer seems to have a firmer consistence than the interior substance; and this may or may not be surrounded by a gelatinous-looking envelope, which is equally far from possessing a membranous firmness, and yet is the only representative of the cellulose-wall. This viscid endochrome consists, as elsewhere, of a colorless protoplasm, through which coloring particles are diffused, sometimes uniformly, sometimes in local aggregations, leaving parts of the protoplasm uncolored. The superficial layer, in particular, is frequently destitute of color; and the "primordial utricle" ap-

pears to be formed by its solidification. In the interior of the viscid mass, are commonly found *vacuoles*, which are distinguished from the surrounding substance by their difference in refracting power; these, however, are not usually void spaces, but are cavities in the protoplasm occupied by fluid of a more watery consistence; and this "vacuolation" of the interior, which increases until the cell-contents have almost entirely lost their original viscosity and are of a more watery character, seems to take place *pari passu* with the consolidation of the exterior into distinct membranous walls; so that the development of a perfect cell out of a rudimentary mass of endochrome, may be stated to consist essentially in the *gradual differentiation* of its substance, which was at first a nearly homogeneous viscid mass, into the solid cell-wall and the liquid cell-contents. It is interesting to observe, at the very outset of our inquiry into the nature of Organization and Vital action, so characteristic an illustration of the great law of Von Bär, already referred to (pp. 52, 55).

149. Now among the *Protophytes* or simplest Plants, on the examination of which we are about to enter, there are many, of which every single cell is not only capable of living in a state of isolation from the rest, but even normally does so; and thus, in the ordinary phraseology, every cell is to be accounted a "distinct individual." There are others, again, of which shapeless masses are made up by the aggregation of contiguous cells, which, though quite capable of living independently, remain attached to each other by the mutual fusion (so to speak) of their gelatinous investments. And there are others, moreover, in which a definite adhesion exists between the cells, and in which regular plant-like structures are thus formed, notwithstanding that every cell is still but a repetition of every other, and is capable of living independently if detached, so as to answer to the designation of a "unicellular" or single-celled plant. These different conditions we shall find to arise out of the mode in which each particular species multiplies by binary subdivision (§ 150): for where the pair of cells that is produced by the segmentation of the previous cell, undergo a *complete* separation from one another, they will henceforth live quite separately; but if, instead of undergoing this complete fusion, they should be held together by the intervening gelatinous envelope, a shapeless mass results from repeated subdivisions not taking place on any determinate plan; and if, moreover, the binary subdivision should always take place in a determinate direction, a long narrow filament (Fig. 104, d), or a broad flat leaf-like expansion (Fig. 104, e), may be generated. To such extended fabrics, the term "unicellular plants" can scarcely be applied with propriety, since they may be built up of many thousands or millions of distinct cells, which have no disposition to separate from each other spontaneously. Still they correspond with those which are strictly unicellular, in the absence of any differentiation, either in structure or in

actions, between their component cells; each one of these being a repetition of the rest, and no relation of mutual dependence existing among them. All such organisms may well be included under the general term of *Protophytes*, by which it is convenient to designate the primitive or elementary forms of Vegetation; and we shall now enter, in such detail as the nature of the present Treatise allows, into the history of those forms of the group, which present most of interest to the Microscopist, or which best serve to illustrate the general doctrines of Physiology.

150. The life-history of one of these Unicellular Plants, in its most simple form, can scarcely be better exemplified than in the *Palmoglaea macrococca* (Kützing); one of those humble kinds of vegetation which spreads itself as a green slime over damp stones, walls, &c.

When this slime is examined with the microscope, it is found to consist of a multitude of green cells (Fig. 67, A), each surrounded by a gelatinous envelope; the cell, which does not seem to have any distinct membranous wall, is filled with granular particles of a green color; and a "nucleus" may sometimes be distinguished through the midst of these. When treated with tincture of iodine, however, the

green contents of the cell are turned to a brownish hue, and a dark-brown nucleus is distinctly shown (G). Other cells are seen (B), which are considerably elongated, some of them beginning to present a sort of hour-glass contraction across the middle; in these is commencing that curious *multiplication by duplicative subdivision*, which is the mode in which increase nearly always takes place throughout the Vegetable kingdom; and when cells in this condition are treated with tincture of iodine, the nucleus is seen to be undergoing the like elongation and constriction (H). A more advanced state of the process of subdivision is seen at C, in which the constriction has proceeded to the extent of completely cutting off the two halves of the cell, as well as of the nucleus (I) from each other, though they still remain in mutual contact; but in a yet later stage, they are found detached from each other (D), though still included within the same gelatinous

FIG. 67.



Various phases of development of *Palmoglaea macrococca*; A, full-grown cell; B, C, D, E, successive stages of binary subdivision; F, row of cells, produced by succession of subdivisions; G, H, I, cells treated by iodine; K, L, M, cells in conjunction.

envelope. Each new cell then begins to secrete its own gelatinous envelope; so that, by its intervention, the two are usually soon separated from one another (ε). Sometimes, however, this is not the case; the process of subdivision being quickly repeated, before there is time for the production of the gelatinous envelope, so that a series of cells (ϕ), hanging on one to another, is produced. There appears to be no definite limit to this kind of multiplication; and extensive areas may be quickly covered, in circumstances favorable to the growth of the plant, by the products of the duplicative subdivision of one primordial cell. This, however, is simply an act of *Growth*, precisely analogous to that by which any one of the higher forms of Vegetation extends itself, and differing only in this, that the cells produced by each act of cell-subdivision in the present case exactly resemble that from which they sprang; whilst in the case of more highly organized Plants, they gradually become differentiated to a greater or less degree, so that special "organs" are evolved, which take upon themselves dissimilar yet mutually dependent parts, in the economy of the entire organism.

151. The process which represents the *Generation* of the higher Plants is here performed in a manner so simple, that it would not be recognized as such, if we were not able to trace it up through a succession of modes of gradually increasing complexity, until we arrive at the elaborate operations which are concerned in the production and fertilization of the seeds of Flowering Plants. For it consists in nothing else than the reunion or fusion together of any pair of cells (κ),—a process which is termed *Conjugation*; and it is characteristic of this humble plant, and shows how imperfect must be the consistence of its cell-membrane, that this seems to enter into the fusion, no less completely than the cell-contents. The communication is at first usually made by a narrow neck or bridge (κ); but before long it extends through a large part of the contiguous boundaries (L); and at last the two cells are seen to be completely fused into one mass (μ); which is termed the *spore*. Each "spore" thus formed is the "primordial cell" of a *new generation*, into which it evolves itself by successive repetitions of the process of binary subdivision. It is curious to observe, that during the conjugating process, a production of oil-particles takes place in the cells; these at first are small and distant, but gradually become larger and approximate more closely to each other, and at last coalesce so as to form oil-drops of various sizes, the green granular matter disappearing; and the color of the conjugated body changes, with the advance of this process, from green to a light yellowish-brown. When the spore begins to vegetate, on the other hand, producing a pair of new cells by binary subdivision, a converse change takes place; the oil-globules disappear, and green granular matter takes their place. Now this is precisely what occurs in the formation of seed among the higher Plants; for starchy

substances are transformed into oil, which is stored up in the seed for the nutrition of the embryo, and is applied, during germination, to the purposes which are at other times answered by starch or chlorophyll. The *growth* of this little plant appears to be favored by cold and damp; its *generation*, on the other hand, is promoted by heat and dryness; and it is obvious that the spore-cell must be endowed with a greater power of resisting this, than the vegetating plant has, since the species would otherwise be destroyed by every drought.


152. If the preceding sketch really comprehends the whole life-history of the humble plant to which it relates, this history is much more simple than that of other forms of vegetation, which, without appearing to possess an essentially higher structure, present themselves under a much greater variety of forms and conditions. One of the most remarkable of these varieties is the *motile* condition, which seems to be common, in some stage or other of their existence, to a very large proportion of the lower forms of aquatic vegetation; and which usually depends upon the extension of the primordial utricle into one or two thread-like filaments (Fig. 68, H-L), endowed with the power of executing rhythmical contractions, whereby the cell is impelled through the water. As an illustration of this peculiar mode of activity, which was formerly supposed to betoken Animal life, a sketch will be given of the history of a plant, the *Protococcus pluvialis*, which is not uncommon in collections of rain-water,¹ and which, in its motile condition, has been very

¹ The Author had under his own observation, about eight years ago, an extraordinary abundance of what he now feels satisfied must have been this plant, in a rain-water cistern, which had been newly cleaned out. His notice was attracted to it, by seeing the surface of the water covered with a green froth, whenever the sun shone upon it. On examining a portion of this froth under the Microscope, he found that the water was crowded with green cells in active motion; and although the only bodies at all resembling them, of which he could find any description, were the so-called *Animalcules*, constituting the genus *Chlamydomonas* of Prof. Ehrenberg, and very little was known at that time of the "motile" conditions of *Plants* of this description, yet of the vegetable nature of these bodies he could not entertain the smallest doubt. They appeared in freshly collected rain-water, and could not, therefore, be deriving their support from organic matter; under the influence of light, they were obviously decomposing carbonic acid and liberating oxygen, and this influence he found to be essential to the continuance of their growth and development, which took place entirely upon the Vegetative plan. Not many days after the Protophyte first appeared in the water, a few Wheel-Animalcules presented themselves; these fed greedily upon it, and increased so rapidly (the weather being very warm) that they soon became almost as crowded as the cells of the *Protococcus* had been; and it was probably due in part to their voracity, that the plant soon became less abundant, and before long disappeared altogether. Had the Author been then aware of its assumption of the "still" condition, he might have found it at the bottom of the cistern, after it had ceased to present itself at the surface. The account of this Plant given above, is derived from that of Dr. Cohn, in the "Nova Acta Acad. Nat. Curios." (Bonn, 1850), torn. xxii; of which an abstract by Mr. George Bask is contained in the "Botanical and Physiological Memoirs," published by the Ray Society for 1853. This excellent observer states that he kept his plants for observation in little glass vessels, having the form of a truncated cone, about two inches deep, and one inch and a quarter in diameter, with a flat bottom polished on both sides, and filled with water to the depth of from two to three lines. "It was only in vessels of this kind," he says, "that he was able to follow the development of a number of various

commonly regarded as an Animalcule, its different states having been described under several different names.

153. In the first place, the *color* of these cells varies considerably; since, although they are usually *green* at the period of their most active life, they are sometimes *red*; and their red form has received the distinguishing appellation of *Hæmatococcus*. Very commonly the red coloring matter forms only a central mass of greater or less size, having the appearance of a nucleus (as shown in Fig. 68, E); and sometimes it is reduced to a single granular point, which has been erroneously represented by Prof. Ehrenberg as the *eye* of these so-called Animalcules. It is quite certain that the red coloring substance is very nearly related in

FIG. 68.



Various phases of development of *Protococcus pluvialis*:—A, an encysted cell, which has passed into the “still” condition; B, division of a “still” cell into two; C, another mode of division into two, each primordial vesicle having developed a cellulose envelope around itself, whilst yet within the original cell; D, division of an encysted cell into four; E, division of an encysted cell into eight; F, division of an encysted cell into thirty-two segments; G, motile gonidia (zoospores) after their escape from the original cell; H, a primordial utricle, without cellulose envelope, furnished with two cilia; I, a similar primordial utricle, with distinct cellulose envelope, and threads of protoplasm extending towards it; K, an encysted primordial utricle, pointed at both ends, and furnished with two cilia; L, an encysted primordial utricle, of which nearly half is composed of a colorless granular substance, enclosing a red body resembling a nucleus.

its chemical character to the green, and that the one may be converted into the other; though the conditions under which this conversion may take place are not precisely known. In the *still* form of the cell, with which we may commence the history

cells throughout its whole course.” Probably he would have found the glass tube cells represented in Fig. 65, if he had been acquainted with them, to answer his purpose just as well as these specially constructed vessels.

of its life, we find a mass of endochrome, consisting of a colorless protoplasm, through which red or green colored granules are more or less uniformly diffused; on the surface of this endochrome, the colorless protoplasm is condensed into a more consistent layer, forming an imperfect "primordial utricle;" and this is surrounded by a tolerably firm layer, which seems to consist of cellulose or of some modification of it. Outside this (as shown in Fig. 68, A), when the "still" cell is formed by a change in the condition of a cell that has been previously "motile," we find another envelope, which seems to be of the same nature, but which is separated by the interposition of aqueous fluid; this, however, may be altogether wanting. The multiplication of the "still" cells by self-division takes place as in the previous instance; the endochrome, enclosed in its primordial utricle, first undergoing separation into two halves (as seen at B), and each of these halves subsequently developing a cellulose envelope around itself, and undergoing the same division in its turn. Thus 2, 4, 8, 16 new cells are successively produced; and these are sometimes set free by the complete dissolution of the envelope of the original cell; but they are more commonly held together by its transformation into a gelatinous investment, in which they remain imbedded. Sometimes the contents of the primordial utricle subdivide at once into *four* segments (as at D), of which every one forthwith acquires the characters of an independent cell; but this, although an ordinary method of multiplication among the "motile" cells, is comparatively rare in the "still" condition. Sometimes, again, the cell-contents of the "still" form subdivide at once into eight portions, which, being of small size, and endowed with motile power, may be considered as "zoospores;" it is not quite clear what becomes of these; but there is reason to believe that some of them retain their motile powers, and, after increasing in size, develop an investing cyst, like the free primordial utricles to be presently described; that others produce a firm cellulose envelope, and become "still" cells; and that others (perhaps the majority) perish without any further change.

154. When the ordinary self-division of the "still" cells into two segments has been repeated four times, so as to produce 16 cells—and sometimes at an earlier period—the new cells thus produced assume the "motile" condition; being liberated before the development of the cellulose envelope, and becoming furnished with two long vibratile filaments, which appear to be extensions of the primordial utricle (H). In this condition, it seems obvious that the colorless protoplasm is more developed relatively to the coloring matter, than it is in the "still" cells; it generally accumulates in the part from which the vibratile filaments or cilia proceed, so as to form a sort of transparent beak (H, K, L); and it usually contains "vacuoles," occupied only by clear aqueous fluid, which are sometimes so numerous as to

take in a large part of the cavity of the cell, so that the colored contents seem only like a deposit on its walls. Before long, this "motile" primordial utricle acquires a peculiar saccular investment, which seems to correspond with the cellulose envelope of the "still" cells, but which is not so firm in its consistence (I, K, L). Thread-like extensions of the protoplasm, sometimes containing colored globules, are not unfrequently seen to radiate from the primordial utricle towards the exterior of this enveloping bag (I); these are rendered more distinct by iodine, and can be made to retract by means of reagents; and their existence seems to show, on the one hand, that the transparent space through which they extend themselves is only occupied by a watery liquid, and on the other, that the layer of protoplasm which constitutes the primordial utricle, is far from possessing the tenacity of a completely formed membrane. The vibratile filaments pass through the cellulose envelope, which invests them with a sort of sheath; and in the portion that is within this sheath, no movement is seen. During the active life of the "motile" cells, the vibration of these cilia is so rapid, that it can be recognized only by the currents it produces in the water, through which the cells are rapidly propelled; but when the motion becomes slacker, the filaments themselves are readily distinguishable; and they may be made more obvious by the addition of iodine. The multiplication of these motile cells may take place in various modes, giving rise to a great variety of appearances. Sometimes they undergo a regular binary subdivision, whereby a pair of motile cells is produced (C), each resembling its single predecessor in possessing the cellulose investment, the transparent beak, and the vibratile filaments, before the solution of the original investment. Sometimes, again, the contents of the primordial cell undergo a segmentation in the first instance into four divisions (D); which may either become isolated by the dissolution of their envelope, and may separate from each other in the condition of free primordial utricles (U), developing their cellulose investments at a future time; or may acquire their cellulose investments (as in the preceding case) before the solution of that of the original cell; and sometimes, even after the disappearance of this, and the formation of their own independent investments, they remain attached to each other at their beaked extremities, the primordial utricles being connected with each other by peduncular prolongations, and the whole compound body having the form of a +. This quaternary segmentation appears to be a more frequent mode of multiplication among the "motile" cells, than the subdivision into two; although as we have seen, it is less common in the "still" condition. So, also, a primary segmentation of the entire endochrome of the "motile" cells, into 8, 16, or even 32 parts, may take place (E, F), thus giving rise to as many minute primordial cells. These, when they are set free, and possess active powers of movement, rank

as "zoospores" (a): which may either develop a loose cellulose investment or cyst, so as to attain the full dimensions of the ordinary motile cells (I, K), or may become clothed with a dense envelope, and lose their vibratile cilia, thus passing into the "still" condition (A); and this last transformation may even take place, before they are set free from the envelope within which they were produced, so that they constitute a mulberry-like mass, which fills the whole cavity of the original cell, and is kept in motion by its cilia.

155. All these *varieties*, whose relation to each other has been clearly proved by watching the successional changes that make up the history of this one Plant, have been regarded as constituting, not merely distinct *species*, but distinct *genera* of Animalcules; such as, *Chlamydomonas*, *Euglena*, *Trachelomonas*, *Gyges*, *Gonium*, *Pandorina*, *Botryocystis*, *Uvella*, *Syncrypta*, *Monas*, *Astasia*, *Bodo*, and probably many others.¹ Certain forms, such as the "motile" cells (I, K, L), appear in a given infusion, at first exclusively and then principally; they gradually diminish, become more and more rare, and finally disappear altogether, being replaced by the "still" form. After some time, the number of the "motile" cells again increases, and reaches, as before, an extraordinary amount; and this alteration may be repeated several times in the course of a few weeks. The process of segmentation is often accomplished with great rapidity. If a number of motile cells be transferred from a larger glass into a small capsule, it will be found, after the lapse of a few hours, that most of them have subsided to the bottom; in the course of the day, they will all be observed to be upon the point of subdivision; on the following morning, the divisional brood will have become quite free; and on the next, the bottom of the vessel will be found covered with a new brood of self-dividing cells, which again proceed to the formation of a new brood, and so on. The activity of motion and the activity of multiplication, seem to stand, in some degree, in a relation of reciprocity to each other; for the self-dividing process takes place with greater rapidity in the "still" cells, than in the "motile."

¹ In the above sketch, the Author has presented the facts described by Dr. Cohn, under the relation which they seemed to him naturally to bear; for the membrane which immediately surrounds the primordial utricle of the "still" cells, appears to him to be essentially the same with the sacculated investment of the "motile" form, since it differs only in its greater density, and in the absence of the interposed fluid. It is distinctly stated by Cohn, towards the conclusion of his Memoir, that the one like the other consists of cellulose, since both alike give the characteristic blue color with a very dilute solution of iodine, and with moderately diluted sulphuric acid; yet he speaks of what he terms the "encysted zoospore" (I, K, L), as being formed by *one cell within another*, the outer cell having a true cell-membrane and aqueous contents, but being destitute of primordial utricle; whilst the inner cell has denser, colored contents, but is without the true cell-membrane. Having enjoyed, since the above was written, the opportunity of personally communicating with Dr. Cohn with regard to the question to which it refers, the Author is glad to be able to state, that Dr. Cohn's later observations have led him to adopt a view of the relationship of the "still" and "motile" forms, which is in essential accordance with his own.


156. What are the precise conditions which determine the transition from one state to the other, cannot yet be precisely stated; but the influence of certain agencies can be predicted with tolerable certainty. Thus it is only necessary to pour the water containing these organisms, from a smaller and deeper into a larger and shallower vessel, at once to determine segmentation in numerous cells,—a phenomenon which is observable also in many other Protophytes. The “motile” cells seem to be favorably affected by light, for they collect themselves at the surface of the water and at the edges of the vessel; but when they are about to undergo segmentation, or to pass into the “still” condition, they sink to the bottom of the vessel, or retreat to that part of it in which they are least subjected to light. When kept in the dark, the “motile” cells undergo a great diminution of their chlorophyll, which becomes very pale, and is diffused, instead of forming definite granules; they continue their movement, however, uninterruptedly, without either sinking to the bottom, or passing into the still form, or undergoing segmentation. A moderate warmth, particularly that of the vernal sun, is favorable to the development of the “motile” cells; but a temperature of excessive elevation prevents it. Rapid evaporation of the water in which the “motile” forms of *Protococcus* may be contained, kills them at once; but a more gradual loss, such as takes place in deep glasses, causes them merely to pass into the “still” form; and in this condition,—especially when they have assumed a red hue,—they may be completely dried up, and may remain in a state of dormant vitality for many years. It is in this state that they are wafted about in atmospheric currents, and that, being brought down by the rain into pools, cisterns, &c., they may present themselves where none had been previously known to exist; and there, under favorable circumstances, they may undergo a very rapid multiplication, and may maintain themselves until the water is dried up, or some other change occurs which is incompatible with the continuance of their vital activity. They then very commonly become red throughout, the red coloring substance extending itself from the centre towards the circumference, and assuming an appearance like that of oil-drops; and these red cells, acquiring thick cell-walls and a mucous envelope, float in flocculent aggregations on the surface of the water. This state seems to correspond with the “winter spores” of other Protophytes; and it may continue until warmth, air, and moisture, cause the development of the red cells into the ordinary “still” cells, green matter being gradually produced, until the red substance forms only the central part of the endochrome. After this occurs the cycle of changes which has been already described; and the Plant may pass through a long series of these, before it returns to the state of the red thick-walled cell, in which it may again remain dormant for an unlimited period. Even this cycle, however, cannot be regarded as completing the

history of the species before us; since it does not include the performance of any true Generative act. There can be little doubt that, in some stage of its existence, a "conjugation" of two cells occurs, as in the preceding case; and the attention of observers should be directed to its discovery, as well as to the detection of other varieties in the condition of this interesting little Plant, which will be probably found to present themselves before and after the performance of that act.

157. From the composite motile forms of the preceding type, the transition is easy to the group of *Volvocineæ*,—an assemblage of minute Plants of the greatest interest to the Microscopist, on account both of the Animalcule-like activity of their movements, and of the great beauty and regularity of their forms. The most remarkable example of this group, is the well-known *Volvox globator* (Fig. 69), or "globe-animalcule;" which is not uncommon in fresh-water pools, and which, attaining a diameter of 1-30th of an inch, may be seen with the naked eye, when the drop containing it is held up to the light, swimming through the water which it inhabits.

Its onward motion is usually of a rolling kind; but it sometimes slides smoothly along, without turning on its axis; whilst sometimes, again, it rotates like a top, without changing its position. When examined with a sufficient magnifying power, the *Volvox* is seen to consist of a hollow sphere, composed of a very pellucid material, which is studded at regular intervals with minute green spots, and which is often (but not constantly) traversed by green threads connecting these spots together. From each of the spots proceed two long cilia; so that the entire surface is beset with these vibratile filaments, to whose combined action its movements are due. Within the external sphere, there may generally be seen from two to twenty other globes, of a darker color, and of varying sizes; the smaller of these are attached to the inner surface of the investing sphere, and project into its cavity; but the larger lie freely within the cavity, and may often be observed to revolve by the agency of their own ciliary filaments. After a time, the original sphere bursts, the contained sphericles swim forth and speedily develop themselves into the likeness of that within which they have been evolved; their component particles, which are at first closely aggregated together, being separated from each other by the interposition of the transparent pellicle. It was long supposed that the *Volvox* was a *single* Animal; and it was first shown to be a *composite* fabric, made up of a repetition of organisms in all respects similar to each other, by Prof. Ehrenberg; who, however, considered these organisms as *Monads*, and described

FIG. 69.



Volvox Globator.

them as each possessing a mouth, several stomachs, and an eye! Our present knowledge of their nature, however, leaves no doubt of their Vegetable character; and the peculiarity of their history renders it desirable to describe it in some detail.

158. Each of the so-called "Monads" is in reality a somewhat flask-shaped mass of endochrome, about 1-3000th of an inch in diameter; consisting, as in the previous instances, of chlorophyll-granules, diffused through a colorless protoplasm (Fig. 70, 1, L); and bounded by a layer of condensed protoplasm, which represents a primordial utricle, but is obviously far from having attained a membranous consistence. It is prolonged outwardly (or towards the circumference of the sphere) into a sort of colorless peak or proboscis, from which proceed two long vibratile cilia (L); and it is invested by a pellucid or "hyaline" envelope (1, d) of considerable thickness, the borders of which are flattened against those of other similar envelopes (E, c c), but which does not appear to have the tenacity of a true membrane. It is impossible not to recognize the precise similarity between the structure of this body, and that of the motile "encysted" cell of *Protococcus pluvialis* (Fig. 68, K); there is not, in fact, any perceptible difference between them, save that which arises from the regular aggregation, in *Volvox*, of the cells which normally detach themselves from one another in *Protococcus*. The presence of cellulose in the hyaline substance is not indicated, in the ordinary condition of *Volvox*, by the iodine and sulphuric acid test, though the use of "Schulz's solution" gives to it a faint blue tinge; there can be no doubt of its existence, however, in the hyaline envelope of what has been termed *Volvox aureus*, which is in reality nothing but the "winter spore" of *Volvox globator*. The cilia and endochrome, as in the motile forms of *Protococcus*, are tinged of a deep brown by iodine, with the exception of one or two particles in each cell, which, being turned blue, may be inferred to be starch; and when the contents of the cell are liberated, bluish flocculi, apparently indicative of the presence of cellulose, are brought into view by the action of sulphuric acid and iodine. All these reactions are strictly *vegetable* in their nature. When the cell is approaching maturity, its endochrome always exhibits one or more "vacuoles" (Fig. 70, 1, a a), of a spherical form, and usually about one-third of its own diameter; and these "vacuoles" (which are the so-called "stomachs" of Prof. Ehrenberg) have been observed by Mr. G. Busk to undergo a very curious rhythmical contraction and dilatation at intervals of about 40 seconds; the contraction (which seems to amount to complete obliteration of the cavity of the vacuole) taking place rapidly or suddenly, whilst the dilatation is slow and gradual. This curious action ceases, however, as the cell arrives at its full maturity; a condition which seems to be marked by the greater consolidation of the primordial utricle, by the removal or transformation of some of the chlorophyll (the


greater part of the coloring matter contracting into a small irregular mass, which adheres to the bottom or side of the cell, leaving the rest of the cavity clear and transparent), and the formation of the red spot (*b*), which obviously consists, as in *Protococcus*, of a peculiar modification of chlorophyll.

159. Each mass of endochrome normally communicates with those in nearest proximity with it, by extensions of its own substance, which are sometimes single and sometimes double (*x*, *b*); and these connecting processes necessarily cross the lines of division between their respective hyaline investments. The thickness of these processes varies very considerably; for sometimes they are broad bands, and in other cases mere threads; whilst they are occasionally wanting altogether. This difference seems partly to depend upon the age of the specimen, and partly upon the abundance of nutriment which it obtains; for, as we shall presently see, the connection is most intimate at an early period, before the hyaline investments of the cells have increased so much as to separate the masses of endochrome to a distance from one another (*B*, *C*, *D*); whilst in a mature individual, in which this separation has taken place to its full extent, and the nutritive processes have become less active, the masses of endochrome very commonly assume an angular form, and the connecting processes are drawn out into threads (as seen at *E*), or they retain their globular form, and the connecting processes altogether disappear. The influence of reagents, or the infiltration of water into the interior of the hyaline investment, will sometimes cause the connecting processes (as in *Protococcus*, § 154), to be drawn back into the central mass of endochrome; and they will also retreat on the mere rupture of the hyaline investment; from these circumstances it may be inferred, that they are not enclosed in any definite membrane. On the other hand, the connecting threads are sometimes seen as double lines, which seem like tubular prolongations of a consistent membrane, without any protoplasmic granules in their interior. It is obvious, then, that an examination of a considerable number of specimens, exhibiting various phases of conformation, is necessary to demonstrate the nature of these communications; but this may be best made out by attending to the history of their development, which we shall now describe.

160. The spherical body of the young *Volvox* (Fig. 78, *A*) is composed of an aggregation of somewhat angular masses of endochrome (*b*), separated by the interposition of hyaline substance; and the whole seems to be enclosed in a distinctly membranous envelope, which is probably the distended hyaline investment of the primordial cell, within which, as will presently appear, the entire aggregation originated. In the midst of the polygonal masses of endochrome, one mass (*a*), rather larger than the rest, is seen to present a circular form; and this, as will presently appear, is the originating cell of what is hereafter to become a

new sphere. The growing *Volvox* at first increases in size, not only by the interposition of new hyaline substance between its component masses of endochrome, but also by an increase in these masses themselves (B, a), which come into continuous connection with each other by the coalescence of processes (b) which

FIG. 70.



Various Stages in the Development of *Volvox Globator*.

they severally put forth; at the same time, an increase is observed in the size of the globular cell (c), which is preliminary to its binary subdivision. A more advanced stage of the same developmental process is seen at c; in which the connecting processes (a a) are so much increased in size, as to establish a most intimate union between the masses of endochrome, although the increase of the intervening hyaline substance carries these masses apart from one another; whilst the endochrome of the central

globular cell has undergone segmentation into two halves. In the stage represented at *D*, the masses of endochrome have been still more widely separated by the interposition of hyaline substance; each has become furnished with its pair of ciliary filaments; and the globular cell has undergone a second segmentation. Finally at *E*, which represents a portion of the spherical wall of a mature Volvox, the endochrome masses are observed to present a more scattered aspect, partly on account of their own reduction in size, and partly through the interposition of a greatly increased amount of hyaline substance, which is secreted from the surface of each mass; and that portion which belongs to each cell, standing to the endochrome mass in the relation of the cellulose coat of ordinary cells to their primordial utricle, is frequently seen to be marked out from the rest by delicate lines of hexagonal areolation (*e, e*), which indicate the boundaries of each. Of these it is often difficult to obtain a sight, a nice management of the light being usually requisite with fresh specimens; but the prolonged action of water (especially when it contains a trace of iodine), or of glycerine, will often bring them into clear view. The prolonged action of glycerine, moreover, will often show that the boundary lines are double, being formed by the coalescence of two contiguous cell-walls; and they sometimes retreat from each other so far, that the hexagonal areolæ become rounded.

161. As the primary sphere approaches maturity, the secondary germ, whose origin has been traced from the beginning, also advances in development; its contents undergoing multiplication by successive segmentations, so that we find it to consist of 8, 16, 32, 64, and still more numerous divisions, as shown in Fig. 70, *F, G, H*. Up to this stage, at which first the sphere appears to become hollow, it is retained within the hyaline envelope of the cell within which it has been produced; a similar envelope can be easily distinguished, as shown at *K*, just when the segmentation has been completed, and at that stage the cilia pass into it, but do not extend beyond it; and even in the mature volvox, it continues to form an investment around the hyaline envelopes of the separate cells, as shown at *L*. It seems to be by the adhesion of the hyaline investment of the new sphere to that of the old, that the secondary sphere remains for a time attached to the interior wall of the primary; at what exact period, or in what precise manner, the separation between the two takes place, has not yet been determined. At the time of the separation, the developmental process has generally advanced as far as the stage represented at *A*; the foundation of one or more tertiary spheres being usually distinguishable in the enlargement of certain of its cells. And thus the cycle of development is completed, in so far as regards the increase of cells by subdivision. But, as already pointed out, the life-history of no organism can be considered as complete, unless it

includes an act of "conjugation," or some other form of the true Generative process; and as none such has yet been observed in *Volvox*, there is strong ground to suspect that we are not acquainted with its whole life, but that conjugation may occur in some condition not yet known to us, which may only present itself after a long succession of repetitions of that process of gemmation, by which the *Volvox* spheres are multiplied and reproduced.

162. Certain curious varieties, however, occasionally present themselves in the preceding cycle. Thus, the young *Volvox* globe which has attained the size and has undergone the degree of segmentation indicated at G or H, sometimes becomes at first deep green and then yellow; its hyaline envelope acquires an unusual firmness, and a second coat is formed within the first, the colored contents undergoing some retraction; and eventually a considerable space intervenes between the two coats, which is occupied by a clear fluid. The protoplasmic contents of the inner envelope consist chiefly of starch grains, mixed with a bright yellow and apparently oily fluid. In this and other respects, the body in question corresponds so closely with the "still" or "winter" spore of other Protophytes, which is adapted to resist influences that are fatal to the more actively vegetating forms of these organisms, that it seems most probable (as was first suggested by Mr. G. Busk) to stand in this relation to *Volvox globator*, although the ultimate mode of its development is not known; and thus the *Volvox aureus*, as the kind that produces these golden yellow spheres has been designated, is nothing else than an ordinary *Volvox* preparing its brood for the winter state of inactivity. The *Volvox stellatus* of Ehrenberg, again, seems to be nothing more than a variety of the same specific type; its peculiarity consisting in the presence of numerous conical eminences, formed of the hyaline substance only, on the secondary globules, giving them a stellate aspect. The endochrome segments resemble those of the globules of *Volvox aureus*, both in color and composition; and these two forms of secondary globules, the stellate and the smooth, having been observed by Mr. Busk to coexist in the very same sphere, are certainly to be regarded as varieties of one and the same type.¹ The same excellent observer has also pointed out the probability, that the


¹ "In the month of August last," says Mr. Busk (Op. cit. p. 44), "when, in a certain pond on Blackheath, there was the most incredible abundance of *Volvox*, so great, in fact, as to render the water at the lee side of the pond in certain spots of a deep green color, and to cause it to afford, when collected, a very strong herbaceous or conserved smell, the majority of the plants exhibited the *stellate* form of spores, or rapidly acquired spores of that character, and very many were in, or soon assumed, the form of *V. aureus*. They seemed, in fact, to be entering on their hibernating state. Many among them, however, though all small and starved-looking, were of the common kind; in all these, Prof. Williamson's hexagonal areolation was very distinct. In the month of October, however, upon returning to the same pond, I was able to find very few *Volvooes* at all, and all of the usual kind; in none of these could I detect the least appearance of the same arrangement." Prof. Williamson has noticed analogous variations, in the specimens taken at different seasons from one locality (Op. cit. p. 56).

body designated by Prof. Ehrenberg *Sphærosira volvox*, is an ordinary *Volvox* in a different phase of development. For it does not present any marked feature of dissimilarity, except that a large proportion of the green cells, instead of being single (as in the ordinary form of *Volvox*) save where they are developing themselves into young spheres, are very commonly double, quadruple, or multiple; and the groups of ciliated cells thus produced, instead of constituting a hollow sphere, form by their aggregation discoid bodies, of which the separate fusiform cells are connected at one end, whilst at the other they are free, each being furnished with a single cilium. These clusters separate themselves from the primary sphere, and swim forth freely, under the forms which have been designated as *Uvella* and *Syncrypta* by Prof. Ehrenberg. The further history of these has not been traced; and it does not seem improbable that more than one intermediate stage may be passed through, before a return is made to the type of *Volvox globator*.¹

163. *Desmidiaceæ*.—Among the simplest tribes of Protophytes, there are two which are of such peculiar interest to the Microscopist, as to need a special notice; these are the *Desmidiæ* (or more properly *Desmidiaceæ*), and the *Diatomaceæ*. Both of them have been ranked by Ehrenberg, and by many other Naturalists, as Animalcules; but the fuller knowledge of their life-history, and the more extended acquaintance with the parallel histories of other simple forms of Vegetation, which have been gained during the last ten years, can scarcely be considered by judges who are at once competent and unprejudiced, as otherwise than decisive in regard to their vegetable nature. The *Desmidiaceæ* are minute plants of a green color, growing in fresh water; generally speaking, the cells are independent of each other (Figs. 71, 75, 76); but sometimes those which have been formed by duplicative subdivision from a single primordial cell, remain adherent one to another in linear series, so as to form a filament (Fig. 77); whilst in other instances, they constitute beautiful star-like groups (Figs. 73, 74). This tribe is distinguished by two peculiar features; one of these being the semblance of a subdivision into two symmetrical halves, which is seen in the cells of most species, and which is sometimes so decided as to have led to the belief that the cell is really double (Fig. 75 A),

¹ The doctrine of the vegetable nature of the *Volvox*, which had been suggested by Siebold, Braun, and other German Naturalists, was first distinctly enunciated by Prof. Williamson, on the basis of the history of its development, in the "Transactions of the Philosophical Society of Manchester," vol. ix. Subsequently Mr. G. Busk, whilst adducing additional evidence of the Vegetable nature of *Volvox*, in his extremely valuable Memoir in the "Transactions of the Microscopical Society," 2d Series, vol. i, called in question some of the views of Prof. Williamson, which were justified by that gentleman in his "Further Elucidations" in the same Transactions. The Author has endeavored to state the facts in which both these excellent observers agree (and which he has himself had the opportunity of verifying), with the interpretation that seems to him most accordant with the phenomena presented by other Protophytes; and he believes that this interpretation harmonizes with what is most essential in the doctrines of both, their differences having been to a certain degree reconciled by their mutual admissions.

though in other cases it is merely indicated by a slight notch on one side (as in the marginal cells of Fig. 73, A, E); whilst the other is the frequency of projections from their surface, which are sometimes short and inconspicuous (Fig. 75), but are often elongated into spines, presenting a very symmetrical arrangement (Fig. 71). These projections are generally formed by the outer




Various species of *Staurastrum*:—A, *S. vestitum*; B, *S. aculeatum*; C, *S. paradoxum*; D, E, *S. brachiatum*.

coat alone, which possess an almost horny consistence, so as to retain its form after the discharge of its contents (Figs. 74, E, 75, B), but which does not include any mineral ingredient, either calcareous or siliceous, in its composition. This outer coat is surrounded by a very transparent sheath of gelatinous substance, which is sometimes very distinct (as shown in Fig. 77), whilst in other cases its existence is only indicated by its preventing the contact of the cells. The outer coat encloses an inner membrane, which is not always, however, closely adherent to it; and this immediately surrounds the colored substance which occupies the whole interior of the cell. It is quite certain that the Desmidiaceæ, like the Confervoid Plants in general, grow at the expense of the inorganic elements which surround them, instead of depending upon other living beings for their subsistence; and that they decompose carbonic acid, and give off oxygen, under the influence of sunlight. They have the power of generating from these materials the organic compounds which they require for their own development; and these are such as are formed by other undoubted Protophytes, as is proved by the application of the appropriate tests. Thus the outer coat is colored blue by sulphuric acid and iodine, and is therefore composed of cellulose. The "endochrome" derives its color from the same green substance "chlorophyll," as that which imparts it to Plants generally; and this is mingled with a protoplasmic fluid, in which "vacuoles" are frequently to be seen. At certain stages in the

growth of these plants, as in other Algæ, starch is produced; the presence of which is made obvious by the application of iodine. There is no one essential point, therefore, in which the *Desmidiaceæ* differ from other Protophytes, or really approach the Animal kingdom. Some of the arguments that have been advanced in support of their Animal affinities,—such as their multiplication by transverse subdivision, and their generation by a process of conjugation,—are really, now that the physiology of the unicellular plants is better understood, much more strongly indicative of their Vegetable alliances. The assertion of Prof. Ehrenberg, that *Closterium* possesses organs which it protrudes through apertures in its extremities, and keeps in continual motion, is (like too many of his statements) simply untrue. And although many of these plants have a power of slowly changing their place, so that they approach the light side of the vessel in which they are kept, and will even traverse the field of the microscope under the eye of the observer, yet this faculty is in no respect different from that which many undoubted Protophytes (such as *Oscillatoria*, § 196) possess.

164. A very peculiar feature which has recently attracted much attention, is the *circulation of fluid* which may be seen to take place in the *Closterium*, both between its “cellulose” coat and its “primordial utricle,” and within the latter; and the *ciliary action* by which, according to the testimony of some good observers, this circulation is maintained. It is not difficult to distinguish this movement along the convex and concave edges of the cell of any vigorous specimen of *Closterium*, if it be examined under a magnifying power of 250 or 300 diameters; and a peculiar whirl-

FIG. 72.



Economy of *Closterium lunula*:—A, frond showing central separation at a, in which large globules, & are not seen;—B, one extremity enlarged, showing at a the double row of cilia, at b the internal current, and at c the external current;—C, external jet produced by pressure on the frond;—D, frond in a state of self-division.

ing movement may also be distinguished, in the large rounded space which is left at each end of the cell by the retreat of the

endochrome from the primordial utricle (Fig. 72, A, B). According to Mr. S. G. Osborne, however, by using the "Rainey moderator" (§ 74), with direct sunlight and an achromatic condenser, and by increasing the power to about 500 diameters (Mr. Ross's objective of 1-6th in., with a deep eye-piece, being the combination employed), a very distinct action of cilia may be discerned, both along the inner edge of the primordial utricle, between this membrane and the endochrome, and along its outer edge, beneath the cellulose coat; the action being in opposite directions in these two situations, and producing two opposite currents. By careful focussing, the circulation may be seen in broad streams over the whole surface of the endochrome; and these streams detach and carry with them, from time to time, little oval or globular bodies (A, B), which are put forth from it, and which are carried by the course of the flow to the chambers at the extremities, where they join a crowd of similar bodies (B). In each of these chambers, a current may be seen from the somewhat abrupt termination of the endochrome, towards the obtuse end of the cell (as indicated by the interior arrows); and the globules it contains are kept in a sort of twisting movement by the action of the cilia on the *inner* side of the primordial utricle, which can here be distinctly seen as at *a*. Other currents are seen externally to it, which seem to be kept up by the *outer* set of cilia; these form three or four distinct courses of globules, passing towards and away from *c* (as indicated by the outer arrows), where they seem to encounter a fluid jetted towards them through an aperture in the primordial utricle at the apex of the chamber; and here some communication between the inner and the outer currents takes place. A corresponding aperture seems also to exist in the outer cellulose coat; for if the endochrome be forced by pressure into closer proximity than usual to the obtuse termination of the cell, the current from its extremity may be seen to spread into the surrounding fluid. This circulation is by no means peculiar to Closterium; having been seen by Mr. Osborne in many other Desmidiaceæ, in several of which he has also detected what he believes to be ciliary action.¹

165. When the single cell has come to its full maturity, it commonly multiplies itself by *duplicative subdivision*; but the plan on which this takes place is often peculiarly modified, in order to maintain the symmetry characteristic of the tribe. In a cell of the simple cylindrical form of those of *Didymoprium* (Fig. 77), little more is necessary than the separation of the two halves, and the formation of a partition between them by the infolding of the primordial utricle, according to the plan already

¹ See Mr. S. G. Osborne's communications to the "Quarterly Journal of Microscopical Science," vol. ii, p. 234, and vol. iii, p. 54. Although the *Circulation* is an unquestionable fact, it may be doubted whether the appearance of *Ciliary action* is not an optical illusion, due to the play of the peculiar light employed, among the moving particles of the fluid. See Mr. Wenham's paper on the Circulation in the Leaf-Cells of *Anacharis*, in the same Journal, vol. iii, p. 278.

described (§ 150); and in this manner, out of the lowest cell of the filament A, a double cell B is produced. But it will be observed that each of the simple cells has a bifid wart-like projection of the cellulose coat on either side, and that the half of this projection, which has been appropriated by each of the two new cells, is itself becoming bifid, though not symmetrically; in process of time, however, the increased development of the sides of the cells which remain in contiguity with each other, brings up the smaller projections to the dimensions of the larger, and the symmetry of the cell is restored. In *Closterium* (Fig. 72, D), the two halves of the endochrome first retreat from one another at the middle line, and a constriction takes place round the cellulose coat; this constriction deepens until it becomes an hour-glass contraction, which proceeds until the cellulose coat entirely closes round the primordial utricle of the two segments; in this state, one half commonly remains passive, whilst the other has a motion from side to side, which gradually becomes more active; and at last one segment quits the other with a sort of jerk. At this time, a constriction is seen across the middle of the primordial utricle of each segment; but there is still only a single chamber, which is that belonging to one of the extremities of the original entire frond. The globular circulation, for some hours previously to subdivision, and for a few hours afterwards, runs quite round the obtuse end *a* of the endochrome; but gradually a chamber is formed, like that at the opposite extremity, by a separation between the cellulose coat and the primordial utricle; whilst at the same time, the obtuse form becomes changed to a more elongated and contracted shape. Thus, in five or six hours after the separation, the aspect of each extremity becomes the same, and each half resembles the perfect frond in whose self-division it originated; and the globular circulation within the newly-formed chamber comes into connection with the general circulation, some of the free particles which are moving over the surface of the primordial utricle, being drawn into its vortex and tossed about in its eddies. The process is seen to be performed after nearly the same method in *Staurastrum* (Fig. 71, D, E); the division taking place across the central constriction, and each half gradually acquiring the symmetry of the original. In such forms as *Cosmarium*, however, in which the cell consists of two lobes united together by a narrow isthmus (Fig. 75), the division takes place after a different method; for the central region of the isthmus expands, and displays two globular enlargements, separated from each other and from the two halves of the original cell (which their interposition carries apart) by a narrow neck; and these enlargements increase, until they assume the appearance of the half-segments of the original cell. In this state, therefore, the plant consists of a row of four segments, lying end to end, the two old ones forming the extremes, and the two new ones (which do not usually acquire the full size or the characteristic markings of the original, before the


division occurs) occupying the intermediate place. At last the central fusion becomes complete, and two bipartite fronds are formed, each having one old and one young segment; the young segment, however, soon acquires the full size and characteristic aspect of the old one; and the same process, the whole of which may take place within twenty-four hours,¹ is repeated ere long. In *Sphaerosoma*, the cells thus produced remain connected in rows within a gelatinous sheath, like those of *Didymoprium* (Fig. 77); and different stages of the process may commonly be observed in the different parts of any one of the filaments thus formed. In any such filament, it is obvious that the two oldest segments are found at its opposite extremities, and that each subdivision of the intermediate cells must carry them further and further from each other. This is a very different mode of increase from that of the *Confervaceæ*, in which the terminal cell alone undergoes subdivision (§ 198), and is consequently the last formed.

166. Many of the *Desmidiaceæ* multiply after another method; namely, by the subdivision of their endochrome into a multitude of granular particles, termed *gonidia*; which are set free by the rupture of the cell-wall, and of which every one may develop itself into a new cell. These "gonidia" may be endowed with cilia, and may possess an active power of locomotion, in which case they are known as "zoospores;" or they may be destitute of any such power, and may become enclosed in a firm cyst or envelope that seems destined for their long-continued preservation, in which case they are designated as "resting-spores." The movement of the zoospores, first within the cavity of the cell that gives origin to them, and afterwards externally to it, has frequently been observed in the various species of *Cosmarium*; and has been described under the title of "the swarming of the granules," from the extraordinary resemblance which the mass of moving particles bears to a swarm of bees. The subsequent history of their development, however, has not been fully traced out; and this is a point to which the attention of Microscopists should be specially directed. In *Pediastrum*, a plant whose frond normally consists of a cluster of cells, the zoospores are not emitted separately, but those formed by the subdivision of the endochrome of one cell, which may be 4, 8, 16, 32, or 64 in number, escape from the parent frond still enclosed in the inner tunic of the cell; and it is within this that they develop themselves into a cluster, resembling that in which they originated. This is well shown in the accompanying series of illustrations of the developmental history of *Pediastrum granulatum* (Fig. 73, A-F), a species whose frond normally consists of 16 cells, but may be

¹ See the observations of Mrs. Herbert Thomas on *Cosmarium margaritiferrum*, in "Microscopical Transactions," Second Series, vol. iii, pp. 33-36. Several varieties in the mode of subdivision are described in this short record of long-continued observations, as of occasional occurrence.

composed of either of the just mentioned multiples or sub-multiples of that number. At A is seen an old disk, of irregular shape, nearly emptied by the emission of its gonidia, which had

FIG. 73.


Various phases of development of *Pediatrum granulatum*.

been seen to take place, within a few hours previously, from the cells *a, b, c, d, e*; most of the empty cells exhibit the cross slit through which their contents had been discharged; and where this does not present itself on the side next the observer, it occurs on the other. Three of the cells still possess their colored contents, but in different conditions. One of them exhibits an early stage of the subdivision of the endochrome, namely, into two halves, one of which already appears halved again. Two others are filled by sixteen very closely crowded gonidia, only half of which are visible, as they form a double layer. Besides these, one cell is in the very act of discharging its gonidia; nine of which have passed forth from its cavity, though still enveloped in a vesicle formed by the extension of its innermost membrane; whilst seven yet remain in its interior. The new-born family, as it appears immediately on its complete emersion, is shown at B; the gonidia are actively moving within the vesicle; and they do not as yet show any indication either of symmetrical arrangement, or of the peculiar form which they are subsequently to assume. Within a quarter of an hour, however, the gonidia are observed to settle down into one plane, and to assume some kind of regular arrangement, most commonly that seen at C, in which there is a single central body, surrounded by a circle of five, and this again by a circle of ten; they do not, however, as yet adhere firmly together. The gonidia now begin to develop themselves into new cells, increase in size, and come into closer approximation (D); and the edge of each, especially in the marginal row, presents a notch, which foreshadows the production of its charac-

teristic "horns." Within about four or five hours after the escape of the gonidia, the cluster has come to assume much more of the distinctive aspect of the species, the marginal cells having grown out into horns (E); still, however, they are not very closely connected with each other; and between the cells of the inner row, considerable spaces yet intervene. It is in the course of the second day, that the cells become closely applied to each other, and that the growth of the horns is completed, so as to constitute a perfect disk, like that seen at F, in which, however, the arrangement of the interior cells does not follow the typical plan.

167. The varieties which present themselves, indeed, both as to the number of cells in each cluster, and the plan on which they are disposed, are such as to baffle all attempts to base specific distinctions on such grounds: and the more attentively the life-history of any one of these plants is studied, the more evident does it appear that many reputed species have no real existence. Some of these, indeed, are nothing else than mere transitory forms; thus it can scarcely be doubted that the specimen represented in Fig. 74, D, under the name of *Pediastrum pertusum*, is in reality nothing else than a young frond of *P. granulatum*, in the stage represented in Fig. 73, E, but consisting of 32 cells. On the other hand, in Fig. 74, E, we see an emptied frond of *P. granulatum*, exhibiting the peculiar surface-marking from which the name of the species is derived, but composed of no more

FIG. 74.



Various species (?) of *Pediastrum*:—A, *P. tetras*; B, C, *P. biradiatum*; D, *P. pertusum*; E, empty frond of *P. granulatum*.

than 8 cells. And instances every now and then occur, in which the frond consists of only 4 cells, each of them presenting the two-horned shape. So, again, in Fig. 74, B and C, are shown two varieties of *Pediastrum biradiatum*, whose frond is normally com-

posed of sixteen cells; whilst at A is figured a form which is designated as *P. tetras*, but which may be strongly suspected to be merely a 4-celled variety of B and C.


168. Many similar cases might be cited; and the Author would strongly urge those Microscopists who have the requisite time and opportunities, to apply themselves to the determination of the *real species* of this group, by studying the entire life-history of whatever forms may happen to lie within their reach, noting all the varieties which present themselves among the offsets from any one stock. It must not be forgotten that this process of *multiplication* is analogous to the propagation of the higher Plants by budding, and to the subsequent separation of the buds, either spontaneously, or by the artificial operations of grafting, layering, &c.; and just as in all these cases, the particular *variety* is propagated, whilst only the characters of the *species* are transmitted by the true generative operation to the descendants raised from seed, so does it come to pass that the characters of any particular variety which may arise among these unicellular Plants, are diffused by the process of duplicative subdivision, amongst vast multitudes of so-called individuals. Thus it happens that, as Mr. Ralfs has remarked, "one pool may abound with individuals of *Staurastrum dejectum* or *Arthrodesmus incus*, having the mucro curved outwards; in a neighboring pool, every specimen may have it curved inwards; and in another, it may be straight. The cause of the similarity in each pool no doubt is, that all its plants are offsets from a few primary fronds." Hence the universality of any particular character, in all the plants of one gathering, is by no means sufficient to entitle these to take rank as a distinct species; since they are, properly speaking, but repetitions of the same form by a process of simple multiplication, really representing, in their entire aggregate, the one plant or tree that grows from a single seed.¹ In the genus *Cælastrum*, the frond of which, like that of *Pediastrum*, is composed of clusters of cells, the endochrome subdivides into segments, as if for the formation of zoospores; but no motion takes place. These segments acquire cellulose coats, and arrange themselves within the parent-cells according to the typical pattern; and then the wall of the parent-cell splits and peels off, leaving them as the foundation of a new cluster (Pringsheim). A somewhat parallel phenomenon has been observed and figured in *Closterium* by Focke; the entire endochrome being retracted from the walls, and breaking up into a number of globules, every one of which acquires a very firm envelope, resembling that of the "resting-spores" or "winter-spores" of many other Protophytes. Probably the following observation of Mr. Jenner's refers to a more limited production of these "resting-spores:"—"In all the Desmidiaceæ,

¹ For a discussion of the question what really constitutes *individuality* in this and similar cases, see the Author's "Principles of Comparative Physiology," Chap. XI, Section 2.—*Am. Ed.*

but especially in *Closterium* and *Micrasterias*, small, compact, seed-like bodies of a blackish color are at times to be met with. Their situation is uncertain, and their number varies from one to four. In their immediate neighborhood the endochrome is wanting, as if it had been required to form them; but in the rest of the frond it retains its usual color and appearance." It seems likely that, when thus enclosed in a firm cyst, the gonidia are more capable of preserving their vitality, than they are when destitute of such a protection; and that in this condition they may be taken up and wafted through the air, so as to convey the species into new localities.

169. The proper Generative process in the *Desmidiaceæ* is always accomplished by the act of "conjugation;" and this takes place after a manner very different from that which we have

FIG. 75



Cosmarium botrytis.—A, mature frond; B, empty frond; C, transverse view; D, sporangium with empty fronds.

seen to occur in *Palmoglaea*. For each cell here possesses, it will be recollected, a firm external envelope, which cannot enter into coalescence with that of any other; and this membrane dehisces more or less completely, so as to separate each of the conjugating cells into two valves (Fig. 75, c, d; Fig. 76, c). The contents of each cell, being thus set free, without (as it appears) any distinct investment, blend with those of the other; and a mass is formed by their union, which so acquires a truly membranous envelope.¹ This envelope is at first very delicate, and is filled with green and granular contents; by degrees the envelope acquires increased thickness, and the contents of the spore-cell become


brown or red. The surface of the *sporangium*, as this body is now termed, is sometimes smooth, as in *Closterium* and its allies (Fig. 76), and in the *Desmidiæ* proper (Fig. 77); but in the *Cosmariæ*, it acquires a granular, tuberculated, or even spinous surface (Fig. 75), the spines being sometimes simple and sometimes forked at their extremities.²

170. The mode in which conjugation takes place in the filamentous species constituting the *Desmidiæ* proper, is, however, in many respects different. The filaments first separate into their component joints; and when two cells approach in con-

¹ In *Closterium lineatum*, as in many of the *Diatomaceæ* (§ 178), the act of conjugation gives origin to two sporangial cells.

² Bodies precisely resembling these, and almost certainly to be regarded as of this kind, are often found fossilized in flint, and have been described by Ehrenberg under the name of *Xanthidia*.


jugation, the outer cell-wall of each splits or gapes at that part which adjoins the other cell, and a new growth takes place, which forms a sort of connecting tube, uniting the cavities of the two cells (Fig. 77, D, E). Through this tube the entire endochrome of one cell passes over into the cavity of the other (D), and the two are commingled so as to form a single mass (E), as is the case in many of the *Conjugateæ* (§ 199). The joint which contains the sporangium can scarcely be distinguished at first (after the separation of the empty cell), save by the greater density of its contents; but



Closterium striatolum.—A, ordinary frond; B, empty frond; C, two fronds in conjugation.

the proper coats of the sporangium gradually become more distinct, and the enveloping cell-wall disappears. The subsequent history of the sporangia is still obscure; since, although it cannot be doubted that they give origin to new plants resembling those by whose conjugation they are formed, it is not known whether each sporangium in the first instance develops a single cell, or a brood of cells. The latter seems, from the observations of Jenner and Focke, to be the case with *Closterium*;

whilst those of Mrs. H. Thomas (loc. cit.) indicate that it is likewise in *Cosmarium*, whose sporangium has been seen by her to emit large numbers of bodies resembling zoospores. This part of the history of the group is yet involved in much mystery; more espe-



Didymoprium Grevillii.—A, portion of filament, surrounded by gelatinous envelope; B, dividing joint; C, single joint viewed transversely; D, two cells in conjugation; E, formation of sporangium.

cially since, according to the observations of Mr. Ralfs, there

are several Desmidiaceæ which never make their appearance in the same pools for two years successively, although their sporangia are abundantly produced,—a circumstance which would seem to indicate that their sporangia give origin to some different forms. It is a subject, therefore, to which the attention of Microscopists cannot be too sedulously directed.

171. The Desmidiaceæ are not found in running streams, unless the motion of the water be very slow; but are to be looked for in standing, but not in stagnant waters. Small shallow pools that do not dry up in summer, especially in open exposed situations, such as boggy moors, are most productive. The larger and heavier species commonly lie at the bottom of the pools, either spread out as a thin gelatinous stratum, or collected into finger-like tufts. By gently passing the fingers beneath these, they may be caused to rise towards the surface of the water, and may then be lifted out by a tin box or scoop. Other species form a greenish or dirty cloud upon the stems and leaves of other aquatic plants; and these also are best detached by passing the hand beneath them, and “stripping” the plant between the fingers, so as to carry off upon them what adhered to it. If, on the other hand, the bodies of which we are in search should be much diffused through the water, there is no other course than to take it up in large quantities by the box or scoop, and to separate them by straining through a piece of linen. At first nothing appears on the linen but a mere stain or a little dirt; but by the straining of repeated quantities, a considerable accumulation may be gradually made. This should be then scraped off with a knife, and transferred into bottles with fresh water. If what has been brought up by hand be richly charged with these forms, it should be at once deposited in a bottle; this at first seems only to contain foul water; but by allowing it to remain undisturbed for a little time, the Desmidiaceæ will sink to the bottom, and most of the water may then be poured off, to be replaced by a fresh supply. If the bottles be freely exposed to solar light, these little plants will flourish, apparently as well as in their native pools; and their various phases of multiplication and reproduction may be observed during successive months or even years. If the pools be too deep for the use of the hand and the scoop, a collecting-bottle attached to a stick (§ 148) may be employed in its stead. The “ring-net” may also be advantageously employed, especially if it be so constructed as to allow of the ready substitution of one piece of muslin for another (§ 143). For by using several pieces of previously-wetted muslin in succession, a large number of these minute organisms may be separated from the water; the pieces of muslin may be brought home, folded up in wide-mouthed bottles, separately, or several in one, according as the organisms are obtained from one or from several waters; and they are then to be opened out in jars of filtered river-water, and exposed to the light, when the Desmidiaceæ will detach themselves from it.

172. *Diatomaceæ*.—Notwithstanding the very close affinity, which, as will be presently shown, exists between this group and the preceding, many Naturalists who do not hesitate in regarding the Desmidiaceæ as Plants, persist in referring the Diatomaceæ to the Animal kingdom. For this separation, no valid reason can be assigned; the curious movements which the Diatomaceæ exhibit, being certainly not of a nature to indicate the possession of any truly animal endowment; and all their other characters being unmistakably vegetable. Like the Desmidiaceæ, they are *simple cells*, having a firm external coating, within which is included a mass of endochrome whose superficial layer seems to be consolidated into a sort of primordial utricle. The external coat, however, though it seems to have a basis of organic membrane,¹ is consolidated by *silix*; the presence of which in this situation is one of the most distinctive characters of the group. The endochrome, instead of being bright green, is of a yellowish-brown; and its peculiar color seems to be in some degree dependent upon the presence of *iron*, which is assimilated by the plants of this group, and which may be detected even in their colorless silicified envelopes. The coloring substance appears to be a modification of ordinary chlorophyll; it takes a green or greenish-blue tint with sulphuric acid; and often assumes this hue in drying. The endochrome consists, as in other Plants, of a viscid protoplasm, in which float the granules of coloring matter. In the ordinary condition of the cell, these granules are diffused through it with tolerable uniformity, except in the central spot which is occupied by a *nucleus*; round this nucleus they commonly form a ring, from which radiating lines of granules may be seen to diverge into the cell-cavity. At certain times, oil-globules are observable in the protoplasm; these seem to represent the starch-granules of the Desmidiaceæ (§ 163) and the oil-globules of other Protophytes (§ 151). A distinct movement of the granular particles of the endochrome, closely resembling the circulation of the cell-contents of the Desmidiaceæ (§ 164), has been noticed by Prof. W. Smith in some of the larger species of Diatomaceæ, such as *Surirella biseriata*, *Nitzschia scalaris*, and *Campylodiscus spiralis*; and although this movement has not the regularity so remarkable in the preceding group, yet its existence is important, as confirming the conclusion that each Diatom is a single cell (the endochrome moving freely from one part of its interior to another), and that it does not contain in its

¹ A membrane bearing all the markings of the siliceous envelope has been found by Prof. Bailey to remain, after the removal of the silix by hydrofluoric acid; and this membrane seems to have been presumed by him, as also by Prof. W. Smith, to lie beneath the siliceous envelope, and to *secrete* this on its surface as a sort of epidermis. The Author agrees, however, with the authors of the "Micrographic Dictionary" (p. 200), in considering it much more likely that this membrane is the proper "cellulose coat" *interpenetrated* by silix; especially since it has been found by Schmidt, that after removing the protoplasm of *Frustrulia salina* by potash, and the oil by ether, a substance remained identical in composition with the cellulose of Lichens.

interior the aggregation of separate organs which have been imagined to exist in it.

173. The Diatomaceæ seem to have received their name from the readiness with which those forms that grow in coherent masses (which were those with which Naturalists first became acquainted) may be *cut* or *broken through*; hence they have been also designated by the vernacular term "brittle-works." Of this we have an example in the common *Diatoma* (Fig. 94), whose component cells (which in this tribe are usually designated as *frustules*) are sometimes found adherent side by side (as at *b*) so as to form filaments, but are more commonly met with in a state of partial separation, remaining connected at their angles only (usually the *alternate* angles of the contiguous frustules) so as to form a zigzag chain. A similar cohesion at the angles is seen in the allied genus *Grammatophora* (Fig. 95), in *Isthmia* (Fig. 96), and in many other Diatoms; in *Biddulphia* (Fig. 81), there even seems to be a special organ of attachment at these points. In some Diatoms, however, the cells produced by successive acts of binary subdivision, habitually remain adherent one to another; and thus are produced filaments or clusters of various shapes. Thus it is obvious that, when each cell is a short cylinder, an aggregation of such cylinders, end to end, must form a rounded filament, as in *Meloseira* (Figs. 97, 98); and whatever may be the form of the sides of the cells, if they be parallel one to the other, a *straight* filament will still be produced, as in *Achnanthes* (Fig. 93). But if, instead of being parallel, the sides be somewhat inclined towards each other, a *curved* band will be the result; this may not continue entire, but may so divide itself as to form fan-shaped expansions, as those of *Lichmophora flabellata* (Fig. 91); or the cohesion may be sufficient to occasion the band to wind itself (as it were) round a central axis, and thus, not merely to form a complete circle, but a spiral of several turns, as in *Meridion circulare* (Fig. 92, *v*). Many Diatoms, again, possess a *stipes* or stalk-like appendage, by which they are attached to other plants or to stones, pieces of wood, &c., and this may be a simple foot-like appendage, as in *Achnanthes longipes* (Fig. 93), or it may be a composite plant-like structure, as in *Lichmophora* (Fig. 91), *Gomphonema* (Fig. 89), and *Mastogloia* (Fig. 99). Little is known respecting the nature of this stipes; it is, however, quite flexible; and may be conceived to be an extension of the cellulose coat unconsolidated by silex, analogous to the prolongations which have been seen in the *Desmidiaceæ* (§ 163), and to the filaments which sometimes connect the cells of the *Palmellaceæ* (§ 194). Some Diatoms, again, have a mucous or gelatinous investment, which may even be so substantial that they lie as it were in a bed of it, as in *Mastogloia* (Figs. 99, 100), or which may form a sort of tubular sheath, as in *Schizonema*. In a large proportion of the group, however, the frustules are always met with entirely free; neither remaining in the least degree coherent


one to another, after the process of duplicative subdivision has once been completed; nor being in any way connected, either by a stipes, or by a gelatinous investment. This is the case, for example, with *Triceratium* (Fig. 79), *Pleurosigma* (Fig. 80), *Actinocyclus* (Figs. 84, 101, *b b*), *Heliopelta* (Fig. 85), *Arachnoidiscus* (Fig. 86), *Campylodiscus* (Fig. 87), *Surirella* (Fig. 88), *Coscinodiscus* (Fig. 101, *a, a, a*), and many others. The discoid forms, however, when obtained in their living state, are commonly found cohering to the surface of seaweeds.

174. We have now to examine more minutely into the curious structure of the siliceous envelope, which constitutes the characteristic feature of the Diatomaceæ, and the presence of which imparts a peculiar interest to the group, not merely on account of the elaborately marked pattern which it often exhibits, but also through the perpetuation of the minutest details of that pattern, in the specimens obtained from fossilized deposits (Figs. 101, 102). The siliceous envelope of every Diatomaceous "frustule" or cell, consists of two *valves* or plates, usually of the most perfect symmetry, closely applied to each other, like the two valves of a Mussel or other bivalve shell, along a line of fracture or *suture*; and each valve being more or less concavo-convex, a cavity is left between the two, which is occupied by the cell-contents. The form of this cavity, however, differs very greatly; for sometimes each valve is hemispherical, so that the cavity is globular; sometimes it is a smaller segment of a sphere, resembling a watch-glass, so that the cavity is lenticular; sometimes the central portion is completely flattened, and the sides abruptly turned up, so that the valve resembles the cover of a pill-box, in which case the cavity will be cylindrical; and these and other varieties may coexist with any modifications of the contour of the valves, which may be square, triangular (Fig. 79), heart-shaped (Fig. 87), boat-shaped (Fig. 88, *A*), or very much elongated (Fig. 80, *A*), and may be furnished (though this is rare among the Diatomaceæ) with projecting outgrowths (Fig. 81). In all instances, the frustule is considered to present its "front" view, when its suture is turned towards the eye, as in Fig. 88, *B, C*; whilst its "side" view is seen, when the centre of either valve is directly beneath the eye (*A*). Although the two valves meet along the suture, in those newly formed frustules which have been just produced by binary subdivision (as shown in Fig. 81, *A, e*), yet as soon as they begin to undergo any increase, the valves separate from one another, and the cell-membrane which is thus left exposed, immediately becomes consolidated by siliceous matter, and thus forms a sort of *hoop* that intervenes between the valves (as seen at *B*); this hoop becomes broader and broader with the increase of the cell in length; and it sometimes attains a very considerable width (Fig. 81, *A, b*). As growth and self-division are continually going on when the frustules are in a healthy vigorous condition, it is rare to find a specimen in which the valves are not in some degree separated by the interposition of

the "hoop." The impermeability of the siliceous envelope renders necessary some special aperture, through which the surrounding water may communicate with the contents of the cell. Such apertures are found along the whole line of suture in disk-like frustules; but when the Diatom is of an elongated form, they are found at the extremities of the frustules only. They do not appear to be absolute perforations in the envelope, but are merely points at which its siliceous impregnation is wanting; and these are usually indicated by slight depressions of its surface. In some Diatoms, as *Surirella* (Fig. 88) and *Campylodiscus* (Fig. 87), these interruptions are connected with what seem to be minute canals hollowed out between the siliceous envelope and the membrane investing the endochrome. In many genera, the surface of each valve is distinguished by the presence of a longitudinal band, on which the usual markings are deficient; and this is widened into small expansions at the extremities, and sometimes at the centre also, as we see in *Pleurosigma* (Fig. 80) and *Gomphonema* (Fig. 89). This band seems to be merely a portion in which the siliceous envelope is thicker than it is elsewhere, forming a sort of rib that seems designed to give firmness to the valve; and its expansions are solid nodules of the same substance.¹

175. The nature of the delicate and regular markings, with which probably every Diatomaceous valve is beset, has been of late years a subject of much discussion among Microscopists, and cannot be said to be even yet settled, although (in the Author's opinion) the weight of evidence now decidedly preponderates on

FIG. 78.



Portion of Cell of *Isthmia nervosa*, highly magnified.


one side. In the first place it may be remarked, that there is a much greater uniformity in the general character of these markings, than was supposed when attention was first directed to them; for what were at first supposed to be *lines*, are now resolved by objectives of large angular aperture into *rows of dots*; and these dots, when sufficiently magnified, are found to bear a close resemblance to the coarser

markings on the larger species. It is to the latter, therefore,

¹ These *nodules* were mistaken by Prof. Ehrenberg for *apertures*; and in this error he has been followed by Kützing. There cannot any longer, however, be a doubt as to their real nature. As Prof. W. Smith has justly remarked:—"The internal contents of the frustule never escape at these points when the frustule is subjected to pressure, but invariably at the suture or at the extremities, where the foramina already described exist. Nor does the valve, when fractured, show any disposition to break at the expansions of the central line, as would necessarily be the case were such points perforations and not nodules."

that we should have recourse for the determination of the nature of these markings; and we cannot resort to better illustrations, than those which are afforded by *Isthmia* (Fig. 78), *Triceratium* (Fig. 79), and *Biddulphia* (Fig. 81), in all of which the structure of the valve can be distinctly seen under a low magnifying power and with ordinary light. In each of these instances, we see a number of *areolæ*, rounded, oval, or hexagonal, with intervening spaces symmetrically disposed; and the idea at once suggests itself, that these *areolæ* are portions of the surface either *elevated above* or *depressed below* the rest. That the latter is their true condition, is suggested by the appearances they present when the light is obliquely directed; and it may also be inferred from the aspect presented by these objects when viewed by the black ground illumination (§ 61), since the *areolæ* are then less bright than the intervening spaces, less light being stopped by their thinner substance. Moreover, when a valve is broken, the line of fracture corresponds to what, on this supposition, is its weakest portion; since it passes through the *areolæ*, instead of through the intervening spaces, which last would be the weaker portions if the *areolæ* were prominences. But the most satisfactory proof that the *areolæ* are depressions, is perhaps that which is afforded by a *side view* of them, such as may be obtained by examining the curved edges of the valves in *Isthmia*; this, it may be safely affirmed, can leave no doubt in the mind of any competent and unprejudiced observer, as to the nature of the markings in *that* genus; and analogy would seem to justify the extension of the same view to the other cases in which the microscopic appearances correspond.¹ Now, it would not be difficult to bring together a connected series of Diatomaceæ, in which the markings, still exhibiting the same general aspect, become more and more minute, requiring for their resolution the use of oblique


FIG. 79.



Triceratium favus.—A, side view; B, front view.

¹ It is considered by Prof. W. Smith, that this areolation is indicative of a *cellular* structure in the siliceous envelope. But when it is borne in mind that the entire frustule constitutes a single cell, such an hypothesis seems altogether inadmissible. The Author would rather consider the markings in question as analogous to those which are presented by the surface of many pollen-grains (Fig. 189), of whose single-celled nature no doubt can exist; and in his researches on the Foraminifera, he has met with several instances, in which the calcareous investments of those segments of sarcode which must be considered as the representatives of single cells, are marked with a like areolation, the *areolæ* being here unquestionably depressions, formed by the thinning away of the envelope at those parts.

light or of stops with a central diaphragm, and of objectives of larger and larger angular aperture; until we come to those species which present the greatest difficulty, and the nature of whose markings seem most obscure. The more perfectly these markings can be defined, however, in any case, the more decidedly are they found to correspond with what has been already seen. Thus, if we examine *Pleurosigma angulatum*, one of the easier tests (§ 102), with an objective of 1-4th inch focus and 75° aperture, we shall see very much what is represented in Fig. 80, A; namely, a double series of somewhat interrupted lines, crossing each other at an angle of 60 degrees, so as to have between them imperfectly defined lozenge-shaped spaces.¹ When, however,



Pleurosigma angulatum:—A, entire frustule, as seen under a power of 500 diam.; B, hexagonal areolation, as seen under a power of 1300 diam.; C, the same, as seen under a power of 15,000 diam.


the valve is examined with an objective of 1-12th inch focus, having an angular aperture of 130°, and is illuminated by oblique rays, its hexagonal areolation becomes very distinct, as shown at B. And if a photographic representation obtained by such a power be itself enlarged by photography, as has been accom-

¹ This representation is taken from a Photograph by Mr. Delves, which is imitated as closely as Wood-engraving can imitate on a scale of such minuteness, but with a reversal of the lights and shades.

plished by Mr. Wenham, the appearance represented at *c* is obtained; which is in all respects comparable with that presented under a low power by the valve of *Triceratium* or *Isthmia*. At the upper part of this figure, which represents a portion of the object that was accurately in focus, the hexagonal areas are seen to be light, and the intervening spaces dark; the reverse is the case with the lower portion, which was out of focus; and a curious transition from one condition to the other is seen in the intermediate part.¹

176. The process of multiplication by self-division takes place among the Diatomaceæ on the same general plan as in the Desmidiaceæ, but with some modifications incident to the peculiarities of the structure of the former group. The first stage consists in the elongation of the cell, and the increase in the breadth of the "hoop," which is well seen in Fig. 81; for in the newly-

formed cell *e*, the two valves are in immediate apposition, in *d* a hoop intervenes, in *a* this hoop has become much wider, and in *b* the increase has gone on until the original form of the cell is completely changed. At the same time, the endochrome separates into two halves so that its granules form two layers, applied to the opposite sides of the frustule; the nucleus also subdivides, in the manner formerly shown (Fig. 67, *g*, *h*, *i*); and (although the process has not been clearly made out in this group) it may be pretty certainly concluded that the primordial utricle folds in, first forming a mere constriction, then an hour-glass contraction, and finally a complete double partition, as in other instances (§ 165). From each of these two surfaces a new siliceous valve is formed, as shown at Fig. 81, *A*, *c*, just as a new cellulose wall is generated in the subdivision of other cells; and this valve is usually



Biddulphia pulchella;—*A*, chain of cells in different states; *a*, full size; *b*, elongating preparatory to subdivision; *c*, formation of two new cells; *d*, *e*, young cells,—*B*, end view;—*c*, side view of a cell more highly magnified.

¹ The Author does not think it necessary to go more in full into the discussion of the nature of these markings, which some have represented to be due to hemispherical elevations on the valves; as he thinks that no argument is likely to convince those, whose minds are prepossessed with a conception which influences their interpretation of what they see. To those who come fresh to the question, he would strongly recommend the education of their judgment upon the larger and more closely marked Diatoms, as above described, before they commit themselves to an opinion on either side. A fuller statement of the question will be found in the "Micrographic Dictionary," "Introduction," p. xxxiii, and *Arts.* "Angular Aperture" and "Diatomaceæ."

the exact counterpart of the one to which it is opposed, and forms with it a complete cell, so that the original frustule is replaced by two frustules. Sometimes, however, the new valves seem to be a little larger than their predecessors; so that, in the filamentous species, there may be an increase sufficient to occasion a gradual widening of the filament, although not perceptible when two contiguous frustules are compared; whilst, in the free forms, frustules of different sizes may be met with, of which the larger are more numerous than the smaller, the increase in number having taken place in geometrical progression, whilst that of size was uniform. It is not always clear what becomes of the "hoop." In *Biddulphia* and *Isthmia* (Fig. 96) the two young cells slip out of it, and the hoop at last becomes completely detached; and the same thing happens with many other Diatoms; so that the "hoops" are to be found in large numbers, in the settlings of water in which they have been growing for some time. In *Meloseira* (Figs. 97, 98), and perhaps in the filamentous species generally, on the other hand, the "hoops" appear to keep the new frustules united together for some time. But in some other cases, all trace of it is lost; and it may be questioned whether it has ever been properly silicified, and whether it does not become fused (as it were) into the gelatinous envelope. During the healthy life of the Diatom, the process of self-division is continually being repeated; and a very rapid multiplication of frustules thus takes place, all of which (as in the cases already cited, §§ 150, 165) must be considered to be repetitions of one and the same individual. Hence it may happen (as among the Desmidiaceæ, § 168) that myriads of frustules may be found in one locality, uniformly distinguished by some peculiarity of form, size, or marking; which may yet have had the same remote origin as another collection of frustules, found in some different locality, and alike distinguished by some peculiarity of its own. For there is strong reason to believe, that such differences spring up among the progeny of any true *generative* act (§ 178); and, when that progeny is dispersed by currents into different localities, each will continue to multiply its own special type, so long as the process of self-division goes on.

177. It is uncertain whether the Diatomaceæ also multiply by the breaking up of their endochrome into "gonidia," and by the liberation of these, either in the *active* condition of "zoospores," or in the state of "still" or "resting" spores. Certain recent observations by Focke,¹ however, taken in connection with the analogy of other Protophytes, and with the fact that the sporangial frustules undoubtedly thus multiply by gonidia (§ 178), seem to justify the conclusion that such a method of multiplication does obtain in this group. And it is not at all improbable, that very considerable differences in the size, form,

¹ "Physiologisch. Studien," Heft ii, 1853; quoted in "Micrographical Dictionary," p. 201.

and markings of the frustules, such as many consider sufficient to establish a diversity of species, have their origin in this mode of propagation. It is probable that, so long as the vegetating processes are in full activity, multiplication takes place in preference by self-division; and that it is when deficiency of warmth, of moisture, or of some other condition, gives a check to these, that the formation of encysted gonidia, having a greater power of resisting unfavorable influences, will take place; whereby the species is maintained in a dormant state, until the external conditions are favorable to a renewal of active vegetation (§ 156).

178. The process of "conjugation," or true Generation, has been observed to take place among the ordinary Diatomaceæ, almost exactly as among the Desmidiaceæ. Thus in *Surirella* (Fig. 88), the valves of two free and adjacent frustules separate from each other at the sutures, and the two endochromes (probably included in their primordial utricle) are discharged; these coalesce, and form a single sporangial mass, which becomes enclosed in a gelatinous envelope; and in due time this mass shapes itself into a frustule resembling that of its parent, but of larger size. In *Epithemia* (Fig. 82, A, B), however,—the first Diatom in which the conjugating process was observed, by Mr. Thwaites,—the endochrome of each of the conjugating frustules (C, D) appears to divide at the time of its discharge, into two halves; each half coalesces with half of the other endochrome; and thus two sporangial frustules (E, F) are formed (as in *Closterium lineatum*, § 169, note), which, as in the preceding case, become invested with a gelatinous envelope, and gradually assume the form and markings of the parent frustules, but grow to a very much larger size, the sporangial masses having obviously a power of self-increase up to the time when their envelopes are consolidated. This double conjugation seems to be the ordinary type of the process among the Diatoms. A curious departure from the usual plan is observed in some of the filamentous species; for their component cells, instead of conjugating with those of another filament (as is the case with the filamen-





FIG. 82.
Conjugation of *Epithemia turgida*:—A, front view of single frustule; B, side view of the same; C, two frustules with their concave surfaces in close apposition; D, front view of one of the frustules, showing the separation of its valves along the suture; E, F, side and front views after the formation of the sporangia.

with those of another filament (as is the case with the filamen-

tous *Desmidiaceæ*, § 170, and usually but not invariably with the *Zygnemaceæ*, § 199), conjugate with each other; and this may take place even before they have been completely separated by self-division. Thus in *Meloseira* (§ 188) and its allies, the endochrome of particular frustules, after separating as if for the formation of a pair of new cells, moves back from the extremities towards the centre, rapidly increasing in quantity and aggregating into a sporangial mass (Fig. 83, 2, a, b, c), and around

FIG. 83.



Self-Conjugation of *Aulacoseira crenulata*:—1, simple filament; 2, filament developing sporangia; a, b, c, successive stages in the formation of sporangia; 3, embryonic frustules, in successive stages a, b, c, of multiplication.

this a new envelope is developed, which may or may not resemble that of the ordinary frustules, but which remains in continuity with them, giving rise to a strange inequality in the size of the different parts of the filaments (Figs. 97, 98). Of the subsequent history of the sporangial frustule, much remains to be learned; and it is probably not the same in all cases. It has been already shown that the sporangial frustule, even where it precisely resembles its parent in form and marking, greatly exceeds it in size; and this excess seems to render it improbable that it should reproduce the race by ordinary self-division. Appearances have been seen, which make it probable that the contents of each sporangial frustule break up into "gonidia;" and that it is from these that the new generation originates. These gonidia, if each be surrounded (as in many other cases) by a distinct cyst, may remain undeveloped for a considerable period; and they must augment considerably in size, before they attain the dimensions of the parent frustule. It is in this stage of the process, that the modifying influence of external agencies is most likely to exert its effects; and it may be easily conceived that (as in higher Plants and Animals) this influence may give rise to various diversities among the respective individuals of the same brood; which diversities (as we have seen) are transmitted to all the repetitions of each, that are produced by the

self-dividing process. Hence a very considerable latitude is to be allowed to the limits of species, when the different forms of Diatomaceæ are compared; and here, as in many other cases, a most important question arises as to what *are* those limits,—a question which can only be answered by such a careful study of the entire life-history of every single type, as may advantageously occupy the attention of many a Microscopist, who is at present devoting himself to the mere detection of differences, and to the multiplication of reputed species.¹

179. Most of the Diatoms which are not fixed by a stipes, possess some power of spontaneous movement; and this is especially seen in those, whose frustules are of a long narrow form, such as that of the *Naviculæ* generally. The motion is of a peculiar kind, being usually a series of jerks, which carry forward the frustule in the direction of its length, and then carry it back through nearly the same path. Sometimes, however, the motion is smooth and equable; and this is especially the case with the curious *Bacillaria paradoxa* (Fig. 92, B), whose frustules slide over each other in one direction, until they are all but detached, and then slide as far in the opposite direction, repeating this alternate movement at very regular intervals.² In either case, the motion is obviously quite of a different nature from that of beings possessed of a power of self-direction. "An obstacle in the path," says Prof. W. Smith, "is not avoided, but pushed aside; or, if it be sufficient to avert the onward course of the frustule, the latter is detained for a time equal to that which it would have occupied in its forward progression, and then retires from the impediment as if it had accomplished its full course." The character of the movement is obviously similar to that of those motile forms of Protophyta which have been already described; but it has not yet been definitely traced to any organ of impulsion; and the cause of it is still obscure.³ By Prof. W. Smith it is referred to

¹ See on this subject a valuable Paper by Prof. W. Smith "On the Determination of Species in the *Diatomaceæ*," in the "Quart. Journ. of Microsc. Science," vol. iii, p. 130: a Memoir by Prof. W. Gregory "On shape of Outline as a specific character of *Diatomaceæ*," in "Trans. of Microsc. Soc.," 2d Series, vol. iii; and the Author's Presidential Address in the same volume, pp. 44-50.

² This curious phenomenon, the Author has himself more than once had the opportunity of witnessing.

³ Prof. Smith says:—"Among the hundreds of species which I have examined in every stage of growth and phase of movement, aided by glasses which have never been surpassed for clearness and definition, I have never been able to detect any semblance of a motile organ; nor have I, by coloring the fluid with carmine or indigo, been able to detect in the colored particles surrounding the Diatom, those rotary movements, which indicate, in the various species of true Infusorial animalcules, the presence of cilia." (Synopsis of British Diatomaceæ, Introduction, p. xxiv.)—Mr. Jabez Hogg, however, has recently stated ("Quart. Journ. of Microsc. Science," vol. iii, p. 235) that by the employment of the same mode of illumination as that by which the ciliary action may be discerned in *Closterium*, &c. (§ 164), a ciliary movement may be detected at the orifices which have already been described as existing in the siliceous envelope of the Diatomaceous frustule (§ 174). It may be questioned, however, whether this be anything else than an optical illusion, arising from the existence of *currents* at these orifices, produced by the vital actions going on within the cell, as noticed above.

forces operating within the frustule, and originating in the vital operations of growth, &c., which may cause the surrounding fluid to be drawn in through one set of apertures, and expelled through the other.¹ "If," as he remarks, "the motion be produced by the exosmose taking place alternately at one and the other extremity, while endosmose is proceeding at the other, an alternating movement would be the result in frustules of a linear form; whilst in others of an elliptical or orbicular outline, in which foramina exist along the entire line of suture, the movements, in any, must be irregular or slowly lateral. Such is precisely the case. The backward and forward movements of the *Navicula* have been already described; in *Surirella* (Fig. 88) and *Campylo-discus* (Fig. 87), the motion never proceeds further than a languid roll from one side to the other; and in *Gomphonema* (Fig. 89), in which a foramen, fulfilling the nutritive office, is found at the larger extremity only, the movement (which is only seen when the frustule is separated from its stipes) is a hardly perceptible advance in intermitted jerks in the direction of the narrow end."


180. The principles upon which this interesting group should be classified, cannot be properly determined, until the history of the Generative process,—of which nothing whatever is yet known in a large proportion of Diatoms, and very little in any of them,—shall have been thoroughly followed out. As already stated, there is a strong probability that many of the forms which are at present considered as distinct from each other, would prove to be but different states of the same, if their *whole* history were ascertained. On the other hand, it is by no means impossible that some which appear to be nearly related in the structure of their frustules and in their mode of growth, may prove to have quite different modes of reproduction. At present, therefore, *any* classification must be merely provisional; and in the notice now to be taken of some of the most interesting forms of the *Diatomaceæ*, the method of Prof. W. Smith, which is based upon the degree of connection that remains between the several frustules after self-division, will be adopted, since it possesses the advantage of being in accordance with the general "physiognomies" of these organisms, as it brings together those forms which correspond most closely in plan of growth; but it cannot be regarded as a truly natural classification, since it often separates genera which are closely allied in the structure of the individual frustules, as, for example, *Coscinodiscus* and *Meloseira*. The whole order is thus grouped, in the first instance, under two Tribes; the *first* including those which have the frustules naked, that is, neither

¹ It has been objected to this view, by the authors of the "Micrographic Dictionary," that, if such were the case, the like movements would be frequently met with in other minute unicellular organisms. They seem to have forgotten, however, that there are no other such organisms, in which the cell is almost entirely enclosed in an impermeable envelope, which limits the imbibition and expulsion of fluid to a small number of definite points, instead of allowing it to take place equally (as in other unicellular organisms) over the entire surface.

imbedded in gelatine, nor enclosed in a membranaceous tube; whilst the *second* is composed of those forms, whose frustules have a gelatinous or membranaceous envelope. The frustules of the Diatoms belonging to the first tribe, however, may have various degrees of connection; for whilst, in some (*a*), the union is dissolved almost immediately upon the completion of self-division, there are others (*b*) in which a gelatinous cushion or a stipes, to which the frustules are attached by a small portion of their surface, maintains a partial connection between the divided frustules, and others (*c*), again, in which the frustules remain in more or less complete cohesion, and form filaments, which, if the cohesion be limited to the angles of the frustules, are mere zigzag chains, but, if the cohesion extend to the entire surfaces of their sides, are continuous filaments, either flattened or cylindrical.

181. That section of the first tribe, in which the frustules are entirely disconnected from each other after the completion of their self-division, includes a number of beautiful discoidal forms, which seem to constitute a natural group, and may therefore be appropriately noticed in connection with each other. The genus *Coccinodiscus* is one of great interest, from the vast abundance of its valves in certain fossil deposits (Fig. 101, *a a a*), especially the Infusorial earth of Richmond in Virginia, of Bermuda, and of Oran, as also in Guano. Each frustule is of discoidal shape, being composed of two nearly flattened valves, united by a hoop; so that, if the frustules remained in adhesion, they would form a filament resembling that of *Meloseira* (Fig. 97). The regularity of the hexagonal divisions on the valves, renders them beautiful microscopic objects; in some species, the areolæ are smallest near the centre, and gradually increase in size towards the margin; in others, a few of the central areolæ are the largest, and the rest are of nearly uniform size. Most of the species are either marine, or are inhabitants of brackish water; when living, they are most commonly found adherent to sea-weeds or zoophytes; but when dead, the valves fall as a sediment to the bottom of the water. In both these conditions, they were found by Prof. J. Quekett in connection with Zoophytes which had been brought home from Melville Island by Sir E. Parry; and the species seemed to be identical with those of the Richmond earth. Nearly allied to the preceding is the beautiful genus *Actinocyclus* (Fig. 84), of which also the frustules are discoidal in form, but of which each valve, instead of being flat, has an undulating surface, as is seen in front view (B); giving to the side view (A)

FIG. 84.




Actinocyclus undulatus:—A, side view;
B, front view.

the appearance of being marked by radiating bands. Owing to this peculiarity of shape, the whole surface cannot be brought into focus at once, except with a lower power; and the difference of aspect which the different radial divisions present in Fig. 84, is simply due to the fact, that one set is out of focus, whilst the other is in it, since the appearances are reversed by merely altering the focal adjustment. The *number* of radial divisions has been considered a character of sufficient importance to serve for the distinction of species; but this is probably subject to variation; since we not unfrequently meet with disks, of which one has (say) 8 and another 10 such divisions, but which are so precisely alike in every other particular, that they can scarcely be accounted as specifically different. The valves of this genus are very abundant in the infusorial earths of Richmond, Bermuda, and Oran (Fig. 101, *b, b, b*); and many of the same species have been found recent in guano, and in the seas of various parts of the world. The frustules in their living state appear to be generally attached to sea-weeds or zoophytes.

182. The Bermuda earth also contains the very beautiful form (Fig. 85), which, though scarcely separable from *Actinocyclus*


FIG. 85.

*Heliopelta.*

except by its marginal spines, has received from Prof. Ehrenberg the distinctive appellation of *Heliopelta* (sun-shield). In the representation here given, the object is delineated as seen by the parabolic illuminator (§ 61), which brings into view certain features that can scarcely be seen by ordinary transmitted light. Four of the radial divisions are seen to be marked out into circular areolæ; but in the four which alternate with them, a minute granular structure is observable. This may be shown by careful adjustment of the focus, to exist over the whole of the valve, even on the divisions in which the circular areolation

is here displayed; and it hence appears that this marking is *superficial*, and that the circular areolation exists in a deeper layer of the siliceous lorica. In the alternating divisions whose surface is here displayed, the subjacent areolation, when brought into view by focussing down to it, is seen to be formed of equilateral triangles; it is not, however, nearly so well marked as the circular areolation of the first-mentioned divisions. The dark spots seen at the ends of the rays, like the dark centre, appear to be solid tubercles of silex, not traversed by markings, as in many other Diatoms; most assuredly they are *not* orifices, as supposed by Prof. Ehrenberg. Of this type, again, specimens are found presenting 6, 8, or 10 radial divisions, but in other respects exactly similar; on the other hand, two specimens agreeing in their number of divisions, may exhibit minute differences of other kinds; in fact it is rare to find two that are *precisely* alike. It seems probable, then, that we must allow a considerable latitude of variation in these forms, before attempting to separate any of them as distinct species. Another very

FIG. 86.

*Arachnoidiscus Ehrenbergii.*


beautiful discoidal Diatom, which occurs in guano, and is also found attached to sea-weeds from different parts of the world (especially to a species employed by the Japanese in making soup), is the *Arachnoidiscus* (Fig. 86), so named from the resemblance which the beautiful markings on its disk cause it to bear to a spider's web. According to Mr. Shadbolt,¹ who has carefully examined its structure, each valve consists of two layers; the outer one, a thin flexible horny membrane, indestructible by

¹ "Transactions of Microscopical Society," 1st Series, vol. iii, p. 49.

boiling nitric acid; the inner one, siliceous. It is the former which has upon it the peculiar spider's web-like markings; whilst it is the latter that forms the supporting framework, which bears a very strong resemblance to that of a circular Gothic window. The two can occasionally be separated entire, by first boiling the disks for a considerable time in nitric acid, and then carefully washing them in distilled water. Even without such separation, however, the distinctness of the two layers can be made out by focussing for each separately under a 1-4th or 1-5th in. objective; or by looking at a valve as an opaque object (either by the Parabolic illuminator, or by the Lieberkühn, or by a side light) with a 4-10ths in. objective, first from one side, and then from the other.

188. Nearly allied to the preceding in general characters, but differing in the triangular shape of its valves, is the *Triceratium*; of which striking form a considerable number of species are met with in the Bermuda and other Infusorial earths, while others are inhabitants of the existing ocean and of tidal rivers. The *T. favus* (Fig. 79), which is one of the largest and most regularly marked of any of these, occurs in the mud of the Thames and in various other estuaries on our own coast; it has been found, also, on the surface of the large sea-shells from various parts of the world, such as those of *Hippopus* and *Halio-tis*, before they have been cleaned; and it presents itself likewise in the infusorial earth of Petersburg (U. S.) Although the *triangular* form, when the frustule is looked at sideways, is that which is characteristic of the genus, yet in some of the species there seems a tendency to produce *quadrangular* and even *pentagonal* forms; these being marked as *varieties*, by their exact correspondence in sculpture, color, &c., with the normal triangular forms.¹ This departure is extremely remarkable, since it

breaks down what seems at first to be the most distinctive character of the genus; and its occurrence is an indication of the degree of latitude which we ought to allow in other cases. The genus *Campylo-discus* (Fig. 87) is distinguished by its saddle-shaped curvature, and by its ribbed markings, which




Campylo-discus costatus.—A, front view; B, side view.

seem to indicate the presence of canals excavated in or beneath

¹ See Mr. Brightwell's excellent memoir "On the genus *Triceratium*," in "Quart. Microsc. Journal," vol. i, p. 245.

the valves, for the passage of fluid between the orifices in the siliceous envelope and the soft cell-membrane beneath. The form of the valves, in most of the species, is circular, or nearly so; some are nearly flat, whilst in others the twist is greater than in the species here represented. Some of the species are marine, whilst others occur in fresh water; a very beautiful form, the *C. clypeus*, exists in such abundance in the Infusorial stratum discovered by Prof. Ehrenberg at Soos near Ezer in Bohemia, that the earth seems almost entirely composed of it. Some of the forms of the last genus lead towards *Surirella* (Fig. 88), which, like it, presents the appearance of possessing a canalicular system, though this is by no means equally conspicuous in all the species. The distinctive character of the genus, in addition to the presence of the canaliculi, is derived from the longitudinal line down the centre of each valve (A), and the prolongation of the margins into "alæ." Numerous species are known, which are mostly of a somewhat ovate form, some being broader and others narrower than *S. con-*




Surirella constricta:—A, side view; B, front view; C, binary subdivision.

stricta; the greater part of them are inhabitants of fresh or brackish water, though some few are marine; and several occur in those infusorial earths, which seem to have been deposited at the bottoms of lakes, such as that of the Mourne Mountains in Ireland (Fig. 102, b, c, k).

184. We now come to that interesting series of forms, which has been ranked under the genus *Navicula*, until its recent subdivision by Prof. W. Smith into the three genera *Navicula*, *Pinnularia*, and *Pleurosigma*, each of which still comprehends a very large number of species. They are all distinguished by the oblong or lanceolate form of their valves, by the convexity of their surfaces, by the presence of a longitudinal line along the middle line of each valve, dilating into nodules at the centre and extremities, and by the more or less conspicuous marking of the valves with transverse or oblique striæ, which, under a sufficient magnifying power, are resolvable into a regular areolation, resembling, though on a very minute scale, that of the more coarsely marked Diatoms (§ 175). The genus *Navicula*, as now constituted, is distinguished by these striæ, and by the appearance of rows of circular dots which they present, when sufficiently magnified. In *Pinnularia* (Fig. 102, h), the striæ are not resolvable into dots, and are so strongly marked as almost to resemble the "ribs" of

Surirella. The genus *Pleurosigma* is at once distinguished by the peculiar curvature of its valves (Fig. 80); and it is further remarkable for the extreme closeness of their striation, and for the consequent "difficulty" which attends its resolution into a regular areolation. The species of the first two of these genera are for the most part inhabitants of fresh water; and they constitute a large part of most of the Infusorial Earths which were deposited at the bottoms of lakes. Among the most remarkable of such deposits, are the substances largely used in the arts for the polishing of metals, under the names of Tripoli and rottenstone; these consist in great part of the frustules of *Naviculæ* and *Pinnulariæ*. The Polierschiefer or polishing-slate of Bilin in Bohemia, the powder of which is largely used in Germany for the same purpose, and which also furnishes the fine sand used for the most delicate castings in iron, occurs in a series of beds averaging fourteen feet in thickness; and these present appear-

FIG. 89.



Gomphonema geminatum: its frustules connected by stipes.

ances which indicate that they have been at some time exposed to a high temperature. The well-known Turkey-stone, so generally employed for the sharpening of edge tools, seems to be essentially composed of a similar aggregation of frustules of *Naviculæ*, &c., which has been consolidated by heat. The species of *Pleurosigma*, on the other hand, are for the most part either marine, or are inhabitants of brackish water; and they comparatively seldom present themselves in a fossilized state. The genus *Stauroneis*, which belongs to the same group, differs from all the preceding forms, in having the central nodule of each valve dilated laterally into a band free from striæ, which forms a cross with the longitudinal band; of this very beautiful form, some species are fresh water, others marine; and the former present themselves frequently in certain infusorial earths.¹

185. The group we have next to notice, consists of those genera which have the frustules,

¹ For some very curious examples of the extent to which variation in form, size, and distance of striæ, may take place in this group, among individuals which must be accounted as of the same species, see the memoirs of Profra. W. Smith and W. Gregory already referred to (p. 287, note).

after self-division, attached by a gelatinous cushion, or by a dichotomous stipes (Fig. 89). Many of these present a strong resemblance to some of the preceding, so far as the structure of their frustules is itself concerned; so that when, as frequently happens, they are found unattached, the difference is not apparent. In *Synedra* (Fig. 102, *l*), which is not unlike a long narrow *Navicula* (an imperfect longitudinal line, with central and terminal dilatations, being often, but not always, apparent), the frustules are at first invariably attached to larger Algæ, or other aquatic plants, by a cushion-like gelatinous basis; and when they remain adherent to this after repeated subdivision, they sometimes form a fan-like band of frustules, not unlike that of *Licmophora* (Fig. 91), or even a stellate cluster, the appearance of which is extremely characteristic. So again, the frustules of *Gomphonema* (Fig. 90) in a side view, are not unlike those of *Navicula*; but they are distinguished in front view by their cuneate (wedge-like) form, which arises chiefly from the unequal development of the membrane connecting the valves, at the upper and lower ends. The stipes seems to be formed by an exudation from the frustule, which is secreted only during the process of self-division; hence when this process has been completed, the extension of the single filament below the frustule ceases; but when it recommences, a sort of joint or articulation is formed, from which a new filament begins to sprout for each of the half-frustules; and when these separate, they carry apart the peduncles which support them, as far as their divergence can take place; and it is in this manner that the dichotomous character is given to the entire stipes (Fig. 89). The species of *Gomphonema* are, with scarcely an exception, inhabitants of fresh water; and are among the commonest forms

of Diatomaceæ. In *Licmophora* (Fig. 91) we meet with a different mode of growth; for the newly formed part of the stipes, instead of itself becoming double with each act of self-division of the frustule, increases in breadth, while the frustules themselves remain coherent; so that a beautiful fan-like arrangement is produced. A splitting away of a few frustules seems occasionally to take place, from one side or the other, before the elongation of the stipes; so that the entire plant presents us with a more or less complete *flabella* or fan upon the summit of the branches, with imperfect *flabellæ* or single frustules irregularly scattered throughout the entire length of the footstalk. This beautiful plant is marine, and is parasitic upon sea-weeds and zoophytes.





FIG. 90.
Gomphonema geminatum.—A, side view of frustule more highly magnified; B, front view; C, frustule in the act of self-division.

186. In the group at which we now arrive, there is more or less of permanent connection between the frustules themselves; and this may depend merely upon the cohesion of their surfaces, or may

FIG. 91.




Licmophora flabellata.

be occasioned by the persistence of the connecting membrane of the valves after the completion of the self-division of the frustules. To the former division belong several genera, the form of whose frustules is more or less elongated, so that the filament formed by their cohesion is a flattened band. If the two extremities of the frustule be of equal breadth, as in *Bacillaria*, the band will be straight; but if one be broader than the other, so that the frustule in front view has a cuneate or wedge-like form, the filament will be curved, as in the beautiful *Meridion circulare* (Fig. 92, A). Although these, when gathered and placed

under the microscope, present the appearance of circles overlying one another, they really grow in a helical (screw-like) form, making several continuous

FIG. 92.



A, *Meridion circulare*.—B, *Bacillaria paradoxa*.

turns. This Diatom abounds in many localities in this country; but there is none in which it presents itself in such rich luxuriance, as in the mountain brooks about West Point in the United States, the bottoms of which, according to Prof. Bailey,

“are literally covered in the first warm days of spring with a ferruginous colored mucous matter, about a quarter of an inch thick, which on examination by the microscope, proves to be filled with millions and millions of these exquisitely beautiful siliceous bodies. Every submerged stone, twig and spear of grass is enveloped by them; and the waving plume-like appearance of a filamentous body covered in this way is often very elegant.” The genus *Bacillaria*, so named from the staff-like form of its frustules, is now limited to the species *B. paradoxa* (Fig. 92, b), whose remarkable movements have been already described (§ 179). Owing to this displacement of the frustules, its filaments seldom present themselves with straight parallel sides, but nearly always in forms more or less oblique, such as those represented above. This curious object is an inhabitant of salt or of brackish water. Many of the species formerly ranked under this genus are now referred to the genus *Diatoma* (§ 187); to which also the genus *Fragillaria* is nearly allied, the difference between them lying chiefly in the mode of adhesion of the frustules. These in *Fragillaria* form long straight filaments with parallel sides; the filaments, however, as the name of the genus implies, very readily break up into their component frustules, often separating at the slightest touch. Its various species are very common in pools and ditches. Among the numerous genera belonging to this group, we may stop to notice *Achnanthes*; some of the species of which, particularly *A. longipes* (Fig. 93), are furnished with a single nearly straight stipes, attached to one end of the lower margin of the first frustule of the filament. There is a curious difference in the markings of the valves of the upper and lower frustules; for while both are traversed by striæ, which are resolvable under a sufficient power into rows of dots, as well as by a longitudinal line, which sometimes has a nodule at each end (as in *Navicula*), the lower valve (*a*), has also a transverse line, forming a *stauros* or cross, which is wanting in the upper valve (*e*). A persistence of the connecting membrane may sometimes be observed in this genus, so as to form an additional connection between the cells; thus, in Fig. 93, it not only holds together the two new frustules resulting from the subdivision of the lowest cell, *a*, which are not yet completely separated the one from the other, but it may be observed to invest the two frustules *b* and *c*, which have not merely separated, but are themselves beginning to undergo binary subdivision; and it may also

FIG. 93.



Achnanthes longipes :—
a, b, c, d, e, successive
frustules in different
stages of self-division.

be observed to invest the frustule *d*, from which the frustule *e*, being the terminal one, has more completely freed itself.

187. It is by the persistence of the connecting membrane, notwithstanding the dissolution of the adhesion between the surfaces of the valves, that the frustules are so held together after self-division as to form zigzag chains in the next group; of which the genus *Diatoma* (Fig. 94), that gives the name ("cutting through") to the whole order, is a typical example. Its

FIG. 94.




FIG. 95.




FIG. 96.




Fig. 94. *Diatoma vulgare*:—*a*, side view of frustule; *b*, frustule undergoing self-division.

Fig. 95. *Grammatophora serpentina*:—*a*, front and side views of single frustule; *b*, *b*, front and end views of divided frustule; *c*, a frustule about to undergo self-division; *d*, a frustule completely divided.

Fig. 96. *Lathmia nervosa*.

valves, when turned sideways, are seen to be strongly marked by transverse striæ, which extend into the front view. The proportion between the length and the breadth of each valve is found to vary so considerably, that, if the extreme forms only were compared, there would seem adequate ground for regarding them as belonging to different species. This genus inhabits fresh water, preferring gently running streams, in which it is sometimes very abundant. The genus *Grammatophora* (Fig. 95) is nearly allied to the preceding; but its transverse striæ are extremely faint and difficult of detection, so that its valves serve as "test-objects" (§ 102, III); and the frustules, when seen in front view, are marked by peculiar bands, usually sinuous, which are termed *vittæ*. The curious *Biddulphia*, whose self-division has already been described (§ 176), belongs to this group; its frustules have a very peculiar external form (Fig. 81), and they are believed to be divided internally by partitions which correspond

to the external ribbings. So far as is yet known, it is exclusively marine. Nearly allied to it, is the beautiful genus *Isthmia* (Fig. 96), which is also marine, and which grows attached to larger sea-weeds, the basal frustule being very commonly attached by a stipes. In this, as in the preceding genus, the areolated structure of the surface is very conspicuous (Fig. 78), both in the valves and in the connecting "hoop;" and this hoop, being silicified, not only connects the two new frustules (as at *b*), until they have separated from each other, but, after such separation, remains for a time round one of the frustules, so as to give it a truncated appearance (*a, c*).

188. It is by the complete persistence of the siliceous hoop, that the frustules are held together in that curious group of Diatomaceæ, which consists of a few genera whose cylindrical filaments bear a close external resemblance to those of *Confer-vaceæ*. The most important of these is *Meloseira* (Figs. 97, 98), long since characterized as a plant by the Swedish algologist Agardh, but taken from

the Vegetable kingdom with other Diatoms by Prof. Ehrenberg, who included its species in his genus *Gallionella*. Some of its species are marine, others fresh water; one of the latter, the *M. ochracea*, seems to grow best in boggy pools containing a ferruginous impregnation; and it is stated by Prof. Ehrenberg to take up from the water, and to incorporate with its own substance, a considerable quantity of iron. The filaments of *Meloseira* very commonly fall apart at the slightest touch; and in the infusorial earths, in which

some species abound, the frustules are always found detached (Fig. 102, *a a, d d*). The meaning of the remarkable difference in the sizes and forms of the frustules of the same filaments (Fig. 97, 98) has not yet been fully ascertained; but it seems to be related to the curious process of self-conjugation already described (§ 178). The sides of the valves are often marked with radiating striæ (Fig. 102, *d d*); and in some species they have toothed or serrated margins, by which the frustules lock together.

189. The Second tribe of Diatomaceæ, in which the frustules are completely enveloped by a gelatinous or membranaceous

FIG. 97.

FIG. 98.

*Meloseira subflexilis.**Meloseira varians.*

envelope, does not contain many forms so interesting as those which have been already described; and it will not be requisite to dwell long upon it. This envelope forms a *frond*, that holds together all the frustules which have originated in the self-division of one individual; and it may either consist of an indefinite gelatinous mass, or rather aggregation of masses, in which the frustules are imbedded (Fig. 100), or of an indefinite cluster

FIG. 99.

FIG. 100.




Fig. 99. *Mastogloia Smithii*:—A, entire stipes; B, frustule in its gelatinous envelope; C—F, different forms of frustule as seen in front view; G, side view; H, frustule undergoing subdivision.

Fig. 100. *Mastogloia lanceolata*.

of such masses, supported by a stipes (Fig. 99), as in the genus *Mastogloia*; or it may have a definite shape, either globular, compressed, or filamentous, within which the frustules either lie scattered, as in *Dickieia* and *Berkeleyia*, or in rows, as in *Schisonema*, or in bundles, as in *Homæocladia*. These are all, or nearly all, marine forms; and many of them present so strong a resemblance to the smaller filamentous Algæ, that they might easily be mistaken for such. Their frustules, however, when taken out of their containing tubes, frequently present so near an approximation, both in structure and markings, to members of the preceding groups of Diatomacæ, that their close relationship to them cannot be questioned. Very strongly marked varieties sometimes present themselves within the limits of a single species; thus the valves C, D, E, F (Fig. 99), would scarcely have been supposed to belong to the same specific type, were they not found upon the same stipes. The careful study of these varieties, in every instance in which any disposi-

tion to variation shows itself, so as to *reduce* the enormous number of species with which our systematic treatises are loaded, is a pursuit of far greater real value, than the *multiplication* of species by the detection of such minute differences as may be presented by forms discovered in newly explored localities; such differences, as already pointed out, being probably, in a large proportion of cases, the result of the multiplication of some one form, which, under modifying influences that we do not yet understand, has departed from the ordinary type. The more faithfully and comprehensively this study is carried out, in *any* department of Natural History, the more does it prove that the range of variation is far more extensive than had been previously imagined; and this is especially likely to be the case with such humble organisms as those we have been considering; since they are obviously more influenced than are those of higher types, by the conditions under which they are developed; whilst, from the very wide geographical range through which the same forms are diffused, they are subject to very great diversities of such conditions.

190. The general habits of this most interesting group cannot be better stated than in the words of Prof. W. Smith, from whose admirable Monograph the Author has drawn the greater part of his materials for the foregoing account of it.¹ "The Diatomacæ inhabit the sea, or fresh water; but the species peculiar to the one, are never found in a living state in any other locality; though there are some which prefer a medium of a mixed nature, and are only to be met with in water more or less brackish. The latter are often found in great abundance and variety in districts occasionally subject to marine influences, such as marshes in the neighborhood of the sea, or the deltas of rivers, where, on the occurrence of high tides, the freshness of the water is affected by percolation from the adjoining stream, or more directly by the occasional overflow of its banks. Other favorite habitats of the Diatomacæ are stones of mountain streams or waterfalls, and the shallow pools left by the retiring tide at the mouths of our larger rivers. They are not, however, confined to the localities I have mentioned,—they are, in fact, most ubiquitous, and there is hardly a roadside ditch, water-trough, or cistern, which will not reward a search, and furnish specimens of the tribe." Such is their abundance in some rivers and estuaries, that their multiplication is affirmed by Prof. Ehrenberg to have exercised an important influence in blocking up harbors and diminishing the depth of channels!

¹ The Author has great pleasure in here acknowledging the liberality of Messrs. Smith and Beck, who have allowed him, not only to make free use of this volume, but also to copy as many as he desired of the admirable series of illustrations which have been executed for it by Mr. Tuffen West, many of them still unpublished. All the figures of Diatomacæ, given in this Manual, except Figs. 80 and 85, which are drawn from nature, and Figs. 101, 102, which are copied from Prof. Ehrenberg, are as exact copies of Mr. West's lithographs as the best Wood-engraving can produce.

Of their extraordinary abundance in certain parts of the ocean, the best evidence is afforded by the observations of Dr. W. J. Hooker upon the Diatomaceæ of the southern seas; for within the Antarctic Circle, they are rendered peculiarly conspicuous by becoming enclosed in the newly-formed ice, and by being washed up in myriads by the sea on to the "pack" and "bergs," everywhere staining the white ice and snow of a pale ochreous brown. A deposit of mud, chiefly consisting of the siliceous loriceæ of Diatomaceæ, not less than 400 miles long and 120 miles broad, was found at a depth of between 200 and 400 feet, on the flanks of Victoria Land in 78° south latitude; of the thickness of this deposit no conjecture could be formed; but that it must be continually increasing is evident, the silex of which it is in a great measure composed being indestructible. A fact of peculiar interest in connection with this deposit, is its extension over the submarine flanks of Mount Erebus, an active volcano of 12,400 feet elevation; since a communication between the ocean-waters and the bowels of a volcano, such as there are other reasons for believing to be occasionally formed, would account for the presence of Diatomaceæ in volcanic ashes and pumice, which was discovered by Prof. Ehrenberg. It is remarked by Dr. Hooker, that the universal presence of this in-

FIG. 101.



Fossil Diatomaceæ, &c., from Oran:—*a, a, a*, *Coscinodiscus*; *b, b, b*, *Actinocyclus*; *c*, *Dictyochya fibula*; *d*, *Lithasteriscus radiatus*; *e*, *Spongolithis acicularis*; *f, f*, *Grammatophora parallela* (side view); *g, g*, *Grammatophora angulosa* (front view).

visible vegetation throughout the South Polar Ocean, is a most important feature; since there is a marked deficiency, in this region, of higher forms of vegetation; and were it not for them,

there would neither be food for aquatic animals, nor (if it were possible for these to maintain themselves by preying on one another) could the ocean-waters be purified of the carbonic acid which animal respiration and decomposition would be continually imparting to it. It is interesting to observe, that some species of marine Diatomaceæ are found through every degree of latitude between Spitzbergen and Victoria Land; whilst others seem limited to particular regions. One of the most singular instances of the preservation of Diatomaceous forms, is their existence in Guano; into which they must have passed from the intestinal canals of the birds of whose accumulated excrement that substance is composed, those birds having received them, it is probable, from shell-fish, to which these minute organisms serve as ordinary food (§ 192.)

191. The indestructible nature of the epiderms of *Diatomaceæ* has also served to perpetuate their presence in numerous localities, from which their living forms have long since disappeared; for the accumulation of sediment formed by their successive production and death, either on the bed of the ocean, or on the bottoms of fresh-water lakes, gives rise to deposits which may attain considerable thickness, and which, by subsequent changes of level, may come to form part of the dry land. Thus very extensive siliceous strata, consisting almost entirely of marine *Diatomaceæ*, are found to alternate, in the neighborhood of the Mediterranean, with calcareous strata, chiefly formed of *Foraminifera* (Chap. X); the whole series being the representative of the Chalk formation of Northern Europe, in which the silex that was probably deposited at first in this form, has undergone conversion into *flint*, by agencies hereafter to be considered (Chaps. X, XIX). Of the Diatomaceous composition of these strata, we have a characteristic example in Fig. 101, which represents the fossil Diatomaceæ of Oran, in Algeria. The so-called "infusorial earth" of Richmond, in Virginia, and that of Bermuda, also marine deposits, are very celebrated among Microscopists for the number and beauty of the forms they have yielded; the former constitutes a stratum of 18 feet in thickness, underlying the whole city, and extending over an area whose limits are not known. Several deposits of more limited extent, and apparently of fresh-water origin, have been found in our own islands; as for instance at Dolgelly, in North Wales, at Lough Mourne, in Ireland (Fig. 102), and in the island of Mull, in Scotland. Similar deposits in Sweden and Norway are known under the name of *berg-mehl* or mountain flour; and in times of scarcity, the inhabitants of those countries are accustomed to mix these substances with their dough in making bread. This has been supposed merely to have the effect of giving increased bulk to their loaves, so as to render the really nutritive portion more satisfying. But as the *berg-mehl* has been found to lose from a quarter to a third of its weight by exposure to a red-heat, there seems a strong proba-

bility that it contains organic matter, which renders it nutritious in itself. When thus occurring in strata of a fossil or sub-fossil character, the Diatomaceous deposits are generally distinguishable as white or cream-colored powders of extreme fineness.

192. For collecting fresh *Diatomaceæ*, three general methods are to be had recourse to, which have been already described (§ 148, 171). "Their living masses," says Prof. W. Smith, "pre-

FIG. 102.



Fossil Diatomaceæ, &c., from Mourne Mountain, Ireland:—*a, a, a*, *Gallonella (Melosira) proceræ* and *G. granulata*; *d, d, d*, *G. biseriata* (side view); *b, b*, *Suirella plicata*; *e*, *S. craticula*; *k*, *S. caldonica*; *g*, *Gomphonema gracile*; *f*, *Cocconeum fusidium*; *h*, *Tabellaria vulgaris*; *l*, *Pinnularia dactylus*; *i*, *P. nobilis*; *j*, *Synedra ulna*.

sent themselves as colored fringes attached to larger plants, or forming a covering to stones or rocks in cushion-like tufts—or spread over their surface as delicate velvet—or depositing themselves as a filmy stratum on the mud, or intermixed with the scum of living or decayed vegetation floating on the surface of the water. Their color is usually a yellowish-brown of a greater or less intensity, varying from a light chestnut, in individual specimens, to a shade almost approaching black in the aggregated masses. Their presence may often be detected without the aid of a microscope, by the absence, in many species, of the fibrous tenacity which distinguishes other plants; when removed from their natural position, they become distributed through the water, and are held in suspension by it, only subsiding after some little time has elapsed." Notwithstanding every care, the collected specimens are liable to be mixed with much foreign matter; this may be partly got rid of by repeated washings in pure water, and, by taking advantage, at the same time, of the different specific gravities of the Diatoms and of the intermixed

substances, to secure their separation. Sand, being the heaviest, will subside first; fine particles of mud, on the other hand, will float after the Diatoms have subsided. The tendency which the Diatomaceæ have, to make their way towards the light, will afford much assistance in procuring the free forms in a tolerably clean state; for if the gathering which contains them be left undisturbed for a sufficient length of time, in a shallow vessel exposed to the sunlight, they may be skimmed from the surface. The marine forms must be looked for upon sea-weeds, and in the fine mud or sand of soundings or dredgings; they are frequently found also, in considerable numbers, in the stomachs of the oyster, scallop, whelk, and other Mollusks, especially the "bivalves," in those of the crab and lobster, and even in those of the sole, turbot, and other "flat-fish." "Several species," says Prof. W. Smith, "rarely or never occurring in my usual haunts, have been supplied in abundance by the careful dissection of the above microphagists." The separation of the Diatoms from the other contents of these stomachs, must be accomplished by the same process as that by which they are obtained from Guano or the calcareous Infusorial Earths; of this, the following are the most essential particulars. The guano or earth is first to be washed several times in pure water, which should be well stirred, and the sediment then allowed to subside for some hours before the water is poured off, since, if it be decanted too soon, it may carry the lighter forms away with it. Some kinds of earth have so little impurity, that one washing suffices; but in any case it is to be continued so long as the water remains colored. The deposit is then to be treated, in a flask or test tube, with hydrochloric (muriatic) acid; and after the first effervescence is over, a gentle heat may be applied. As soon as the action has ceased, and time has been given for the sediment to subside, the acid should be poured off, and another portion added; and this should be repeated as often as any effect is produced. When hydrochloric acid ceases to act, strong nitric acid should be substituted; and after the first effervescence is over, a continued heat of about 200° should be applied for some hours. When sufficient time has been given for subsidence, the acid may be poured off, and the sediment treated with another portion; and this is to be repeated, until no further action takes place. The sediment is then to be washed, until all trace of the acid is removed; and if there have been no admixture of siliceous sand in the earth or guano, this sediment will consist almost entirely of Diatomaceæ, with the addition, perhaps, of Sponge-spicules. The separation of siliceous sand, and the subdivision of the entire aggregate of Diatoms into the larger and the finer kinds, may be accomplished by stirring the sediment in a tall jar of water, and then, while it is still in motion, pouring off the supernatant fluid as soon as the coarser particles have subsided; this fluid should be set aside, and, as soon as a finer sediment has subsided,

it should again be poured off; and this process may be repeated three or four times at increasing intervals, until no further sediment subsides after the lapse of half an hour. The first sediment will probably contain all the sandy particles, with, perhaps, some of the largest Diatoms, which may be picked out from among them; and the subsequent sediments will consist almost exclusively of Diatoms, the sizes of which will be so graduated, that the earliest sediments may be examined with the lower powers, the next with the medium powers, while the latest will require the higher powers,—a separation which is attended with great convenience.¹

193. The mode of mounting specimens of Diatomaceæ, will depend upon the purpose which they are intended to serve. If they can be obtained quite fresh, and it be desired that they should exhibit, as closely as possible, the appearance presented by the living plants, they should be put up in distilled water within cement cells (§ 134); but if they are not thus mounted within a short time after they have been gathered, about a sixth part of alcohol should be added to the water. If it be desired to exhibit the stipitate forms in their natural parasitism upon other aquatic plants, the entire mass may be mounted in Deane's gelatine (§ 131), in a deeper cell; and such a preparation is a very beautiful object for the black-ground illumination. If, on the other hand, the minute structure of the siliceous envelopes is the feature to be brought into view, the fresh Diatoms must be boiled in nitric or hydrochloric acid, which must then be poured off (sufficient time being allowed for the deposit of the residue), and the sediment, after repeated washings, is to be either mounted in balsam in the ordinary manner (§ 128), or, if the species have markings that are peculiarly difficult of resolution, is to be set up dry between two pieces of thin glass (§ 122). In order to obtain a satisfactory view of these markings, object-glasses of very wide angle of aperture are required, and all the refinements which have recently been introduced into the methods of Illumination, need to be put in practice. (Chaps. III, IV.) It will often be convenient to mount certain particular forms of Diatomaceæ separately from the general aggregate; but on account of their minuteness, they cannot be selected and removed by the usual means. The larger forms, which may be readily distinguished under a simple microscope, may be taken up by a camel-hair pencil, which has been so trimmed as to leave two or three hairs projecting beyond the rest. But the smaller can only be dealt with by a single fine bristle or stout sable-hair, which may be inserted into the cleft end of a slender wooden handle; and if the bristle or hair should be split at its extremity in a brush-like manner, it will be particularly useful.


¹ A somewhat more complicated method of applying the same principle, is described by Mr. Okeden in the "Quart. Journ. of Microsc. Science," vol. iii, p. 158. The Author believes, however, that the method above described will answer every purpose.

Such split hairs (as Dr. Redfern first noticed) may always be found in a shaving brush which has been for some time in use. Those should be selected, which have thin split portions so closely in contact, that they appear single until touched at their ends. When the split extremity of such a hair touches the glass slide, its parts separate from each other to an amount proportionate to the pressure; and, on being brought up to the object, first pushed to the edge of the fluid on the slide, may generally be made to seize it. Supposing that we wish to select certain particular forms, from a Diatomaceous sediment which has been obtained by the preceding processes; either of the two following modes may be put in practice. A small portion of the sediment being taken up in the dipping-tube, and allowed to escape upon the slide, so as to form a long narrow line upon it, this is to be examined with the lowest power with which the object we are in search of can be distinguished (the erector and draw-tube, §§ 43, 44, will here be very useful); and when one of the specimens has been found, it may be taken up, if possible, on the point of the hair, and transferred to a new slide, to which it may be made to adhere by first breathing on its surface. But if it be found impracticable thus to remove the specimens, on account of their minuteness, they may be pushed to one side of the slide on which they are lying; all the remainder of the sediment which it is not desired to preserve, may be washed off; and the objects may then be pushed back into the middle of the slide, and mounted in any way that may be desired.

194. *Palmellaceæ*.—To the little group of Plants which is ranked under this designation, those two genera belong, which have been already cited as illustrations of the humblest types of vegetation (§§ 150, 152); and the other forms which are associated with these, are scarcely less simple in their essential characters, though sometimes attaining considerable dimensions. They all grow either on damp surfaces, or in fresh or salt water; and they may either form (1) a mere powdery layer, of which the component particles have little or no adhesion to each other, or they may present themselves (2) in the condition of an indefinite slimy film, or (3) in that of a tolerably firm and definitely bounded membranous "frond." The *first* of these states we have seen to be characteristic of *Palmoglæa* and *Protococcus*: the new cells which are originated by the process of duplicative subdivision, usually separating from each other after a short time; and even where they remain in cohesion, nothing like a frond or membranous expansion being formed. The "red snow," which sometimes colors extensive tracts in Arctic or Alpine regions, penetrating even to the depth of several feet, and vegetating actively at a temperature which reduces most plants to a state of torpor, is generally considered to be a species of *Protococcus*; but as its cells are connected by a tolerably firm gelatinous investment, it would rather seem to be a *Palmella*. The

second is the condition of the genus *Palmella*; of which one species, the *P. cruenta*, usually known under the name of "gory dew," is common on damp walls and shady places, sometimes extending itself over a considerable area as a tough gelatinous mass, of the color and general appearance of coagulated blood. A characteristic illustration of it is also afforded by the *Hæmatococcus sanguineus* (Fig. 103), which chiefly differs from *Palmella*

FIG. 103.



Hamatococcus sanguineus, in various stages of development:—*a*, single cells, enclosed in their mucous envelope; *b*, *c*, clusters formed by subdivision of parent cell; *d*, more numerous cluster, its component cells in various stages of division; *e*, large mass of young cells, formed by the continuance of the same process, and enclosed within a common mucous envelope.

in the partial persistence of the walls of the parent cells, so that the whole mass is subdivided by partitions, which enclose a larger or smaller number of cells originating in the subdivision of their contents. Besides increasing in the ordinary mode of binary multiplication, the *Palmella* cells seem occasionally to rupture and diffuse their granular contents through the gelatinous stratum, and thus to give origin to a whole cluster at once, as seen at *e*, after the manner of other simple Plants to be

presently described (§ 195), save that these minute segments of the endochrome, having no power of spontaneous motion, cannot be ranked as Zoospores. The gelatinous masses of the *Palmelleæ* are frequently found to contain parasitic growths, formed by the extension of other plants through their substance; but numerous branched filaments sometimes present themselves, which, being traceable into absolute continuity with the cells, must be considered as properly appertaining to them. Sometimes these filaments radiate in various directions from a single central cell, and must at first be considered as mere extensions of this; their extremities dilate, however, into new cells; and, when these are fully formed, the tubular connections close up, and the cells become detached from each other.¹ This is obviously an additional mode of increase by *gemmation*; analogous to that which we shall meet with among the *Conferveæ* (§ 198).

¹ Although the Authors of the "Micrographic Dictionary" throw doubt upon this fact, yet the writer, having had the opportunity of verifying the observations of that most accurate Algologist, Mr. Thwaites, can entertain no doubt of their correctness. (See "Ann. of Nat. Hist." N. S. vol. ii, p. 313.)

Of the *third* condition, we have an example in the curious *Palmodyctyon* described by Kützing; the frond of which appears to the naked eye like a delicate network consisting of anastomosing branches, each composed of a single or double row of large vesicles, within every one of which is produced a pair of elliptical cellules that ultimately escape as "zoospores." The alternation between the "motile" form and the "still" or resting form, which has been described as occurring in *Protococcus* (§§ 153-155), has been observed in several other forms of this group; and it seems obviously intended, like the production of "zoospores," to secure the dispersion of the plant, and to prevent it from choking itself by overgrowth in any one locality. No other form of the true Generative process, than the "conjugation" of two cells, and the formation of a spore by the reunion of their contents, has yet been observed among the members of this family; and as this has only been witnessed in a few of them, it is probable that we do not yet know by any means the whole of their life-history. And from the close resemblance which many reputed *Palmellaceæ* bear to the early stages of higher Plants (Fig. 104, A, B, c), there is considerable doubt whether they ought to be regarded in the light of distinct and complete organisms, or whether they are anything else than embryonic forms of more elevated types. Here, again, therefore, there is an ample field for investigation to the Microscopist who desires to employ himself in extending the boundaries of Science, and in perfecting our very imperfect acquaintance with these humble but most interesting and instructive types of Vegetation.

195. Notwithstanding the very definite form and large size attained by the fronds or leafy expansions of the *Ulvaceæ*, to which group belong the grass-green seaweeds (or "lavers") found on every coast, yet their essential structure differs but very little from that of the preceding group; and the principal advance is shown in this, that the cells, when multiplied by


Fig. 104.



Successive stages of development of *Ulva*.

binary subdivision, not only remain in firm connection with each other, but possess a very regular arrangement (in virtue of the determinate plan on which the subdivision takes place), and form a regular membranous stratum. The mode in which this frond is produced, is best understood by studying the history of its development, some of the principal phases of which are seen in Fig. 104; for the isolated cells (A), in which it originates, resembling in all points those of a *Protococcus*, give rise, by their successive subdivisions in determinate directions, to such regular clusters as those seen at B and C, or to such confervoid filaments as that shown at D. A continuation of the same regular mode of subdivision, taking place alternately in two directions, may at once extend the clusters B and C into leaf-like expansions; or, if the filamentous stage be passed through (different species presenting variations in the history of their development), the filament increases in breadth as well as in length (as seen at E), and finally becomes such a frond as is shown at F, G. In the simple membranous expansions thus formed, there is no approach to a "differentiation" of parts, by even the semblance of a formation of root, stem, and leaf, such as the higher Algæ present; every portion is the exact counterpart of every other; and every portion seems to take an equal share in the operations of growth and reproduction. Each cell is very commonly found to exhibit an imperfect partitioning

FIG. 105.



Formation of Zoospores in *Phycoseris gigantea* (*Ulva latissima*);—a, portion of the ordinary frond; b, cells in which the endochrome is beginning to break up into segments; c, cells from the boundary between the colored and colorless portion, some of them containing zoospores, others being empty; d, ciliated zoospores, as in active motion; e, subsequent development of the zoospores.

into four parts, preparatory to multiplication by double subdivision; and the entire frond usually shows the groups of cells arranged in clusters containing some multiple of four. Besides this continuous increase of the individual frond, however, we

find in most species of *Ulva* a provision for extending the plant by the dispersion of "zoospores;" for the endochrome (Fig. 105, *a*) subdivides into numerous segments (as at *b* and *c*), which at first are seen to lie in close contact within the cell that contains them, then begin to exhibit a kind of restless motion, and at last pass forth through an aperture in the cell-wall, acquire four or more cilia (*d*), and swim freely through the water for some time. At last, however, they come to rest, attach themselves to some fixed point, and begin to grow into clusters or filaments (*e*), in the manner already described. The walls of the cells which have thus discharged their endochrome, remain as colorless spots on the frond; sometimes these are intermingled with the portions still vegetating in the usual mode; but sometimes the whole endochrome of one portion of the frond may thus escape in the form of zoospores, thus leaving behind it nothing but a white flaccid membrane. If the Microscopist who meets with a frond of an *Ulva* in this condition should examine the line of separation between its green and its colored portion, he may not improbably meet with cells in the very act of discharging their zoospores, which "swarm" around their points of exit, very much in the manner that *Animalcules* are often seen to do around particular spots of the field of view, and which might easily be taken for true *Infusoria*; but on carrying his observations further, he would see that similar bodies are moving *within* cells a little more remote from the dividing line, and that a little further still, they are obviously but masses of endochrome in the act of subdivision.¹ Of the true Generative process in the *Ulvaceæ*, nothing whatever is known; and it is consequently altogether uncertain whether it takes place by simple conjugation, or according to that more truly sexual method which will be presently described. Here, again, therefore, is an unsolved problem of the greatest physiological interest, which probably requires nothing more for its solution, than patient and discriminating study. And the Author would point out, that it is by no means unlikely that the generative process may not be performed in the complete plant, but, as in the Ferns (§ 219), in the early product of the development of the zoospore. Although the typical *Ulvaceæ* are marine, yet there are several fresh-water species; and there are some which can even vegetate on damp surfaces, such as those of rocks or garden-walls kept moist by the percolation of water.

196. The *Oscillatoriaceæ* constitute another tribe of simple Plants, of great interest to the Microscopist, on account both of the extreme simplicity of their structure, and of the peculiar animal-like movements which they exhibit. They are continuous tubular filaments, formed by the elongation of their pri-

¹ Such an observation the Author had the good fortune to make in the year 1842, when the emission of zoospores from the *Ulvaceæ*, although it had been described by the Swedish algologist Agardh, had not been seen (he believes) by any British naturalist.

mordial cells, usually lying together in bundles or in strata, sometimes quite free, and sometimes invested by a gelatinous matrix. The endochrome which they contain usually exhibits some degree of transverse striation, as if breaking up into short segments by the division of the tube into cells; but this division is never perfected by the formation of complete partitions; and the filaments ultimately break up into distinct joints, the fragments of endochrome, which are to be regarded as *gonidia*, usually escaping from their sheaths, and giving origin to new filaments. These plants are commonly of some shade of green, often mingled, however, with blue; but not unfrequently they are of a purplish hue, and are sometimes so dark, as, when in mass, to seem nearly black. They occur not only in fresh, stagnant, brackish, and salt waters (certain species being peculiar to each), but also in mud, on wet stones, or on damp ground. Their very curious movements constitute the most remarkable feature in their history. These are described by Dr. Harvey¹ as of three kinds; first, a pendulum-like movement from side to side, performed by one end, whilst the other remains fixed so as to form a sort of pivot; second, a movement of flexure of the filament itself, the oscillating extremity bending over first to one side and then to the other, like the head of a worm or caterpillar seeking something on its line of march; and third, a simple onward movement of progression. "The whole phenomenon," continues Dr. H., "may perhaps be resolved into a spiral onward movement of the filament. If a piece of the stratum of an *Oscillatoria* be placed in a vessel of water, and allowed to remain there for some hours, its edge will first become fringed with filaments, radiating as from a central point, with their tips outwards. These filaments by their constant oscillatory movements, are continually loosened from their hold on the stratum, cast into the water, and at the same time propelled forward; and as the oscillation continues after the filament has left its nest, the little swimmer gradually moves along, till it not only reaches the edge of the vessel, but often, as if in the attempt to escape confinement—continues its voyage up the sides, till it is stopped by dryness. Thus in a very short time, a small piece of *Oscillatoria* will spread itself over a large vessel of water." This rhythmical movement, impelling the filaments in an undeviating onward course, is evidently of a nature altogether different from the truly spontaneous motions of Animals; and must be considered simply as the expression of certain vital changes taking place in the interior of the cells. It is greatly influenced by temperature and light, being much more active in warmth and sunshine than in cold and shade; and it is checked by any strong chemical agents. The true Generation of *Oscillatoriaceæ* is as yet completely unknown; and it does not seem at all unlikely that these plants may be the "motile" forms of some others, which, in their "still"

¹ "Manual of British Marine Algae," p. 220.

condition, present an aspect altogether different. Nearly allied to the preceding, is the little tribe of *Nostochaceæ*; which consists of distinctly beaded filaments, lying in firmly gelatinous fronds of definite outline. The filaments are usually simple, though sometimes branched; and are almost always curved or twisted, often taking a spiral direction. The masses of jelly in which they are imbedded are sometimes globular or nearly so, and sometimes extend in more or less regular branches; they frequently attain a very considerable size; and as they occasionally present themselves quite suddenly (especially in the latter part of the autumn, on damp garden walks), they have received the name of "fallen stars." They are not always so suddenly produced, however, as they appear to be; for they shrink up into mere films in dry weather, and expand again with the first shower. These plants multiply themselves, like the *Oscillatoria*, by the subdivision of their filaments, the portions of which escape from the gelatinous mass wherein they are imbedded, and move slowly through the water in the direction of their length; after a time they cease to move, and a new gelatinous envelope is formed around each piece, which then begins not only to increase in length by the transverse subdivision of its segments, but also to double itself by longitudinal fission, so that each filament splits lengthways (as it were) into two new ones. By the repetition of this process, a mass of new filaments is produced, the parts of which are at first confused, but afterwards become more distinctly separated by the interposition of the gelatinous substance developed between them. Besides the ordinary cells of the beaded filaments, two other kinds are occasionally observable; namely, "vesicular cells" of larger size than the rest (sometimes occurring at one end of the filaments, sometimes in their centre, and sometimes at intervals along their whole length), which are destitute of endochrome, and are sometimes furnished with cilia; and "sporangial cells," which seem like enlarged forms of the ordinary cells, and which are usually found in the neighborhood of the preceding. There is very strong evidence from analogy, that the "vesicular cells" are "antheridia" or sperm-cells, and that the "sporangial cells" contain germs, which, when fertilized by the antherozoids, and set free, become "resting-spores," as in certain members of the family to be next noticed.


197. Although many of the plants belonging to the family *Siphonaceæ* attain a considerable size, and resemble the higher Sea-weeds in their general mode of growth, yet they retain a simplicity of structure so extreme, that it apparently requires them to be ranked among the Protophytes. They are inhabitants both of fresh water and of the sea, and consist of very large tubular cells, which commonly extend themselves into branches, so as to form an arborescent frond. These branches, however, are seldom separated from the stem by any intervening partition; but the whole frond is composed of a simple continuous tube,

the entire contents of which may be readily pressed out through an orifice made by wounding any part of the wall. The *Vaucheria*, named after the Genevese botanist whose admirable researches on the fresh-water Confervæ have been already referred to (p. 38), may be selected as a particularly good illustration of this family; its history having been pretty completely made out. Most of its species are inhabitants of fresh water; but some are marine, and they commonly present themselves in the form of cushion-like masses, composed of irregularly branching filaments, which, although they remain distinct, are densely tufted together, and variously interwoven. The formation of moving gonidia or "zoospores" may be readily observed in these plants, the whole process usually occupying but a very short time. The extremity of one of the filaments usually swells up in the form of a club, and the endochrome accumulates in it, so as to give it a darker hue than the rest; a separation of this part from the remainder of the filament, by the interposition of a transparent space, is next seen; a new envelope is then formed around the mass thus cut off; and at last the membranous wall of the investing tube gives way, and the "zoospore" escapes, not, however, until it has undergone marked changes of form, and exhibited curious movements. Its motions continue for some time after its escape, and are then plainly seen to be due to the action of the cilia with which its whole surface is clothed. If it be placed in water in which some carmine or indigo has been rubbed, the colored granules are seen to be driven in such a manner as to show that a powerful current is produced by their propulsive action, and a long tract is left behind it. When it meets with an obstacle, the ciliary action not being arrested, the zoospore is flattened against the object; and it may thus be compressed, even to the extent of causing its endochrome to be discharged. The cilia are best seen, when their movements have been retarded or entirely arrested by means of opium, iodine, or other chemical reagents. The motion of the spore continues for about two hours; but after the lapse of that time it soon comes to an end, and the spore begins to develop itself into a new plant. It has been observed by Unger, that the escape of the zoospores generally takes place towards 8 A.M.; to watch this phenomenon, therefore, the plant should be gathered the day before, and its tufts examined early in the morning. It is stated by Dr. Hassall, that he has seen the same filament give off two or three zoospores successively. Their emission is obviously to be regarded as a method of increase by gemmation, rather than as a generative act; and recent discoveries have shown that there exists in this humble plant a true process of Sexual Generation, as was, indeed, long ago suspected by Vaucher, though upon no sufficient grounds. The branching filaments are often seen to bear at their sides peculiar globular or oval capsular protuberances, sometimes separated by the interposition of a stalk, which are filled with dark endochrome;

and these have been observed to give exit to large bodies covered with a firm envelope, from which, after a time, new plants arise. In the immediate neighborhood of these "capsules" are always found certain other projections, which, from being usually pointed and somewhat curved, have been named "horns;" and these were supposed by Vaucher to fulfil the function of the anthers of flowering-plants. The recent observations of Pringsheim have shown that such is really the case; for the "horns" are "antheridia," which, like those of the *Characeæ* (§ 202), produce antherozoids in their interior; whilst the capsules are "germ-cells," whose aggregate mass of endochrome is destined to become, when fertilized, the primordial cell of a new generation. The antherozoids, when set free from the antheridium, swarm over the exterior of the capsule, and have actually been seen to penetrate its cavity, through an aperture which opportunely forms in its wall, and to come into contact with the surface of its endochrome-mass, over which they diffuse themselves; there they seem to undergo dissolution, their contents mingling themselves with those of the germ-cell; and the endochrome mass, which had previously no proper investment of its own, soon begins to form an envelope, which increases in thickness and strength, until it has acquired such a density as enables it to afford a firm protection to its contents. This body, possessing no power of spontaneous movement, is known as a "resting-spore," in contradistinction to the "zoospores" already described; and it answers the purpose of a seed, in laying the foundation for a new generation, whilst the "zoospores" merely multiply the individual by a process analogous to budding. The Microscopist who wishes to study the development of zoospores, as well as several other phenomena of this low type of vegetation, may advantageously have recourse to the little plant termed *Achlya prolifera*, which grows parasitically upon the bodies of dead flies lying in the water, but also not unfrequently attaches itself to the gills of fish, and is occasionally found on the bodies of frogs. Its tufts are distinguishable by the naked eye, as clusters of minute colorless filaments; and these are found, when examined by the microscope, to be long tubes devoid of all partitions, extending themselves in various directions. The tubes contain a colorless slightly granular protoplasm, the particles of which are seen to move slowly in streams along the walls, as in *Chara*, the currents occasionally anastomosing with each other (Fig. 106, c). Within about thirty-six hours after the first appearance of the parasite on any body, the protoplasm begins to accumulate in the dilated ends of the filaments, each of which is cut off from the remainder by the formation of a partition; and within this dilated cell, the movement of the protoplasm continues for a time to be distinguishable. Very speedily, however, its endochrome shows the appearance of being broken up into a large number of distinct masses, which are at first in close contact with each other and

with the walls of the cell (Fig. 106, A), but which gradually become more isolated, each seeming to acquire a proper cell-wall; they then begin to move about within the parent cell; and, when quite mature, they are set free by the rupture of its wall (B), to go forth and form new attachments, and to develop themselves into tubiform cells resembling those from which they sprang. Each of these "motile gonidia" is possessed of only two cilia; their movements are not so powerful as those of the zoospores of *Vaucheria*; and they come to an end sooner. This plant forms "resting-spores" also, like those of *Vaucheria*; and there is every probability that they are generated by a like sexual process.

They may remain unchanged for a long time in water, when no suitable *nidus* exists for them; but will quickly germinate, if a dead insect or other suitable object be thrown in. One of the most curious forms of this group, is the *Hydrodictyon utriculatum*, which is found in fresh-water pools in the midland and southern counties of England. Its frond consists of a green open network of filaments, acquiring, when full grown, a length of from four to six inches, and composed of a vast number of cylindrical tubular cells, which attain the length of four lines or more, and adhere to each other by their rounded extremities, the points of junction corresponding to the knots or intersections of the network. Each of these cells may form within itself an enormous multitude (from 7000 to 20,000) of



Development of *Achlya prolifera*:—A, dilated extremity of a filament, b, separated from the rest by a partition, a, and containing young cells in progress of formation;—B, conceptacle discharging itself, and setting free young cells, a, b, c;—C, portion of filament, showing the course of the circulation of granular protoplasm.


"gonidia;" which, at a certain stage of their development, are observed in active motion in its interior; but of which groups are afterwards formed by their mutual adhesion, that are set free by the dissolution of their envelopes, each group giving origin to a new plant-net. Besides these bodies, however, certain cells produce from 30,000 to 100,000 more minute bodies of longer shape, each furnished with four long cilia and a red spot, which are termed by Braun "microgonidia:" these escape from the cell in a swarm, move freely in the water for some time, and

then come to rest and sink to the bottom, where they remain heaped in green masses. What is their future history, has not yet been ascertained; but there is evidence from analogy that they are "antheridial" cells, which have for their office to fecundate the true germ-cells by contained antherozoids. No development of "resting-spores," however, has yet been observed. The rapidity of the growth of this curious plant, is not one of the least remarkable parts of its history. The individual cells of which the net is composed, at the time of their emersion as gonidia, measure no more than 1-2500th of an inch in length; but in the course of a few weeks, they grow to a length of from 1-12th to 1-3d of an inch.

198. Almost every pond and ditch contains some members of the family *Confervaceae*; but they are especially abundant in moving water; and they constitute the greater part of those green threads, which are to be seen attached to stones, with their free ends floating in the direction of the current, in every running stream, and upon almost every part of the sea-shore, and which are commonly known under the name of "silk-weeds" or "crow-silk." Their form is usually very regular; for each thread is a long cylinder, made up by the union of a single file of short cylindrical cells united to each other by their flattened extremities; sometimes these threads give off lateral branches, which have the same structure. The endochrome, though usually green, is occasionally of a brown or purple hue; it is sometimes distributed uniformly throughout the cell (as in Fig. 107), whilst in other instances it is arranged in a pattern of some kind, as a network or a spiral; but this may be only a transitional stage in its development. The plants of this order are extremely favorable subjects for the study of the method of cell-multiplication by binary subdivision. This process usually takes place only in the *terminal* cell; and it may be almost always observed there, in some one of its stages. The first step is seen to be the subdivision of the endochrome, and the inflection of the primordial utricle around it (Fig. 107, A, a); and thus there is gradually formed a sort of hour-glass contraction across the cavity of the parent cell, by which it is divided into two equal halves (B). The two surfaces of the infolded utricle produce a double layer of cellulose membrane between them; this is not confined, however, to the contiguous surfaces of the young cell, but takes place over the whole exterior of the primordial utricle, so that the new septum is continuous with new layers that are formed throughout the interior of the cellulose wall of the original cell (C). Sometimes, however, as in *Conferva glomerata* (a common species), new cells may originate as branches from any part of the surface, by a process of budding; which, notwithstanding its difference of mode, agrees with that just described in its essential character, being the result of the subdivision of the original cell. A certain portion of the primordial utricle seems to undergo

increased nutrition, for it is seen to project, carrying the cellulose envelope before it, so as to form a little protuberance; and this sometimes attains a considerable length, before any separation of its cavity from that of the cell which gave origin to it, begins

Fig. 107.



Process of cell-multiplication in *Conferva glomerata*:—A, portion of filament with incomplete separation at *a*, and complete partition at *b*; B, the separation completed, and new cellulose partition being formed at *a*; C, formation of additional layers of cellulose wall, *c*, beneath the mucous investment, *d*, and around the primordial utricle, *e*, which encloses the endochrome, *b*.

to take place. This separation is gradually effected, however, by the infolding of the primordial utricle, just as in the preceding case; and thus the endochrome of the branch cell is completely severed from that of the stock. The branch then begins to elongate itself by the subdivision of its first-formed cell; and this process may be repeated for a time in all the cells of the filament, though it usually comes to be restricted at last to the terminal cell. The *Confervaceæ* multiply themselves by “zoospores,” which are produced within their cells, and are then set free, just as in the *Ulvaceæ* (§ 195); in most of the genera, the endochrome of each cell divides into numerous zoospores, which are of course very minute; but in *Edogonium*,—a fresh-water genus distinguished by the circular markings which form rings round the extremities of many of the cells, and by many interesting peculiarities of growth and reproduction,¹—only a single large zoospore is set free from each cell; and its liberation is accomplished by the almost complete fission of the wall of the cell through

one of these rings, a small part only remaining unleft, which serves as a kind of hinge, whereby the two parts of the filament are prevented from being altogether separated. Sometimes the zoospore does not completely extricate itself from the parent cell; and it may begin to grow in this situation, the root-like processes which it puts forth being extended into the cavity. A true sexual generation has been observed in several *Confervaceæ*; and is probably universal throughout the group. Thus in *Sphæroplea annulina*, according to the recent observations of Dr. Cohn,² the ring-like masses of endochrome, of which several are found in each cell, resolve themselves in certain of the cells into minute bodies resembling the antherozoids of *Chara* (Fig. 112, u), which, after moving freely within these cells, escape through apertures in their walls, and then penetrate

¹ See the account of these processes in the “Micrographic Dictionary,” p. 468.


² “Monatsberichte der Königl. Akad. der Wissenschaften,” Mai, 1855.

apertures in the walls of certain other cells, within each of which the endochrome has coalesced into a globular mass; over this mass the antherozoids spread themselves, and seem to dissolve away upon its surface; and after this process has taken place, the mass of endochrome acquires a firm envelope, and becomes a "resting-spore," which, when set free by the rupture of the parent cell-wall, germinates into a new plant. A curious variation of this process is seen in *Edogonium*; for instead of the antherozoids escaping freely from the "sperm-cells" which produced them, they are discharged *en masse*, included within a capsule which is furnished with cilia, and which so resembles a "zoospore" as to be easily mistaken for it; and it is only when this has attached itself, and has set free its contents by the falling off of a sort of lid, that the antherozoids are enabled to perform their fertilizing office. The same thing happens in some other Confervaceæ; in which, however, the antheridial capsules, being smaller than the zoospores, are distinguished as *microgonidia*, whilst the latter are known as *macrogonidia*. The offices of these different classes of reproductive bodies are only now beginning to be understood; and the inquiry is one so fraught with Physiological interest, and, from the facility of growing these plants in artificial Aquaria, may be so easily pursued, that it may be hoped that Microscopists will apply themselves to it so zealously, as not long to leave any part of it in obscurity.

199. The family *Conjugateæ* agrees with that of the *Confervaceæ* in its mode of growth, but differs from it in the plan in which its generative process is performed; this being accomplished by an act of "conjugation," resembling that which has been described in the simplest Protophytes. These plants are not found so much in running streams, as in waters that are perfectly still, such as those in ponds, reservoirs, ditches, or marshy grounds; and they are for the most part unattached, floating freely at or near the surface, especially when buoyed up by the bubbles of gas which are liberated from the midst of them under the influence of solar light and heat. In an early stage of their growth, whilst as yet the cells are undergoing multiplication by subdivision, the endochrome is commonly diffused pretty uniformly through their cavities (Fig. 109, A); but as they advance towards the stage of conjugation, the endochrome ordinarily arranges itself in regular spirals (B), but occasionally in some other forms. The act of "conjugation" usually occurs between the cells of two distinct filaments, that happen to lie in proximity to each other; and all the cells of each filament generally take part in it at once. The adjacent cells put forth little protuberances, which come into contact with each other, and then coalesce by the breaking down of the intervening partitions, so as to establish a free passage between the cavities of the conjugating cells. In some genera of this family (such as *Mesocarpus*), the conjugating

cells pour their endochromes into a dilatation of the passage that has been established between them; and it is there that they commingle, so as to form the spore or the embryo-cell. But in the *Zygnema* (Fig. 108), which is among the commonest and best known forms of Conjugatæ, the endochrome of one cell passes over entirely into the cavity of the other; and it is within the latter that the spore is formed (c), the two endochromes co-

FIG. 108.




Various stages of the history of *Zygnema quininum*.—A, three cells, a, b, c, of a young filament, of which b is undergoing subdivision; B, two filaments in the first stage of conjugation, showing the spiral disposition of their endochromes, and the protuberances from the conjugating cells; C, completion of the act of conjugation, the endochromes of the cells of the filament a having entirely passed over to those of filament b, in which the sporangia are formed.

lescening into a single mass, around which a firm envelope gradually makes its appearance. Further, it may be generally observed, that all the cells of one filament thus empty themselves, whilst all the cells of the other filaments become the recipients; here, therefore, we seem to have a foreshadowing of the sexual distinction of the generative cells into "sperm-cells" and "germ-cells," which we have just seen to exist in the Confervaceæ (§ 198). And this transition will be still more complete, if (as Itzigsohn has affirmed) the endochrome of certain filaments of *Spirogyra* breaks up before conjugation into little spherical aggregations, which are gradually converted into nearly colorless spiral filaments, having an active spontaneous motion, and therefore corresponding precisely to the antherozoids of the truly sexual Protophytes.¹

¹ This group of plants seems to serve as the connecting link between those simple Protophytes in which the sexes are not yet differentiated, and those higher forms in which the distinction between the "sperm-cells" and "germ-cells" is very apparent. For let it be supposed that in *Sphaeroplea* (§ 198) a conjugation of two adjacent cells were to take place, at that stage in their development in which the endochrome is uniformly arranged in rings, no differentiation of sexes yet showing itself,—the process would in all respects correspond with that of the ordinary Conjugatæ. Again, whilst in *Mesocarpus*, the two conjugating cells appear to take (as in the *Desmidiæ*, § 169) a precisely similar share in the formation of their product, the first stage of differentiation

200. The *Chætophoraceæ* constitute another beautiful and interesting little group of Confervoid plants, of which some species inhabit the sea, whilst others are found in fresh and pure water, rather in that of gently moving streams, however, than in strongly flowing currents. Generally speaking, their filaments put forth lateral branches, and extend themselves into arborescent fronds; and one of the distinctive characters of the group is afforded by the fact, that the extremities of these branches are usually prolonged into bristle-shaped processes (Fig. 109). As in many preceding cases, these plants multiply themselves by the conversion of the endochrome of certain of their cells into "zoospores;" and these, when set free, are seen to be furnished with four large cilia. "Resting-spores" have also been seen in many species; and it is probable that these, as in Confervaceæ, are true generative products of the fertilization of the contents of "germ-cells" by "antherozoids" developed within "sperm-cells" (§ 198). Nearly allied to the preceding are the *Batrachospermææ*, whose name is indicative of the strong resemblance which their beaded filaments bear to frog-spawn; these exhibit a somewhat greater complexity of structure, and afford objects of extreme beauty to the Microscopist. The plants of this family are all inhabitants of fresh water, and they are chiefly found in that which is pure and gently flowing. "They are so extremely flexible," says Dr. Hassall, "that they obey the slightest motion of the fluid which surrounds them; and nothing can surpass the ease and grace of their movements.



Branches of *Chætophora elegans*, in the act of discharging ciliated zoospores, which are seen, as in motion, on the right.

into sperm-cells and germ-cells is manifested in *Zygnema*, by the passage of the whole endochrome of those of one filament into the cavities of the other, and by the formation of the spores within the latter. In *Spirogyra*, moreover, the endochrome of one set of cells becomes converted into antherozoids before conjugation, whilst that of the other aggregates into a sporangial mass; thus exhibiting the second stage of differentiation. Further, there are certain species which agree with the ordinary Conjugatæ in their general habit, and which form "resting-spores" like theirs, but in which no conjugation has been observed; and it seems not improbable that in these, as in Sphæroplea, the antherozoids make their way out of the sperm-cell by minute apertures in its wall, and swim freely about before finding their way into the germ-cell through the apertures in its wall;—still, however, performing by this means the very same act, as that which is accomplished by the more direct process of conjugation,—viz., the introduction of the contents of the sperm cell into the interior of the germ-cell.

When removed from the water, they lose all form, and appear like pieces of jelly, without trace of organization; on immersion, however, the branches quickly resume their former disposition." Their color is for the most part of a brownish-green; but sometimes they are of a reddish or bluish purple. The central axis of each plant is originally composed of a single file of large cylindrical cells laid end to end; but this is subsequently invested by other cells, in the manner to be presently described. It bears, at pretty regular intervals, whorls of short radiating branches, each of them composed of rounded cells arranged in a bead-like row (Fig. 110), and sometimes subdividing again into two, or themselves giving off lateral branches. Each of the primary branches originates in a little protuberance from the primitive cell of the central axis, precisely after the manner of the lateral cells of *Conferva glomerata* (§ 198); as this protuberance

FIG. 110.


*Batrachospermum moniliforme.*

increases in size, its cavity is cut off by a septum, so as to render it an independent cell; and by the continual repetition of the process of duplicative subdivision, this single cell becomes converted into a beaded filament. Certain of these branches, however, instead of radiating from the main axis, grow downwards upon it, so as to form a closely fitting investment, that seems properly to belong to it. Some of the radiating branches grow out into long transparent points, like those of *Chætophoraceæ*; and it does not seem by any means improbable, that these, like the "horns" of *Vaucheria* (§ 197), are really antheridia. For within certain cells of other branches, "resting-spores" are formed; by the agglomeration of which are produced the large dark bodies, that are seen in the midst of the whorls of branches (Fig. 110).

201. This seems the most appropriate place to consider a group of humble plants, having a peculiar interest for Microscopists,—that, namely, of *Characeæ*; in which we have a vegetative apparatus as simple as that of the *Protophytes* already described, whilst their reproductive apparatus is even more highly developed than that of the proper *Algæ*. They are for the most part inhabitants of fresh waters, and are found rather in such as are still, than in those which are in motion; one species, however, may be met with in ditches whose waters are rendered salt by communication with the sea. They may be easily grown for

the purposes of observation, in large glass jars exposed to the light; all that is necessary being to pour off the water occasionally from the upper part of the vessel (thus carrying away a film that is apt to form on its surface), and to replace this by fresh water. Each plant is composed of an assemblage of long tubiform cells, placed end to end; with a distinct central axis, around which the branches are disposed at intervals with great regularity (Fig. 111, A). In one of the genera, *Nitella*, the stem and

FIG. 111.



Nitella flexilis:—A, stem and branches of the natural size; a, b, c, d, four verticils of branches issuing from the stem; e, f, subdivision of the branches;—B, portion of the stem and branches enlarged; a, b, joints of stem; c, d, verticils; e, f, new cells sprouting from the sides of the branches; g, h, new cells sprouting at the extremities of the branches.

branches are simple cells, which sometimes attain the length of several inches; whilst in the true *Chara*, each central tube is surrounded by an envelope of smaller ones, which is formed as in *Batrachospermæ*, save that the investing cells grow upwards as well as downwards from each joint, and meet each other on the stem half-way between the joints. Some species have the power of secreting carbonate of lime from the water in which they grow, if this be at all impregnated with calcareous matter; and by the deposition of it beneath their teguments, they have gained their popular name of “stone-worts.” These humble plants have attracted much attention, in consequence of the facility with which the “rotation,” or movement of fluid in the interior of the individual cells, may be seen in them. Each cell,

in the healthy state, is lined by a layer of green oval granules, which cover every part, except two longitudinal lines that remain nearly colorless (Fig. 111, B); and a constant stream of semi-fluid matter, containing numerous jelly-like globules, is seen to flow over this green layer, the current passing up one side, changing its direction at the extremity, and flowing down the other side, the ascending and descending spaces being bounded by the transparent lines just mentioned. That the currents are in some way directed by the layer of granules, appears from the fact noticed by Mr. Varley,¹ that if accident damages or removes them near the boundary between the ascending and descending currents, a portion of the fluid of the two currents will intermingle by passing the boundary; whilst, if the injury be repaired by the development of new granules on the part from which they had been detached, the circulation resumes its regularity, no part of either current passing the boundary. In the young cells, however, the rotation may be seen, before their granular lining is formed. The rate of the movement is affected by anything which influences the vital activity of the plant; thus, it is accelerated by moderate warmth, whilst it is retarded by cold; and it may be at once checked by a slight electric discharge through the plant. The moving globules, which consist of starchy matter, are of various sizes; being sometimes very small, and of definite figure, whilst in other instances they are seen as large irregular masses, which appear to be formed by the aggregation of the smaller particles.² The production of new cells, for the extension of the stem or branches, or the origination of new whorls, is not here accomplished by the subdivision of the parent cell, but takes place by the method of out-growth (Fig. 111, B, e, f, g, h), which, as already shown (§ 198), is nothing but a modification of the usual process of cell-multiplication; in this manner, the extension of the individual plant is effected with considerable rapidity. When these plants are well supplied with nutriment, and are actively vegetating under the influence of light, warmth, &c., they not unfrequently develop "bulbels," or gonidia of a peculiar kind, which serve the same purpose in multiplying the individual, as is answered by the zoospores of


¹ "Transactions of the Microscopical Society" (First Series), vol. ii, p. 99.

² This interesting phenomenon may be readily observed, by taking a small portion of the plant out of the water in which it is growing, and either placing it in a large aquatic box (§ 68) or in the zoophyte-trough (§ 69), or laying it on the glass stage-plate (§ 67) and covering it with thin glass. The modification of the stage-plate which is termed the "growing-slide," will enable the Microscopist to keep a portion of *Chara* under observation for many days together; and this is a much simpler and more convenient arrangement than the method devised by Mr. Varley for growing *Chara* in bottles; since the bottle requires a special "phial-holder" for fixing it on the Microscope stage, and the convexity of its surface produces some distortion of the image. The latter method, however, has its advantages for those who wish to make a special study or a frequent exhibition of the phenomenon in question; and such should consult Mr. Varley's memoir in the "Transactions of the Society of Arts," vols. xlviii, xlix, 1; some parts of which are cited by Mr. Quekett in his "Practical Treatise on the Microscope," Third Ed. pp. 166, 397, *et seq.*

the simpler Protophytes; these are little clusters of cells, filled with starch, which sprout from the sides of the central axis, and then, falling off, evolve the long tubiform cells characteristic of the plant from which they were produced.¹ The *Characeæ* may also be multiplied by artificial subdivision; the separated parts continuing to grow, under favorable circumstances, and developing themselves into the typical form.

202. The generative apparatus of *Characeæ* consists of two sets of bodies, both of which grow at the bases of the branches (Fig. 112, A, B); one set is known by the designation of "globules,"

FIG. 112.



Antheridia of *Chara fragilis*:—A, antheridium or "globule" developed at the base of pistillidium or "nucule";—B, nucule enlarged, globule laid open by the separation of its valves; C, one of the valves, with its group of antheridial filaments, each composed of a linear series of cells, within every one of which an antherozoid is formed;—in D, E, and F, the successive stages of this formation are seen;—and at G is shown the escape of the mature antherozoids, H.

the other by that of "nucules." The globules are really antheridia; whilst the nucules contain the germ-cells. The "globules," which are nearly spherical, have an envelope made up of eight

¹ This multiplication by bulbels was described by Amici in 1827; but his observations seem to have been forgotten by Botanists, until the rediscovery of the fact by M. Montague.


triangular valves (B, C), often curiously marked, which enclose a nucleus of a light reddish color; this nucleus is principally composed of a mass of filaments rolled up compactly together; and each of these filaments (C) consists, like a *Conferva*, of a linear succession of cells. In every one of these cells there is formed, by a gradual change in its contents (the successive stages of which are seen at D, E, F), a spiral thread of two or three coils, which, at first motionless, after a time begins to move and revolve within the cell; and at last the cell-wall gives way, and the spiral thread makes its way out (G), partially straightens itself, and moves actively through the water for some time (H), in a tolerably determinate direction, by the lashing action of two long and very delicate filaments with which they are furnished. The exterior of the "nucule" (A, B) is formed by five spirally twisted tubes, that give it a very peculiar aspect; and these enclose a central sac containing protoplasm, oil, and starch-globules. At a certain period, the spirally twisted tubes, which form a kind of crown around the summit, separate from each other, leaving a canal that leads down to the central cell; and it is probable that through this canal the antherozoids make their way down, to perform the act of fertilization. Ultimately the nucule falls off like a seed, and gives origin to a single new plant by a kind of germination. The complete *specialization* of the Generative apparatus which we here observe (the organs of which it is composed being distinctly separated from the ordinary vegetating structure of the plant), as well as the complex structure of the organs themselves, mark out this group, in spite of the simplicity of the rest of its structure, as belonging to a grade very much above that of the other families that have been treated of in this chapter; but as scarcely any two Botanists agree upon the exact place which ought to be assigned to it, the convenience of associating it with other forms of vegetation of which the Microscopist especially takes cognizance, is a sufficient reason for so arranging it in a work like the present.

CHAPTER VII.

MICROSCOPIC STRUCTURE OF HIGHER CRYPTOGAMIA.

203. FROM those simple Protophytes, whose minuteness causes their entire fabrics to be fitting objects for Microscopic examination, we pass to those higher forms of Vegetable life, whose larger dimensions require that they should be analyzed (so to speak) by the examination of their separate parts. And in the present chapter, we shall bring under notice some of the principal points of interest to the Microscopist, which are presented by the *Cryptogamic* series; commencing with those simpler Algæ, which scarcely rank higher than some of the Protophytes already described; and ending with the Ferns and their allies, which closely abut upon the *Phanerogamia* or Flowering Plants. In ascending this series, we shall have to notice a *gradual differentiation* of organs; those set apart for Reproduction being in the first place separated from those appropriated to Nutrition (as we have already seen them to be in the Characæ); and the principal parts of the Nutritive apparatus, which are at first so blended together that no real distinction exists between root, stem, and leaf, being progressively evolved on types more and more peculiar to each respectively, and having their functions more and more limited to themselves alone. Hence we find a differentiation, not merely in the external form, but also in the intimate structure of organs; its degree bearing a close correspondence to the degree in which their functions are respectively *specialized* or limited to particular actions. Thus in the simple *Ulva* (Fig. 104), whatever may be the extent of the frond, every part has exactly the same structure, and performs the same actions, as every other part; living *for* and *by* itself alone. In *Batrachospermum* (Fig. 110), we have seen a definite arrangement of branches upon an axis of growth; and while the branches are formed of simple necklace-like rows of rounded cells, the cells of the stem are elongated and adhere to one another by flattened ends. This kind of dif-

FIG. 113.



Mesogloia vermicularis.

ferentiation is seen to be carried to a still greater extent in *Mesogloia* (Fig. 113); a plant which may be considered as one of the connecting links between such Protophytes as *Batrachospermæ*—which it resembles in general plan of structure,—and the Fucoid Algæ, which it resembles in fructification.

204. When we pass to the higher Sea-weeds, such as the common *Fucus* and *Laminaria*, we observe a certain foreshadowing of the distinction between root, stem, and leaf; but this distinction is but very imperfectly carried out, the root-like and stem-like portions serving for little else than the mechanical attachment of the leaf-like part of the plant, and each still absorbing and assimilating its own nutriment, so that no transmission of fluid takes place from one portion of the fabric to another. Hence we find that there is not yet any departure from the simple *cellular* type of structure; the only modification being, that the several layers of cells, where many exist, are of different sizes and shapes, the texture being usually closer on the exterior and looser within; and that the texture of the stem and roots is denser than that of the expanded fronds. This simple cellular type of structure is maintained through all but the highest Cryptogamia; for it is not until we come to the Mosses, that the differentiation of stem, root, and leaf is established; and even in these it is not so fully carried out, as to require a provision for the free transmission of fluid from one part to another; whilst the scale of their fabrics is not such as to render it necessary that their softer parts should be supported by a tissue of peculiar density. But in the group of Ferns, which, notwithstanding their complete adhesion to the Cryptogamic type of Reproduction, have the general form of the higher plants, and even attain the size and bearing of trees, we find the leaves separated from the roots by the intervention of a stem; and in this stem, as also in the leaf-stalks prolonged from it, we find, interposed in the midst of the cellular tissue which forms their principal substance, two new forms of structure,—namely, *woody fibre* which serves to give strength and support to the stem and to the organs it bears, and *ducts* through which the liquid absorbed by the roots may be readily conveyed to the leaves.

205. The group of *Melanospermous* or olive-green Sea-weeds, which, in the family *Fucaceæ*, exhibits the highest type of Algal structure, presents us with the lowest in the family *Ectocarpaceæ*; which, notwithstanding, contains some of the most elegant and delicate structures that are anywhere to be found in the group, the full beauty of which can only be discerned by the microscope. Such is the case, for example, with the *Sphacelaria*, a small and delicate sea-weed, which is very commonly found parasitic upon larger Algæ, either near low-water mark, or altogether submerged; its general form being remarkably characterized by a symmetry that extends also to the individual branches (Fig. 114), the ends of which, however, have a decayed

look, that seems to have suggested the name of the genus (from the Greek *σφακελος*, gangrene). The study of the higher and larger members of this group, has recently come to present a new and very attractive source of interest to the Microscopist, in consequence of the discovery of the truly sexual nature of their fructification; and we shall take that of a common species of *Fucus* as the type of that of the order generally. The "receptacles" which are borne at the extremities of the fronds, here contain both "sperm-cells" and "germ-cells;" in some other species, however, they are disposed in different receptacles on the same plant; whilst in the commonest of all, *F. vesiculosus* (bladder-wrack), they are limited to different individuals.¹ When a section is made through one of the flattened receptacles of *F. platycarpus*, its interior is seen to be a nearly globular cavity (Fig. 115), lined with filament-

FIG. 115.




FIG. 114.



Fig. 114. Terminal portion of branch of *Sphaecelaria cirrhosa*.

Fig. 115. Vertical Section of receptacle of *Fucus platycarpus*, lined with filaments, among which lie the antheridial cells, and the sporangia containing octospores.

ous cells, some of which are greatly elongated, so as to project through the pore by which the cavity opens on the surface. Among these are to be distinguished, towards the period of their

¹ It was at first stated by MM. Thuret and Decaisne, that this species was sometimes dioecious, sometimes hermaphrodite; but they now consider the hermaphrodite form to be a distinct species, the *F. platycarpus* described above.

maturity, certain filaments (Fig. 116, A) whose granular contents acquire an orange hue, and gradually shape themselves into oval bodies (B), each with an orange-colored spot, and two long

FIG. 116.



Antheridia and antherozoids of *Fucus platycarpus*:—A, branching articulated hairs, detached from the walls of the receptacle, bearing antheridia in different stages of development; B, antherozoids, some of them free, others still included in their antheridial cells.

thread-like appendages, which, when discharged by the rupture of the containing cell, have for a time a rapid undulatory motion, whereby these antherozoids are diffused through the surrounding liquid. Lying amidst the filamentous mass, near the walls of the cavity, are seen (Fig. 115) numerous dark pear-shaped bodies, which are the *sporangia*, or parent cells of the "germ-cells." Each of these sporangia gives origin, by duplicative subdivision, to a cluster of eight cells, which is thence known as an "octospore;" and these are liberated from their envelopes, before the act of fertilization takes place. This act consists in the swarming of the antherozoids over the surface of the germ-cells, to which they communicate a rotatory motion by the vibration of their own filaments; it takes place within the receptacles in the hermaphrodite *Fuci*, so that the spores do not make their exit from the cavity until after they have been fecundated; but in the monœcious and dicœcious species, each kind of receptacle separately discharges its contents, which come into mutual contact on their exterior. The antheridial cells are usually ejected entire, but soon rupture, so as to give exit to their filaments; the sporangia of the female receptacles discharge their globular octospores within the receptacle; and these, soon after passing forth, liberate their separate spores, which speedily meet with antherozoids and are fecundated by them. The spores, when fertilized, soon acquire a new and firmer envelope; and under favorable circumstances they speedily begin to develop themselves into new plants. The first change that is seen in them, is


the projection and narrowing of one end into a kind of footstalk, by which the spore attaches itself, its form passing from the globular to the pear-shaped; a partition is speedily observable in its interior, its single cell being subdivided into two; and by a continuation of a like process of duplication, first a filament, and then a frondose expansion, is produced, which gradually evolves itself into the likeness of the parent plant. The whole of this process may be watched without difficulty, by obtaining specimens of *F. vesiculosus* at the period at which the fructification is shown to be mature, by the recent discharge of the contents of the conceptacles in little gelatinous masses on their orifices; for if some of the spores, which have been set free from the olive-green (female) receptacles, be placed in a drop of sea-water in a very shallow cell, and a small quantity of the mass of antherozoids set free from the orange-yellow (male) receptacles, be mingled with the fluid, they will speedily be observed, with the aid of a magnifying power of 200 or 250 diameters, to go through the actions just described; and the subsequent processes of germination may be watched by means of the "growing-slide."¹ The winter months, from December to March, are the most favorable for the observation of these phenomena; but where the Fuci abound, some individuals will usually be found in fructification at almost any period of the year. Even in the *Fucaceæ*, according to recent observations, a multiplication by zoospores, like that of the *Ulvaceæ* (§ 195), still takes place; these bodies being produced within certain of the cells that form the superficial layer of the frond, and swimming about freely for a time after their emission, until they fix themselves and begin to grow. That they are to be considered *gemmæ*, and not as generative products, appears certain from the fact that they will vegetate without the assistance of any other bodies; whereas the antherozoids of themselves never come to anything, and the octospores undergo no further changes, but decay away (as M. Thuret has experimentally ascertained), if not fecundated by the antherozoids.

206. Among the *Rhodospirææ*, or red Sea-weeds, also, we find various simple but most beautiful forms, which connect this group with the more elevated Protophytes, especially with the family *Chaetophoraceæ*; such delicate feathery or leaf-like fronds belong, for the most part to the family *Ceramiceæ*, some members of which are found upon every part of our coasts, attached either to rocks or stones, or to larger Algæ, and often themselves affording an attachment to Zoophytes and Bryozoa. They chiefly live in deeper water than the other sea-weeds; and their richest tints are only exhibited, when they grow under the shade of projecting rocks or of larger dark-colored Algæ. Hence in growing

¹ If a cell be not employed, the drop should not be covered, unless some precaution be taken to keep the pressure of the thin glass from the minute bodies beneath, whose movements it will otherwise impede.

them artificially in *Aquaria*, it is requisite to protect them from an excess of light; since otherwise they become unhealthy. The nature of the fructification of the *Rhodospirææ* (or *Floridææ*) is less perfectly understood than that of the *Fucoid Algæ*. It is certain, however, that antheridia exist among them; these being developed in individuals that do not produce spores, and in pretty much the same situations. The products of these antheridia, however, do not exhibit the spontaneous motion of ordinary antherozoids. Of the spores there are two kinds, of which one set are probably "gemmæ," whilst the other are "germ-cells;" but it is not yet determined to which of the two these characters respectively belong. The "tetraspores,"—which are peculiarly characteristic of the group, being found in every one of its subdivisions,—are usually imbedded in the general substance of the frond, though they sometimes congregate in particular parts, or are restricted to a special branch. Each group (Fig. 117, B) seems

FIG. 117.



Arrangement of tetraspores, in *Carpocaulon mediterraneum*:—A, entire plant; B, longitudinal section of branch. (N. B. Where only three tetraspores are seen, it is merely because the fourth did not happen to be so placed as to be seen at the same view.)

to be evolved within one of the ordinary cells of the frond, which undergoes a duplicative subdivision; the four secondary cells, however, remain enclosed within their primary cell until the period of maturity, a new envelope, the "perispore," being formed around them. In the *Corallines*, which are sea-weeds whose tissue is consolidated by calcareous deposit, the tetraspores are included within hollow conceptacles; but generally speaking, it is the simple spores only which are thus specially

protected. These are never scattered through the frond, like the tetraspores; and are commonly developed within a *ceramidium*, which is an urn-shaped case, furnished with a pore at its summit, and containing a tuft of pear-shaped spores arising from the base of its cavity. The resemblance of these bodies in position to the "octospores" of Fuci, would seem to justify the conclusion that *they* are the true generative spores, whilst the tetraspores are *gemmae*, as Harvey and Thwaites consider them; but a different view is taken by Decaisne, Agardh, and other eminent Algologists, who regard the tetraspores as the true generative spores, and consider the simple spores to be *gemmae*. It is, therefore, a point of much interest to determine by careful observation and experiment which is the right view; and Microscopists who have the opportunity of studying these plants, either in their native haunts, or in artificial Aquaria, can scarcely apply themselves to a better subject of investigation.

207. The class of *Lichens*, which consists of plants that closely correspond with *Algæ* in simplicity of organization, but differ from them widely in habit, does not present so many objects of attractive interest to the Microscopist; and the peculiar density which usually characterizes their structure, renders a minute examination of it more than ordinarily difficult. Lichens are commonly found growing upon the trunks or branches of trees, upon rocks or stones, upon hard earth, or in other situations in which they are sparingly supplied with moisture, but are freely exposed to light and air. In the simpler forms of this group, the primordial cell gives origin, by the ordinary process of subdivision, to a single layer of cells, which may spread itself over the surface to which it is attached, in a more or less circular form; and one or more additional layers being afterwards developed upon its free surface, a *thallus* is formed, which has no very defined limit, and which, in consequence of the very slight adhesion of its component cells, is said to be "pulverulent." Sometimes, however, the cells of the thallus are rather arranged in the form of filaments, which penetrate the superficial layers of the bark whereon such Lichens grow, and which are sometimes also so interwoven at the outer surface, as to form a sort of cuticle. Interposed among the ordinary cells of the thallus, we very commonly find certain green globular cells, arranged in single bead-like filaments; these, which are termed *gonidia*, being found to be capable of reproducing the plant when detached, must be considered as *gemmae*. In the higher tribes of Lichens, we find the interlacing filaments forming a tough cortical envelope to both surfaces; whilst in the interior of the firm "crustaceous" thallus, the gonidial cells are found in regular layers. Sometimes these increase in particular spots, and make their way through the upper cortical layer, so as to appear on the surface as little masses of dust, which are called *soredia*. Besides these, Lichens contain proper Generative organs, by which a true sexual

reproduction seems to be effected. In addition to the "fructification" which is commonly recognized by its projection from the surface of the thallus, the researches of M. Tulasne have detected a set of peculiar organs of much smaller size, not unlike the male receptacles of *Fuci* (§ 205), to which he has given the appellation of *spermogonia*. From the exterior of the cellular filaments which line these cavities, a vast number of minute oval bodies termed *spermatia* are budded off, which, when mature, escape in great numbers from the orifices of the spermogonia. They differ from ordinary antherozoids in being destitute of any power of spontaneous movement; but in this respect they are paralleled by the spermatoid bodies of the *Florideæ* (§ 206). As their participation in the production of fertile spores has not yet been demonstrated, we cannot yet indubitably assign to them the character of "sperm-cells;" although various considerations concur to render their possession of this attribute highly probable. The female portion of the generative apparatus, though sometimes dispersed through the thallus, is usually collected into special aggregations, which form projections of various shapes; these, although they have received a variety of designations according to their particular conformation, may all be included under the general term *apothecia*. When divided by a vertical section, these bodies at their maturity are found to contain a number of *asci* or spore-cases, arranged vertically in the midst of straight elongated cells or filaments, which are termed *paraphyses*. Each of the *asci* contains a definite number of spores (usually eight, but always a multiple of two), which are projected from the *apothecia* with some force, the emission being kept up continuously for some time; this discharge seems to be due to the different effect of moisture upon the different layers of the apothecium. When and how the act of fecundation is accomplished, is a matter still hidden in obscurity; and the problem is one which, owing to the difficulties arising out of the dense structure of the organs, will only be resolved by a combination of sagacity, manipulative skill, and perseverance, on the part of Microscopic observers who may devote themselves to the study.

208. In the simplest forms of *Fungi*, we again return to the lowest type of Vegetable existence, namely, the single cell; and such, if perfect plants, would properly take rank among the lowest Protophytes. But there is good reason for regarding many—perhaps all—of those which *seem* most simple, as the imperfectly developed states of other Plants, which, if they attained their full evolution, would present a much more complex structure. This is the case, for example, with the *Torula cerevisiæ* or Yeast-plant, which so abounds in Yeast, that this substance may be said to be almost entirely made up of it. When a small quantity of yeast is placed under the Microscope, and is magnified 300 or 400 diameters, it is found to be full of globules, which are clearly cells; and these cells vegetate, when placed in a fer-

mentable fluid containing some form of albuminous matter in addition to sugar, in the manner represented in Fig. 118. Each

FIG. 118.




Torula Cerevisia, or Yeast-Plant, as developed during the process of fermentation:—a, b, c, d, successive stages of Cell-multiplication.

cell puts forth one or two projections, which seem to be young cells developed as buds or offsets from their predecessors; these, in the course of a short time, become complete cells, and again perform the same process; and in this manner the single cells of yeast develop themselves, in the course of a few hours, into rows of four, five, or six, which remain in continuity with each other whilst the plant is still growing, but which separate if the fermenting process be checked, and return to the isolated condition of those which originally constituted the yeast. Thus it is, that the quantity of yeast first introduced into the fermentable fluid, is multiplied six times or more, during the changes in which it takes part. The full development of the Plant, however, and the evolution of its apparatus of fructification, only occur, when the fermenting process is allowed to go on without check; and it seems capable of producing a considerable variety of forms, whose precise relationship to each other has not yet been made clear. In fact, with regard to the Fungi generally, it has been made apparent by recent observations, that different individuals of the very same species may not only develop themselves according to a great number of very dissimilar modes of growth, but that they may even bear very dissimilar types of fructification; and further, that even the same individual may put forth, at different periods of its life, those two kinds of fructification,—the *basidio-sporous*, in which the spores are developed by out-growth from free points (*basidia*), and the *theca-sporous*, in which they are developed in the interior of cases (*thecæ* or *asci*, Fig. 125),—which had been previously considered as separately characterizing the two principal groups, into which the class is primarily divided.

209. Many of the simpler forms of Fungi are inhabitants of the interior of the bodies of other animals, and are only known as living in these situations. Among these may first be mentioned the *Sarcina ventriculi* (Fig. 119), which is most frequently found in the matters vomited by persons suffering under disorder of the stomach, but has also been met with in other diseased parts of the body. The plant has been detected in the contents of the stomach, however, under circumstances which seem to indicate that it is not an uncommon tenant of that organ even in health, and that it may accumulate there to a

considerable amount without producing any inconvenience; it seems probable, therefore, that its presence in disease is




Sarcina ventriculi.

rather to be considered as favored by the changed state of the fluids which the disease induces (either an acid or a fermentable state of the contents of the stomach having been generally found to exist in the cases in which the plant has been most abundant), than to be itself the occasion of the disease, as some have supposed. The *Sarcina* presents itself in the form of clusters of adherent cells arranged in squares, each square containing from

4 to 64, and the number of cells being obviously multiplied by duplicative subdivision in directions transverse to each other. In fact, its general mode of growth would indicate a near relationship to *Gonium*, one of the Volvocinæ, which presents itself in similar quadripartite aggregations; and many Botanists, looking to this circumstance, and to the residence of the plant in liquid, regard it as belonging to the group of Algæ. It agrees with the Fungi, however, in not living elsewhere than in liquids containing organic matter; and there can be little doubt that, as no fructification has yet been seen in it, only its earlier and simpler condition is yet known to us. Its true place cannot be determined, until its whole life-history shall have been followed out. There is a form of Fungous vegetation that is prone to develope itself within the living body, which is of great economic importance, as well as of scientific interest; this is the *Botrytis bassiana* (Fig. 120), a kind of "mould," the growth of which is the real source of a disease termed *Muscardine*, that sometimes carries off Silk-worms in large numbers, just when they are about to enter the chrysalis state, to the great injury of their breeders. The sporules of this fungus, floating in the air, enter the breathing-pores which open into the tracheal system of the silk-worm (Chap. XVII); they first develope themselves within the air-tubes, which are soon blocked up by their growth; and they then extend themselves through the fatty mass beneath the skin, occasioning the destruction of this tissue, which is very important as a reservoir of nutriment to the animal, when it is about to pass into a state of complete inactivity. The disease invariably occasions the death of the silk-worm which it attacks; but it seldom shows itself externally until afterwards, when it rapidly shoots forth from beneath the skin, especially at the junction of the rings of the body. Although it spontaneously attacks only the larva, yet it may be communi-

cated by inoculation to the Chrysalis and the Moth, as well as to the worm; and it has been also observed to attack other Lepidopterous Insects. A careful investigation of the circumstances which favor the development of this disease, was made by Audouin, who first discovered its real nature; and he showed that its spread is favored by the overcrowding of the worms in the breeding establishments, and particularly by the practice of throwing the bodies of such as die, into a heap in the immediate neighborhood of the living worms; this heap speedily becomes covered with this kind of "mould," which finds upon it a most congenial soil; and it keeps up a continual supply of sporules, which, being diffused through the atmosphere of the neighborhood, are drawn into the breathing pores of individuals previously healthy. Wherever the precautions obviously suggested by the knowledge of the nature of the disease thus afforded by

FIG. 120.



Botrytis bassiana.—A, the fungus as it first appears at the orifices of the stigmata; B, tubular filaments bearing short branches, as seen two days afterwards; C, magnified view of the same; D, appearance of filaments on the fourth and sixth days; E, masses of mature spores falling off the branches, with filaments proceeding from them.

the Microscope have been duly put in force, its extension has been kept within comparatively limited bounds. The plant pre-

sents itself (Fig. 120) under a considerable variety of forms; all of which, however, are of extremely simple structure, consisting of elongated or rounded cells, connected in necklace-like filaments, very nearly as in the ordinary "bead-moulds."

210. Again, it is not at all uncommon in the West Indies to see individuals of a species of *Polistes* (the representative of the Wasp of our own country) flying about with plants of their own length projecting from some part of their surface, the germs of which have been probably introduced (as in the preceding case) through the breathing-pores at their sides, and have taken root in their substance, so as to produce a luxuriant vegetation. In time, however, this fungous growth spreads through the body, and destroys the life of the insect; it then seems to grow more rapidly, the decomposing tissue of the dead body being still more adapted than the living structure to afford it nutriment. A similar growth of different species of the genus *Sphaeria* takes place in the bodies of certain caterpillars in New Zealand, Australia, and China; and being thus completely pervaded by a dense substance, which, when dried, has almost the solidity of wood, these Caterpillars come to present the appearance of twigs, with long slender stalks that are formed by the projection of the fungus itself. The Chinese

species is valued as a medicinal drug. The stomachs and intestines of many Worms and Insects are infested with Entophytic Fungi, which grow there with great luxuriance. In the accompanying illustrations (Figs. 121, 122) are shown some of the forms of the *Enterobryus*,¹ which has been found by Dr. Leidy to be so constantly present in the stomach of certain species of *Iulus* (gally-worm), that it is extremely rare to meet with individuals whose stomachs do not contain it. The *Enterobryus* originally consists of a single long tubular cell, which sometimes grows in

FIG. 121.




Growth of *Enterobryus spiralis* from mucous membrane of stomach of *Iulus*:—a, epithelial cells of mucous membrane; b, spiral thallus of *Enterobryus*; c, primary cells; d, e, secondary cells.

a spiral mode (Fig. 121), sometimes straight and tapering (Fig. 122, A); in its young state, the cell contains a transparent protoplasma, with granules and globules of various sizes; but in its more advanced condition, the tube of the filament is occupied by cells in various stages of development; these distend the

¹ This plant, also, has much affinity to Algae in its general type of structure, and is referred to that group by many Botanists; but the conditions of its growth, as in the case of *Sarcina*, seem rather to indicate its affinity to the Fungi; and until its proper fructification shall have been made out, its true place in the scale must be considered as undetermined.

terminal part of the cell (Fig. 122, B), and press so much against each other that their walls become flattened; whilst nearer the middle of the same filament (c), we find them retaining their rounded form, and merely lying in contact with each other; and

FIG. 122.



Structure of *Enterobryus*.—A, growth of *E. attenuatus*, from mucous membrane of the stomach of *Passeris*; B, dilated extremity of primary cell of *E. elegans*, filled with secondary cells, which, near its termination, become mutually flattened by pressure; C, lower portion of the same filament, containing cells mingled with granules; D, base of the same filament, containing globules interspersed among granules.

at the base (D) they lie detached in the midst of the granular protoplasm. In *E. spiralis*, the primary cells (Fig. 121, b, c) very commonly have secondary and even ternary cells (d) developed at their extremities; but this is rarely seen in *E. attenuatus* (Fig. 122). It may be considered as next to certain that the tubular filaments rupture, when the contained cells have arrived at maturity, and give them exit; and that these cells are developed, under favorable circumstances, into tubular filaments like those from which they sprang; but the process has not yet been thoroughly made out. This is obviously not the true Generation of the plant, but is analogous to the development of zoospores in *Achlya* (§ 197). It is not a little curious, moreover, that the Entozoa or parasitic worms infesting the alimentary canal of these animals, should be frequently clothed *externally* with an abundant growth of such plants; in one instance Dr. Leidy found an *Ascaris* bearing twenty-three filaments of *Enterobryus* "which appeared to cause no inconvenience to the animal, as it moved and wriggled about with all the ordinary activity of the species." The presence of this kind of vegetation seems to be related to the peculiar food of the animals in whose stomachs it is found; for Dr. Leidy could not discover a trace of these or of any other parasitic plants in the alimentary canal of the *carnivorous* Myriapods which he examined; whilst he met with a constant and most extraordinary profusion of vegetation (Fig.

123) in the stomach of an herbivorous beetle, the *Passulus cornutus*,

FIG. 123.



Fungoid Vegetation, clothing membrane of stomach of *Passulus*, intermingled with brush-like hairs.

which lives, like the Iuli, in stumps of old trees, and feeds as they do on decaying wood. Of this vegetation, some parts present themselves in tolerably definite forms, which have been described under various names; whilst other portions have the indefiniteness of imperfectly developed organisms, and can scarcely be characterized in the present state of our knowledge of them. With regard to several forms, indeed, Dr. Leidy expresses a doubt whether they are parasitic plants, or whether they are outgrowths of the membrane itself.


There are various diseased conditions of the Human skin and mucous membranes, in which there is a combination of fungoid vegetation and morbid growth of the animal tissues; this is the case, for example, with the *Tinea favosa*, a disease of the scalp, in which yellow crusts are formed, that consist almost entirely of the mycelium, receptacles, and sporules of a fungus; and the like is true also of those white patches (*Aphthæ*) on the lining membrane of the mouth of children, which are known as *Thrush*. In these and similar cases, two opinions are entertained as to the relation of the fungi to the diseases in which they present themselves; some maintaining that their presence is the essential condition of these diseases, which originate in the introduction of the vegetable germs; and others considering their presence to be secondary to some morbid alteration of the parts wherein the fungi appear, which alteration favors their development. The first of these doctrines derives a strong support from the fact, that the diseases in question may be communicated to healthy individuals, through the introduction of the germs of the fungi by inoculation; whilst the second is rather consistent with general analogy, and especially with what is known of the conditions under which the various kinds of fungoid "blights" develop themselves in or upon growing Plants (§ 212).

211. There are scarcely any Microscopic objects more beautiful, than some of those forms of "mould" or "mildew," which are so commonly found growing upon the surface of jams and

other preserves; especially when they are viewed with a low magnifying power, by reflected light. For they present themselves as a forest of stems and branches of extremely varied and elegant forms (Fig. 124), loaded with fruit of singular delicacy of conformation, all glistening brightly on a dark ground. In removing a portion of the "mould" from the surface whereon it grows, for the purpose of microscopic examination, it is desirable to disturb it no more than can be helped, in order that it may be seen as nearly as possible in its natural condition; and it is therefore preferable to take up a portion of the membrane-like substance whereon it usually rests, which is, in fact, a *mycelium* composed of interlacing filaments of the *vegetative* part of the plant, the stems and branches being its *reproductive* portion (§ 213). The universality of the appearance of these simple forms of Fungi upon all spots favorable to their development, has given rise to the belief that they are spontaneously produced by decaying substances; but there is no occasion for this mode of accounting for it; since the extraordinary means adopted by Nature for the production and diffusion of the germs of these plants, adequately suffices to explain the facts of the case. The number of sporules which any one Fungus may develop, is almost incalculable; a single individual of the puff-ball tribe has been computed to send forth no fewer than ten millions. And their minuteness is such, that they are scattered through the air in the condition of the finest possible dust; so that it is difficult to conceive of a place from which they should be excluded. This mode of explanation has received further confirmation from the facts recently ascertained, in regard to the great number of forms under which a single germ may develop itself; the particular form being determined, it seems likely, by the soil whereon each germ happens to grow. Hence we are not obliged to suppose that distinct germs are floating about in the atmosphere, for all the forms of fungous vegetation which appear to be of different species, and which are only found in particular situations,—the *Puccinia rosæ*, for example, only upon rose-bushes, *Isaria felina* only upon the dung of cats deposited in humid and obscure situations, and *Onygena exigua* upon the hoofs of dead horses;—but are warranted in believing that the real variety of germs is comparatively small, and that the facts just stated, with others of the same order, only indicate the modifying influence of the circumstances under which they are developed.

212. The parasitic Fungi which infest some of the Vegetables most important to Man, as furnishing his staple articles of food,

FIG. 124.

*Stysanus cupul-medusa.*

constitute a group of special interest to the Microscopist; of which a few of the chief examples may here be noticed. The *mildew*, which is often found attacking the straw of Wheat, shows itself externally in the form of circular clusters of pear-shaped spore-cases (Fig. 125), each containing two compartments filled with sporules; these (constituting the

FIG. 125.

*Puccinia graminis.*


Puccinia graminis) arise from a filamentous tissue constituting the *mycelium*, the threads of which interweave themselves with the tissue of the straw; and they generally make their way to the surface through the "stomata" or breathing-pores of its epidermis. The *rust*, which makes its appearance on the leaves and chaff-scales of Wheat, has a fructification that seems essentially distinct from that just described, consisting of oval spore-cases, that grow without any regularity of arrangement from the threads of the mycelium; and hence it has been considered to belong to a different genus and species, *Uredo rubigo*. But from the observations of Prof. Henslow, it seems cer-

tain that the "rust" is only an earlier form of the "mildew;" the one form being capable of development into the other, and the fructification characteristic of the two supposed genera having been evolved on one and the same individual. Another reputed species of *Uredo* (the *U. segetum*) it is, which, when it attacks the flower of the wheat, reducing the ears to black masses of sooty powder, is known as "smut" or "dust-brand." The corn-grains are entirely replaced by aggregations of spores; and these being of extreme minuteness, they are very easily and very extensively diffused. The "bunt" or "stinking rust" is another species of *Uredo* (the *U. foetida*), which is chiefly distinguished by its disgusting odor. The prevalence of these "blights" to any considerable extent, seems generally traceable to some seasonal influences unfavorable to the healthy development of the wheat plant; but they often make their appearance in particular localities, through careless cultivation, or want of due precaution in the selection of seed. It may be considered as certain that an admixture of the spores of any of these fungi with the grains, will endanger the plants raised from them; but it is equally certain that the fungi have little tendency to develop themselves in plants that are vegetating with perfect healthfulness. The wide prevalence of such blights in bad seasons is not difficult to account for, if it be true (as the observations of Mr. John Marshall, a few years since, rendered probable) that there are really *very few* wheat-grains, near the points of which one or two sporules of Fungi may not be found, entangled among their minute hairs;

and it may be fairly surmised that these sporules remain dormant, unless an unfavorable season should favor their development, by inducing an unhealthy condition of the wheat-plant. The same general doctrine probably applies to the *Botrytis*, which, from 1847 to the present time, has had a large share in the production of the "Potato-disease;" and to the *Oidium*, which has a like relation to the "Vine-disease" that has been extending itself for some years past through the south of Europe. There seems no doubt that, in the fully developed disease, the Fungus is always present; and that its growth and multiplication have a large share in the increase and extension of the disorder, just as the growth of the Yeast-plant excites and accelerates fermentation, and its reproduction enables this action to be indefinitely extended through its instrumentality. But just as the Yeast-plant will not vegetate save in a fermentable fluid, that is, in a solution which, in addition to sugar, contains some decomposable albuminous matter,—so does it seem probable, on a consideration of all the phenomena of the Potato and Vine diseases, that neither the *Botrytis* of the one, nor the *Oidium* of the other will vegetate in perfectly healthy plants; but that a disordered condition, induced either by forcing and therefore unnatural systems of cultivation, or by unfavorable seasons, or by a combination of both, is necessary as a "predisposing" condition.

213. In those lower forms of this class to which our notice of it has hitherto been chiefly restricted, there is not any very complete separation between its nutritive or vegetative, and its Reproductive portions; every cell, as in the simplest Protophytes, being equally concerned in both. But such a separation makes

FIG. 126.



Oidium tussilaginis.—A, portion of the plant magnified; B, section of one of the conceptacles with its spores.

itself apparent in the higher; and this in a very curious mode. For the ostensible Fungi of almost every description (Fig. 126) consist, in fact, of nothing else than the organs of *fructification*;

the nutritive apparatus of these plants being composed of an indefinite *mycelium*, which is a filamentous expansion (Fig. 127), composed of elongated branching cells (*a*), interlacing amongst each other, but having no intimate connection; and this "mycelium" has such an indefiniteness of form, and varies so little in

FIG. 127.



Clavaria crispula:—*a*, portion of the mycelium magnified.

the different tribes of Fungi, that no determination of species, genus, or even family, could be certainly made from it alone. The recent observations of Tulasne render it probable that a true sexual generation exists among the Fungi; since he has ascertained that the presence of bodies resembling the *spermatia* of Lichens (§ 207) may be regarded as universal in the organs of fructification, at an early period of their development. These are budded off (so to speak) from ramifying filaments, which are sometimes developed in the midst of those that bear the spores, and are sometimes found on other parts of the plant, being occasionally included in distinct conceptacles or *spermatogonia*, as in Lichens. The whole history of the development of the Fungi, and the question of the relationship of its different forms to each other, is one that most urgently calls for re-examination at the present time, under the guidance of our recently acquired knowledge, and with the assistance of improved instruments of Microscopic investigation; and whilst there is a wide field for the labors of those who possess only instruments of very moderate capacity, there are several questions which can only be worked out by means of the highest powers and the most careful appliances which the practised Microscopist can bring to bear upon them.

214. The little group of *Hepaticæ* or "Liverworts," which is intermediate between Lichens and Mosses,—rather agreeing with the former in its general mode of growth, whilst approaching the latter in its fructification,—presents numerous objects of great interest to the Microscopist; and no species is richer in these, than the very common *Marchantia polymorpha*, which may often be found growing between the paving-stones of damp court-yards, but which particularly luxuriates in the neighborhood of springs or water-falls, where its lobed fronds are found covering extensive surfaces of moist rock or soil, adhering by the radical (root) filaments which arise from their lower surface.


At the period of fructification, these fronds send up stalks, which carry at their summit either round shield-like disks, or radiating bodies that bear some resemblance to a wheel without its tire (Fig. 128); the former carry the male organs or antheridia, and the latter, at an early period, the female organs or archegonia, which afterwards give place to the *sporangia* or spore-cases.¹ But besides these, the frond usually bears upon its surface (as shown in Fig. 128) a number of little open basket-shaped "conceptacles," whose nature and purpose will be presently explained. The green surface of the frond of this Liverwort is seen under a low magnifying power to be divided into minute diamond-shaped spaces (Fig. 129, A, *a, a*) bounded by raised bands (*c, c*); every one of these spaces has in its centre a curious brownish-colored body (*b, b*), with an opening in its middle, which allows a few small green cells to be seen through it. When a thin vertical section is made of the frond (B), it is seen that each of the lozenge-shaped divisions of its surface corresponds with an air-chamber in its interior; which is bounded below by a floor (*a, a*) of closely set cells (from whose under surface the radical fila-

FIG. 128.



Frond of *Marchantia polymorpha*, with gemmiferous conceptacles, and lobed receptacles bearing pistillidia.

FIG. 129.



A, Portion of frond of *Marchantia polymorpha* seen from above; *a, a*, lozenge-shaped divisions; *b, b*, stomata seen in the centre of the lozenges; *c, c*, greenish bands separating the lozenges:—B, vertical section of the frond, showing *a, a*, the dense layer of cellular tissue forming the floor of the cavity *d, d*; *b, b*, cuticular layer, forming its roof; *c, c*, its walls; *f, f*, loose cells in its interior; *g*, stoma divided perpendicularly; *h*, rings of cells forming its wall; *i*, cells forming the obturator ring.


ments arise), at the sides by walls (*c, c*) of similar solid parenchyma, the projection of whose summits forms the raised bands on the surface, and above by a cuticle (*b, b*) formed of a single layer of cells; whilst its interior is occupied by a very loosely

¹ In some species, the same shields bear both sets of organs; and in *Marchantia androgyna*, we find the upper surface of one half of the pelta developing antheridia, whilst the under surface of the under half bears archegonia.

arranged parenchyma, composed of branching rows of cells (*f, f*) that seem to spring from the floor,—these cells being what are seen from above, when the observer looks down through the central aperture just mentioned. If the vertical section should happen to traverse one of the peculiar bodies which occupies the centres of the divisions, it will bring into view a structure of remarkable complexity. Each of these *stomata* (as they are termed, from the Greek *στομα*, mouth) forms a sort of shaft (*g*), composed of four or five rings (like the “courses” of bricks in a chimney) placed one upon the other (*h*) every ring being made up of four or five cells; and the lowest of these rings (*i*) appears to regulate the aperture, by the contraction or expansion of the cells which compose it, and it is hence termed the “obturator ring.” In this manner, each of the air-chambers of the frond is brought into communication with the external atmosphere; the degree of that communication being regulated by the limitation of the aperture. We shall hereafter find (§ 245) that the leaves of the higher plants contain intercellular spaces, which also communicate with the exterior by “stomata;” but that the structure of these organs is far less complex in them, than it is in this humble Liverwort.

215. The basket-shaped “conceptacles” which are borne upon the surface of the frond (Fig. 130, *A*), and which may often be

FIG. 130.



Gemmiparous conceptacles of *Marchantia polymorpha*:—*A*, conceptacle fully expanded, rising from the surface of the frond, *a, a*, and containing disks already detached;—*B*, first appearance of conceptacle on the surface of the frond, showing the formation of its fringe by the splitting of the cuticle.


found in all stages of development, are structures of singular beauty. They contain, when mature, a number of little green round or oblong disks, each composed of two or more layers of cells; and their wall is surmounted by a glistening fringe of “teeth,” whose edges are themselves regularly fringed with minute outgrowths. This fringe is at first formed by the splitting up of the epidermis, as seen at *B*, at the time when the conceptacle and its contents are first making their way above the surface. The little disks (sometimes termed “bulbels,” from their analogy to the bulbels or detached buds of Flowering

Plants) are at first evolved as single globular cells, supported upon other cells which form their footstalks; these single cells gradually undergo multiplication by duplicative subdivision, until they evolve themselves into the disks; and these disks, when mature, spontaneously detach themselves from their footstalks, and lie free within the cavity of the conceptacle. Most commonly they are at last washed out by rain, and are thus carried to different parts of the neighboring soil, on which they grow very rapidly when well supplied with moisture; sometimes, however, they may be found growing whilst still contained within the conceptacles, forming natural grafts (so to speak) upon the stock from which they have been developed and detached; and many of the irregular lobes which the frond of the *Marchantia* puts forth, seem to have this origin. When this plant vegetates in damp shady situations, which are favorable to the nutritive processes, it does not readily produce the true fructification, which is to be looked for rather in plants growing in more exposed places. Each of the stalked peltate (shield-like) disks contains a number of flask-shaped cavities opening upon its upper surface, which are brought into view by a vertical section; and in each of these cavities is lodged an "antheridium," composed of a mass of "sperm-cells," within which are developed "antherozoids" like those of *Chara* (§ 202), surmounted by a long neck that projects through the mouth of the flask-shaped cavity. The wheel-like receptacles (Fig. 128) on the other hand, bear on their under surface, at an early stage, concealed between membranes that connect the origins of the lobes with one another, a set of "archegonia," shaped like flasks with elongated necks (Fig. 131); each of these has in its interior a "germ-cell," to which a canal

leads down from the extremity of the neck; and there is every reason to believe that, as in Ferns, the germ-cell is fertilized by the penetration of the antheridia through this canal, until they reach it. Instead, however, of at once evolving itself into a new plant resembling its parent, the fertilized germ-cell, or embryo-cell, develops itself into a mass of cells enclosed within a capsule, which is termed a "sporangium;" and thus the mature receptacle, in place of archegonia, bears capsules or sporangia, which finally burst open, and discharge their contents. These contents

consist of "spores," which are isolated cells, enclosed in firm yellow envelopes; and of "elaters," which are ovoidal cells, each containing a double spiral fibre coiled up in its interior. This fibre is so elastic, that, when the surrounding pressure is withdrawn by the bursting of the sporangium, the spires extend

FIG. 131.

Archegonia of *Marchantia polymorpha*, in successive stages of development.


themselves (Fig. 132), tearing apart the cell-membrane; and they do this suddenly, so as to jerk forth the spores which may be adherent to their coils, and thus to assist in their dispersion. The spores, when subjected to moisture, with a moderate amount of light and warmth, develop themselves into little collections of cells, which gradually assume the form of a flattened frond; and thus the species is very extensively multiplied, every one of the mass of spores, which is the product of a single germ-cell, being capable of giving origin to an independent individual.

Fig. 132.

Eliater and Spores of *Marchantia*.


216. The tribe of *Mosses* is as remarkable from the delicacy and minuteness of all the plants composing it, as other orders of the Vegetable Kingdom are for the majesty of their forms, the richness of their foliage, or the splendor of their blossoms. There is not one of this little tribe, whose external organs do not serve as beautiful objects, when viewed with low powers of the Microscope; while their more concealed wonders are admirably fitted for the detailed scrutiny of the practised observer. The Mosses always possess a distinct axis of growth, commonly more or less erect, on which the minute and delicately formed leaves are arranged with great regularity. The stem shows some indication of the separation of a *cortical* or bark-like portion, from the *medullary* or pith-like, by the intervention of a circle of bundles of elongated cells, which seem to prefigure the woody portion of the stem of higher plants, and from which prolongations pass into the leaves, so as to afford them a sort of midrib. The leaf usually consists of either a single or a double layer of cells, having flattened sides by which they adhere one to another; they rarely present any distinct epidermic layer; but such a layer, perforated by stomata of simple structure, is commonly found on the *setæ* or bristle-like footstalks bearing the fructification, and sometimes on the midribs of the leaves. The leaf-cells of the *Sphagnum* (bog moss) exhibit a very curious departure from the ordinary type; for instead of being small and polygonal, they are large and elongated (Fig. 133); they contain spiral fibres loosely coiled in their interior; and their membranous walls have large rounded apertures, by which their cavities freely communicate with one another, as is sometimes curiously evidenced by the passage of Wheel Animalcules, that make their habitation in these chambers. Between these coarsely spiral cells, are some thick-walled narrow elongated cells, which give to the leaf its firmness; these, in the very young leaf (as Mr. Huxley has pointed out) do not differ much in appearance from the others; the peculiarities of both being evolved by a gradual process of "differentiation." The chief interest of the Mosses, however, to the Microscopist, lies

in their fructification; which recent discoveries have invested with a new character. What has been commonly regarded in that light, namely, the "capsule" or "urn," borne at the top of a long footstalk, which springs from the centre of a cluster of leaves (Fig. 134, A), is not the real fructification, but its product; for Mosses, like Liverworts, possess both antheridia and pistillidia, although these are by no means conspicuous. These organs are sometimes found in the same envelope (or perigone), sometimes on different parts of the same plants, sometimes only on different individuals; but in either case, they are usually situated close to the axis, among the bases of the leaves. The antheridia are globular, oval, or elongated bodies (Fig. 135, A) composed of aggregations of cells, of which the exterior form a sort of capsule, whilst the interior are "sperm-cells," each of which, as it comes to maturity, develops within itself an "antherozoid" (B, C, D); and the antherozoids, set free by



Portion of the leaf of *Sphagnum*; showing the large cells, a, a, a, with spiral fibres and communicating apertures; and the intervening bands, b, b, b, composed of small elongated cells.

FIG. 134.



Structure of Mosses:—A, Plant of *Funaria hygrometrica*, showing *f*, the leaves, *u*, the urns supported upon the setae or footstalks, *s*, closed by the operculum, *o*, and covered by the calyptra, *c*:—B, urn of *Encalypta vulgaris*, one of them closed and covered with the calyptra, the other open; *u*, *u*, the urns; *o*, *o*, the opercula; *c*, calyptra; *p*, peristome; *s*, *s*, setae:—C, longitudinal section of very young urn of *Splachnum*; *a*, solid tissue forming the lower part of the capsule; *c*, columella; *l*, loculus or space around it for the development of the spores; *e*, epidermic layer of cells, thickened at the top to form the operculum, *o*; *p*, two intermediate layers, from which the peristome will be formed; *s*, inner layer of cells forming the wall of the loculus.

the rupture of the cells within which they are formed, make

their escape by a passage that opens for them at the summit of the antheridium. The antheridia are generally surrounded by a

cluster of hair-like filaments, composed of cells joined together (Fig. 134, A), and called "paraphyses." The archegonia bear a general resemblance to those of *Marchantia* (Fig. 131); and there is every reason to believe that the fertilization of their contained germ-cells is accomplished in the manner already described; for antherozoids have been observed swimming about around the archegonia within their involucre,¹ and the evolution of capsules from archegonia has been ascertained not to take place in those Mosses which bear the two sets of organs on separate individuals, unless an antheridial plant be in the neighborhood. The fertilized embryonal cell becomes gradually developed by cell division into a conical body elevated upon a stalk; and this at length tears across the walls of the flask-shaped archegonium by a circular fissure, carrying the higher part upwards as a *calyptra* or hood (Fig. 135, B, c) upon its summit, while the lower part remains to form a kind of collar round the base of the stalk.




Antheridia and Antherozoids of *Polytrichum commune*: A, group of antheridia, mingled with hairs and sterile filaments (paraphyses); of the three antheridia, the central one is in the act of discharging its contents; that on the left is not yet mature, while that on the right has already emptied itself, so that the cellular structure of its walls becomes apparent;—B, cellular contents of an antheridium, previously to the development of the antherozoids;—C, the same, showing the first appearance of the antherozoids;—D, the same, mature and discharging the antherozoids.

217. The "urns" or spore-capsules of Mosses, which are thus the immediate product of the Generative act, and which must really be considered as the offspring of the plants that bear them

¹ The detection of the antherozoids *within* the canal of the archegonium, and upon the surface of the germ-cell, is a point well worthy of Microscopic research.


(although grafted on to these, and drawing their nourishment from them), are closed at their summit by *opercula* or lids (Fig. 135, B, o, o), which fall off when the contents of the capsules are mature, so as to give them free exit; and the mouth thus laid open is surrounded by a beautiful toothed fringe, which is termed the *peristome*. This fringe, as seen in its original undisturbed position, is shown in Fig. 136; whilst in Figs. 137–139 are shown

FIG. 136.



Mouth of capsule of *Funaria*, showing the Peristome *in situ*.


FIG. 137.



Double Peristome of *Fontinalis antipyretica*.

three different forms of it, spread out and detached, illustrating the varieties which it exhibits in different genera of Mosses,—varieties whose existence and readiness of recognition render them characters of extreme value to the systematic Botanist, whilst they furnish objects of great interest and beauty for the

FIG. 138.



Double Peristome of *Bryum intermedium*.

FIG. 139.




Double Peristome of *Cinctidium arcticum*.

Microscopist. The peristome seems always to be originally double, one layer springing from the outer, and the other from the inner, of two layers of cells which may be distinguished in the immature capsule (Fig. 135, c, p); but frequently, at the time of maturity, one or other of these is wanting, and sometimes

both are obliterated, so that there is no peristome at all. The number of the "teeth" is always a multiple of 4, varying from 4 to 64; sometimes they are prolonged into straight or twisted hairs. The spores are contained in the upper part of the capsule, where they are clustered round a central pillar, which is termed the *columella*. In the young capsule, the whole mass is nearly solid (Fig. 135, o), the space (*l*) in which the spores are developed being very small; but this gradually augments, the walls becoming more condensed; and at the time of maturity, the interior of the capsule is almost entirely occupied by the spores, in the dispersion of which the peristome seems in some degree to answer the same purpose as the elaters of Hepaticæ. The development of the spores into new plants, commences with the rupture of their outer walls, and a protrusion of their inner coats; and from the projecting extremity new cells are put forth by a process of outgrowth, which form a sort of confervoid filament (as in Fig. 145, c). At certain points of this filament, its component cells multiply by subdivision, so as to form rounded clusters, from every one of which an independent plant may arise; so that several individuals may be evolved from a single spore. A numerous aggregation of spores may be developed, as we have seen, from a single germ-cell; so that the immediate product of each act of fertilization does not consist (as in the higher Plants) of a single seed, that afterwards develops itself into a composite fabric, whence are put forth a multitude of leaf buds, every one of which is capable (under favorable circumstances) of evolving itself into a complete Plant; but divides itself at once into a mass of isolated cells (spores), of which every one may be considered in the light of a bud or *gemma* of the simplest possible kind, and one of the first acts of which is to put forth other buds, whereby the rapid extension of these plants is secured, although no separate individual ever attains more than a very limited size.

FIG. 140.



Oblique section of footstalk of *Fern-leaf*, showing bundle of scalariform ducts.

218. In the *Ferns* we have in many respects a near approximation to Flowering plants; but this approximation does not extend to their Reproductive apparatus, which is formed upon a type essentially the same as that of Mosses, but is evolved at a very different period of life. As the component tissues of which their fabrics are composed, are essentially the same as those which will be described in the next chapter, it will not be requisite here to dwell upon them. The stem (where it exists)

is for the most part made up of cellular parenchyma, which is separated into a cortical and a medullary portion, by the inter-

position of a circular series of fibro-vascular bundles containing true woody tissue and ducts. These bundles form a kind of irregular network, from which prolongations are given off that pass into the leaf-stalks, and thence into the midrib and its lateral branches; and it is their peculiar arrangement in the leaf-stalks, which gives to the transverse section of these the figured marking commonly known as "King Charles in the oak." A thin section, especially if somewhat oblique (Fig. 140), displays extremely well the peculiar character of the

FIG. 141.

Leaflet of *Polypodium*, with sori.

FIG. 142.

Portion of Frond of *Hamionites*, with sori.

ducts of the Fern; which are termed "scalariform," from the resemblance of the regular markings on their walls to the rungs of a ladder. What is usually considered the "fructification" of the Ferns, affords a most beautiful and readily prepared class of opaque objects for the lowest powers of the Microscope; nothing more being necessary, than to lay a fragment of the frond that bears it on its under surface, upon the glass stage-plate, or to hold it in the stage-forceps, and to throw an adequate light upon it by the side-condenser. It usually presents itself in the form of isolated spots, termed *sori*, as in the common *Polypodium* (Fig. 141), and in the *Aspidium* (Fig. 143); but sometimes these "sori" are elongated into bands, as in the common *Scolopendrum* (Harts tongue): and these bands may coalesce with each other, so as almost to cover the surface of the frond with a network, as in *Hamionites* (Fig. 142); or they may form merely a single band along its borders, as in the common *Pteris* (brake-fern). The sori are sometimes naked on the under surface of the fronds;

but they are frequently covered with a delicate membrane, termed the *indusium*, which may either form a sort of cap upon the summit of each sorus, as in *Aspidium* (Fig. 143), or a long fold, as in *Scolopendrum* and *Pteris*, or a sort of cup, as in *Deparia* (Fig. 144). Each of these sori, when sufficiently magnified, is found to be made up of a multitude of capsules or *thecae* (Figs. 143, 144), which are sometimes closely attached to the surface of the leaf, but more commonly spring from it by a pedicel or foot-stalk. The wall of the capsule is composed of flattened cells, applied to each other by their edges; but there is generally one row of these, thicker and larger than the rest, which springs from the pedicel, and is continued over the summit of the capsule, so as to form a projecting ring, which is known as the *annulus*. This ring has an elasticity superior to that of all the rest of the capsular wall, causing it to split across, when mature, so that the contained spores may escape; and in many instances carrying the two halves of the capsule widely apart from each other (Fig. 141), the fissure extending to such a depth as to separate them completely. It will frequently happen, that specimens of Fern-fructification gathered for the Microscope, will be found to have all the capsules burst and the spores dispersed, whilst in others, less advanced, the capsules may all be closed; others, however, may often be met with, in which some of the capsules are closed and others are open; and if these be watched with sufficient attention, the rupture of some of the thecae and the dispersion of the spores may be observed to take place, whilst the specimen is under observation in the field of the microscope. In sori whose capsules have all burst, the annuli connecting their two halves are the most conspicuous objects, looking, when a strong light is thrown upon them, like strongly banded worms of a bright brown hue. This is particularly the case in *Scolopendrum*,

FIG. 143.


Sorus and indusium of *Aspidium*.

FIG. 144.


Sorus and cup-shaped indusium of *Deparia prolifera*.

whose elongated sori are remarkably beautiful objects for the microscope in all their stages; until quite mature, however, they need to be brought into view by turning back the two indusial folds that cover them. The commonest Ferns, indeed, which are found in almost every hedge, furnish objects of no less beauty

than those yielded by the rarest exotics; and it is in every respect a most valuable training to the young, to teach them how much there may be found to interest, when looked for with intelligent eyes, even in the most familiar and therefore disregarded specimens of Nature's handiwork.

219. The spores (Fig. 145, A) set free by the bursting of the thecae, usually have a somewhat angular form, and are invested by a yellowish or brownish outer coat, which is marked, very much in the manner of pollen-grains (Fig. 187), with points, streaks, ridges, or reticulations. When placed upon a damp surface, and exposed to a sufficiency of light and warmth, the spore begins to "germinate," the first indication of its vegetative activity being a slight enlargement, which is manifested in the

FIG. 145.




Development of Prothallium of *Pteris serrulata*.—A, spore set free from the theca;—B, spore beginning to germinate, putting forth the tubular prolongation *a* from the principal cell, *b*;—C, first formed linear series of cells;—D, prothallium taking the form of a leaf-like expansion; *a* first, and *b* second radical fibre; *c, d*, the two lobes, and *e* the indentation between them; *f, f*, first formed part of the prothallium; *g*, external coat of the original spore; *h, h*, antheridia.

rounding off of its angles; this is followed by the putting forth of a tubular prolongation (B, *a*) of the internal cell-wall, through an aperture in the outer spore-coat; and by the absorption of moisture through this root-fibre, the inner cell is so distended, that it bursts the external unyielding integument, and soon begins to elongate itself in a direction opposite to that of the root-fibre. A production of new cells by subdivision then takes place from its growing extremity; this at first proceeds in a single series, so as to form a kind of confervoid filament (C); but the multiplication of cells by subdivision soon takes place transversely as well as longitudinally, so that a flattened leaf-like expansion (D) is produced, so closely resembling that of a young *Marchantia* as to be readily mistaken for it. This expansion, which is termed the *prothallium*, varies in its configuration in different species;

but its essential structure always remains the same. From its under surface are developed, not merely the root-fibres (*a, b*) which serve to fix it in the soil, and at the same time to supply it with moisture, but also the antheridia and archegonia, which constitute the true representatives of the essential parts of the flower of higher Plants. Some of the *antheridia* may be distinguished at an early period of the development of the prothallium (*h, h*); and at the time of its complete evolution these bodies are seen in considerable numbers, especially about the origins of the root-fibres. Each has its origin in a peculiar protrusion that takes place from one of the cells of the prothallium (Fig. 146, *A, a*);

FIG. 146.




Development of the Antheridia and Antherozoids of *Pteris serrulata*.—*A*, projection of one of the cells of the prothallium, showing the antheridial cell, *b*, with its sperm cells, *c*, within the cavity of the original cell, *a*;—*B*, antheridium completely developed; *a*, wall of antheridial cell; *c*, sperm-cells, each enclosing an antherozoid;—*C*, one of the antherozoids more highly magnified, showing *a*, its large extremity, *b*, its small extremity, *d, d*, its cilia.

this is at first entirely filled with chlorophyll-granules; but soon a peculiar free cell (*b*) is seen in its interior, filled with mucilage and colorless granules. This cell gradually becomes filled with another brood of young cells (*c*), and increases considerably in its dimensions, so as to fill the projection which encloses it; this part of the original cavity is now cut off from that of the cell of which it was an offshoot, and the antheridium henceforth ranks as a distinct and independent organ. Each of the secondary cells (*B*) contained within its primary cell, is seen, as it approaches maturity, to contain a spirally coiled filament; and when they have been set free by the bursting of the antheridium, they themselves burst and give exit to their "antherozoids" (*c*), which execute rapid movements of rotation on their axes, partly dependent on the six long cilia with which they are furnished. The *archegonia* are fewer in number, and are found upon a different part of the prothallium. Each of them at its origin presents itself only as a slight elevation of the cellular layer of the prothallium, within which is a large intercellular space containing a peculiar cell (the "germ-cell"), and opening externally by an orifice at the summit of the projection; but when fully developed (Fig. 147), it is composed of from ten to twelve cells built up in layers of four cells each, one upon another, so as to form a kind of chimney or shaft, having a central passage that leads down to the cavity at its base, wherein the germ-cell is contained. Into this cavity the antherozoids penetrate, so as to come into contact with

the "germ-cell;" and, by the softening of the membrane at its apex, they are even enabled to enter its cavity, within which a minute "embryonic vesicle" was previously distinguishable. This embryonic vesicle, when fertilized by the antherozoids which

FIG. 147.



Archegonium of *Pteris serrulata*:—A, as seen from above; a, a, a, cells surrounding the base of the cavity; b, c, d, successive layers of cells, the highest enclosing a quadrangular orifice:—B, side view, showing, A, A, cavity containing the germ-cell; B, B, walls of the archegonium, made up of the four layers of cells, b, c, d, e, and having an opening at the summit; c, c, antherozoids within the cavity; g, large extremity; h, thread-like portion; i, small extremity in contact with the germ-cell and dilated.


move actively round it, becomes the primordial cell of a new plant, the development of which speedily commences.¹ By the usual process of duplicative subdivision a globular homogeneous mass of cells is at first formed; but rudiments of special organs soon begin to make their appearance; the germ grows at the expense of the nutriment prepared for it by the prothallium; and it soon bursts forth from the cavity of the archegonium, which organ in the meantime is becoming atrophied. In the very beginning of its development, the tendency is seen in the cells of one extremity to grow upwards, so as to evolve the stem and leaves, and in those of the other extremity to grow downwards,

¹ See Hofmeister, in "Ann. of Nat. Hist.," 2d Ser. vol. xiv, p. 272. The study of the development of the spores of Ferns, and of the act of fertilization and of its products, may be conveniently prosecuted as follows:—Let a frond of a Fern, whose fructification is mature, be laid upon a piece of fine paper, with its spore-bearing surface downwards; in the course of a day or two, this paper will be found to be covered with a very fine brownish dust, which consists of the discharged spores. This must be carefully collected, and should be spread upon the surface of a smoothed fragment of porous sandstone; the stone being placed in a saucer, the bottom of which is covered with water, and a glass tumbler being inverted over it, the requisite supply of moisture is insured, and the spores will germinate luxuriantly. Some of the prothallia soon advance beyond the rest; and at the time when the advanced ones have long ceased to produce antheridia, and bear abundance of archegonia, those which have remained behind in their growth are beginning to be covered with antheridia. If the crop be now kept with little moisture for several weeks, and then suddenly watered, a large number of antheridia and archegonia simultaneously open; and in a few hours afterwards, the surface of the larger prothallia will be found almost covered with moving antherozoids. Such prothallia as exhibit freshly opened archegonia, are now to be held by one lobe between the forefinger and thumb of the left hand, so that the upper surface of the prothallium lies upon the thumb; and the thinnest possible sections are then to be made, with a thin narrow-bladed knife, perpendicularly to the surface of the prothallium. Of these sections, which, after much practice, may be made no more than 1-15th of a line in thickness, some will probably lay open the canals of the archegonia; and within these, when examined with a power of 200 or 300 diameters, antherozoids may be occasionally distinguished.

to form the root; and when these organs have been sufficiently developed to absorb and prepare the nutriment which the young plant requires, the prothallium, whose functions as a "nurse" is now discharged, decays away.

220. The little group of *Equisetaceæ* (Horsetails), which seem nearly allied to the Ferns in the type of their generative apparatus, though that of their vegetative portion is very different, affords certain objects of considerable interest to the Microscopist. The whole of their structure is penetrated to so extraordinary a degree by *silex*, that, even when the organic portion has been destroyed by prolonged maceration in dilute nitric acid, a consistent skeleton still remains. This mineral, in fact, constitutes in some species not less than 13 per cent. of the whole solid matter, and 50 per cent. of their inorganic ash. The cuticle, which is used by cabinet-makers for smoothing the surface of wood, becomes, through the peculiar arrangement of its siliceous particles, an extremely beautiful object under polarized light. Of these particles (each of which has a regular axis of double refraction), some are distributed in two lines, parallel to the axis: others, however, are grouped into oval forms, connected with each other, like the jewels of a necklace, by a chain of particles forming a sort of curvilinear quadrangle; and these (which are, in fact, the particles occupying the cells of the stomata) are arranged in pairs. What is usually designated as the fructification of the *Equisetaceæ*, forms a cone or spike at the extremity of certain of the stem-like branches (the real stem being a horizontal rhizoma);

FIG. 148.

Spores of *Equisetum*.

and consists of a cluster of shield-like disks, each of which carries a circle of *thecae* or spore-cases, that open by longitudinal slits to set free the spores. Each of these bodies has, attached to it, a pair of elastic filaments (Fig. 148), that are originally formed as spiral fibres on the interior of the wall of the primary cell within which the spore is generated, and are set free by its rupture; these are at first coiled up closely around the spore, in the manner represented at A, though more closely applied to the surface; but, on the slightest application of moisture, they suddenly extend themselves in the manner shown at B; and this motion, like the extension of the spiral elaters of *Marchantia*, probably serves to assist in the dispersion of the spores. If a number of the spores be spread out on a slip of glass under the field of view, and, whilst the observer watches them, a bystander

breathe gently upon the glass, all the filaments will be instantaneously put in motion,—thus presenting an extremely curious spectacle,—and will almost as suddenly return to their previous condition, when the effect of the moisture has passed off. These spores are to be regarded in the same light as those of Ferns; namely, as *gemmæ* or rudimentary buds, not as seeds. They evolve themselves after the like method into a prothallium; and this developes antheridia and archegonia, by the conjoint action of which an embryo is produced.

221. In ascending, as we have now done, from the lower to the higher Cryptogamia, we have seen a gradual change in the general plan of structure, so that the superior forms present a close approximation to the Flowering Plant, which is undoubtedly the highest type of Vegetation. But we have everywhere encountered a mode of Generation, which, whilst essentially the same throughout the series, is essentially distinct from that of the Phanerogamia; the fertilizing material of the “sperm-cell” being embodied, as it were, in self-moving filaments, which find their way to the germ-cells by their own independent movements; and the “embryo-cell” being destitute of that store of prepared nutriment, which surrounds it in the true seed, and serves as the pabulum for its early development. In the lower Cryptogamia, we have seen that the embryo-cell, after fertilization, is thrown at once upon the world (so to speak) to get its own living; but in the Liverworts, Mosses, and Ferns, the embryo-cell is nurtured by the parent-plant, for a period that varies in each case according to the nature of the fabric into which it evolves itself. While the true reproduction of the species is effected by the proper Generative act, the multiplication of the individual is accomplished by the production and dispersion of spores; and this production, as we have seen, takes place at very different periods of existence in the several groups, dividing the life of each into two separate epochs, in which it presents itself under two very distinct phases that contrast remarkably with each other. Thus, the frond of the *Marchantia*, bearing its antheridia and archegonia, is that which seems naturally to constitute *the* plant; but that which represents it in the Ferns, is the minute *Marchantia*-like prothallium. On the other hand, the product into which the fertilized embryo-cell evolves itself in the Ferns, is that which is commonly regarded as *the* plant; and this is represented in the Liverworts and Mosses by the spore-capsules alone. We shall notice, in the next Chapter (§ 256), the representation of these two phases in the life of the Flowering Plant, which is traceable by means of the study of the *Lycopodiaceæ* and *Coniferæ*; two groups that form the link of transition between these two great divisions of the Vegetable kingdom, the former being probably to be regarded as the highest of the Cryptogamia, and the latter as the lowest of the Phanerogamia.¹

¹ See the Author's "Principles of Comparative Physiology," Am. ed. 1854, §§ 499-505.

CHAPTER VIII.

OF THE MICROSCOPIC STRUCTURE OF PHANEROGAMIC PLANTS.

222. *Elementary Tissues.*—In passing from the Cryptogamic division of the Vegetable Kingdom, to that larger and more ostensibly important province which includes the Flowering Plants, we do not meet with so wide a departure from those simple types of structure we have already considered, as the great differences in general aspect and external conformation might naturally lead us to expect. For a very large proportion of the fabric of even the most elaborately formed Tree, is made up of components of the very same kind with those which constitute the entire organisms of the simplest Cryptogamia; and that proportion always includes the parts most actively concerned in the performance of the vegetative functions. For although the stems, branches, and roots, of trees and shrubs, are principally composed of *woody* tissue, such as we do not meet with in any but the highest Cryptogamia, yet the special office of this is to afford mechanical support; when it is once formed, it takes no further share in the vital economy, than to serve for the conveyance of fluid from the roots, upwards through the stem and branches, to the leaves; and even in these organs, not only the pith and the bark, with the “medullary rays” which serve to connect them, but that “cambium-layer” intervening between the bark and the wood (§ 240), in which the periodical formation of the new layers both of bark and wood takes place, are composed of *cellular* substance. This tissue is found, in fact, wherever *growth* is taking place; as, for example, in the spongioles or growing points of the root-fibres, in the leaf-buds and leaves, and in the flower-buds and sexual parts of the flower; it is only when these organs attain an advanced stage of development, that woody structure is found in them,—its purpose (as in the stem) being merely to give support to their softer textures; and the small proportion of their substance which it forms, being at once seen in those beautiful skeletons, which, by a little skill and perseverance, may be made of leaves, flowers, and certain fruits. All the softer and more pulpy tissue of these organs is composed of *cells* more or less compactly aggregated together,


and having forms which approximate more or less closely to the globular or ovoidal, which may be considered as their original type.

223. As a general rule, the rounded shape is preserved only when the cells are but loosely aggregated, as in the parenchymatous (or fleshy) substance of leaves (Fig. 149); and it is then only, that the distinctness of the walls becomes evident. When the tissue becomes more solid, the sides of the vesicles are pressed against each other, so as to flatten them and to bring them into close apposition; and they then adhere to one another in such a manner, that the partitions appear, except when carefully examined, to be single, instead of being (as they really are) double. Frequently it happens that the pressure is exerted more in one direction than in another, so that the form presented by the outline of the cell varies according to the direction in which the section is made.

This is well shown in the pith of the young shoots of Elder, Lilac, or other rapidly growing trees; the cells of which, when cut transversely, generally exhibit circular outlines, whilst, when the section is made vertically, their borders are straight, so as to make them appear like cubes or elongated prisms, as in Fig. 152. A very good example of such a cellular parenchyma is to be found in the substance known as "Rice-paper;" which is made by cutting the herbaceous stem of a Chinese plant termed *Aralia papyrifera*,¹ vertically round and round, with a long sharp knife, so that its tissue may be (as it were) unrolled in a sheet. The shape of the cells, as seen in the rice-paper thus prepared, is irregularly prismatical, as shown in Fig. 150, B; but if the stem be cut transversely, their outlines are seen to be circular or nearly so (A). When, as often happens, the cells have a very elongated form, this elongation is in the direction of their growth, which is that, of course, wherein there is least resistance. Hence their greatest length is nearly always in the direction of the axis; but there is one remarkable exception,—that, namely, which is afforded by the "medullary rays" of Exogenous stems (§ 239), whose cells are greatly elongated in the horizontal direction (Fig. 161, a), their growth being from the centre of the stem

¹ The *Eschynomene*, which is sometimes named as the source of this article, is an Indian plant, employed for a similar purpose.


FIG. 149.



Section of leaf of *Agave*, treated with dilute nitric acid, showing the primordial utricle contracted in the interior of the cells:—a, epidermic cells; b, boundary cells of the stoma; c, cells of parenchyma; d, their primordial utricles.

towards its circumference. It is obvious that fluids will be more readily transmitted in the direction of greatest elongation, being that in which they will have to pass through the least number of partitions; and whilst their ordinary course is in the direction of


FIG. 150.



Sections of Cellular Parenchyma of *Aralia*, or Rice-paper Plant;—A, transversely to the axis of the stem; B, in the direction of the axis.


the length of the roots, stem, or branches, they will be enabled, by means of the medullary rays, to find their way in the transverse direction. One of the most curious varieties of form which Vegetable cells present is that represented in Fig. 151, which constitutes the *stellate* cell. This modification, to which we have already seen an approximation in the *Volvox* (§ 159), is found in the spongy parenchymatous substance where lightness is an object; as in the stems of many

FIG. 151.



Section of Cellular parenchyma of *Rush*.

FIG. 152.



Cubical parenchyma, with stellate cells, from petiole of *Nuphar lutea*.

aquatic plants, such as the *Rush*, which need to be furnished with air-spaces. In other instances, these air-spaces are large cavities, which are altogether left void of tissue; such is the case

in the *Nuphar lutea* (Yellow water-lily), the footstalks of whose leaves contain large air-chambers, the walls of which are built up of very regular cubical cells, whilst some curiously formed large stellate cells project into the cavity which they bound (Fig. 152). The dimensions of the component vesicles of Cellular tissue are extremely variable; for although their diameter is very commonly between 1-300th and 1-500th of an inch, they occasionally measure as much as 1-30th of an inch across, whilst in other instances they are not more than 1-3000th.

224. The component cells of Cellular tissue are usually held together by an intercellular substance, which may be considered analogous to the "gelatinous" layer that intervenes between the cells of the Algae. There are many cellular substances, however, in which, in consequence of the loose aggregation of their component cells, these may be readily isolated, so as to be prepared for separate examination without the use of reagents which alter their condition; this is the case with the pulp of ripe fruits, such as the Strawberry or Currant (the Snowberry is a particularly favorable subject for this kind of examination), and with the parenchyma of many fleshy leaves, such as those of the Carnation (*Dianthus caryophyllus*) or the London Pride (*Saxifraga crassifolia*). Such cells usually contain evident nuclei, which are turned brownish-yellow by iodine, whilst their membrane is only turned pale-yellow; and in this way the nucleus may be brought into view, when, as often happens, it is not previously distinguishable. If a drop of the iodized solution of chloride of zinc be subsequently added, the cell-membrane becomes of a beautiful blue color, whilst the nucleus and the granular protoplasm that surrounds it, retain their brownish-yellow tint. The use of dilute nitric or sulphuric acid, of alcohol, of syrup, or of several other reagents, serves to bring into view the "primordial utricle" of Mohl; its contents being made to coagulate and shrink, so that it detaches itself from the cellulose wall with which it is ordinarily in contact, and shrivels up within its cavity, as shown in Fig. 149.

225. It is probable that all cells, at some stage or other of their growth, exhibit, in a greater or less degree of intensity, that curious movement of "rotation," which has been already described as occurring in the *Characæ* (§ 201), and which consists in the steady flow of one or of several currents of protoplasm over the inner wall of the cell; this being rendered apparent by the movement of the particles which the current carries along with it. The best examples of it are found among submerged plants, in the cells of which it continues for a much longer period than it usually does elsewhere; and among these there are two, the *Vallisneria spiralis* and the *Anacharis alismastrum*, which are peculiarly fitted for the exhibition of this interesting phenomenon. The former is an aquatic plant that grows abundantly in the rivers of the south of Europe, but is not a native

of this country; it may, however, be readily grown in a tall glass jar having at the bottom a couple of inches of mould, which, after the roots have been inserted into it, should be closely pressed down, the jar being then filled with water, of which a portion should be occasionally changed.¹ The jar should be freely exposed to light, and should be kept in as warm but equal a temperature as possible. The long grass-like leaves of this plant are too thick to allow the transmission of sufficient light through them for the purpose of this observation; and it is requisite to make a thin slice or shaving with a sharp knife. If this be taken from the surface, so that the section chiefly consists of the superficial layer of cells, these will be found to be small, and the particles of chlorophyll, though in great abundance, will rarely be seen in motion. But if, after the removal of this layer, a deeper stratum be sliced off, this will be found to consist of larger cells, some of them greatly elongated, with particles of chlorophyll in smaller number, but carried along in active rotation by the current of protoplasm; and it will often be noticed, that the rotation takes place in contiguous cells in opposite directions. If the movement (as is generally the case) be checked by the shock of the operation, it will be revived again by a gentle warmth; and it may continue under favorable circumstances, in the separated fragment, for a period of weeks or even of months. Hence when it is desired to exhibit the phenomenon, the preferable method is to make the sections a little time before they are likely to be wanted, and to carry them in a small vial of water in the waistcoat pocket, so that they may receive the gentle and continuous warmth of the body. In summer, when the plant is in its most vigorous state of growth, the section may be taken from any one of the leaves; but in winter, it is preferable to select those which are a little yellow.² The *Anacharis alsinastrum* is a water-weed, which, having been accidentally introduced into this country about twelve years ago; has since spread itself with such rapidity through our canals and rivers, as, in many instances, seriously to impede the navigation. It does not require to root itself in the bottom, but floats in any part of the water it inhabits; and it is so tenacious of life, that even small fragments are sufficient for the origination of new plants. The leaves have no distinct cuticle, but are for the most part composed of two layers of cells, and these are elongated and colorless in the centre, forming a kind of midrib; towards the margins of the leaves, however, there is but a single layer. Hence no preparation whatever is required for the exhibition of

¹ Mr. Quekett has found it the most convenient method of changing the water in the jars in which *Chara*, *Vallisneria*, &c., are growing, to place them occasionally under a water-tap, and allow a very gentle stream to fall into them for some hours; for by the prolonged overflow thus occasioned, all the impure water, with the conferva that is apt to grow on the sides of the vessel, may be readily got rid of.

² An object-glass of $\frac{1}{4}$ th inch focus affords the best power for the observation of this interesting phenomenon.


this interesting phenomenon; all that is necessary being to take a leaf from the stem (one of the older yellowish leaves being preferable), and to place it with a drop of water either in the aquatic box, or on a slip of glass beneath a thin glass cover. A higher magnifying power is required, however, than that which suffices for the examination of the rotation in *Chara* or *Vallisneria*; the $\frac{1}{3}$ th inch object-glass being here preferable to the $\frac{1}{4}$ th, and the assistance of the Achromatic condenser being desirable. With this amplification, the phenomenon may be best studied in the single layer of marginal cells; although, when a lower power is used, it is most evident in the elongated cells forming the central portion of the leaf. The number of chlorophyll granules in each cell varies from three or four to upwards of fifty; they are somewhat irregular in shape, some being nearly circular flattened disks, whilst others are oval; and they are usually from 1-3000th to 1-5000th of an inch in diameter. When the rotation is active, the greater number of these granules travel round the margin of the cells, a few, however, remaining fixed in the centre; their rate of movement, though only 1-40th of an inch per minute, being sufficient to carry them several times round the cell within that period. As in the case of the *Vallisneria*, the motion may frequently be observed to take place in opposite directions in contiguous cells. The thickness of the layer of protoplasm in which the granules are carried round, is estimated by Mr. Wenham at no more than 1-20,000th of an inch. A peculiar undulating appearance is seen in this, under certain modes of illumination, which suggests the idea of ciliary action; but this appearance is decidedly affirmed by Mr. Wenham to be an optical illusion. It is a curious circumstance, first remarked by Dr. Branson, that the elongated cells along the margin of the leaf and forming the midrib, contain a large quantity of silicex; as may be readily seen by polarized light, especially after the leaf has been boiled for a few minutes in equal parts of nitric acid and water, which removes part of the vegetable substance, and thus renders the siliceous portion more distinct, without destroying the form of the leaf.¹

226. The phenomena of "rotation," however, is by no means restricted to submerged Plants; for it has been witnessed by numerous observers in so great a variety of other species, that it may fairly be presumed to be universal. It is especially observable in the *hairs* of the epidermic surface; and according to Mr. Wenham, who has recently given much attention to this subject, "the difficulty is to find the exceptions, for hairs taken alike from the loftiest Elm of the forest, to the humblest weed that we trample beneath our feet, plainly exhibit this circulation." Such hairs are furnished by various parts of plants; and what is chiefly necessary is, that the part from which the hair is gathered

¹ See Dr. Branson in "Quart. Journ. of Microsc. Science," vol. ii, p. 131, and vol. iii, p. 374; and Mr. Wenham, *Op. cit.* vol. iii, p. 277.

should be in a state of vigorous growth. The hairs should be detached by tearing off, with a pair of fine-pointed forceps, the portion of the cuticle from which they spring; care being taken not to grasp the hair itself, whereby such an injury would be done to it, as to check its circulation. The hair should then be placed with a drop of water under thin glass; and it will generally be found advantageous to use a 1-8th in. objective, with an achromatic condenser having a series of diaphragms. The nature

FIG. 153.



Circulation of fluid in hairs of *Tradescantia virginica*:—A, portion of cuticle with hair attached; a, b, c, successive cells of the hair; d, cells of the cuticle; e, stoma;—a, joint of a beaded hair, showing several currents; a, nucleus.

of the movement in the hairs of different species of plants is far from being uniform. In some instances, the currents pass in single lines along the entire length of the cells, as in the hairs from the filaments of the *Tradescantia virginica* or Virginian spider-wort (Fig. 153, A); in others, there are several such currents which retain their distinctness, as in the jointed hairs of the calyx of the same plant (B); in others, again, the streams coalesce into a network, the reticulations of which change their position at short intervals, as in the hairs of *Glaucium luteum*; whilst, lastly, the current may flow in a sluggish uniformly moving sheet or layer. Where several distinct currents exist in one cell, they are all found to have one common point of departure and return, namely the "nucleus" (B, a); from which it seems fairly to be inferred, that this body is the centre of the vital activity of the cell.¹ Mr. Wenham states that in all cases in which the sap-motion is seen in the hairs of a plant, the cells of the cuticle also display it, provided that their walls are not so opaque or so strongly marked, as to prevent the rotation from


being distinguished.² The cuticle may be most readily torn off

¹ The above statement is called in question by Mr. Wenham, who affirms that "whenever he has observed such a 'nucleus,' it has either been formed by an accidental conglomeration of some of the cell-contents, or by morbid conditions." The Author is satisfied, however, from the constancy with which the "nucleus" is the centre of the diverging lines of protoplasm, in those cells which have several streams radiating from one point, that it can neither be an accidental nor a morbid conglomeration.

² Op. cit. vol. iv. p. 44.

from the stalk or the midrib of the leaf; and must then be examined as speedily as possible, since it loses its vitality, when thus detached, much sooner than do the hairs. Even where no obvious movement of particles is to be seen, the existence of a rotation may be concluded from the peculiar arrangement of the molecules of the protoplasm, which are remarkable for their high refractive power, and which, when arranged in a "moving train," appear as bright lines across the cell; and these lines, on being carefully watched, are seen to alter their relative positions. The leaf of the common *Plantago* (plantain or dock) furnishes an excellent example of "rotation;" the movement being distinguishable at the same time, both in the cells and in the hairs of the cuticle torn from its stalk or midrib. It is a curious circumstance, that when a plant (such as the *Anacharis*) which exhibits the "rotation," is kept in a cold dark place for one or two days, not only is the movement suspended, but the moving particles collect together in little heaps; these being again broken up by the separate motion of their particles, when the stimulus of light and warmth occasions a renewal of the circulatory action. It is well to collect the specimens about mid-day, that being the time when the rotation is most active; and the movement is usually quickened by artificial warmth, which, indeed, is a necessary condition in some instances to its being seen at all. The most convenient method of applying this warmth, while the object is on the stage of the microscope, is to blow a stream of air upon the thin glass cover, through a glass or metal tube previously heated in a spirit lamp.

227. The walls of the cells of Plants are frequently thickened by internal deposits, which may present very different appearances according to the manner in which they are arranged. In its simplest condition, such a deposit forms a thin uniform layer over the whole internal surface of the cellulose wall (probably on the outside of the primordial utricle), scarcely detracting at all from its transparency, and chiefly distinguishable by the "dotted" appearance which the membrane then presents (Fig. 150, A). These dots, however, are not pores, as their aspect might naturally suggest; but are merely points at which the deposit is wanting, so that the original cell-wall there remains unthickened. When the cellular tissue is required to possess unusual firmness, a deposit of *sclerogen* (a substance which, when separated from the resinous and other matters that are



Tissue of the Testa of the seed-coat of *Narantia*:—A, as seen in section; B, as seen on the surface.

commonly associated with it, is found to be allied in chemical composition to cellulose) is formed in successive layers, one within another (Fig. 154, A), which present themselves as concentric rings when the cells containing them are cut through; and these layers are sometimes so thick and numerous, as almost to obliterate the original cavity of the cell. By a continuance of the same arrangement as that which shows itself in the single layer of the dotted cell,—each deposit being deficient at certain points, and these points corresponding with each other in the successive layers,—a series of passages is left, by which the cavity of the cell is extended at some points to its membranous wall; and it commonly happens that the points at which the deposit is wanting on the walls of two contiguous cells, are coincident, so that the membranous partition is the only obstacle to the communication between their cavities (Figs. 154–156). It is of such tissue that the “stones” of fruit, the gritty substance

FIG. 155.

Section of *Cherry Stone*, cutting the cells transversely.

FIG. 156.

Section of *Coquilla Nut*, in the direction of the long diameters of the cells.

which surrounds the seeds and forms little hard points in the fleshy substance of the pear, the shell of the cocoa-nut, and the albumen of the seed of *Phytelphas* (known as “vegetable ivory”), are made up; and we see the use of this very curious arrangement, in permitting the cells, even after they have attained a considerable degree of consolidation, still to remain permeable to the fluid required for the nutrition of the parts which such tissue encloses and protects.


228. The deposit sometimes assumes, however, the form of definite *fibres*, which lie coiled up in the interior of cells, so as to form a single, double, or even a triple or quadruple spire

(Fig. 157). Such "spiral cells" are found most abundantly in the leaves of certain Orchideous plants, immediately beneath the cuticle, where they are brought into view by ver-

FIG. 157.

Spiral cells of leaf of *Oncidium*.

FIG. 158.

Spiral fibres of Seed-coat of *Collomia*.

tical sections, and they may be obtained in an isolated state, by macerating the leaf and peeling off the cuticle, so as to expose the layer beneath, which is then easily separated into its components. In an Orchideous plant named *Saccolabium guttatum*, the spiral cells are unusually long, and have spires winding in opposite directions; so that, by their mutual intersection, a series of diamond-shaped markings is produced. Spiral cells are often found upon the surface of the *testa* or outer coat of seeds; and in the *Collomia grandiflora*, the *Salvia verbenaca* (wild clary), and some other plants, the membrane of these cells is so weak, and the elasticity of their fibres so great, that when the membrane is softened by the action of water, the fibres suddenly uncoil and elongate themselves (Fig. 158), springing out, as it were, from the surface of the seed, to which they give a peculiar flocculent appearance. This very curious phenomenon, which greatly preplexes those who are ignorant of its real nature, may be best observed in the following manner:—A very thin transverse slice of the seed should first be cut, and laid upon the lower glass of the aquatic box; the cover should then be pressed down, and the box placed upon the stage, so that the body of the microscope may be exactly "focussed" to the object, the power employed being the 1 inch, 2-3ds inch, or the $\frac{1}{2}$ inch objective. The cover of the aquatic box being then removed, a small drop of water should be placed on that part of its internal surface, with which the slice of the seed had been in contact; and the cover being replaced, the object should be immediately looked at. It is important that the slice of the seed should be very thin, for two reasons; first that the view of the spires may not be confused by their aggregation in too great numbers; and second, that the drop of water should be held in its place by


capillary attraction, instead of running down and leaving the object, as it will do if the glasses be too widely separated.

229. In some part or other of most Plants, we meet with cells containing granules of *Starch*. These granules are sometimes minute and very numerous, and are so closely packed together as to fill the cavity (Fig. 159); in other instances they are of

FIG. 159.

Cells of *Peany*, filled with Starch.

FIG. 160.

Granules of *Starch*, as seen under polarized light.

much larger dimensions, so that only a small number of them can be included in any one cell; while in other cases, again, they are both few and minute, so that they form but a small proportion of the cell-contents. Their nature is at once detected by the addition of a solution of iodine, which gives them a beautiful blue color. Each granule exhibits a peculiar spot termed the *hilum*, which marks the point at which, in its early state, it is attached to the cell-wall; and it also presents, when highly magnified, a set of circular lines, which are for the most part concentric (or nearly so) with the hilum. When viewed by polarized light (§ 63), each grain exhibits a beautiful dark cross, the point of intersection being at the hilum (Fig. 160). Regarding the internal structure of the starch grain, opinions are very much divided; for whilst some affirm the concentric lines to indicate the existence of a number of concentric lamellæ, one enclosing another, others consider that they are due to the peculiar plaiting or involution of a single vesicular wall;¹ and among those who consider it to be concentrically lamellated, some hold that each lamella is formed *outside* or *upon* that which preceded it, while others consider that each is formed *inside* or *within* its predecessor. The centre of the granule is often occupied by starchy matter in an unconsolidated state; and it is the appearance arising from its different refractive power, that has caused some observers to describe the starch-grain as possessing

¹ The first of these opinions is the one which had been generally received until recently, when Mr. G. Busk supported the latter by new observations made upon the unfolding of the starch-granule by dilute sulphuric acid; since when, Prof. Allman, after repeating Mr. Busk's observations, has been led to affirm them to be fallacious, and to revert to the first of the above-mentioned doctrines. See Mr. Busk's memoir in "Trans. of Microsc. Soc." ser. 2, vol. i, p. 58; and that of Prof. Allman in "Quart. Journ. of Microsc. Sci." vol. ii, p. 163.

a nucleus. Although the dimensions of the starch-grains produced by any one species of Plant are by no means constant, yet there is a certain average for each, from which none of them depart very widely; and by reference to this average, the starch-grains of different plants that yield this product in abundance, may be microscopically distinguished from one another,—a circumstance of considerable importance in commerce. The largest starch-grains in common use are those of the plant (a species of *Canna*) known as *Tous les mois*; the average diameter of those of the *Potato* is about the same as the diameter of the smallest of the “*tous les mois*”; and the size of the ordinary starch-grains of *Wheat* and of *Sago* is about the same as that of the smallest grains of potato-starch; whilst the granules of *Rice*-starch are so very minute, as to be at once distinguished from any of the preceding.

230. Deposits of mineral matter in a crystalline condition, known as *Raphides*, are not unfrequently found in the cells of Plants; where they are at once brought into view by the use of polarized light. Their designation (derived from *ῥαψίς*, a needle) is very appropriate to one of the most common states in which these bodies present themselves, that, namely, of bundles of needle-like crystals, lying side by side in the cavity of the cell; such bundles are well seen in the cells lying immediately beneath the cuticle of the bulb of the medicinal Squill. It does not apply, however, to other forms which are scarcely less abundant; thus, instead of bundles of minute needles, single large crystals, octohedral or prismatic, are frequently met with; and the prismatic crystals are often aggregated in beautiful stellate groups. One of the most common materials of raphides, is oxalate of lime, which is generally found in the stellate form; and no plant yields these stellate raphides so abundantly as the common *Rhubarb*, the best specimens of the dry medicinal root containing as much as 35 per cent. of them. In the cuticle of the bulb of the *Onion*, the same material occurs under the octohedral or the prismatic form. In other instances, the calcareous base is combined with tartaric, citric, or malic acid; and the acicular raphides are said to consist usually of phosphate of lime. Some raphides are as long as 1-40th of an inch, while others measure no more than 1-1000th. They occur in all parts of plants,—the wood, pith, bark, root, leaves, stipules, sepals, petals, fruit, and even in the pollen. They are always situated in cells, and not, as some have stated, in the intercellular passages; the cell-membrane, however, is often so much thinned away, as to be scarcely distinguishable. Certain plants of the *Cactus* tribe, when aged, have their tissues so loaded with raphides as to become quite brittle; so that when some large specimens of *C. senilis*, said to be a thousand years old, were sent to Kew Gardens, from South America, some years since, it was found necessary for their preservation during transport, to pack them in cotton, like jewelry.


It is not yet known what office the raphides fulfil in the economy of the Plant; or whether they are to be considered in any other light, than as non-essential results of the vegetative processes. For as all these processes require the introduction of mineral bases from the soil, and themselves produce organic acids in the substance of the plant, it may be surmised that the accidental union of the components will occasion the formation of raphides wherever such union may occur; and this view is supported by the fact, that the late Mr. E. Quekett succeeded in artificially producing raphides within the cells of "rice paper" (§ 223), by first filling these with lime water by means of the air-pump, and then placing the paper in weak solutions of phosphoric and oxalic acids. The artificial raphides of phosphate of lime were rhombohedral; while those of oxalate of lime were stellate, exactly resembling the natural raphides of the Rhubarb.¹

231. A large proportion of the denser parts of the fabric of the higher Plants, is made up of the substance which is known as *Ligneous Tissue*, or *Woody Fibre*. This, however, can only be regarded as a very simple variety of the "cellular;" for it is composed of peculiarly elongated cells (Fig. 171, *a a*), usually pointed at their two extremities so as to become spindle-shaped, whose walls have a special tendency to undergo consolidation by the internal deposit of sclerogen. It is obvious that a tissue consisting of elongated cells, adherent together by their entire length, and strengthened by internal deposit, must possess much greater tenacity than any tissue in which the cells depart but little from the primitive spherical form; and we accordingly find *Woody Fibre* introduced, wherever it is requisite that the fabric should possess not merely density, but the power of resistance to tension. In the higher classes of the Vegetable kingdom, it constitutes the chief part of the stem and branches, where these have a firm and durable character; and even in more temporary structures, such as the herbaceous stems of annual plants, and the leaves and flowers of almost every tribe, this tissue forms a more or less important constituent, being especially found in the neighborhood of the spiral vessels and ducts, to which it affords protection and support. Hence the bundles or fasciculi composed of these elements, which form the skeletons of leaves, and which give "stringiness" to various esculent vegetable substances, are commonly known under the name of *fibro-vascular tissue*. In their young and unconsolidated state, the ligneous cells seem to conduct fluids with great facility in the direction of their length; and in the *Coniferous* tribe whose stems and branches are desti-

¹ The materials of the above paragraph are derived from the excellent section on this subject in Prof. Quekett's "Lectures on Histology." Besides the vegetables therein named as affording good illustrations of different kinds of Raphides, may be mentioned, the parenchyma of the leaf of *Agave*, *Aloe*, *Cycas*, *Encephalartos*, &c., the cuticle of the bulb of the *Hyacinth*, *Tulip*, and *Garlic* (and probably of other bulbs), the bark of the *Apple*, *Cascarilla*, *Cinchona*, *Lime*, *Lorust*, and many other trees, the pith of *Eleagnus*, and the testa of the seeds of *Anagallis* and the *Elm*.

tute of vessels, they afford the sole channel for the ascent of the sap. But after their walls have become thickened by internal deposit, they are no longer subservient to this function; nor, indeed, do they then appear to fulfil any other purpose in the vegetable economy, than that of affording mechanical support. It is this which constitutes the difference between the *alburnum* or "sap-wood," and the *duramen* or "heart-wood," of Exogenous stems (§ 238). A peculiar set of markings, seen on the woody fibres of the *Coniferae*, and of some other tribes, is represented in Fig. 161; in each of these spots, the inner circle appears to mark a deficiency of the lining deposit, as in the porous cells of other plants; whilst the outer circle indicates the boundary of a lenticular cavity, which intervenes between the adjacent cells at this point, and which contains a small globular body that may be sometimes detached. Of the purpose of these minute bodies interposed between the wood-cells, nothing is known; there can be no doubt, however, from the definiteness and constancy of their arrangement, that they fulfil some important object in the economy of the plants in which they occur; and there are varieties in this arrangement so characteristic of different tribes, that it is sometimes possible to determine, by the microscopic inspection of a minute fragment, even of a fossil wood, the tribe to which it belonged. The woody fibre thus marked, is often designated as "glandular."

FIG. 161.



Section of *Coniferous Wood* in the direction of the fibres, showing their "glandular" dots; a, a, a, medullary rays crossing the fibres.


232. All the more perfect forms of Phanerogamia contain, in some part of their fabric, the peculiar structures which are known as *Spiral Vessels*.¹ These have the elongated shape of woody fibres; but the internal deposit, as in the "spiral cells" (§ 228), takes the form of a spiral fibre winding from end to end, remaining distinct from the cell-wall, and retaining its elasticity; this fibre may be single, double, or even quadruple,—this last character presenting itself in the very large elongated fibre-cells of the *Nepenthes* (Chinese pitcher-plant). These cells are especially found in the delicate membrane ("medullary sheath") surrounding the pith of Exogens, and in the midst of the woody bundles occurring in the stem of Endogens; thence they proceed in each case to the leaf-stalks, through which they are distributed to the leaves. By careful dissection under the microscope, they may be separated entire; but their structure may be more easily dis-

¹ So long, however, as they retain their original cellular character, and do not coalesce with each other, these fusiform spiral cells cannot be regarded as having any more claim to the designation of *vessels*, than have the elongated cells of the ligneous tissue.

played, by cutting *round*, but not *through*, the leaf-stalk of the Strawberry, Geranium, &c., and then drawing the parts asunder. The membrane composing the tubes of the vessels will thus be broken across; but the fibres within, being elastic, will be drawn out and unrolled. Spiral vessels are sometimes found to convey liquid, whilst in other cases they contain air only; the conditions of this difference are not yet certainly known.

233. Although fluid generally finds its way with tolerable facility through the various forms of Cellular tissue, especially in the direction of the greatest length of its cells, a more direct means of connection between distant parts is required for an active circulation. This is afforded by what has been termed *Vasiform tissue*, which consists merely of cells laid end to end, the partitions between them being more or less obliterated, so that a continuous *Duct* is formed. The origin of these ducts in cells is occasionally very evident, both in the contraction of their calibre at regular intervals, and in the persistence of the remains of their partitions (Fig. 175, *b b*); but in most cases it can only be ascertained by studying the history of their development, neither of these indications being traceable. The component cells appear to have been sometimes simply membranous, but more commonly to have possessed the fibrous type (§ 228).

FIG. 162.



Longitudinal section of stem of Italian Reed:—*a*, cells of the pith; *b*, fibro-vascular bundle, containing, 1, annular duct; 2, spiral duct; 3, dotted duct, with woody fibre; *c*, cells of the integument.

Some of the ducts formed from the latter (Fig. 162, *2*) are so like continuous spiral vessels, as to be scarcely distinguishable from them, save in the want of elasticity in the spiral fibre, which causes it to break when the attempt is made to draw it out. This would seem to have taken place, in some instances, from the natural elongation of the cells by growth; the fibre being broken up into rings, which sometimes lie close together, but more commonly at considerable intervals; such a duct is said to be *annular* (Fig. 162, *1*). Intermediate forms between the spiral and annular ducts, which show the derivation of the latter from the former, are very frequently to be met with. The spires are sometimes broken

up still more completely, and the fragments of the fibre extend in various directions, so as to meet and form an irregular net-

work lining the duct, which is then said to be *reticulated*. The continuation of the deposit, however, gradually contracts the meshes, and leaves the walls of the duct marked only by pores like those of porous cells (§ 227); and canals upon this plan, commonly designated as *dotted* ducts, are among the most common forms of vasiform tissue, especially in parts of most solid structure and least rapid growth (Fig. 162, 3). The "scalariform" ducts of Ferns (§ 218) are for the most part of the spiral type; but spiral ducts are frequently to be met with also in the rapidly growing leaf-stalks of Flowering plants, such as the Rhubarb. Not unfrequently, however, we find all forms of ducts in the same bundle, as seen in Fig. 162. The size of these ducts is occasionally so great, as to enable their openings to be distinguished by the unaided eye. They are usually largest in stems, whose size is small in proportion to the surface of leaves which they support, such as the common Cane, or the Vine; and generally speaking they are larger in woods of dense texture, such as Oak or Mahogany, than in those of which the fibres, being softer, can themselves be subservient to the conveyance of fluid. They are entirely absent in the Coniferæ.

234. The Vegetable Tissues, whose principal forms have been now described, but among which an immense variety of detail is found, may be either studied as they present themselves in thin *sections* of the various parts of the plant under examination, or in the isolated condition in which they are obtained by *dissection*. The former process is the most easy, and yields a large amount of information; but still it cannot be considered that the characters of any tissue have been properly determined, until it has been dissected out. Sections of some of the hardest vegetable substances, such as "vegetable ivory," the "stones" of fruit, the "shell" of the cocoa-nut, &c. (§ 227), can scarcely be obtained except by slicing and grinding (§ 108); and these may be mounted either in Canada balsam or in weak spirit. In cases, however, in which the tissues are of only moderate firmness, the section may be readily and most effectually made with the "Section-Instrument" (§ 107); and there are few parts of the Vegetable fabric which may not be advantageously examined by this means, any very soft or thin portions being placed in it between two pieces of cork. In certain cases, however, in which even this compression would be injurious, the sections must be made with a sharp knife, the substance being laid upon a slip of glass. In dissecting the Vegetable tissues, scarcely any other instrument will be found really necessary, than a pair of needles (in handles), one of them ground to a cutting edge. The adhesion between the component cells, fibres, &c., is often sufficiently weakened by a few hours' maceration, to allow of their readily coming apart, when they are torn asunder by the needle points beneath the simple lens of a dissecting microscope. But if this should not prove to be the case, it is desirable to employ some

other method, for the sake of facilitating their isolation. None is so effectual as the boiling of a thin slice of the substance under examination, either in dilute nitric acid, or in a mixture of nitric acid and chlorate of potass. This last method (which was devised by Schultz) is the most rapid and effectual, requiring only a few minutes for its performance; but as oxygen is liberated with such freedom as to give an almost explosive character to the mixture, it should be put in practice with extreme caution. After being thus treated, the tissue should be boiled in alcohol, and then in water; and it will then be found very easy to tear apart the individual cells, ducts, &c., of which it may be composed. These may be preserved by mounting in weak spirit.

235. *Structure of the Stem and Root.*—It is in the stems and roots of Plants, that we find the greatest variety of tissues in combination, and the most regular plans of structure; and sections of these, viewed under a low magnifying power, are objects of peculiar beauty, independently of the scientific information which they afford. The Axis (under which term is included the Stem with its branches, and the Root with its ramifications) always has for the basis of its structure a dense cellular parenchyma; though this, in the advanced stage of development, may constitute but a small proportion of it. In the midst of this parenchyma we generally find fibro-vascular bundles; that is, fasciculi of woody fibre, with ducts of various kinds, and (very commonly) spiral vessels. It is in the mode of arrangement of these bundles, that the fundamental difference exists between the stems that are commonly designated as *Endogenous*, and those which are (more appropriately) termed *Exogenous*; for in the former, the bundles are dispersed throughout the whole diameter of the axis, without any peculiar plan, the intervals between them being filled up by cellular parenchyma; whilst in the latter they are arranged side by side, in such a manner as to form a hollow cylinder of *wood*, which includes within it that portion of the cellular substance which is known as *pith*, whilst it is itself enclosed in an envelope of the same substance, that forms the *bark*. These two plans of axis formation,—respectively characteristic of those two great groups into which the Phanerogamia are subdivided, namely the *Monocotyledonous*, and the *Dicotyledonous*,—will now be more particularly described.

236. When a transverse section (Fig. 163) of a *Monocotyledonous* stem is examined microscopically, it is found to exhibit a number of fibro-vascular bundles, disposed without any regularity in the midst of the mass of cellular tissue, which forms (as it were) the matrix or basis of the fabric. Each bundle contains two, three, or more large ducts, which are at once distinguished by the size of their openings; and these are surrounded by woody fibre, and spiral vessels, the transverse diameter of which is so extremely small, that the portion of the bundles which they form is at once distinguished in transverse section, by the *close-*

ness of its texture (Fig. 164). The bundles are least numerous in the centre of the stem, and become gradually more approximated towards the circumference: but it frequently happens that the portion of the area in which they are most compactly

FIG. 163.



Transverse section of stem of young *Palm*.

FIG. 164.

Portion of transverse section of stem of *Waghie Cane*.

arranged, is not absolutely at its exterior, this portion being itself surrounded by an investment composed of cellular tissue only; and sometimes we find the central portion, also, completely destitute of fibro-vascular bundles; so that a sort of indication of the distinction between pith, wood, and bark is here presented. This distinction, however, is very imperfect; for we do not find either the central or the peripheral portions ever separable, like pith and bark, from the intermediate woody layer. In its young state, the centre of the stem is always filled up with cells; but these not unfrequently disappear after a time, except at the nodes, leaving the stem hollow, as we see in the whole tribe of Grasses. When a vertical section is made of a woody stem (as that of a Palm) of sufficient length to trace the whole extent of the fibro-vascular bundles, it is found that whilst they pass at their upper extremity into the leaves, they pass at the lower end, also, towards the surface of the stem, and assist, by their interlacement with the outer bundles, in forming that extremely tough investment, which the lower ends of these stems present. The fibro-vascular bundles once formed, receive no further additions; and the augmentation of the stem in diameter depends upon the development of new woody bundles, in continuity with the leaves which are successively evolved at its summit. It was formerly supposed that these successively

formed bundles descend in the interior of the stem through its entire length, until they reach the roots; and as the successive development of leaves involves a successive development of new bundles, the stem was imagined to be continually receiving additions to its interior, whence the term *Endogenous* was given to this type of stem-structure. From the fact just stated, however, regarding the course of the fibro-vascular bundles, it is obvious that such a doctrine cannot be any longer admitted; for those which are most recently formed only pass into the centre of the stem during the higher part of their course, and usually make their way again to its exterior at no great distance below; and thus the lower and older portions of a Palm-stem really do receive very little augmentation in diameter, while a rapid elongation is taking place at its summit. In fact, the dense unyielding nature of the fabric, which is formed by the interlacement of the fibro-vascular bundles at or near the surface of the trunk, would prevent any such augmentation by expanding pressure from within.

237. In the stems of *Dicotyledonous* Phanerogamia, we find a method of arrangement of the several parts, which must be re-




Diagram of the first formation of an Exogenous stem; a, pith; b b, bark; c c, plates of cellular tissue (medullary rays) left between the woody bundles, d d.


garded as the highest form of the development of the axis, being that in which the greatest *differentiation* exists. A distinct division is always seen in a transverse section, between the three concentric areas of the *pith*, the *wood*, and the *bark*; the first (*a*) being central and the last (*b*) peripheral, and these having the wood interposed between them, its circle being made up of wedge-shaped bundles (*d, d*), kept apart by the bands (*c, c*) that pass between the pith and the bark. The *Pith* (Fig. 165, *a*) is almost invariably composed of cellular tissue only, which usually presents (in transverse section) a hexagonal areola-

tion. When newly formed, it has a greenish hue, and its cells are filled with fluid; but it gradually dries up and loses its color; and not unfrequently its component cells are torn apart by the rapid growth of their envelope, so that irregular cavities are found in it; or, if the stem should increase with extreme rapidity, it becomes hollow, the pith being reduced to fragments which are found adhering to its interior wall. The pith is immediately surrounded by a delicate membrane, consisting almost entirely of spiral vessels, and termed the "medullary sheath."

238. The *Woody* portion of the stem (Fig. 165, *b, b*) is made up of woody fibres, usually with the addition of ducts of various kinds; these, however, are absent in one large group, the *Coni-*

feræ or Fir tribe with its allies (Figs. 166, 170–172), in which the woody fibres are of unusually large diameter, and have the peculiar “glandular” markings already described (§ 231). In any stem or branch of more than one year’s growth, the woody struc-


FIG. 165.



Transverse section of young stem of *Clematis*:—*a*, pith; *b, b, b*, woody bundles; *c, c, c*, medullary rays.

ture presents a more or less distinct appearance of division into concentric rings, the number of which varies with the age of the tree (Fig. 167). The composition of the several rings, which are the sections of so many cylindrical layers, is uniformly the same, however different their thickness; the arrangement of the two principal elements, however,—namely the woody fibre and the ducts,—varies in different species; the ducts being sometimes

FIG. 166.



Portion of transverse section of stem of *Cedar*:—*a*, pith; *b, b*, wood; *c*, bark.

almost uniformly diffused through the whole layer, but in other instances being confined to its inner part; while in other cases, again, they are dispersed with a certain regular irregularity (if such an expression may be allowed), so as to give a curiously figured appearance to the transverse section (Figs. 167, 168). The general fact, however, is, that the ducts predominate towards the inner side of the ring (which is the part of it first formed), and

that the outer portion of each layer is almost exclusively composed of woody tissue; such an arrangement is seen in Fig. 165. This alternation of ducts and woody fibre frequently serves to mark the succession of layers, when, as is not uncommon, there is no very distinct line of separation between them.


FIG. 167.



Transverse section of stem of *Buckthorn* (*Rhamnus*).

The number of layers is usually considered to correspond with that of the years during which the stem or branch has been growing; and this is, no doubt, generally true in regard to the trees of temperate climates. There appears strong reason to believe, however, that such is not the universal rule;

FIG. 168.



Portion of the preceding figure, more highly magnified.

and that we should be more correct in stating that each layer indicates an "epoch of vegetation;" which, in temperate climates, is usually (but not invariably) a year, but which is commonly much less in the case of trees flourishing in tropical regions. For example, we not unfrequently meet with stems, in which the place of a layer of the ordinary breadth is occupied by two narrow layers;

the line of demarcation between them having apparently been formed by a temporary interruption to the process of growth, in the middle of the period through which the formation of wood extends. Such an interruption might occur from heat

FIG. 169.



a b c


Portion of transverse section of stem of *Hazel*, showing, in the portion a, b, c, six narrow layers of wood.

and drought, in a tree that flourishes best in a cold damp atmosphere, or from a fall of temperature in a tree that requires heat; and in a variable season, it might recur several times. Something of this kind would appear to have been the cause of the

peculiar appearance presented by a section of Hazel-stem (in the Author's possession), of which a portion is represented in Fig. 169; for between two layers of the ordinary thickness, there intervenes a band whose breadth is altogether less than that of either of them, and which is yet composed of no fewer than six layers, four of them (*c*) being very narrow, and each of the other two being about as wide as these four together. The inner layers of wood are the oldest, and the most solidified by matters deposited within their component cells and vessels; hence they are spoken of collectively under the designation *duramen* or "heart-wood." On the other hand, it is through the cells and ducts of the outer and newer layers, that the sap rises from the roots towards the leaves; and these are consequently designated as *alburnum* or "sap-wood." The line of demarcation between the two is sometimes very distinct, as in *Lignum-vitæ* and *Cocos* wood; and as a new layer is added every year to the exterior of the alburnum, an additional layer of the innermost part of the alburnum is every year consolidated by internal deposit, and is thus added to the exterior of the *duramen*. More generally, however, this consolidation is gradually effected, and the alburnum and *duramen* are not separated by any abrupt line of division.

239. The *Medullary Rays* which cross the successive rings of wood connecting the cellular substance of the pith with that of

FIG. 170.



Portion of transverse section of large stem of *Coniferous Wood* (fossil), showing part of two annual layers, divided at *a, a*, and traversed by very thin but numerous *Medullary Rays*.

of the bark, and dividing each ring of wood into wedge-shaped segments, are thin plates of cellular tissue (Fig. 165, *c, c*), not usually extending to any great depth in the vertical direction. It is not often, however, that their character can be so clearly seen in a transverse section, as it is in that just referred to; for they are usually compressed so closely, as to appear darker than the wedges of woody tissue between which they intervene (Figs. 166–169); and their real nature is best understood by a

comparison of *longitudinal* sections made in two different directions,—namely, *radial* and *tangential*,—with the transverse. Three such sections of a fossil *Coniferous wood* in the Author's possession are shown in Figs. 170–172. The stem was of such large size, that, in so small a part of the area of its transverse section as is represented in Fig. 170, the medullary rays seem to run parallel to each other, instead of radiating from a common centre. They are very narrow; but are so closely set together, that only two or three rows of woody fibres (no ducts being here

present) intervene between any pair of them. In the longitudinal section taken in a radial direction (Fig. 171), and conse-

FIG. 171.



FIG. 172.




Fig. 171. Portion of vertical section of the same wood, taken in a radial direction, showing the glandular woody fibres, without ducts, crossed by the Medullary Rays, *a, a*.

Fig. 172. Portion of vertical section of the same wood, taken in a tangential direction, so as to cut across the Medullary Rays.

quently passing in the same course with the medullary rays, these are seen as thin plates (*a, a, a*) made up of superposed cells

FIG. 173.

Vertical section of *Mahogany*.

very much elongated, and crossing, in a horizontal direction, the glandular woody fibres which lie parallel to one another vertically. And in the tangential section (Fig. 172), which passes in a direction at right angles to that of the medullary rays, and therefore cuts them across, we see that each of the plates thus formed has a very limited depth from above downwards, and is composed of no more than one thickness of horizontal cells. A section of the stem of *Mahogany*, taken in the same direction as the last (Fig. 173), gives a very good view of the cut ends of the medullary rays, as they pass between the woody fibres; and they are seen to be here of somewhat greater thickness, being composed of two or three rows of cells, arranged side by side. In another fossil wood, whose transverse section is shown in Fig. 174, and its tangential section in Fig. 175, the medullary rays are seen to occupy a much larger part of the substance of the stem; being seen in the transverse section as broad bands (*a a, a a*), intervening between the closely set woody fibres, among which the large ducts are scattered; whilst in the tangential, they are observed to be not

only deeper than the preceding from above downwards, but also to have a much greater thickness. This section also gives an excellent view of the ducts *b b*, *b b*, which are here plainly seen

FIG. 174.



FIG. 175.



Fig. 174. Portion of transverse section of Fossil Wood, showing the Medullary Rays, *a a, a a, a a*, running nearly parallel to each other, and the openings of the large Ducts in the midst of the woody fibres.

Fig. 175. Portion of vertical (tangential) section of the same Wood, showing the woody fibres separated by the Medullary Rays, and by the large Ducts, *b b, b b*.

to be formed by the coalescence of large cylindrical cells, lying end to end. In another fossil wood in the Author's possession, the medullary rays constitute a still larger proportion of the

FIG. 176.



FIG. 177.



Fig. 176. Portion of transverse section of a Fossil Wood, remarkable for the very large size of the Medullary Rays, *b, b*, which separate the woody plates, *a, a*.

Fig. 177. Vertical (tangential) section of the same wood, showing, *a, a*, the woody bundles, and *b, b*, the Medullary Rays.

stem; for in the transverse section (Fig. 176), they are seen as very broad bands (*b, b*), alternating with plates of woody struc-

ture (*a, a*), whose thickness is often less than their own; whilst in the tangential section (Fig. 177), the cut extremities of the medullary rays occupy a very large part of the area, having apparently determined the sinuous course of the woody fibres; instead of looking, as in Fig. 172, as if they had forced their way between the woody fibres, which there hold a nearly straight and parallel course on either side of them.

240. The *Bark* may be usually found to consist of three principal layers; the external, or *epiphlæum*, also termed the *suberous* (or corky) layer; the middle, or *mesophlæum*, also termed the "cellular envelope;" and the internal, or *endophlæum*, which is more commonly known as the *liber*. The two outer layers are entirely cellular; and are chiefly distinguished by the form, size, and direction of their cells. The *epiphlæum* is generally composed of one or more layers of colorless or brownish cells, which usually present a cubical or tabular form, and are arranged with their long diameters in the horizontal direction: it is this which, when developed to an unusual thickness, forms *Cork*, a substance which is by no means the product of one kind of tree exclusively, but which exists in greater or less abundance in the bark of every exogenous stem. The *mesophlæum* consists of cells, usually of green color, prismatic in their form, and disposed with their long diameters parallel to the axis; it is more loosely arranged than the preceding, and contains "intercellular passages," which often form a network of canals, that have been termed "laticiferous vessels;" and although usually less developed than the suberous layer, it sometimes constitutes the chief thickness of the bark. The *liber* or inner bark, on the other hand, usually contains woody fibre in addition to the cellular tissue and laticiferous vessels of the preceding; and thus approaches more nearly in its character to the woody layers, with which it is in close proximity on its inner surface. The *liber* may generally be found to be made up of a succession of thin layers, equalling in number those of the wood, the innermost being the last formed; but no such succession can be distinctly traced in the cellular envelope, or in the suberous layer; although it is certain that they, too, augment in thickness by additions to their interior, whilst their external portions are frequently thrown off in the form of thickish plates, or detach themselves in smaller and thinner flakes. The bark is always separated from the wood by the *cambium layer*, which is the part wherein all new growth takes place: this seems to consist of mucilaginous semifluid matter; but it is really made up of cells of a very delicate texture, which gradually undergo transformations, whereby they are for the most part converted into woody tissue, ducts, spiral vessels, &c. These materials are so arranged, as to augment the fibro-vascular bundles of the wood on their external surface; thus forming a new layer of alburnum, which *encloses* all those that preceded it; whilst they also form a new layer of *liber*, on

the *interior* of all those which preceded it ; they also extend the medullary rays, which still maintain a continuous connection between the pith and the bark ; but a portion remains unconverted, so as always to keep apart the liber and alburnum. This type of stem structure is termed *Exogenous* ; a designation which applies very correctly to the mode of increase of the woody layers, although (as we have just seen) the liber is formed upon a truly endogenous plan.

241. Numerous departures from the normal type are found in particular tribes of Exogens. Thus in some, the wood is not marked by concentric circles, their growth not being interrupted by any seasonal change. In other cases, again, each woody zone is separated from the next, by the interposition of a thick layer of cellular substance. Sometimes wood is formed in the bark (as in *Calycanthus*), so that several woody columns are produced, which are quite independent of the principal woody axis, but cluster around it. Occasionally the woody stem is divided into distinct segments, by the peculiar thickness of certain of the medullary rays ; and in the stem of which Fig. 178 represents a transverse section, these cellular plates form four large segments, disposed in the manner of a Maltese cross, and alternating with the four woody segments, which they equal in size.

242. In its first-developed state, the Exogenous stem consists, like the so-called endogenous, of cellular tissue only ; but

FIG. 178.



FIG. 179.



Fig. 178. Transverse section of the stem of a Climbing-plant (*Aristolochia?*) from New Zealand.
Fig. 179. Portion of transverse section of *Burdock* (*Arctium*), showing one of the fibro-vascular bundles, that lies beneath the cellular integument.

after the leaves have been actively performing their functions for a short time, we find a circle of fibro-vascular bundles, as represented in the diagram, p. 378, interposed between the central (or medullary) and the peripheral (or cortical) portions of the cellular matrix ; these fibro-vascular bundles being themselves

separated from each other by plates of cellular tissue, which still remain to connect the central and the peripheral portions of the matrix. This first stage in the formation of the Exogenous axis, in which its principal parts—the pith, wood, bark, and medullary rays—are marked out, is seen even in the stems of herbaceous plants, which are destined to die down at the end of the season (Fig. 179); and sections of these, which are very easily prepared, are most interesting Microscopic objects. In such stems, the difference between the “Endogenous” and the “Exogenous” types is manifested in little else than the disposition of the fibro-vascular layers; which are scattered through nearly the whole of the cellular matrix (although most abundant towards its exterior), in the former case; but are limited to a circle within the peripheral portion of the cellular tissue, in the latter. It is in the further development which takes place during succeeding years in the woody stems of perennial Exogens, that those characters are displayed, which separate them most completely from the Ferns and their allies, whose stems contain a cylindrical layer of fibro-vascular bundles, as well as from (so-called) Endogens. For whilst the fibro-vascular layers of the latter, when once formed, undergo no further increase, those of Exogenous stems are progressively augmented by the metamorphosis of the cambium layer; so that each of the bundles which once lay as a mere series of parallel cords beneath the cellular investment of a first year’s stem, may become in time the small end of a wedge-shaped mass of wood, extending continuously from the centre to the exterior of a trunk of several feet in diameter, and becoming progressively thicker as it passes outwards. The fibro-vascular bundles of Exogens are therefore spoken of as “indefinite;” whilst those of Exogens and Acrogens (Ferns, &c.) are said to be “definite” or “closed.”

243. The structure of the *Roots* of Endogens and Exogens is essentially the same in plan with that of their respective stems. Generally speaking, however, the roots of Exogens have no pith, although they have medullary rays; and the succession of distinct rings is less apparent in them, than it is in the stems from which they diverge. In the delicate radical filaments which proceed from the larger root-fibres, a central bundle of vessels will be seen, enveloped in a sheath of cellular substance; and this investment also covers in the end of the fibril, which is usually somewhat dilated, and composed of peculiarly succulent tissue, forming what is termed the *spongiole*. The structure of the radical filaments may be well studied in the common *Duckweed*, every floating leaf of which has a single fibril hanging down from its lower surface.

244. The structure of Stems and Roots cannot be thoroughly examined in any other way than by making sections in different directions with the Section-instrument. The general directions already given (§ 107) leave little to be added respecting this

special class of objects; the chief points to be attended to being the preparation of the stems, &c., for slicing, the sharpness of the knife and the dexterity with which it is handled, and the method of mounting the sections when made. The wood, if green, should first be soaked in strong alcohol for a few days, to get rid of the resinous matter; and it should then be macerated in water for some days longer, for the removal of its gum, before being submitted to the cutting process. If the wood be dry, it should first be softened by soaking for a sufficient length of time in water, and then treated with spirit and afterwards with water, like green wood. Some woods are so little affected even by prolonged maceration, that boiling in water is necessary to bring them to the degree of softness requisite for making sections. No wood that has once been dry, however, yields such good sections, as that which is cut fresh. When a piece, of the appropriate length, has been placed in the grasp of the Section-instrument (wedges of deal or other soft wood being forced in with it, if necessary for its firm fixation), a few thick slices should first be taken, to reduce its surface to an exact level; the surface should then be wetted with spirit, the micrometer-screw moved through a small part of a revolution, and the slice taken off with the razor, the motion given to which should partake both of *drawing* and *pushing*. A little practice will soon enable the operator to discover, in each case, *how thin* he may venture to cut his sections, without a breach of continuity; and the micrometer-screw should be turned so as to give the required elevation. If the surface of the wood has been sufficiently wetted, the section will not curl up in cutting, but will adhere to the surface of the razor, from which it is best detached by dipping the razor in water, so as to float away the slice of wood, a camel-hair pencil being used to push it off, if necessary. All the sections that may be found sufficiently thin and perfect, should be put aside in a bottle of weak spirit, until they be mounted. For the minute examination of their structure, it is generally much better to preserve them in fluid, than to mount them either dry or in Canada balsam; and no fluid answers better than weak spirit. Where a mere general view only is needed, the dry mounting answers the purpose sufficiently well. It is only in the case of the section being unusually opaque, that mounting it in Canada balsam can be of any service whatever; and in general it is rather injurious than useful, making the section so transparent that its features can scarcely be discerned. Transverse sections, however, when charred by heating between two plates of glass until they turn brown, may be mounted with advantage in Canada balsam, and are then very showy specimens for the solar or gas-microscope. The number of beautiful and interesting objects which may be thus obtained, at the cost of a very small amount of trouble, can scarcely be conceived save by those who have made a special study of these wonderful struc-

tures. Even the commonest trees, shrubs, and herbaceous plants, yield specimens that exhibit a varied elaboration of design, which cannot but strike with astonishment even the most cursory observer; and there is none in which a careful study of sections, made in different parts of the stem, and especially in the neighborhood of the "growing point," will not reveal to the eye of the scientific Physiologist, some of the most important phenomena of Vegetation. *Fossil Woods*, when well preserved, are almost invariably *silicified*, and require, therefore, to be cut and polished by a Lapidary. Should the Microscopist be fortunate enough to meet with a portion of a *calcified* stem, in which the organic structure is preserved, he should proceed with it after the manner of other hard substances which need to be reduced by grinding (§§ 108–110).

245. *Structure of the Cuticle and Leaves*.—On all the softer parts of the higher Plants, save such as grow under water, we find a superficial layer, different in its texture from the parenchyma beneath, and constituting a distinct membrane, known as the

FIG. 180.


Cuticle of Leaf of *Yucca*.

FIG. 181.


Cuticle of Leaf of *Indian Corn* (*Zea Mays*).

Cuticle.¹ This membrane is composed of cells, the walls of which are flattened above and below, whilst they adhere closely to each other laterally, so as to form a continuous stratum (Fig. 184, *a, a*). Their shape is different in almost every tribe of Plants; thus in the cuticle of the *Yucca* (Fig. 180), *Indian Corn* (Fig. 181), *Iris* (Fig. 183), and most other Monocotyledons, the cells are elongated, and present an approach to a rectangular contour; their margins being straight in the *Yucca* and *Iris*, but minutely

¹ The term *epidermis* is applied to this membrane by many Vegetable Physiologists, on account of the analogy it seems to present to the epidermis of Animals; but as epidermis means a membrane that lies upon the *derm* or "true skin," and as no such subjacent layer exists in the Plant, the transference of the designation is altogether inappropriate. It would be much more correct to designate by the name *cutis* or *derm* what is ordinarily denominated the Cuticle; and to reserve the term *epidermis* for the thin pellicle which may be sometimes detached from it (§ 247).

sinuous or crenated in the Indian Corn. In most Dicotyledons, on the other hand, the cells of the cuticle depart less from the form of circular disks; but their margins usually exhibit large irregular sinuosities, so that they seem to fit together like the pieces of a dissected map, as is seen in the cuticle of the *Apple* (Fig. 182, *b, b*). Even here, however, the cells of the portion of

FIG. 182.




Portion of the Cuticle of the inferior surface of the Leaf of the *Apple*, with the layer of parenchyma in immediate contact with it:—*a, a*, elongated cells of the cuticle overlying the veins or nerves of the leaf; *b, b*, ordinary cuticle cells overlying the parenchyma; *c, c*, stomata; *d, d*, green cells of the parenchyma, forming a very open network near the lower surface of the leaf.

the cuticle (*a, a*) that overlies the veins of the leaf, have an elongated form, approaching that of the wood-cells of which these veins are chiefly composed; and it seems likely, therefore, that the elongation of the ordinary cuticle-cells of Monocotyledons has reference to that parallel arrangement of the veins, which their leaves almost constantly exhibit. The cells of the cuticle are colorless, or nearly so, no chlorophyll being formed in their interior; and their walls are generally thickened by secondary deposit, especially on the side nearest the atmosphere. This deposit is of a waxy nature, and consequently renders the membrane very impermeable to fluids; the retention of which within the soft tissue of the leaf is obviously the purpose to be answered by the peculiar organization of the cuticle. In most European Plants, the cuticle contains but a single row of cells, which are usually, moreover, thin-sided; whilst in the generality of tropical species, there exist two, three, or even four layers of thick-sided cells; this last number being seen in the *Oleander*, the cuticle of which, when separated, has an almost leathery firmness. The difference in conformation is obviously adapted to the conditions of growth under which these plants respectively exist; since the cuticle of a plant indigenous to temperate climates, would not afford a sufficient protection to the interior structure against the rays of a tropical sun; whilst the diminished heat of this country would scarcely overcome the resistance presented by the dense and non-conducting tegument of a

species formed to exist in tropical climates. A very curious modification of the cuticle is presented by the *Rochea falcata*, commonly known as the "ice-plant;" a designation it owes to


FIG. 183.



Portion of the Cuticle of the upper surface of the leaf of *Rochea falcata*, as seen at A from its inner side, and at B from its outer side:—a, a, small cells forming the inner layer of the cuticle; b, b, large prominent cells of the outer layer; c, c, stomata, disposed between the latter.

the peculiar appearance of its surface, which looks as if it were covered with frozen dewdrops. This appearance is occasioned by the presence of a layer of very large oval cells (Figs. 183, 184, b, b), which lie detached one from another upon the surface

FIG. 184.



Portion of a vertical section of the same Leaf, showing the small cells, a, a, of the inner layer of cuticle; the large cells, b, b, of the outer layer; c, one of the stomata; d, d, cells of the parenchyma; l, lacuna between the parenchymatous cells, into which the stoma opens.

of the ordinary cuticle, a, a. In other instances, the cuticle is partially invested by a layer of scales, which are nothing else than flattened cells, often having a very peculiar form; whilst in numerous cases, again, we find the surface beset with hairs, which occasionally consist of single elongated cells, but are more commonly made up of a linear series, attached end to end, as in Fig. 153. Sometimes these hairs

bear little glandular bodies at their extremities, by the secretion of which a peculiar viscosity is given to the surface of the leaf, as in the Sundew (*Drosera*); in other instances, the hair has a glandular body at its base, with whose secretion it is moistened; so that when this secretion is of an irritating quality, as in the *Nettle*, it constitutes a "sting." A great variety of such organs may be found, by a microscopic examination of the surface of the leaves of Plants having any kind of superficial investment to the cuticle. Many connecting links are found between hairs and scales, such as the "stellate hairs" of the *Deutzia scabra*, which a good deal resemble those within the air-chambers of the Yellow Water-lily (Fig. 152).


246. The Cuticle in many Plants, especially those belonging to the *Grass* tribe, has its cell-walls impregnated with silex, like that of the *Equisetum* (§ 220); so that, when all the organic matter has been got rid of by heat or by acids, the forms of the cuticle-cells, hairs, stomata, &c., are still marked out in silex, and are most beautifully displayed by Polarized light. Such silicified cuticles are found on the husks of the "grains" yielded by these plants: and there is none in which a larger proportion of mineral matter exists, than that of *Rice*, which contains some curious elongated cells with toothed margins. The hairs with which the *paleæ* (chaff-scales) of most Grasses are furnished, are strengthened by the like siliceous deposit; and in the *Festuca pratensis*, one of the common meadow grasses, the paleæ are also beset with longitudinal rows of little cup-like bodies formed of silica. The cuticle and scaly hairs of *Deutzia* also contain a large quantity of silex; and are remarkably beautiful objects for the Polaroscope.

247. Externally to the cuticle, there usually exists a very delicate transparent pellicle, without any decided traces of organization, though occasionally somewhat granular in appearance, and marked by lines that seem to be impressions of the junctions of the cells with which it was in contact. When detached by maceration, it not only comes off from the surface of the cuticle, but also from that of the hairs, &c., which this may bear. This membrane, the proper *Epidermis* (p. 388, note), is obviously formed by the agency of the cells of the cuticle; and it seems to consist of the external layers of their thickened cellulose walls, which have coalesced with each other, and have separated themselves from the subjacent layers, by a change somewhat analogous to that which occurs in the *Palmelleæ* (§ 194), the outer walls of whose original cells seem to melt away into the gelatinous investment, that surrounds the "broods" which have originated in their subdivision.

248. In nearly all Plants which possess a distinct Cuticle, this is perforated by the minute openings termed *Stomata* (Figs. 182, 183, c, c); which are bordered by cells of a peculiar form, distinct from those of the cuticle, and more resembling in character those of the tissue beneath. These boundary-cells are usually somewhat kidney-shaped, and lie in pairs (Fig. 185, b, b), with an oval opening between them; but by an alteration in their form, the opening may be contracted or nearly closed. In the cuticle of *Yucca*, however, the opening is bounded by two pairs of cells, and is somewhat quadrangular (Fig. 180); and a like doubling of the boundary-cells, with a narrower slit between them, is seen in the cuticle of the *Indian Corn* (Fig. 181). In the stomata of no Phanerogamic Plant, however, do we meet with any conformation at all to be compared in complexity with that which has been described as existing in the humble *Mar-*

chantia (214). Stomata are usually found most abundantly (and

FIG. 185.



Portion of the Cuticle of the leaf of the *Iris germanica*, torn from its surface, and carrying away with it a portion of the parenchymatous layer in immediate contact with it:—*a*, *a*, elongated cells of the cuticle; *b*, *b*, cells of the stomata; *c*, *c*, cells of the parenchyma; *d*, *d*, impressions formed by their contact, on the epidermic cells; *e*, *e*, lacunæ in the parenchyma, corresponding to the stomata.

sometimes exclusively) in the cuticle of the *lower* surfaces of leaves, where they open into the air-chambers that are left in the parenchyma which lies next the inferior cuticle; in leaves which float on the surface of water, however, they are found in the cuticle of the upper surface only; whilst, in leaves that habitually live entirely submerged, as there is no distinct cuticle, so there are no stomata. In the erect leaves of Grasses, the *Iris* tribe, &c., they are found equally (or nearly so) on both surfaces. As a general fact, they are least numerous in succulent Plants, whose moisture, obtained in a scanty supply, is destined to be retained in the system; whilst

they abound most in those which exhale fluid most readily, and therefore absorb it most quickly. It has been estimated that no fewer than 160,000 are contained in every square inch of the under surface of the leaves of *Hydrangea* and of several other plants; the greatest number seeming always to present itself in species, the upper surface of whose leaves is entirely destitute of these organs. In *Iris germanica*, each surface has nearly 12,000 stomata in every square inch; and in *Yucca*, each surface has 40,000. In *Oleander*, *Banksia*, and some other plants, the stomata do not open directly upon the lower surface of the cuticle, but lie in the deepest part of little pits or depressions which are excavated in it, and which are lined with hairs; the mouths of these pits, with the hairs that line them, are well brought into view by taking a thin slice from the surface of the cuticle with a sharp knife; but the form of the cavities, and the position of the stomata, can only be well made out in vertical sections of the leaves.

249. The internal structure of *Leaves* is best brought into view by making vertical sections, that shall traverse the two layers of cuticle and the intermediate cellular parenchyma; portions of such sections are shown in Figs. 184, 186, and 187. In close apposition with the cells of the upper cuticle (Fig. 186, *a*, *a*), which may or may not be perforated with stomata (*c*, *c*, *d*, *d*), we find a layer of soft thin-walled cells, containing a large quantity of chlorophyll; these cells usually press so closely against one another, that their sides become mutually flattened, and no spaces are left, save where there is a definite air-chamber into which the stoma opens (Fig. 184, *L*); and the compactness of

this superficial layer is well seen, when, as often happens, it adheres so closely to the cuticle, as to be carried away with this when it is torn away (Fig. 185, *c, c*). Beneath this first layer of leaf-cells, there are usually several others, rather less compactly arranged; and the tissue gradually becomes more and more lax, its cells not being in close apposition, and large intercellular passages being left amongst them, until we reach the lower cuticle, which the parenchyma only touches at certain points, its lowest layer forming a sort of network (Fig. 182, *d, d*), with large interspaces, into which the stomata open. It is to this arrangement that the darker shade of green almost invariably presented by the superior surfaces of leaves, is principally due; the color of the component cells of the parenchyma not being deeper in one part of the leaf than in another. In those plants, however, whose leaves are erect instead of being horizontal, so that their two surfaces are equally exposed to light, the parenchyma is arranged on both sides in the same manner, and their cuticles are furnished with an equal number of stomata. This is the case, for example, with the leaves of the common Garden *Iris* (Fig. 187); of which, moreover, we find a central portion (*d, d*) formed by


FIG. 186.



Vertical section of the Cuticle, and of a portion of the subjacent parenchyma, of a Leaf of *Iris germanica*, taken in a transverse direction:—*a, a*, cells of the cuticle; *b, b*, cells at the sides of the stomata; *c, c*, small green cells placed within these; *d, d*, openings of the stomata; *e, e*, lacunae of the parenchyma, corresponding to the stomata; *f, f*, cells of the parenchyma.

thick-walled colorless tissue, very different either from ordinary leaf-cells or from woody fibre. The explanation of its presence is probably to be found in the peculiar conformation of the leaves; for if we pull one of them from its origin, we shall find that what appears to be the flat expanded blade really exposes but half its surface; the blade being doubled together longitu-

FIG. 187.



Portion of a vertical longitudinal section of the leaf of *Iris*, extending from one of its flattened sides to the other:—*a, a*, elongated cells of the epidermis; *b, b*, stomata cut through longitudinally; *c, c*, green cells of the parenchyma; *d, d*, colorless tissue, occupying the interior of the leaf.

thick-walled colorless tissue, very different either from ordinary leaf-cells or from woody fibre. The explanation of its presence is probably to be found in the peculiar conformation of the leaves; for if we pull one of them from its origin, we shall find that what appears to be the flat expanded blade really exposes but half its surface; the blade being doubled together longitu-

dinally, so that what may be considered its under surface is entirely concealed. The two halves are adherent together at their upper part; but at their lower they are commonly separated by a new leaf, which comes up between them; and it is from this arrangement, which resembles the position of the legs of a man on horseback, that the leaves of the Iris tribe are said to be *equitant*. Now by tracing the middle layer of colorless cells, *d, d*, down to that lower portion of the leaf, where its two halves diverge from one another, we find that it there becomes continuous with the cuticle, to the cells of which (Fig. 185, *a*) these bear a strong resemblance, in every respect save the greater proportion of their breadth to their length. Another interesting variety in leaf-structure is presented by the Water-Lily, and other plants whose leaves float on the surface; for here the usual arrangement is entirely reversed, the closely-set layers of green leaf-cells being found in contact with the lower surface, whilst all the upper part of the leaf is occupied by a loose spongy parenchyma, containing a very large number of air-spaces that give buoyancy to the leaf; and these spaces communicate with the external air through the numerous stomata, which, contrary to the general rule (§ 248), are here found in the upper cuticle alone.

250. The examination of the foregoing structures is attended with very little difficulty. Many cuticles may be torn off, by the exercise of a little dexterity, from the surfaces of the leaves they invest, without any preparation; this is especially the case with Monocotyledonous plants, the "veins" of whose leaves run parallel, and with such Dicotyledons as have very little woody structure in their leaves; in those, on the other hand, whose leaves are furnished with reticulated veins, to which the cuticle adheres (as is the case in by far the larger proportion), this can only be detached by first macerating the leaf for a few days in water; and if the texture of the cuticle should be particularly firm, the addition of a few drops of nitric acid to the water will render them more easily separable. If it be desired to preserve them, they may be advantageously mounted in weak spirit. Very good sections of most leaves may be made by a sharp knife, handled by a careful manipulator; but it is generally preferable to use Valentin's knife (§ 106) or the section instrument (§ 107); taking care in the former case to cut down upon a piece of fine cork; and in the latter not to crush the leaf between the two pieces of cork that hold it, and to use very soft cork whenever the delicacy of the leaf renders this desirable. In order to study the structure of leaves with the fulness that is needed for scientific research, numerous sections should be made in different directions; and slices taken parallel to the surfaces, at different distances from them, should also be examined. There is no known liquid, in which such sections can be preserved altogether without change; but water with a small dash of spirit,

seems to answer best, provided the cell be air-tight, and the specimen fresh.

251. *Structure of Flowers.*—Many of the smaller Flowers are, when looked at entire, with a low magnifying power, very striking Microscopic objects; and the interest of the young in such observations can scarcely be better excited, than by directing their attention to the new view they thus acquire of the “composite” nature of the humble down-trodden *Daisy*, or to the beauty of the minute blossoms of many of those *Umbelliferous* plants, which are commonly regarded only as rank weeds. The scientific Microscopist, however, looks more to the organization of the separate parts of the flower; and among these he finds abundant sources of gratification, not merely to his love of knowledge, but also to his taste for the beautiful. The general structure of the *Sepals* and *Petals* which constitute the “perianth” or “floral envelopes,” closely corresponds with that of leaves; the chief difference lying in the peculiar changes of hue which the chlorophyll almost invariably undergoes in the latter class of organs, and very frequently in the former also. There are some petals, however, whose cells exhibit very interesting peculiarities, either of form or marking, in addition to their distinctive coloration; such are those of the *Geranium* (*Pelargonium*), of which a small portion is represented in Fig. 188. The different portions of the petal,—when it has been dried after stripping it of its cuticle, immersed for an hour or two in oil of turpentine, and then mounted in Canada balsam,—exhibit a most beautiful variety of vivid coloration, which is seen to exist chiefly in the thickened partitions of the cells; whilst the surface of each cell presents a very curious opaque spot with numerous diverging prolongations, which looks as if formed by a deposit of




FIG. 188.

Cells from the Petal of the *Geranium* (*Pelargonium*).

sclerogen upon its interior. This method of preparation, however, does not give a true idea of the structure of the cells; for each of them has a peculiar mammillary protuberance, the base of which is surrounded by hairs; and this it is which gives the velvety appearance to the surface of the petal, and which, when altered by drying and compression, occasions the peculiar spots represented in Fig. 188. The real character may be brought into view by Dr. Inman's method; which consists in drying the petal (when stripped of its cuticle) on a slip of glass, to which it adheres, and then placing on it a little Canada balsam diluted with turpentine, which is to be boiled for an instant over the

spirit-lamp, after which it is to be covered with a thin glass. The boiling "blisters" it, but does not remove the color; and on examination, many of the cells will be found showing the mammilla very distinctly, with a score of hairs surrounding its base, each of these slightly curved, and pointing towards the apex of the mammilla. The petal of the common Scarlet Pimpernel (*Anagallis arvensis*), that of the common Chickweed (*Stellaria media*), together with many others of a small and delicate character, are also very beautiful microscopic objects; and the two just named are peculiarly favorable subjects for the examination of the spiral vessels in their natural position. For the "veins" which traverse these petals are entirely made up of spiral vessels, none of which individually attain any great length; but one follows or takes the place of another, the conical commencement of each somewhat overlapping the like termination of its predecessor; and where the veins seem to branch, this does not happen by the bifurcation of a spiral vessel, but by the "splicing on" (so to speak) of one to the side of another, or by the "splicing on" of two new vessels diverging from one another, to the end of that which formed the principal vein.¹

252. The *Anthers* and *Pollen-grains*, also, present numerous objects of great interest, both to the scientific Botanist and to the amateur Microscopist. In the first place, they afford a good opportunity of studying that "free" cell-development, which seems peculiar to the parts concerned in the Reproductive process, and which consists in the development of a new cell-wall round an isolated mass of protoplasm forming part of the contents of a "parent-cell;" so that the new cell lies free within its cavity, instead of being developed in continuity with it, as in the ordinary methods of multiplication (§§ 150, 198). If the Anther be examined, by thin sections, at an early stage of its development within the young flower-bud, it will be found to be made up of ordinary cellular parenchyma, in which no peculiarity anywhere shows itself; but a gradual "differentiation" speedily takes place, consisting in the development of a set of very large cells in two vertical rows, which occupy the place of the *loculi* or pollen-chambers that afterwards present themselves; and these cells give origin to the pollen-grains, whilst the ordinary parenchyma remains to form the walls of the pollen-chambers. The first change consists in the multiplication of the cells of the primary row, by cell-division, in correspondence with the general increase in the size of the anther; until at length they form masses of considerable size, composed of large squarish cells, filled with granular contents, well defined as constituting a distinct tissue from the walls of the pollen-chambers. The history of the development of the pollen-grains in their interior is thus described by Mr. Henfrey, who has made a special study of it. "The cou-

¹ See Mr. R. H. Solly's description and figure of the petal of the *Anagallis*, in "Trans. of Society of Arts," vol. xlviii.


tents of each of these cells secrete a layer of cellulose, which does not adhere to the wall of the parent-cell to form a layer of secondary deposit, but lies free against it, so that a new free cell is formed within each old one, nearly filling it. The walls of the old cells then dissolve, so that the free cells become free, no longer in their parent-cells, but in a cavity which is to constitute the pollen-chamber or loculus of the anther. These free cells are the 'parent-cells of the pollen' of authors. A new phenomenon soon occurs in these. These parent-cells divide into four by ordinary cell-division; either by one or two successive partings, by septa at right angles to each other, but both perpendicular to an imaginary axis (as when an orange is quartered); or by simultaneously formed septa, which cut off portions in such a manner, that the new cells stand in the position of cannon-balls piled into a pyramid (tetrahedrally). These new cells are the 'special parent-cells of the pollen;' and in each of these the entire protoplasmic contents secrete a series of layers, which, in the ordinary course, by the solution of the primary walls of the special parent-cells upon which they were applied, become the walls of free cells, which constitute the simple ordinary pollen-cells. These subsequently increase in size, and their outer coat assumes its characteristic form and appearance, while free in the chamber of the anther."¹ This history bears a very close parallel with that of the development of the spores within the "theca" of the Mosses (§ 217); and it is not a little curious that the layer of cells which lines the pollen-chambers, should exhibit, in a considerable proportion of plants, a strong resemblance in structure, though not in form, to the elaters of the *Marchantia* (Fig. 132). For they have in their interior a fibrous deposit; which sometimes forms a continuous spiral (like that in Fig. 157), as in *Narcissus* and *Hyoscyamus*; but is often broken up, as it were, into rings, as in the *Iris* and *Hyacinth*; in many instances, forms an irregular network, as in the *Violet* and *Saxifrage*; in other cases, again, forms a set of interrupted arches, the fibres being deficient on one side, as in the yellow *Water-lily*, *Bryony*, *Primrose*, &c.; whilst a very peculiar stellate aspect is often given to these cells, by the convergence of the interrupted fibres towards one point of the cell-wall, as in the *Cactus*, *Geranium*, *Madder*, and many other well-known plants. Various intermediate modifications exist; and the particular form presented, often varies in different parts of the wall of one and the same anther. It seems probable that, as in *Hepaticæ*, the elasticity of these spiral cells may have some share in the opening of the pollen-chambers and the dispersion of the pollen-grains.

253. The form of the Pollen-grains seems to depend in part upon the mode of division of the cavity of the parent-cell into quarters; generally speaking it approaches the spheroidal, but it is sometimes elliptical, and sometimes tetrahedral. It varies

¹ "Micrographic Dictionary," p. 516.

more, however, when the pollen is dry, than when it is moist; for the effect of the imbibition of fluid, which usually takes place when the pollen is placed in contact with it, is to soften down angularities, and to bring the cell nearer to the typical sphere. The pollen-cell (save in a few submerged plants) has a thick outer coat, surrounding a thin interior wall; and this often exhibits very curious markings, which seem due to an increased thickening at some points, and a thinning away at others. Sometimes these markings give to the surface-layer so close a resemblance to a stratum of cells (Fig. 189, B, C, D), that only a very careful examination can detect the difference. The roughening of the surface by spines or knobby protuberances, as shown at A, is a very common feature; and this seems to answer the purpose of enabling the pollen-grains more readily to hold to the surface whereon they may be cast. Besides these and other inequalities of the surface, most pollen-grains have what appear to be pores or slits in the outer coat, varying in number in different species, through which the inner coat protrudes itself, when the bulk of its contents has been increased by imbibition; it seems probable, however, that the outer coat is not absolutely deficient at these points, but is only thinned away. Sometimes the pores are covered by little disk-like pieces, or lids, which fall off when the pollen-tube is protruded. This action takes place naturally, when the pollen-grains fall upon the surface of the "stigma," which is moistened with a viscid secretion; and the pollen-tubes, at first

mere protrusions of the inner coat of their cell, insinuating themselves between the loosely packed cells of the stigma, grow downwards through the "style," sometimes even to the length of several inches, until they reach the ovarium. The first change,—namely, the protrusion of the inner membrane through the pores of the exterior,—may be made to take place artificially, by moistening the pollen with water, thin syrup, or dilute acids (different kinds of pollen-grains requiring a different mode of treatment); but the subsequent extension by



growth will only take place under the natural conditions.

254. The darker kinds of pollen may be best mounted for the Microscope in Canada balsam; but this renders the more transparent kinds too faintly distinguishable; and it is better to

mount them either dry, or (if they will bear it without rupturing) in fluid. The most delicate and interesting forms are found, for the most part, in plants of the Natural families *Amarantaceæ*, *Cichoraceæ*, *Cucurbitaceæ*, *Malvaceæ*, and *Passifloreæ*; others are furnished also by *Convolvulus*, *Campanula*, *Oenothera*, *Pelargonium* (*Geranium*), *Polygonum*, *Sedum*, and many other Plants. It is frequently preferable to lay down the entire anther with its adherent pollen-grains (where these are of a kind that hold to it), as an opaque object; this may be done with great advantage in the case of the common Mallow (*Malva sylvestris*) or of the Hollyhock (*Althæa rosea*); the anthers being picked soon after they have opened, whilst a large proportion of their pollen is yet undischarged, and before they have begun to wither, being laid down as flat as possible between two pieces of smooth blotting-paper, then subjected to moderate pressure, and finally mounted upon a black surface. They are then, when properly illuminated, most beautiful objects for the 2 in. objective.

255. The structure and development of the *Ovules* that are produced within the ovarium at the base of the pistil, and the operation in which their fertilization essentially consists, are subjects of investigation which have a peculiar interest for scientific Botanists, but which, in consequence of the special difficulties that attend the inquiry, are not commonly regarded as within the province of amateur Microscopists. The ovule, in its earliest condition, is, like the anther, a mass of cells, in which no part is differentiated from the rest; gradually, this body, which is termed the *nucleus*, is found to be enveloped in one, two, or three coats, which are formed by the multiplication of cells that at first constitute merely an annular enlargement at its base; these coats, however, do not entirely close in around the nucleus, at the point of which a small aperture always remains, that is called the *micropyle*. In the interior of the nucleus a large cavity is formed, apparently by the enlargement of one of its cells at the expense of those which surround it; and this cavity, which is called the *embryo sac*, is at first filled only with a liquid protoplasm. Some little time before fecundation, however, there are seen in it a certain number of free cell-nuclei, rarely fewer than three, and frequently more; around these, free cells of a spheroidal form are developed, which are "germ-cells," of which one only, the *embryonal vesicle*, is ordinarily destined to be fertilized. This act is accomplished by the penetration of the pollen-tube, which, when it has made its way down to the ovarium, enters the "micropyle" of the ovule, and impinges upon the apex of the "embryo sac," which it sometimes pushes before it, in such a manner as to have given origin to the idea that the tube enters its cavity: no such penetration, however, really exists. By Prof. Schleiden, and his disciple Schacht, it is affirmed that the embryo makes its first appearance within the dilated extremity of the pollen-tube, which speedily becomes filled with a mass of cells;

but nearly all other Vegetable Physiologists who have examined the question with sufficient care, have come to a different conclusion,—namely, that the embryo is the product of a cell-multiplication within the “embryonal vesicle,” which seems to be fertilized by the transudation of the contents of the pollen-tube, just as the embryonal vesicle within the archegonium of the Ferns is fertilized by the contact of the antherozoids (§ 219).

256. The early processes of development, too, correspond closely with those which have been described as taking place through the whole of the inferior tribes; for the primordial cell that is formed within the embryonal vesicle as the result of its fecundation, gives origin by transverse fission to a pair; this again, to four; and so on, it being usually in the terminal cell of the filament so generated, that the process of multiplication chiefly takes place, as in the *Confervæ* (§ 198). The filament then begins to enlarge at its lower extremity, where its cells are often multiplied into a somewhat globular mass; of this mass, by far the larger proportion is destined to be evolved into the “cotyledons,” or seed-leaves, whose function is limited to the earliest part of the life of the young plant; the small remainder is the rudiment of the “plumula,” which is to be developed within the stem and leaves; while the prolonged extremity of the embryonic filament, which is directed towards the micropyle, is the original of the “radicle” or embryonic root. The mucilaginous protoplasm filling the “embryo sac,” in which the “embryonal vesicle” was imbedded, becomes converted, by the formation of free cells, soon after fecundation, into a loose cellular tissue, which constitutes what is known as the “endosperm;” this, however, usually deliquesces again, as the embryonic mass increases in bulk and presses upon it; and its development is of interest, chiefly because it may be shown, by the curious intermediate phase presented by the *Lycopodiaceæ* and *Coniferæ* (§ 221), that this endosperm is the equivalent of the “prothallium” of the higher Cryptogamia.¹

257. In tracing the origin and early history of the Ovule, very thin sections should be made through the flower-bud, both vertically and transversely; but when the ovule is large and distinct enough to be separately examined, it should be placed on the thumb-nail of the left hand, and very thin sections made with a sharp razor; the ovule should not be allowed to dry up, and the sections should be removed from the blade of the razor by a wetted camel-hair pencil. The tracing downwards the pollen-tubes through the tissue of the style, may be accomplished by


¹ For a more detailed account of the Generative process in ordinary Phanerogamia and in *Coniferæ*, as well as for references to the principal original sources of information on this controverted question, the Author may be permitted to refer to his “Principles of Comparative Physiology,” 4th Ed. §§ 501–505. Some recent discussion on the same subject will be found in the “Ann. des Sci. Nat.” 4ième Sér. tom. iii, p. 188 et seq.

sections (which, however, will seldom follow one tube continuously for any great part of its length), or, in some instances, by careful dissection with needles. Plants of the *Orchis* tribe are the most favorable subjects for this kind of investigation, which is best carried on by artificially applying the pollen to the stigma of several flowers, and then examining one or more of the styles daily. "If the style of the flower of an *Epipactis* (says Schacht), to which the pollen has been applied about eight days previously, be examined in the manner above mentioned, the observer will be surprised at the extraordinary number of pollen-tubes, and he will easily be able to trace them in large strings, even as far as the ovules. *Viola tricolor* (heartsease) and *Ribes nigrum* and *rubrum* (black and red currant) are also good plants for the purpose; in the case of the former plant, withered flowers may be taken, and branched pollen-tubes will not unfrequently be met with." The entrance of the pollen-tube into the micropyle may be most easily observed in *Orchideous* plants and in *Euphrasia*; it being only necessary to tear open with a needle the ovary of a flower which is just withering, and to detach from the placenta the ovules, almost every one of which will be found to have a pollen-tube sticking in its micropyle. These ovules, however, are too small to allow of sections being made, whereby the origin of the embryo may be discerned; and for this purpose, *Ænothera* (evening primrose) has been had recourse to by Hoffmeister, whilst Schacht recommends *Lathrœa squamaria*, *Pedicularis palustris*, and particularly *Pedicularis sylvatica*. There is no kind of investigation that requires nicer management, and none which is just now of greater interest to Botanists. Such Microscopists, therefore, as have qualified themselves for the inquiry, by their acquirement of the knowledge which is requisite to guide their dissections, and of the manipulative skill by which alone these dissections can be successfully made, cannot do a greater service to science, than by applying themselves perseveringly to it. The use of high magnifying powers is not at all needed. Much may be done, in the preparation of the objects, under the Simple microscope; and for the examination of the preparations, a power of 200 diameters with a shallow eye-piece is generally sufficient. The assistance of the Binocular Microscope would probably be found peculiarly valuable in this inquiry; since the right interpretation of the appearances presented, mainly depends upon a precise knowledge of the exact relative position of the pollen-tube, embryo sac, &c., such as this instrument is peculiarly fitted to convey.

258. We have now, in the last place, to notice the chief points of interest to the Microscopist, which are furnished by mature *Seeds*. Many of the smaller kinds of these bodies are very curious, and some are very beautiful, objects, when looked at in their natural state, under a low magnifying power. Thus the seed of

the *Poppy* (Fig. 190, A) presents a regular reticulation upon its

FIG. 190.



Seeds, as seen under a low magnifying power:—A, *Poppy*; B, *Amaranthus* (Prince's feather); C, *Antirrhinum majus*? (Snapdragon); D, *Caryophyllum* (Clove-pink); E, *Bignonia*.

surface, pits, for the most part hexagonal, being left between projecting walls; that of *Caryophyllum* (D) is regularly covered with curiously jagged divisions, every one of which has a small bright black hemispherical knob in its middle; that of *Amaranthus hypochondriacus* has its surface traced with extremely delicate markings (B); that of *Antirrhinum* (?) is strangely irregular in shape (C), and looks almost like a piece of furnace-slag; and that of *Bignonia* (E) is remarkable for the beautiful structure of the translucent membrane which sur-

rounds it, the radiating lines shown in the figure being found under a higher magnifying power to consist of rows of elongated spiral cells. Such are seen, too, in the like delicate membrane that surrounds several other seeds, as those of *Sphenogyne speciosa* and *Lophospermum erubescens*, which, from possessing this appendage, are spoken of as "winged." The most remarkable development of this structure is said by Mr. Quekett to exist in a seed of *Calosantes Indica*, an East Indian plant, in which the wing extends more than an inch on either side of the seed. Some seeds are distinguished by a peculiarity of form, which, although readily discernible by the naked eye, becomes much more striking when they are viewed under a very low magnifying power; this is the case, for example, with the seeds of the *Carrot*, whose long radiating processes make it bear, under the Microscope, no trifling resemblance to some kinds of star-fish; and with those of *Cyanthus minor*, which bear about the same degree of resemblance to shaving-brushes. In addition to the preceding, the following may be mentioned as seeds easily to be obtained, and as worth mounting for opaque objects:—*Anagallis*, *Anethum graveolens*, *Antirrhinum*, *Begonia*, *Carum carui*, *Coriopsis tinctoria*, *Datura*, *Delphinium*, *Digitalis*, *Elatine*, *Erica*, *Gentiana*, *Gesnera*, *Hyoscyamus*, *Hypericum*, *Lepidium*, *Limnocharis*, *Linaria*, *Lychnis*, *Mesembryanthemum*, *Nicotiana*, *Orobanche*, *Petunia*, *Roseda*, *Saxifraga*, *Scrophularia*, *Sedum*, *Sempervivum*, *Silene*, *Stellaria*, and *Verbena*. The following may be mounted as transparent objects in Canada balsam:—*Drosera*, *Hydrangea*, *Mono-*

*tropa, Orchis, Parnassia, Pyrola, Saxifraga.*¹ The seeds of Umbelliferous plants generally are remarkable for the peculiar *vittæ*, or receptacles for essential oil, which are found in their coats. Various points of interest respecting the structure of the *testæ* or envelopes of seeds,—such as the fibre-cells of *Cobæa* and *Collomia*, the stellate cells of the *Star-Anise*, and the densely consolidated tissue of the “shells” of the *Coquilla-nut, Cocoa-nut, &c.*,—having been already noticed, we cannot here stop to do more than advert to the peculiarity of the constitution of the “husk” of the *Corn-grains*. In these, as in other Grasses, the ovary itself continues to envelope the seed, forming a covering to it, that surrounds its own *testa*; this covering (which forms the “bran” that is detached in grinding) is composed of hexagonal cells of remarkable regularity and density; and these are so little altered by a high temperature, as still to be readily distinguishable when the grain has been ground after roasting,—thus enabling the Microscopist to detect even a very small admixture of roasted Corn with Coffee or chicory, without the least difficulty.²

¹ These lists have been chiefly derived from the “Micrographic Dictionary,” p. 572.

² In a case in which the Author was called upon to make such an investigation, he found as many as *thirty* distinctly recognizable fragments of this cellular envelope, in a *single grain* of a mixture consisting of Chicory with only 5 per cent. of roasted Corn.

CHAPTER IX.

MICROSCOPIC FORMS OF ANIMAL LIFE:—PROTOZOA; ANIMALCULES.


259. *Protozoa*.—Passing on, now, to the Animal Kingdom, we begin by directing our attention to those minute and simple forms, which correspond, in the Animal series, with the Protophyta in the Vegetable (Chap. VI); and this is the more desirable, since the formation of a distinct group, to which the name of *Protozoa* (first proposed by Siebold) may be appropriately given, is not merely one of the most interesting results of recent Microscopic inquiry, but is a subject on which it is particularly important that the Microscopic observer should know what the Physiologist believes himself to have ascertained. This group, which must be placed at the very base of the animal scale, beneath the great subkingdoms marked out by Cuvier, is characterized by the extreme simplicity that prevails in the structure of the beings composing it; these being either *isolated cells*, or *aggregations of cells* wherein no such differentiation of parts exhibits itself, as constitutes the “organs” of even the simplest Zoophyte or Worm. We have in the first place to consider, therefore, what are the essential characters of the Animal cell; and what are the precise relations of the Protozoa to the Protophyta, to which they seem to bear so close an affinity.

260. The Animal cell, in its most complete form, is comparable in most parts of its structure to that of the Plant; but differs from it in the entire absence of the “cellulose-wall,” or of anything that represents it, the cell contents being enclosed in only a single limitary membrane, the chemical composition of which (being albuminous) indicates its correspondence with the primordial utricle (§ 147). In its young state, it seems always to contain a semi-fluid *plasma*, which is essentially the same as the “protoplasm” of the Plant, save that it does not include chlorophyll-granules; and this may either continue to occupy its cavity (which is the case in cells whose entire energy is directed to growth and multiplication), or may give place, either wholly or in part, to the special product which it may be the function of the cell to prepare. Like the Vegetable cell, that of Animals very commonly multiplies by duplicative subdivision; and it also (especially among Protozoa) may give origin to new cells,

by the breaking up of its contents into several particles; but new cells are not unfrequently to be met with, especially in the nutritive fluids of such Animals as possess a distinct circulation, which have not directly originated in either of these modes from a previously existing cell, but which have been developed by a process of *free* cell-formation, namely, by the aggregation of organic molecules, floating in these fluids, into little masses, of which the external particles coalesce into a membranous cell-wall, whilst the interior liquefy into cell-contents. This can only take place, however, in a liquid which has undergone elaboration in the interior of a highly-organized living body; and we find no traces of such free cell-formation among the members of the group we are first to investigate.

261. As we have seen (§ 150) that, among the lowest Proto-phytes, the general attributes of a cell may exist in a minute mass of protoplasm which is not bounded by a limitary membrane,—the differentiation between cell-wall and cell-contents not having yet manifested itself,—so, among the lowest *Protozoa*, we find the power of maintaining an independent existence to be possessed by similar particles of that peculiar *blastema*, or formative substance, to which the name of *sarcode* has been given by Dujardin (who first drew attention to its extraordinary endowments), and which may be considered as the basis, not merely of the entire organisms of Protozoa, but of a large part of that of higher animals. The properties of this may be most fully understood by the careful study of a creature, which is by no means unfrequently to be met with in fresh and stagnant waters, vegetable infusions, &c., and which, from the great variety of forms it assumes, has received the designation of *Pro-*

FIG. 191.

*Amœba princeps*, in different forms, A, B, C.

teus. This name, however, having been assigned to an animal of far higher organization, that with which we are now concerned is properly known as the *Amœba* (Fig. 191). It may be


described as a minute mass of "sarcode," presenting scarcely any evidence of distinct organization, even of the simplest kind; for, as in the lowest forms of Vegetable cell (§ 148), there is not even a complete differentiation between the cell-wall and the cell-contents; the former not being composed of a distinct membrane, though obviously possessing more consistence than the latter, which are semi-fluid. A contractile vesicle (or rather, perhaps, a "vacuole") may be observed in some part of the body, which pulsates at tolerably regular intervals; and "vacuoles" or clear spaces are seen, surrounding the alimentary particles which have been received into the midst of the jelly-like substance. However inert and shapeless this minute body may be when first noticed, its possession of vital activity is soon made apparent by the movements which it executes, and by the changes of form which it undergoes; these being, in fact, part of one and the same set of actions. For the shapeless mass puts forth one or more finger-like prolongations, which are simply extensions of its gelatinous substance in those particular directions; and a continuation of the same action, first distending the prolongation, and then (as it were) carrying the whole body into it, causes the entire mass to change its place. After a short time another prolongation is put forth, either in the same or in some different direction; and the body is again absorbed into it. These changes seem to be connected with a movement of the semi-fluid particles in the interior of the mass, of which a current may be observed to "set" in the direction wherein the protrusion is about to take place, *before* the surface shows any projection. When the creature, in the course of its progress, meets with a particle capable of affording it nutriment, its gelatinous body spreads itself over or around this, so as to envelope it completely; and the substance (sometimes animal, sometimes vegetable) thus taken into this extemporized stomach, undergoes a sort of digestion there, the nutrient material being extracted, and any indigestible part making its way to the surface and finally being (as it were) squeezed out. Of the mode of reproduction of *Amœba*, nothing is yet known, save that it undergoes multiplication by self-division, very much in the manner of the Protophytes, and that portions separated from the jelly-like mass, either by cutting or tearing, can develop themselves into independent beings. Consequently, as we are quite in the dark respecting the sexual operation, which (as all analogy would lead us to believe) must take place at some period of its life, it cannot be said that we are acquainted with more than one phase of its existence; and it is quite possible that, after many repetitions of the process of multiplication by self-division, some entirely new form may present itself, of which the *Amœba* is (as it were) the larva. The completion of the life-history of this curious creature, therefore, is a most worthy object of Microscopic inquiry; and its abundance in many situations should prevent this from being a matter of

any great difficulty to an observer, who can devote sufficient attention to the study.¹

262. Nearly allied to the preceding, is another curious organism, on which the attention of many eminent Microscopists has been recently fixed. This creature, the *Actinophrys* (Fig. 192), consists like the preceding of a homogeneous, jelly-like contractile substance, or *sarcode*, not enclosed in any distinct envelope, though the outer portion seems to be of firmer consistence than the inner. Throughout the body, which is usually nearly spherical in form, but more particularly near its surface, there are observed vacuoles occupied by fluid; these have no definite boundaries, and may be easily made artificially either to coalesce into larger ones, or to subdivide into smaller. A "contractile vesicle" (*o*), pulsating rhythmically with great regularity, is always to be distinguished either in the midst of the jelly-like substance, or (more commonly), near its surface; and the appearance which it presents in this latter position, seems to leave no doubt of its being included within a distinct though very thin membrane. The "sarcode" extends itself into contractile tentacular filaments, which are called *pseudopodia*; and these, in the *Actinophrys sol*, are commonly seen to radiate from the centre, in such a manner as to have suggested the designation of the species.

Their degree of extension, however, is extremely variable, and sometimes they entirely disappear: the creature cannot then be distinguished with certainty from an *Amœba*. For although the form

FIG. 192.



Actinophrys sol, in different states:—A, in its ordinary sun-like form, with a prominent contractile vesicle, *o*; B, in the act of division or of conjugation, with two contractile vesicles, *o, o*; C, in the act of feeding; D, in the act of discharging fecal (?) matters, *a* and *b*.

¹ It has been recently affirmed by Dr. Hartig, that *Amœbæ* may be produced by the transformation of the "antherozoids" of *Chara* (§ 202), *Marchantia*, or *Mosses*; and that, in their turn, they become metamorphosed, first into *Protococci* or other unicellular *Alge*, and then into *Articulated Alge*. But even if jelly like bodies resembling *Amœbæ* in general appearance and in spontaneous change of form, should be thus produced, they cannot be said to be true *Amœbæ*, unless they should feed in the manner described above,—which Dr. Hartig does not appear to have witnessed. (See "Quart. Journal of Microscopic Science," vol. iii, p. 51.) However strange Dr. Hartig's statements may be, they are not more strange than many assertions of the same class first appeared, which are now admitted as unquestionable truths; and they ought not to be set aside without disproof, any more than they should be received without further confirmation.

of the latter is generally flattened, yet it sometimes becomes nearly spherical; so that neither type can be recognized, until the jelly-like spherule flattens itself out as an *Amœba*, or puts forth radiating "pseudopodia" as an *Actinophrys*. Far less activity is exhibited by *Actinophrys*, than by *Amœba*; and the slight change of place which it undergoes from time to time, does not seem attributable either to any change of form of the body, or to any bending of the tentacles. It is by the agency of these, however, that its nourishment is obtained; and this is derived not merely from vegetable particles, but from various small animals, some of them (as the young of Crustaceans) possessing great activity, as well as a comparatively high organization. When any of these happens to come into contact with one of the tentacular filaments, this usually retains it by adhesion, and forthwith begins to retract itself; as it shortens, the surrounding filaments also apply themselves to the captive particle, bending their points together so as gradually to enclose it, and themselves retracting, until the prey is brought close to the surface of the body. The threads of "sarcode" of which the "pseudopodia" are composed, not being invested (any more than the sarcode of the body) by any limiting membrane, coalesce with each other and with it; and thus the particle which has been entrapped by them becomes actually embedded in the gelatinous mass, and gradually passes from the peripheral towards the central part of it, where its digestible portion undergoes solution, the superficial part of the body, with its pseudopodial prolongations, in the meantime recovering their previous condition. Any indigestible portion, as the shell of a Crustacean, or the hard case of a Rotifer, finds its way to the surface of the body; and is extruded from it by a process exactly the converse of that by which it is drawn in (Fig. 192, D). The number as well as the size of the particles included by the *Actinophrys* at any one time is very various; frequently they are more than ten or twelve. They are not usually embraced closely by the sarcode, but are surrounded by fluid in "vacuoles" in its substance; and it was this appearance which led Prof. Ehrenberg to describe the animalcule as possessing numerous stomachs. The *Actinophrys*, like the *Amœba*, multiplies itself by self-division; but a process which seems to resemble the "conjugation" of Protophytes (§ 151), has also been witnessed in it by Prof. Kölliker and Dr. Cohn.¹ Two individuals approximate and coalesce, so as to form what appears to be a single body; but a "nucleus" then makes its appearance, which gradually develops itself into a mass having the characters of its parent; and the young *Actinophrys* thus generated probably escapes before long from the body within which it

¹ It appears probable, from the recent observations of Mr. Weston (Quarterly Journal of Microsc. Science, Jan 1856), and others, that the supposed "conjugation" of *Actinophrys* is a mere fusion of two bodies which may separate again unchanged, and is not a generative phenomenon. The same would appear to be true of it in this respect, as of *Gregarina* (§ 358).

originated. As the condition of an Actinophrys undergoing self-division is to all appearance the same as that of two individuals in incipient conjugation (Fig. 192, B), it cannot be determined in any particular case which operation is in progress, until the creature has been watched sufficiently long for the *tendency* of its changes to become apparent.

263. If, now, we compare the foregoing history with that of *Palmoglaea* or any other simple Protophyte, we shall see that whilst there is a strong analogy between the two sets of phenomena, there are at the same time certain most important differences. One of the most obvious of these differences, lies in the *movements* exhibited by the Amœba and Actinophrys; for although these are by no means sufficient in themselves (as was once supposed) to establish the distinction between the Animal and Vegetable kingdoms, yet, when they do not consist in the mere vibrations of cilia, such as are executed by zoospores, antherozoids, &c., among Plants, but depend upon alterations in a contractile substance forming the entire body, they bear a much closer resemblance to the actions of higher Animals; and we may trace in fact, in the ascending animal scale, a progressive specialization or setting apart of certain portions of the contractile substance more peculiarly endowed with this property, until they take the form of distinct muscular bands. A more positive and easily defined distinction lies in the *nature of the aliment* of the Protophyta and Protozoa respectively, and in the *method of its introduction*. For whilst the Protophyte obtains the materials of its nutrition from the air and moisture that surround it, and possesses the power of detaching oxygen, hydrogen, carbon, and nitrogen, from their previous binary combinations, and of uniting them into ternary, and quaternary organic compounds (chlorophyll, starch, albumen, &c.), the simplest Protozoon, in common with the highest members of the Animal kingdom, seems utterly destitute of any such power, and is dependent for its support upon organic substances previously elaborated by other beings. But further, the Protophyte obtains its nutriment by mere absorption of liquid and gaseous molecules, which penetrate by simple imbibition; whilst the Protozoon, though destitute of any proper stomach, makes (so to speak) a stomach for itself in the substance of its body, into which it ingests the solid particles that constitute its food, and within which it subjects them to a regular process of digestion. Hence these simplest members of the two kingdoms, which can scarcely be distinguished from each other by any *structural* characters, seem to be *physiologically* separable by the mode in which they perform those actions wherein their life most essentially consists; for the Protococcus-cell decomposes carbonic acid under the influence of light, and generates chlorophyll and proteine compounds, in a manner in all respects comparable to that in which the same operation is performed by the leaf-cells of the most perfect Plant; whilst the


Amœba ingests and digests both Vegetable and Animal food, and applies it to the nutrition of its body, no less effectively than an Animal possessing the most complex digestive and circulating apparatus. And in the present state of our knowledge, we seem justified in laying it down as the most ready and certain differential character we are acquainted with, between the most closely related Protophyta and Protozoa, that the former (with the exception of the Fungi) decompose carbonic acid under the influence of light, and acquire a red or green color from the new compounds which they form in their interior; whilst the latter, having no such power,¹ receive animal and vegetable organisms, or particles of such, into the interior of their bodies, where they extract from them the ready prepared nutriment they are fitted to yield.

264. *Rhizopoda*.—The two creatures above described as the types of the simplest form of Protozoic life, are not now regarded as Infusory Animalcules, among which they were ranked by Prof. Ehrenberg; but are considered as the types of a distinct subdivision, to which the name of *Rhizopoda* has been assigned by M. Dujardin; this name very appropriately representing the leading feature in their organization, which consists in the extension of their sarcode-body into long root-like processes, whereby their aliment is drawn into its substance. In by far the larger proportion of the animals included in this group, a carapace or shell is formed, either by the consolidation of the superficial layer of the sarcode-body through impregnation of its substance with mineral matter, or (as appears to be the case in some instances) by the agglutination of particles of sand, &c., with a viscid secretion exuded from its surface. This "carapace," in *Arcella* (Fig. 193, c, d), has for its basis a layer of dense membrane, which seems analogous to the firm envelope of the Desmidiaceæ (§ 164), and to that which constitutes the basis of the "lorica" of Diatomaceæ (§ 172); and in some species it sends out spinous prolongations like those of the former; whilst in others its surface exhibits symmetrical markings that resemble those of the latter. Its material differs, however, from that of the cellulose wall of Vegetable cells, as may be shown by the effects of reagents; and seems to be rather a modification of horny matter, probably resembling the *chitine* which gives solidity to the integuments of Insects. Not unfrequently it contains particles of sand, minute Diatoms, &c., imbedded in its substance. The *Diffugia* (Fig. 193, a, b) differs in no essential particular from *Arcella*, save in the form of its carapace, which is pitcher-shaped, instead of being shield-like or dish-shaped. In the one case, as in the other, the carapace has but a narrow opening, through which alone can the "pseudopodia" be projected. The general nature of these ani-

¹ Many instances have been cited, of *Animalcules* acquiring a green color by the decomposition of carbonic acid under the influence of light; but there can be no doubt in the mind of any one who is familiar with the results of recent microscopic research, that in all these cases, the supposed *Animalcules* were really *Protophytes*.

mals seems to be exactly the same as that of *Amœba* and *Actinophrys*; and their mode of nutrition differs only in this, that their food can only be drawn into that part of the body which is unprotected by the carapace. Nothing positive is yet known as to their Reproduction. Two individuals may not unfrequently be seen with the apertures of their shells in contact with each other, the pseudopodial prolongations being apparently common to

FIG. 193.



Various forms of Simple Rhizopods:—A, *Diffugia proteiformis*: B, *Diffugia oblonga*: C, *Arcella acuminata*: D, *Arcella dentata*.

both; but whether these are in the act of conjugation, or whether (as the younger aspect of one of the shells sometimes indicates) the union results from the production of a bud not yet separated, cannot yet be certainly affirmed. The most remarkable development of this type of organization presents itself among the marine *Foraminifera* and *Sponges*; in which composite structures, often of very large size, are developed by continuous gemmation, and a complex skeleton is produced. These, not being in themselves (for the most part at least) *microscopic* organisms, will be more fitly considered under a separate head (Chap. X).

265. *Animalcules*.—Dismissing the Rhizopods for the present, we have now to apply ourselves to the special subject of this chapter; namely, the assemblage of minute forms of Animal life, which are commonly known under the designation of Animalcules. Nothing can be more vague or inappropriate than this title, since it only expresses the small dimensions of the beings to which it is applied, and does not indicate any of their characteristic peculiarities. In the infancy of Microscopic knowledge, it was natural to associate together all those creatures which could only be discerned at all under a high magnifying power, and whose internal structure could not be clearly made out with the instruments then in use; and thus the most heterogeneous assemblage of Plants, Zoophytes, minute Crustaceans (water-fleas, &c.), larvæ of Worms and Mollusks, &c., came to be aggregated with the true Animalcules under this head. The class was being gradually limited by the removal of all such forms as could be referred to others; but still very little was known of the real nature of those that remained in it, until the study was taken up by Prof. Ehrenberg, with the advantage of instruments which had derived new and vastly improved capabilities from the application of the principle of Achromatism

(p. 48). One of the first and most important results of his study, and that which has most firmly maintained its ground notwithstanding the weakening of Prof. Ehrenberg's authority in other respects, was the separation of the entire assemblage into two distinct groups, having scarcely any feature in common excepting their minute size, one being of very *low*, and the other of comparatively *high* organization. On the lower group he conferred the designation of *Polygastrica* (many-stomached), in consequence of having been led to form an idea of their organization, which the united voice of the most trustworthy observers now pronounces to be erroneous; and he not only assigned to them a complex digestive apparatus, but considered them to be endowed with genital organs, nervous ganglia, and organs of special sense, for none of which can any adequate basis be found in the appearances that the Microscope brings into view within their bodies. Hence it seems desirable to abandon the term "Polygastrica," as conveying an erroneous idea of the structure of these beings; and we may appropriately fall back on the name *Infusoria*, or Infusory Animalcules, which simply expresses their almost universal prevalence in infusions of organic matter. For although this was applied by the older writers to the higher group as well as to the lower, yet as these are now distinguished by an appropriate appellation of their own, and are, moreover, *not* found in infusions when in that state of rapid decomposition which is most favorable to the presence of the inferior kind of Animalcules, it may very well be withdrawn from them, and be restricted to the Polygastrica of Ehrenberg, which is the sense wherein it has been used by many recent writers. To the higher group, Prof. Ehrenberg's name *Rotifera* or *Rotatoria* is on the whole very appropriate, as significant of that peculiar arrangement of their cilia upon the anterior parts of their bodies, which in some of their most common forms, gives the appearance (when the cilia are in action) of wheels in revolution; the group, however, includes many members, in which the ciliated lobes are so formed as not to bear the least resemblance to wheels. In their general organization, these "Wheel-animalcules" must certainly be considered as members of the *Articulated* division of the Animal Kingdom; and they seem to constitute a class in that lower portion of it, to which the designation *Worms* is now commonly given. Notwithstanding this wide zoological separation between the two kinds of Animalcules, it seems most suitable to the plan of the present work, to treat of them in connection with one another; since the Microscopist continually finds them associated together, and almost necessarily ranges them in his own mind under one and the same category.

266. *Infusoria*.—This term, as now limited by the separation of the Rotifera, is applied to a far smaller range of forms than that which was included by Prof. Ehrenberg under the name of "polygastric" animalcules. For a large section of these, in-

cluding the *Desmidiaceæ*, *Diatomaceæ*, *Volvocineæ*, and many other Protophytes, have been transferred by the almost concurrent voice of those Naturalists whose judgment is most to be relied on, to the Vegetable Kingdom. The *Rhizopod* group, again, must be excluded, as being very distinct in its plan of organization from the true Infusoria. And, lastly, it is not improbable that many of the reputed Infusoria may be but larval forms of some higher organisms, instead of being themselves complete animals.¹ Still an extensive group remains, of which no other account can at present be given, than that the beings of which it is composed go through the whole of their lives, so far as we are acquainted with them, in the condition of isolated cells; differing from Vegetable cells on the one hand, and from Rhizopods on the other, in this remarkable particular,—that the alimentary particles by which they are nourished are taken into their cavity through a distinct oral aperture, some of them also having an anal orifice for the ejection of indigestible matters. It has been imagined by Prof. Ehrenberg, that a distinct alimentary canal exists; sometimes returning upon itself (as in *Vorticella*), sometimes proceeding straight from one extremity of the body to the other (as in *Enchelis*), and sometimes passing round and round in a spiral (as in *Leucophrys*), having a number of flask-shaped stomachs connected with it, into which the alimentary particles find their way, and in which they undergo digestion. But as he made the like assertions with regard to beings that have been since undoubtedly proved to belong to the Vegetable Kingdom, their authority must be explicitly denied; and all the best observers of the present time would agree (the Author believes) in the following general account of the organization of Infusoria.

267. Their bodies consist of "sarcode," of which the outer layer possesses considerably more consistence than the internal portion; the process of differentiation having here advanced sufficiently far to establish a clear distinction between the *wall* and its *contents*. Sometimes, as in *Paramecium*, a distinct pellicle may be recognized on the surface of the proper coat of the body; and this, which is studded with regularly arranged markings like those of *Diatomaceæ*, seems to be the representative of the carapace of *Arcella*, &c. (§ 264), as of the cellulose coat of Protophytes. The form of the body is usually much more definite than that of *Amœba* or *Actinophrys*; each species having its characteristic shape, which is only departed from, for the most part, when the animalcule is subjected to pressure from without,

¹ Professor Agassiz, indeed, goes so far as to assert ("Ann. of Nat. Hist." vol. ii, 1850 p. 157) that, as he has satisfied himself by direct observation, *Byrsaria*, *Paramecium*, and other Animalcules which are commonly accounted as types of this group, are germs of fresh-water worms, some of which he has seen hatched from eggs of *Planaria*. No confirmation has yet been given to this statement by other observers; and until details of these observations shall have been published, it cannot be expected that Zoologists should acquiesce in the entire demolition of this class, which Prof. Agassiz thinks himself warranted by them in proposing.

or when its cavity has been distended by the ingestion of any substance above the ordinary size. The body does not seem to possess much contractile power in its own substance, its movements being principally executed by the instrumentality of locomotive appendages; one remarkable instance of contractility, however, is presented by the stalk of *Vorticella* (§ 268). The locomotive appendages, which may all be considered as prolongations of the tegumentary layer, are destitute of any more minute organization, being, in fact, of the nature of *cilia*, though sometimes of much larger dimensions, and employed in a different manner. The vibration of ciliary filaments, which are either disposed along the entire margin of the body, as well as around the oral aperture (Figs. 194, 195), or are limited to some one part of it, this being always in the immediate vicinity of the

FIG. 194.



FIG. 195.




Fig. 194. *Kerona niturus*:—*a*, contractile cavity; *b*, mouth; *c*, *e*, animalcules swallowed by the *Kerona*, after having themselves ingested particles of indigo.

Fig. 195. *Paramecium caudatum*:—*a*, *a*, contractile cavities; *b*, mouth.


mouth (Fig. 196), is the means by far the most frequently employed by the beings of this class, both for progression through the water, and for drawing alimentary particles into the interior of their bodies. In some their vibration is constant, whilst in others it is only occasional, thus conveying the impression that the Animalcule has a voluntary control over them; but there is strong reason for questioning the existence of any such self-directing power. These cilia, like those of the zoospores of *Protophytes*, can usually be distinctly seen only when their movement is very much slackened in its rate, or when it has entirely ceased. Sometimes, however, instead of a multitude of short cilia, we find a small number of long slender filaments, usually proceeding from the anterior part of the body (that nearest the mouth), and strongly resembling the elongated cilia of *Proto-coccus* (Fig. 68, *h*), or of *Volvox* (Fig. 70, *i*, *k*, *l*). But in other

cases, the filaments are comparatively short, and have a bristle-like firmness; and instead of being kept in vibration, they are moved (like the spines of Echini) by the contraction of the substance to which their bases are attached, in such a manner that the animalcule crawls by their means over a solid surface, as we see especially in *Trichoda lynceus* (Fig. 199, p, q). In *Chilodon* and *Nassula*, the mouth is provided with a circlet of these bristles, which have received the designation of "teeth;" their function, however, is rather that of laying hold of alimentary particles by their expansion and subsequent drawing together (somewhat after the fashion of the tentacula of Zoophytes), than of reducing them by any kind of masticatory process.

268. The modes of movement which Infusory Animalcules execute by means of these instruments, are extremely varied and remarkable. Some propel themselves directly forwards, with a velocity which appears, when thus highly magnified, like that of an arrow, so that the eye can scarcely follow them, whilst others drag their bodies slowly along like a leech. Some attach themselves by one of their long filaments to a fixed point, and revolve around it with great rapidity, whilst others move by undulations, leaps, or successive gyrations; in short, there is scarcely any kind of animal movement which they do not exhibit. There is no sufficient reason, however, to regard such actions as indicative of consciousness; indeed, the very fact that they are performed by the instrumentality of *cilia* seems to imply the contrary; since we know that ciliary action takes place to a large extent in our own bodies, without the least dependence upon our consciousness, and that it is also used as a means of dispersion among the zoospores of the lowest Plants, which cannot for a moment be supposed to be endowed with this attribute. We can only regard it, therefore, as indicative of a wonderful adaptation, on the part of these simple ciliated cells, to a kind of life which enables them to go in quest of their own nutriment, and to introduce it, when obtained, into the interior of their bodies. The curious contraction of the footstalk of the *Vorticella*, however, is a movement of a very different nature, and is due to the contractility of the tissue that occupies the interior of the tubular pedicle. This stalk serves to attach the bell-shaped body of the Animalcule to some fixed object, such as the leaf or stem of duckweed; and when the animal is in search of food, with its cilia in active vibration, the stalk is fully extended. If, however, the Animalcule should have drawn to its mouth any particles too large to be received within it, or should be touched by any other that happens to be swimming near it, or should be "jarred" by a smart tap on the stage of the microscope, the stalk suddenly contracts into a spiral, from which it shortly afterwards extends itself again into its previous condition. The central cord to whose contractility this action is due, has been described as muscular; but it does not possess the characteristic structure of either

kind of muscular fibre, and is probably nothing else than a portion of sarcode specially endowed with this property. Nothing but

FIG. 196.



Group of *Vorticella nebulifera*. showing A, the ordinary form; B, the same with the stalk contracted; C, the same with the bell closed; D, E, F, successive stages of fissiparous multiplication.

the rapidity of its contraction and relaxation differentiates it from the pseudopodia of the Rhizopods. There is no reason ~~what~~ ever to believe that these Animalcules possess any organs of special sense. The red spots which may be seen in many of them, and which have been designated as eyes by Prof. Ehrenberg, from their supposed correspondence with the eyespots of *Rotifera* (§ 278), really bear a much greater resemblance to the red spots which are so frequently seen among Protophytes (§ 153). If they are really endowed with consciousness, as their movements *seem* to indicate, though other considerations render it very doubtful, they must derive their perceptions of external things from the impressions made upon their general surface, but more particularly upon their filamentous appendages.

269. The interior of the body does not always seem to consist of a simple undivided cavity, occupied by soft "sarcodé;" for the tegumentary layer appears in many instances to send prolongations across it in different directions, so as to divide it into chambers of irregular shape, freely communicating with each other, which may be occupied either by sarcodé, or by particles introduced from without. The alimentary particles which can be distinguished in the interior of the transparent bodies of Infusoria, are usually Protophytes of various kinds, either entire or in a fragmentary state. The Diatomaceæ seem to be the ordinary food of many; and the insolubility of their *loricæ* enables the observer to recognize them unmistakably. Sometimes entire Infusoria are observed within the bodies of others not much exceeding them in size (Fig. 199, B); but this is only when they have been recently swallowed, since the prey speedily undergoes digestion. It would seem as if these creatures do not feed by any means indiscriminately; since particular kinds of them are attracted by particular kinds of aliment; the crushed bodies and eggs of Entomostracea, for example, are so voraciously consumed by the *Coleps*, that its body is sometimes quite altered in shape by the distension. This circumstance, however, by no means proves, as some have considered it to do, that such creatures possess a sense of taste and a power of determinate


selection; for many instances might be cited, in which actions of the like apparently conscious nature are performed without any such guidance. The ordinary process of feeding, as well as the nature and direction of the ciliary currents, may be best studied by diffusing through the water containing the Animalcules, a few particles of indigo or carmine. These may be seen to be carried by the ciliary vortex into the mouth, and their passage may be traced for a little distance down a short (usually ciliated) oesophagus. There they commonly become aggregated together, so as to form a little pellet of nearly globular form; and this, when it has attained the size of the hollow within which it is moulded, is projected into the "general cavity of the body," where it lies in a vacuole of the sarcode, its place in the oesophagus being occupied by other particles subsequently ingested. This "moulding," however, is by no means universal; the aggregations of colored particles in the bodies of these animals being often destitute of any regularity of form. One after another of such particles being thus introduced into the interior of the body, each aggregation seems to push on its predecessors; and a kind of circulation is thus occasioned in the contents of the cavity. The pellets that first entered make their way out after a time (after yielding up their nutritive materials), generally by a distinct anal orifice, sometimes, however, by any part of the surface indifferently, and sometimes by the mouth. A circumstance which seems clearly to indicate that they cannot be enclosed (as Prof. Ehrenberg maintains) in distinct stomachal cavities, is that, when the pellets are thus moving round the body of the Animalcule, two of them sometimes appear to become fused together, so that they obviously cannot have been separated by any membranous investment. When the Animalcule has not taken food for some time, "vacuoles" or clear spaces, extremely variable both in size and number, filled only with a very transparent fluid, are often seen in its sarcode; their fluid sometimes shows a tinge of color, and this seems to be due to the solution of some of the vegetable chlorophyll upon which they may have fed last.

270. Contractile vesicles (Figs. 194, 195, *a, a*), usually about the size of the "vacuoles," are found, either singly or to the number of from two to sixteen, in the bodies of most Animalcules; and may be seen to execute rhythmical movements of contraction and dilatation at tolerably regular intervals, being so completely obliterated when emptied of their contents, as to be quite indistinguishable, and coming into view again as they are refilled. These vesicles do not change their position in the individual, and they are pretty constant, both as to size and place, in different individuals of the same species; hence they are obviously quite different in character from the "vacuoles." What is their purpose in the economy of these creatures can be only vaguely guessed at; it may be surmised to be the diffusion

through the body of the liquid product of the digestive operation,—a surmise which seems in some degree justified by their unusual complexity in *Paramecium*. For each of its two globular vesicles (Fig. 195, *a, a*), is surrounded by several elongated cavities, arranged in a radiating manner, so as to give to the whole somewhat of a starlike aspect; and the liquid contents are seen to be propelled from the former into the latter, and *vice versa*.

271. Of the Reproduction of the Infusoria, our knowledge is at present very limited; the attention of observers having, until a comparatively recent period, been fixed almost exclusively upon the act of duplicative subdivision, which, though by far the most frequent method of propagation, is not a true generative operation. It is effected in the same general mode as the subdivision of Protophyta; and has been observed in many instances to commence in the "nucleus" which may usually be distinguished in the cell-bodies of the Infusoria. The division takes place in some species longitudinally, that is, in the direction of the greatest length of the body (Fig. 196, *D, E, F*), in other species transversely (Fig. 199, *A, D*), whilst in some, as in *Chilodon cucullulus* (Fig. 197), it seems to occur in either direc-

FIG. 197.




Fissiparous multiplication of *Chilodon cucullulus*:—*A, B, C*, successive stages of longitudinal fission; *D, E, F*, successive stages of transverse fission.

tion indifferently, though there may not improbably be an alternation, as there usually is in the direction of the subdivision of the component cells of masses that are increasing both in length and in breadth (§ 195). This operation is performed with such rapidity, under favorable circumstances, that, according to the calculation of Prof. Ehrenberg, no fewer than 268 millions might be produced in a month by the repeated subdivisions of a single *Paramecium*. When this fission occurs in *Vorticella* (Fig. 196), one of the divisions is usually smaller than the other, sometimes so much so as to look like a bud; and this usually detaches itself when mature from the main body, and swims freely about until it develops a new footstalk for itself. But sometimes the two parts are equal in size, and the fission extends down the stalk, which thus becomes double for a greater or less part of its length; and thus a whole bunch of Vorticellæ may spring (by a repetition of the same process) from one base. In some members of the same family, indeed, an arborescent structure is produced by the like processes of division and gemmation.

272. The recent observations of Stein and other Microscopists have drawn attention to another very curious mode of propagation, the phenomena of which were completely misapprehended by Prof. Ehrenberg. Many Infusoria at certain times undergo an *encysting* process; that is, their body secretes from its surface a sort of gelatinous case, which hardens so as completely to enclose it, the Animalcule, however, still remaining free in the interior of its coffin-like investment. Previously to the formation of this cyst, the Animalcule loses its activity, its form becomes more rounded, and its cilia or other filamentous prolongations are lost or retracted, as is well seen in *Vorticella* (Fig. 198, A);

and it was not, perhaps, very unnatural, that the encysting process should have been considered by Prof. Ehrenberg as the expiring effort of life. If the cysts and their contents, however, be attentively watched, it will be seen that the process is preliminary to a production of new individuals; and this may take place in different modes. For sometimes the substance of the body appears to break up (C, D) into numerous "gemmules," which are analogous to the "zoospores" of Protophytes, and which, like them, are set free by the bursting of the parent cell (E), swimming forth to develop themselves into new individuals of the same kind, though at first perhaps bearing no resemblance to the type of their predecessor. In other instances, however, only a single offspring is developed from the "nucleus" of the original cell-

body, which offspring may have an entirely dissimilar form; and this latter change occurs in *Vorticella*, in conjunction with other




Development and Metamorphosis of *Vorticella microstoma*: —A, full-grown individual in its encysted state; a, retracted oval circlet of cilia; b, nucleus; c, contractile vesicle;—B, a cyst separated from its stalk;—C, the same more advanced, the nucleus broken up into spore-like globules;—D, the same more developed, the original body of the *Vorticella*, d, having become sacculated, and containing many clear spaces;—E, one of the sacculations having burst through the enveloping cyst, a gelatinous mass, e, containing the spores, is discharged;—F, transformation of encysted *Vorticella* (z) into form of *Acineta*; b, nucleus;—G, stalked *Acineta* form of *Vorticella*, enclosing a young one, the result of the transformation of the nucleus;—H, young free *Vorticella*; a, b, c, as in Fig. 1; g, posterior circlet of cilia.

peculiarities of a very remarkable kind. For the encysted Vorticella becomes changed into the form of an *Acineta* (closely resembling that of *Actinophrys*), as shown in Fig. 198, *r*; and this may acquire a new stalk, so as to correspond with a *Podophrys* (*g*). At the same time, its band-like nucleus becomes entirely metamorphosed into a free body of ovate form, which carries at its narrower end a cirlet of long vibrating cilia, while its more obtuse end is perforated by a mouth which communicates with a distinct oral cavity. In the interior of this offspring (*h*), we already observe a long oval nucleus (*b*) and a round contractile vesicle (*c*), and its whole aspect is that of a young Vorticella-bud just ready to quit its stock. This body escapes from the interior of the *Acineta* by a gap formed in some part of its wall, which, however, soon closes again, and the *Acineta* goes on stretching out and retracting its radiating filaments, and after a time produces in its interior a new nucleus for a second Vorticella-bud.¹

278. Another interesting series of phenomena, of the same order, is presented in the development of the Animalcule designated by Müller as *Trichoda lynceus*, which has been carefully studied by M. Jules Haime.² The form which seems most pro-

FIG. 199.



Metamorphoses of *Trichoda lynceus*:—A, larva (*Oxytricha*); B, a similar larva, after swallowing the animalcule represented at *m*; C, a very large individual on the point of undergoing fission; D, another in which the process has advanced further; E, one of the products of such fission; F, the same body become spherical and motionless; G, aspect of this sphere fifteen days afterwards; H, later condition of the same, showing the formation of the cyst; I, incipient separation between living substance and exuvial matter; K, partial discharge of the latter, with flattening of the sphere; L, more distinct formation of the confined animal; M, its escape from the cyst; N, its appearance some days afterwards; O, more advanced stage of the same; P, Q, perfect individuals, one as seen sideways, moving on its bristles, the other as seen from below; these are magnified twice as much as the preceding figures.

perly to be considered as the larval one, is that shown in Fig. 199, A-E, which has been described by Prof. Ehrenberg under

¹ See Prof. Stein's Memoir in "Siebold and Kolliker's Zeitschrift," Bd. iii, and the translation of it in "Ann. of Nat. Hist." 2d Ser. vol. ix, p. 471.

² Annales des Sci. Nat." Sér. 3, tom. xix, p. 109.

the name of *Oxytricha*. This possesses a long, narrow, flattened body, furnished with cilia along the greater part of both margins, and having also at its two extremities a set of larger and stronger hair-like filaments; and its mouth, which is an oblique slit on the right hand side of its fore part, has a fringe of minute cilia on each lip. Through this mouth, large particles are not unfrequently swallowed, which are seen lying in the midst of the gelatinous contents of the general cavity of the body, without any surrounding "vacuole;" and sometimes even an animalcule of the same species, but in a different stage of its life, is seen in the interior of one of these voracious little devourers (B). In this phase of its existence, the *Trichoda* undergoes multiplication by transverse fission, after the ordinary mode (C, D); and it is usually one of the short-bodied "doubles" thus produced (E), that passes into the next phase. This consists in the assumption of the globular form, and the almost entire loss of the locomotive appendages (F); in the escape of successive portions of the granular sarcode, so that "vacuoles" make their appearance (G); and in the formation of a gelatinous envelope or cyst, which, at first soft, afterwards acquires increased firmness (H). After remaining for some time in this condition, the contents of the cyst become clearly separated from their envelope; and a space appears on one side, in which ciliary movement can be distinguished (I). This space gradually extends all round, and a further discharge of granular matter takes place from the cyst, by which its form becomes altered (K); and the distinction between the newly formed body to which the cilia belong, and the effete residue of the old, becomes more and more apparent (L). The former increases in size, whilst the latter diminishes; and at last the former makes its escape through an aperture in the wall of the cyst, a part of the latter still remaining within its cavity (M). The body thus discharged (N) does not differ much in appearance from that of the *Oxytricha* before its encystment (F), though only of about two-thirds its diameter; but it soon develops itself (O, P, Q) into an Animalcule very different from that in which it originated. First it becomes still smaller, by the discharge of a portion of its substance; numerous very stiff bristle-like organs are developed, on which the animalcule creeps, as by legs, over solid surfaces; the external integument becomes more consolidated on its upper surface, so as to become a kind of carapace; and a mouth is formed by the opening of a slit on one side, in front of which is a single hair-like filament, which is made to turn round and round with great rapidity, so as to describe a sort of inverted cone, whereby a current is brought towards the mouth. This latter form has been described by Prof. Ehrenberg under the name of *Aspidisca*. It is very much smaller than the larva; the difference being, in fact, twice as great as that which exists between A and P, Q (Fig. 199), since the two last figures are drawn under a magnifying power twice as great as that employed for

the preceding. How the *Aspidisca*-form in its turn gives origin to the *Oxytricha*-form, has not yet been made out. A sexual process, it may be almost certainly concluded, intervenes somewhere; but other transformations may not improbably take place, before the latter of these types is reproduced.

274. A like succession of phenomena has been observed among several other forms of Infusoria; so that, considering the strong general resemblance in kind and degree of organization which prevails throughout the group, it does not seem unlikely that the "encysting process" occurs at some stage of the life of nearly all these Animalcules, just as the "still" condition alternates with the "motile" in the most active Protophytes (§§ 152-6). And it is not improbably in the "encysted" condition that their dispersion takes place; since they have been found to endure desiccation in this state, although, in their ordinary condition of activity, they cannot be dried up without loss of life. When this circumstance is taken into account, in conjunction with the extraordinary rapidity of multiplication of these Animalcules, and with the fact that a succession of different forms may be presented by one and the same being, the difficulty of accounting for the universality of their diffusion, which has led some Naturalists to believe in their "spontaneous generation," and others to regard them as isolated particles of higher organisms set free in their decomposition so as to constitute an "equivocal generation," is as readily got over as we have seen it to be in the case of the Fungi (§ 211). Although it may be stated as a general fact, that wherever decaying organic matter exists in a liquid state, and is exposed to air and warmth, it speedily becomes peopled with these minute inhabitants, yet it has been experimentally proved by Prof. Schultz, that perfectly *free* access of air to such infusions, is essential to the appearance of Animalcules in them. For having kept infusions of decaying animal and vegetable matter, in air which had been filtered (so to speak) of any floating germs it might contain, by passing through either a red-hot tube or strong sulphuric acid, he found that no Animalcules made their appearance under these circumstances, even after the lapse of several weeks; although they were seen in abundance after the free exposure of the same infusions to the atmosphere for a few hours only. Hence it may be fairly inferred, that, as seems to be the case with the Fungi, the dried cysts or the germs of Infusoria are everywhere floating about in the air, ready to develop themselves wherever the appropriate conditions are presented; and all our knowledge of their history, as well as the strong analogy of the Fungi, seems further to justify the belief, that the same germs may develop themselves into several different forms, according to the nature of the liquid in which they chance to be deposited. This is a subject peculiarly worthy of the attention of Microscopic observers; who can scarcely be better employed than in tracing out the succes-



sion of phases which any particular type may present, and thus in making a most important extension of our knowledge of its life-history, whilst at the same time effecting a most desirable reduction in the number of reputed species. Such a study, if perserveringly as well as carefully pursued, would be almost certain to lead to the discovery of the true generative process in these creatures, of which we cannot at present be said to know anything with certainty. For although an apparent "conjugation" has been observed between the bodies of distinct individuals, both in the ordinary active states of Infusoria, and in their *Acineta* condition, yet this does not seem truly to represent the sexual union of higher animals; for the blending of the two individuals is not so complete as to amount to the "fusion" we have seen to occur in previous instances, the boundaries of each remaining distinctly traceable; and not only two, but three, four, or even more, have been found in continuous union.

275. It is obvious that no Classification of Infusoria can be of any permanent value, until it shall have been ascertained, by the study of their entire life-history, what are to be accounted really distinct forms; and the differences between them, consisting chiefly in the shape of their bodies, the disposition of their cilia, the possession of other locomotive appendages, the position of the mouth, the presence of a distinct anal orifice, and the like, are matters of such trivial importance as compared with those leading features of their structure and physiology on which we have been dwelling, that it does not seem desirable to attempt in this place to give any account of them. The most remarkable departure from the ordinary type is presented by the *Vorticellinæ*, the habit of which is to attach themselves to the stems of aquatic plants or some other supports, either by the apex of their own conical body,—as is the case with *Stentor*, one of the largest of all Infusoria (being visible to the naked eye), which is very common in ponds and ditches, attaching itself to duckweed, decaying reeds, or other floating bodies, round which it forms a sort of slimy fringe, but which is often found swimming freely, its trumpet-shaped body drawn together into the form of an egg,—or by a footstalk several times its own length, as is the case with *Vorticella* (Fig. 196), which also occasionally quits its attachment (the stalk apparently dying and being thrown off), and swims rapidly through the water, being propelled by the fringe of cilia, which, when the body was fixed by its stalk, served to produce a vortex in the surrounding fluid, that brought it both food and air. Another curious departure from the ordinary type is presented by the family *Ophrydinæ*; the animalcules of which, closely resembling some *Vorticellinæ* in their individual structure, are usually found imbedded in a gelatinous mass, of a greenish color, which is sometimes adherent, sometimes free, and may attain the diameter of four or five inches, presenting such a strong general resemblance to a mass of *Nostoc*



(§ 196) or even of Frogs' spawn, as to have been mistaken for such. The mode in which these masses are produced closely resembles that in which the masses of *Mastoloia* (§ 189) or of *Palmella* (§ 194) are formed; since they simply result from the fact, that the multitude of individuals produced by a repetition of the process of self-division, remain connected with each other for a time by a gelatinous exudation from the surface of their cell-bodies, instead of at once becoming completely isolated. From a comparison of the dimensions of the individual *Ophrydia*, each of which is about 1-120th of an inch in length, with those of the composite masses, some estimate may be formed of the number included in the latter; for a cubic inch would contain nearly *eight millions* of them, if they were closely packed; and many times that number must exist in the larger masses, even making allowance for the fact that the bodies of the animalcules are separated from each other by their gelatinous cushion, and that the masses have their central portions occupied only by water. Hence we have, in such clusters, a distinct proof of the extraordinary extent to which multiplication by duplicative subdivision may proceed, without the interposition of any other operation. These animalcules, however, free themselves at times from their gelatinous bed; and have been observed to undergo an "encysting process," corresponding with that of the *Vorticellinæ* (§ 272). It is much to be desired that Microscopic observers should devote themselves systematically to the continuous study of even the commonest and best known forms of these Animalcules; since there is *not a single one*, whose entire life-history, from one Generative act to another, is known to us. And since it cannot be even guessed at, without such knowledge, what, among the many dissimilar forms that have been described by Prof. Ehrenberg and others, are to be accounted as truly distinct species, and what are mere phases in the existence of others that are perhaps very dissimilar to them in aspect, it is obvious that no credit is really to be gained by the discovery of any number of apparently new species, which shall be at all comparable with that to be acquired by the complete and satisfactory elucidation of the life-history of any one.¹

276. As it is among Animalcules that the action of the organs

¹ The recent memoirs of Prof. Stein in "Wiegmann's Archiv," and in "Siebold and Kölliker's Zeitschrift," which have been separately published in a collective form "Die Infusionsthiere," Leipzig, 1854, contain most valuable contributions to the desiderated knowledge; as do also the memoirs of Dr. Cohn in "Siebold and Kölliker's Zeitschrift." The great work of Prof. Ehrenberg, "Die Infusionsthierchen," will always remain a standard of reference, as the basis of all our higher knowledge of these organisms; but its authority has been completely destroyed by the proof which has been gradually accumulating, of the erroneous conceptions on which he based his descriptions; *first*, as to what *are* Infusory Animalcules, a very large proportion of those considered by him as such being undoubtedly Plants;—*second*, as to the organization of the true Infusoria, the complex digestive apparatus, genital organs, nervous system, and organs of sense, which he believed that he had discovered in them, having no existence but in his imagination;—and, *third*, as to the relations of his supposed species one to another, many of these being only different states of the same.

termed *cilia* has the most important connection with the vital functions, it seems desirable to introduce here a more particular notice of them. They are always found in connection with *cells*, of whose substance, as we have seen among the *Protophyta* (§§ 154, 158), they may be considered as extensions. The form of the filaments is usually a little flattened, and tapering gradually from the base to the point. Their size is extremely variable; the largest that have been observed being about 1-500th of an inch in length, and the smallest about 1-13,000th. When in motion, each filament appears to bend from its root to its point, returning again to its original state, like the stalks of wheat when depressed by the wind; and when a number are affected in succession with this motion, the appearance of progressive waves following one another is produced, as when a wheatfield is agitated by successive gusts. When the ciliary action is in full activity, however, little can be distinguished save the whirl of particles in the surrounding fluid; but the *back-stroke* may often be perceived, when the *forward-stroke* is made too quickly to be seen; and the real direction of the movement is then opposite to the apparent. In this back-stroke, when made slowly enough, a sort of "feathering" action may be observed; the thin edge being made to cleave the liquid, which has been struck by the broad surface in the opposite direction. It is only when the rate of movement has considerably slackened, that the shape and size of the cilia, and the manner in which their stroke is made, can be clearly seen. It has been maintained by some, that the action of the cilia is muscular; but they are often too small to contain even the minutest fibrillæ of true muscular tissue, and no such elements can be discerned around their base; their presence in Plants, moreover, seems distinctly to negative such an idea. Hence we must consider them as organs *sui generis*, wherein the contractility of the cell to which they belong, is (as it were) concentrated. We have seen that in the *Rhizopods*, the entire mass of whose sarcode is highly contractile, no cilia are present; whilst in the Infusoria, whose bodies have comparatively little contractility, the movements are delegated to the cilia. Cilia are not confined, however, to Animalcules and Zoophytes, but exist on some of the free internal surfaces, especially the walls of the respiratory passages, of all the higher animals, not excepting Man himself. Our own experience assures us that their action takes place, not only without any exercise of will on our own parts, but even without affecting our *consciousness*; and it has been found to continue for many hours, or even days, after the death of the body at large. How far it is subject to any conscious control on the part of these Animalcules, in which the cilia serve as instruments for locomotion as well as for bringing to them food or oxygen, it is impossible for any one to say with confidence. In this important respect, however, the ciliary movement of Animalcules differs from that which is

observable in the higher Animals,—that whilst in the latter it is *constant*, giving the idea of purely mechanical agency, in the former it is so interrupted and renewed, as almost necessarily to suggest to the observer the notion of choice and direction.

277. *Rotifera, or Wheel-Animalcules.*—We now come to that higher group of Animalcules, which, in point of complexity of organization, is as far removed from the preceding, as Mosses are from the simplest Protophytes; the only point of real resemblance between the two groups, in fact, being the minuteness of size which is common to both, and which was long the obstacle to the recognition of the comparatively elevated character of the Rotifera, as it still is to the precise determination of certain points of their structure. Some of the Wheel-Animalcules are inhabitants of salt water only, but by far the larger proportion are found in collections of fresh water, and rather in such as are free from actively decomposing matter, than in those which contain organic substances in a putrescent state. Hence when they present themselves in vegetable infusions, it is usually after that offensive condition, which is favorable to the development of many of the Infusoria, has passed away; and they are consequently to be looked for, after the disappearance of many successions (it may be) of Animalcules of inferior organization. Rotifera are more abundantly developed in liquids which have been long and freely exposed to the open air, than in such as have been kept under shelter; certain kinds, for example, are to be met with in the little pools left after rain in the hollows of the lead with which the tops of houses are partly covered; and they are occasionally found in enormous numbers in cisterns which are not beneath roofs or otherwise covered over.¹ They are not, however, absolutely confined to collections of liquid; for there are a few species which can maintain their existence in damp earth; and the common *Rotifer* is occasionally found in the interior of the leaf-cells of *Sphagnum* (§ 216). The wheel-like organs from which the class derives its designation, are most characteristically seen in the common form just mentioned (Fig. 201), where they consist of two disk-like lobes or projections of the body, whose margins are fringed with long cilia; and it is the uninterrupted succession of strokes given by these cilia, each row of which nearly returns (as it were) into itself, that gives rise by an optical illusion to the notion of “wheels.” This arrangement, however, is by no means universal; in fact, it obtains in only a small proportion of the group; and by far the more general plan is that seen in Fig. 200, in which the cilia form one continuous line across the body, being disposed upon the sinuous edges of certain lobes or projections which are borne upon its anterior portion. Some of the chief departures from this plan will be noticed hereafter (§ 281). The great transparency of the Rotifera permits their general structure to


¹ See a remarkable instance of this in p. 253, *note*.

be easily recognized. They have usually an elongated form, similar on the two sides; but this rarely exhibits any traces of segmental division. The body is covered with a double envelope; both layers of which are extremely thin and flexible in some species; whilst in others, the outer one seems to possess a horny consistence. In the former case, the whole integument is drawn together in a wrinkled manner when the body is shortened; in some of the latter the sheath has the form of a polyp-cell, and the body lies loosely in it, the inner layer of the integument being separated from the outer by

a considerable space (Fig. 202); whilst in others, the envelope or *lorica* is tightly fitted to the body, and strongly resembles the horny casing of an insect or the shell of a crab, except that it is not jointed, and does not extend over the head and tail, which can be projected from the openings at its extremities, or completely drawn within it for protection (Fig. 203). In those Rotifera in which the flexibility of the body is not interfered with by the consolidation of the external integument, we usually find it capable of great variation in shape, the elongated form being occasionally exchanged for an almost globular one, as is seen especially when the animals are suffering from deficiency of water; whilst by alternative movements of contraction and extension, they can make their way over solid surfaces, after the manner of a worm or a leech, with considerably activity,—some even of the


loricated species being rendered capable of this kind of progression, by the contractility of the head and tail. All these, too, can swim readily through the water, by the action of their cilia; and there are some species which are limited to the latter mode of progression. The greater number have an organ of attachment at the posterior extremity of the body, which is usually prolonged into a tail, by which they can affix themselves to any solid object; and this is their ordinary position, when keeping their wheels in action for a supply of food or of water. They have no difficulty, however, in letting go their hold and moving through the water in search of a new attachment, and may therefore be considered as perfectly free; the polypoid species, however, remain attached by the posterior extremity to the spot on which they have at first fixed themselves, and their cilia are consequently employed for no other purpose than that of creating currents in the surrounding water.

FIG. 200.


Brachionus pala.

278. In considering the internal structure of Rotifera, we shall take as its type the arrangement which it presents in the *Rotifer vulgaris* (Fig. 101); and specify the principal variations exhibited by others. The body of this animal, when fully extended, possesses greater length in proportion to its diameter than that of most others of the class; and the tail is composed of three joints or segments, which are capable of being drawn up, one within another, like the sliding-tubes of a telescope, each having a pair of prongs or points at its extremity. Within the external integument of the body, are seen a set of longitudinal muscular bands (*h*), which serve to draw the two extremities towards each other; and these are crossed by a set of transverse annular bands, which also are probably muscular, and serve to diminish the diameter of the body, and thus to increase its length. Between the wheels is a prominence, bearing two red spots (*b*), supposed to be rudimentary eyes, and having the mouth (*a*) at its extremity; this prominence may be considered, therefore, as a true head, notwithstanding that it is not clearly distinguishable from the body. This head also bears upon its under surface a projecting tubular organ (*d*), which was thought by Professor Ehrenberg to be a siphon for the admission of water to the cavity of the body for the purpose of respiration; this, however, is certainly not the case, the tube being imperforate at its extremity; and there seems much more probability in the idea of Dujardin, that it represents the *antennæ* or *palpi* of higher Articulata, the single organ being replaced in many Rotifera by a pair, of which each is furnished at its extremity with a brush-like tuft of hairs that can be retraced into the tube. The œsophagus, which is narrow in the *Rotifer*, but which is dilated into a crop in *Stephanoceros* (Fig. 202) and in some other genera, leads to the masticating apparatus, which in these animals is placed far behind the mouth, and in close proximity to the stomach. It consists of a pair of stirrup-shaped jaws (Fig. 201, *e*) each having from one to five teeth (in *Rotifer*, two), which appear to contain mineral matter and to be of harder texture than the rest of the fabric; these jaws are put in action by powerful muscles, and are so moved that all the food which passes into the stomach is subject to be divided and torn by their teeth. In many Rotifera, the conformation of this masticating apparatus is extremely complicated. The form of the alimentary canal varies; this being sometimes a simple tube, passing without enlargement or constriction from the masticating apparatus to the anal orifice at the posterior part of the body; whilst in other instances there is a marked distinction between the stomach and intestinal tube, the former being a large globular dilatation immediately below the jaws, whilst the latter is cylindrical and comparatively small. The alimentary canal of *Rotifer* most resembles the first of these types, but presents a dilatation (*l*) close to the anal orifice, which may be considered as a cloaca; that of *Brachionus* (Fig. 200) is rather formed upon

the second. Connected with the alimentary canal are various glandular appendages, more or less developed; sometimes clustering round its walls as a mass of separate follicles, which seems to be the condition of the glandular investment (*g*) of the alimentary canal in *Rotifer*; in other cases having the form of cæcal tubuli. Some of these open into the stomach close to the termination of the œsophagus, and have been supposed to be salivary or pancreatic in their character; whilst others, which discharge their secretion into the intestinal tube, have been regarded, and probably with correctness, as the rudiment of a liver. In a curious animalcule of this class, minutely described by Mr. Dalrymple,¹ although the mouth, masticating apparatus, and stomach are constructed upon the regular type of the genus *Notommata*, to which it seems nearly allied, yet there is neither intestine nor anal orifice, and the indigestible matters are rejected through the mouth. This, so far as is yet known, is a solitary example of the existence of this character of degradation in the class Rotifera. There does not appear to be any special circulating apparatus in these animals; but the fluid which is contained in the general cavity of the body, between the exterior of the alimentary canal and the inner tegumentary membrane, is probably to be regarded as nutritive in its character; and its aeration is provided for by a peculiar apparatus, which seems to be a rudimentary form of the "water vascular system," that attains a high development in the class of Worms. On either side of the body there is usually to be observed a long flexuous tube (Fig. 200), which extends from a contractile vesicle common to both and opening into the cloaca (Fig. 201, *i, i*), towards the anterior region of the body, where it frequently subdivides into branches, one of which may arch over towards its opposite side, and inosculate with a corresponding branch from its tube. Attached to each of these tubes are a number of peculiar organs (usually from two to eight on each side), in which a trembling movement is seen, very like that of a flickering flame; these appear to be pear-shaped sacs, attached



Rotifer vulgaris, as seen at A, with the wheels drawn in, and at B with the wheels expanded; a, mouth; b, eye-spots; c, wheels; d, calcar (antenna?); e, jaws and teeth; f, alimentary canal; g, glandular (?) mass enclosing it; h, longitudinal muscles; i, i, tubes of water vascular system; k, young animal; l, cloaca.

¹ "Philos. Transact." 1849, p. 339.

by hollow stalks to the main tube, having a long cilium in the interior of each, that is attached by one extremity to the interior of the sac, and vibrates with a quick undulatory motion in its cavity; and there can be little doubt that their purpose is to keep up a constant movement in the contents of the aquiferous tubes, whereby fresh water may be continually introduced from without, for the aeration of the fluids of the body.¹ There is much uncertainty with regard to the structures which Prof. Ehrenberg has described as ganglia and nerves; and it seems doubtful if there is more than a single nervous centre in the neighborhood of the single, double, or multiple red spots, which are seen upon the head of the Rotifera, and which, corresponding precisely in situation with those that in the higher Articulata are unquestionably eyes, are probably to be regarded as rudiments of visual organs.

279. The Reproduction of the Rotifera has not yet been completely elucidated. There is no instance, in this group, in which multiplication by gemmation or spontaneous fission is certainly known to take place; but the occurrence of clusters formed by the aggregation of a number of individuals of *Conochilus*, adherent by their tails, and enclosed within a common *lorica*, would seem to indicate that these clusters, like the aggregations of Polygastrica, Bryozoa, and Tunicata, must have been formed by continuous growth from a single individual. The ordinary method of multiplication, however, is commonly supposed to be by a proper generative act; as distinct sexes have been discovered in several individuals, and the act of sexual union has been witnessed. The condition of the male of the remarkable genus described by Mr. Dalrymple (loc. cit.) is a most extraordinary one; for it possesses no mandibles, pharynx, œsophagus, stomach, nor hepatic glands; having, in fact, no other organs fully developed, than those of generation. It would appear, therefore, quite unfit to obtain aliment for itself; and its existence is probably a very brief one, being continued only so long as the store of nutriment supplied by the egg remains unexhausted. In *Rotifer*, however, as in by far the larger proportion of the class, no males have been discovered; and it remains doubtful whether the two sexes are united in the same individual, or whether the males are produced only at certain times. The female organ consists but of a single ovarian sac, which frequently occupies a large part of the cavity of the body, and which opens at its lower end by a narrow orifice into the cloaca. Although the number of eggs in these animals is so small, yet the rapidity with which the whole process of their development and maturation is accomplished, renders the multiplication of the race very rapid. The egg of the *Hydatina* is extruded from the cloaca within a few


¹ See Mr. Huxley's account of these organs, in his description of *Lacinularia socialis*, "Transact of Microsc. Soc." Ser. 2, vol. i. Other observers have supposed that the pyriform sacs communicate with the general cavity of the body; but the Author has much confidence in the correctness of Mr. Huxley's statements on this point.

hours after the first rudiment of it is visible; and within twelve hours more the shell bursts, and the young animal comes forth. In the *Rotifer* and several other genera, the development of the embryo takes place whilst the egg is yet retained within the body of the parent (Fig. 201, *k*), and the young are extruded alive; whilst in some other instances, the eggs, after their extrusion, remain attached to the posterior extremity of the body (Fig. 200), until the young are set free. In general it would seem that, whether the rupture of the egg-membrane takes place before or after the egg has left the body, the germinal mass within it is developed at once into the form of the young animal, which resembles that of its parent; no preliminary metamorphosis being gone through, nor any parts developed which are not to be permanent. The transparency of the egg-membrane, and also of the tissues, of the parent Rotifer, allows the process of development to be watched, even when the egg is retained within the body; and it is curious to observe, at a very early period, not merely the red eye spot of the embryo, but also a distinct ciliary movement. The multiplication of *Hydatina* (in which genus three or four eggs are deposited at once, and their development completed out of the body) takes place so rapidly, that, according to the estimate of Prof. Ehrenberg, nearly *seventeen millions* may be produced within twenty-four days from a single individual. Even in those species which usually hatch their eggs within their bodies, a different set of ova is occasionally developed, which are furnished with a thick glutinous investment: these, which are extruded entire, and are laid one upon another, so as at last to form masses of considerable size in proportion to the bulk of the animals, seem not to be destined to come so early to maturity, but very probably remain dormant during the whole winter season, so as to produce a new brood in the spring. These "winter eggs" are inferred by Mr. Huxley, from the history of their development, to be really *gemmae* produced by a non-sexual operation; while the bodies commonly called ova, he considers to be true generative products. Dr. Cohn has recently informed the Author, however, that he has ascertained by direct experiment upon those species in which the sexes are distinct, that the bodies commonly termed ova (Figs. 200, 201), are really *internal gemmae*, since they are reproduced, through many successions, without any sexual process, just like the external *gemmae* of *Hydra* (§ 301), or the internal *gemmae* of *Entomostraca* and *Aphides* (Chap. XVI). And this view appears to himself to be more accordant with general physiological analogy than that of Mr. Huxley; since, in the other instances referred to, as in the Rotifera, the multiplication by gemmation goes on rapidly whilst food and warmth are abundantly supplied; but gives place to the true generative process, when the nutritive activity is lowered by their withdrawal.

280. Certain Rotifera, among them the common Wheel-Ani-

malcule, are remarkable for their tenacity of life, even when reduced to the state of most complete dryness; for they can be kept in this condition for any length of time, and will yet revive very speedily upon being moistened. Experiments have been carried still farther with the allied tribe of *Tardigrades*; individuals of which have been kept in a vacuum for thirty days, with sulphuric acid and chloride of calcium (thus suffering the most complete desiccation that the Chemist can effect), and yet have not lost their capability of revivification. This fact, taken in connection with the extraordinary rate of increase mentioned in the preceding paragraph, removes all difficulty in accounting for the extent of the diffusion of these animals, and for their occurrence in incalculable numbers in situations where, a few days previously, none were known to exist. For their entire bodies may be wafted in a dry state by the atmosphere, from place to place; and their return to a state of active life, after a desiccation of unlimited duration, may take place whenever they meet with

FIG. 202.

*Stephanoceros Eichornii.*

the requisite conditions,—moisture, warmth, and food. It is probable that the ova are capable of sustaining treatment even more severe than the fully developed animals can bear; and that the race is frequently continued by them, when the latter have perished.

281. The principles on which the various forms that belong to this class should be systematically arranged, have not yet been satisfactorily determined. By Prof. Ehrenberg, the disposition of the ciliated lobes or wheel-organs, and the enclosure or non-enclosure of the body in a *lorica* or case, were taken as the basis of his classification; but as his ideas on both these points are inconsistent with the actual facts of organization, the arrangement founded upon them cannot be received. Another division of the class has been propounded by M. Dujardin, which is based on the several modes of life of

the most characteristic forms. And in a third, more recently put forth by Prof. Leydig, the general configuration of the body,

with the presence, absence, and conformation of the foot (or tail), are made to furnish the characters of the subordinate groups. Either of the two latter is certainly more *natural* than the first, as bringing together for the most part the forms which most agree in general organization, and separating those which differ; and we shall adopt that of M. Dujardin as most suitable to our present purpose.


I. The first group includes those that habitually live attached by the foot, which is prolonged into a pedicle; and it includes two families, the *Floscularians* and the *Melicertians*, both of which bear a certain general resemblance to the *Vorticellinæ* (§ 268) on the one hand, and to Zoophytes (Chap. XI) on the other. For they are commonly found attached to the stems and leaves of aquatic plants, by a long pedicle or footstalk, which bears a somewhat bell-shaped body; and in one of the most beautiful species, the *Stephanoceros Eichornii* (Fig. 202), this body has five long tentacles, beset with tufts of short bristly cilia, reminding us of the ciliated tentacles of the Bryozoa (Chap. XIII), whilst the body seems to be enclosed in a cylindrical cell, resembling that of Hydrozoa and Bryozoa. A comparison of this with other forms, however, shows that these tentacles are only extensions of the ciliated lobes which are common to all the members of these families; and the so-called "cell" is not formed by a thickening and separation of the outer tegument, but by a gelatinous secretion from it; so that, as the rest of the organization is essentially conformable to the Rotiferous type, no such passage is really established by this animal towards other groups, as it is commonly supposed to form. In one respect, *Floscularia* is still more aberrant; for the long bristly filaments with which its lobes are beset, are not capable of rhythmical vibration, and cannot, therefore, be properly termed "cilia." The body of *Melicerta* is protected by a most curious cylindrical tube, composed of little rounded pellets agglutinated together; this is obviously an artificial construction; and Mr. Gosse has been fortunate enough to have an opportunity of watching the animal whilst engaged in building it up. Beneath a projection on its head which he terms the chin, there is observed a small disk-like organ, in which, when the wheels are at work, a movement is seen very much resembling that of a revolving ventilator. Towards this disk, the greater proportion of the solid particles that may be drawn from the surrounding liquid into the vortex of the wheel-organs, are driven by their ciliary movement, a small part only being taken into the alimentary canal; and there they accumulate, until the aggregation (probably cemented by a glutinous secretion furnished by the organ itself) acquires the size and form of one of the globular pellets of the case; the time ordinarily required being about three minutes. The head of the animal then bends itself down, the pellet-disk is applied to the edge of the tube, the newly formed pellet is left attached there, and, the head being

lifted into its former position, the formation of a new pellet at once commences.

II. The next of M. Dujardin's primary groups (ranged by him, however, as the third), consists of the ordinary *Rotifer* and its allies, which pass their lives in a state of alternation between the conditions of those attached by a pedicle, of those which habitually swim freely through the water, and of those which creep or crawl over hard surfaces. As these have already been fully described, it is not requisite to dwell longer upon them.

III. The next group consists of those Rotifers which seldom or never attach themselves by the foot, but habitually swim freely through the water; and putting aside the peculiar aberrant form *Albertia*, which has only been found as a parasite in the intestines of worms, it may be divided into two families, the *Brachionians* and the *Furcularians*. The former are for the most part distinguished by the short, broad, and flattened form of the body (Figs. 200, 203); which is, moreover, enclosed in a sort of cuirass,

FIG. 203.



Notus quadricornis; A, dorsal view; B, side view.

formed by the consolidation of the external integument. This cuirass is often very beautifully marked on its surface, and may be prolonged into extensions of various forms, which are sometimes of very considerable length. The latter (corresponding almost exactly with the *Hydatineæ* of Prof. Ehrenberg) derive their name from the bifurcation of the foot into a sort of two-bladed forceps; their bodies are ovoidal or cylindrical, and are enclosed in a flexible integument, which is often seen to wrinkle itself into longitudinal and transverse folds, at equidistant lines. To this family belongs the *Hydatina senta*, which is one of the largest of the Rotifera, and which was employed by Prof. Ehrenberg as the chief subject of his examination of the internal structure of this group; as does also the *Notommata*, the curious condition of whose male has been already referred to (§ 279).

IV. The fourth of M. Dujardin's primary orders consists of the very curious tribe, first carefully investigated by M. Doyère, to which the name of *Tardigrada* has been given, on account of the slowness of their creeping movement. Their relation to the true Rotifera, however, is not at all clear; and many naturalists regard them as altogether distinct. They are found in the same localities with the Rotifers, and, like them, can be revived after desiccation (§ 280); but they have a vermiform body, divided transversely into five segments, of which one constitutes the head, whilst each of the others bears a pair of little fleshy protuberances, furnished with four curved hooks, and much resembling the pro-legs of a caterpillar. The head is entirely unpossessed of ciliated lobes; and it is only in the presence of a pair of jaws somewhat resembling those of Rotifera, and in the correspondence of their general grade of organization, that they bear any structural relation to the class we have now been considering. They may be pretty certainly regarded as a connecting link between the Rotifera and the Worms; but they should probably be ranked on the worm side of the boundary.

282. Notwithstanding that all the best informed Zoologists are now agreed in ranking the class of Rotifera in the Articulated series, yet there is still a considerable discordance of opinion as to the precise part of that series in which they should stand. For whilst Prof. Leydig, who has recently devoted much attention to the study of the class, regards them as most allied to the *Crustacea*, and terms them "Ciliocrustaceans," Mr. Huxley, with (as it seems to the Author) a clearer insight into their real nature, has argued that they are more connected with the *Annelida*, through the resemblance which they bear to the early larval forms of that class (Fig. 274). Considered in this light, the *Tardigrada* might seem to represent a more advanced phase of the same developmental history.¹

¹ In addition to the classical works of Ehrenberg and Dujardin, the following Memoirs should be consulted (besides those already referred to) by such as wish to acquaint themselves with the best researches upon the Rotifera:—Leydig in "Siebold and Kolliker's Zeitschrift," Band VI, heft 1; Goss on *Meliceria ringens*, in "Transact. of Microsc. Soc." Ser. I, vol. iii, p. 58; and "Quart. Journ. of Microsc. Sci." vol. i, p. 71; Williamson on *Meliceria ringens*, Op. cit. p. 1; and Huxley on *Lacinularia socialis*, in "Transact. of Microsc. Soc." Ser. II, vol. i, p. 1.

CHAPTER X.


FORAMINIFERA, POLYCYSTINA, AND SPONGES.

283. RETURNING now to the lowest or *Rhizopod* type of Animal life (§ 264), we have to direct our attention to three very remarkable series of forms, almost exclusively marine, under which that type manifests itself; all of them being distinguished by a *skeleton* of greater or less density; and this skeleton being generally so consolidated by mineral deposit, as to retain its form and intimate structure, long after the animal to which it belonged has ceased to live, even for those undefined periods in which they have been imbedded as fossils in strata of various geological ages. In the first of these groups, the *Foraminifera*, this skeleton consists of a *calcareous* shell (save in a few exceptional cases, in which the lorica remains flexible), that invests the sarcode body, but is usually perforated with numerous minute apertures. In the second group, also, the *Polycystina*, there is an investing shell, perforated with numerous apertures; but this shell is *siliceous*, instead of calcareous. In the third group, on the other hand,—that of *Porifera*, or *Sponges*,—the skeleton is usually composed of a network of horny fibres, strengthened either by calcareous or siliceous spicules, and having the soft animal body, which is composed of an aggregate of Amœba-like cells, in its interstices: in this group, moreover, we have a departure from the Rhizopod type, in the fact that certain parts of the free surfaces are furnished with cilia, whereby currents of water are sustained, that serve both for nutrition and for respiration.

284. *Foraminifera*.—The beings now known under this designation, possess, for the most part, *polythalamous* or “many-chambered” cells (Fig. 336), so strongly resembling those of *Nautilus*, *Spirula*, and other Cephalopod Mollusks, that it is not surprising that the older Naturalists, to whom the structure of these animals was entirely unknown, ranked them under that class. As such they were described by M. D’Orbigny (to whom we owe much of our knowledge of this group), in all his earlier publications; and they were distinguished from the ordinary Cephalopods that possess a single siphon passing from chamber to chamber, by the designation *Foraminifera*, which originally imported that the communications *between the chambers* are commonly made by

several such apertures, though it is now more commonly understood as applying to the sieve-like structure often presented by the external shell. It was by M. Dujardin, in 1835, that the structure of these animals was first shown to be conformable to the Rhizopod type; and notwithstanding the opposition to his views which has been set up by Prof. Ehrenberg (who, with inexplicable pertinacity, has continued to rank them among his *Bryozoa*, Chap. XIII), they have been confirmed by all subsequent observers, and more especially by the recent researches of Prof. Schulze,¹ who has given admirable descriptions of the animals of several different kinds of Foraminifera, derived from observation of them during their living state. The conformity of the Foraminifera to the ordinary Rhizopod type, is best seen in those forms, such as *Gromia* (Fig. 204), in which there is no multiplication of chambers; and it is made obvious by an examination of the accompanying figure, that there is no other essential difference between *Gromia* and *Arcella* or *Diffugia* (Fig. 193), than that which lies in the greater length and slenderness of the pseudopodial prolongations of the sarcod body in the former, as compared with those of the latter. The food is obtained by the extension of these pseudopodia in various directions from the mouth of the shell; and the absence of any membrane investing them is clearly indicated by their fusion or coalescence when two or more happen to come into contact, as well as by the vagueness of the expansions into which they are occasionally seen to spread out. These instruments entangle and lay hold of the minute bodies which serve as food to the animals, consisting of Diatomaceæ, Desmidiæ, the smaller forms of Confervæ, &c.; and they draw these, by their contraction, into the substance of the body, within which they may be seen through the transparent shell. It is not by any means constantly, that their indigestible *residua* are cast forth

FIG. 204.

*Gromia oviformis*, with its pseudopodia extended.

¹ "Über den Organismus der Polythalamien (Foraminiferen)." Leipzig, 1854.


again, as we have seen them to be from the surface of the body of the Actinophrys (§ 262); for they sometimes accumulate in considerable numbers, so as even to choke up a large part of the cavity. The sarcode body is occasionally seen to extend itself, as shown in Fig. 204, around the exterior of the shell; and pseudopodia are put forth from this extension, as well as from the ordinary outlet. Although nothing is certainly known of the mode of propagation of these animals, yet it is probable, from the analogy of the composite forms, that they multiply themselves by the detachment of gemmæ, composed of portions of sarcode, which in time form an envelope or shell. Nothing has been yet seen, that in the least corresponds to a true Generative process.

285. By far the greater number of Foraminifera are *composite* fabrics, evolved by a process of continuous gemmation, each gemma remaining in connection with the body by which it was put forth; and according to the plan on which this gemmation takes place, will be the configuration of the composite body thereby produced. Thus if the bud should be put forth from the aperture of *Gromia*, in the direction of the axis of its body, and a second shell should be formed around this bud, in continuity with the first, and this process should be successively repeated, a straight rod-like shell will be produced, having many chambers communicating with each other by the openings that originally constituted their mouths, the mouth of the last formed chamber being the only aperture through which the sarcode body, thus composed of a number of segments connected by a peduncle or "stolon" of the same material, can now project itself or draw in its food. The successive segments may be all of the same size, or nearly so, in which case the entire rod will approach the cylindrical form, or will resemble a line of beads; but it often happens that each segment is somewhat larger than the preceding, so that the composite shell has a conical form, the apex of the cone being the original segment, and its base the one last formed. The method of growth now described, is common to a large number of Foraminifera, chiefly belonging to the genera *Nodosaria*, *Dentalina*, *Morginulina*, which M. D'Orbigny has ranked together in an order, under the designation of *Stichotègues*; it is, however, frequently seen also in the advanced stages of growth of such as usually begin upon the spiral plan; and there are various considerations (into which this is not the place to enter) which satisfy the Author, that the mere *direction* of increase is a character of very subordinate importance in the group, instead of being one of such fundamental value as to serve for the basis of classification.

286. If each of the successively formed segments, instead of being developed exactly in the axis of its predecessor, should be directed a little to one side, it is obvious that a curved instead of a straight rod will be produced; and this curvature may increase,

until it becomes a spiral (Fig. 205). The character of this spiral will depend in great degree upon the enlargement or non-enlargement of the successively formed chambers; for sometimes it opens out very rapidly, every whorl being considerably broader than that which it surrounds, in consequence of the great excess of the size of each segment over that of its predecessor, as is generally the case in *Cristellaria*; whilst in other instances there is so little difference between the successive segments, after the spiral has made two or three turns, that the breadth of each whorl scarcely exceeds that of its predecessor, as is well seen in *Faujasina* (Fig. 209), as also in *Nummulite* (Chap. XIX). An intermediate condition is presented by *Rosalina* (Fig. 205), which

FIG. 205.

*Rosalina ornata*, with its pseudopodia extended.

may be taken as an example of a very large group of Foraminifera, composed of those whose plan of growth is *helical* or *spiral*, and ranged by M. D'Orbigny under the designation *Helicostegues*. In this genus, as in a large proportion of its congeners, we find the shell perforated with numerous apertures, through which pseudopodia can be extended from any of the segments that are not enclosed by others, as well as from the mouth or aperture of the outer segment; and when this is the case, it does not appear that the sarcode body is so often extended over the exterior of the shell, as it is when the shell has no perforations for the putting forth of these extensions. There are generally indications, however, in the structure of the shell itself,—new layers being often formed over the innermost whorls of the spiral, or partial

deposits being added, sometimes in the shape of bosses, spines, or other outgrowths,—that a soft substance, capable of originating such new formation, must occasionally spread itself over the whole external surface of the previous segments. And it is not difficult to understand how this may come to pass, when it is borne in mind that the gelatinous sarcode, however fine may be the threads into which it divides itself, may readily form a continuous layer by the coalescence of those threads. The group of Helicostègues is subdivided by M. D'Orbigny into two families, the *Nautiloidæ*, and the *Turbinoidæ*; the first and most important consisting of those in which the successive whorls all lie in the same plane, so that the shell is “equilateral” (like that of a *Nautilus*), as is the case with *Nummulites* and their allies (such as *Nonionina*, *Assilina*, *Operculina*, &c., between which and *Nummulites* the differences are but very slight); whilst the second contains those in which the spiral passes obliquely round an axis, so that the shell becomes “inequilateral” (like that of a snail or periwinkle), as is the case with *Rotalia* (Fig. 335), *Faujasina* (Fig. 209) and *Rosalina* (Fig. 205).

287. Putting aside less important variations in the plan of gemmation, we have now to notice one that seems essentially distinct from the preceding; that, namely, in which the new segments are added in concentric rings, each surrounding its predecessors, so as to form flattened disks. As an example of this curious type of Foraminiferous structure, the *Orbitolite* may be cited; which, long known as a very abundant fossil in the early tertiary of the Paris basin, has lately proved to be scarcely less abundant in certain parts of the existing ocean. The largest specimens of it, sometimes attaining the size of a sixpence, have hitherto been obtained only from the coast of New Holland and various parts of the Polynesian Archipelago; but disks of comparatively minute size (from the diameter of an ordinary pin's head, to that of a small pea) and of simpler organization, are to be found in almost all Foraminiferous sands and dredgings from the shores of the warmer regions of the globe, being especially abundant in those of some of the Philippine Islands, of the Red Sea, of the Mediterranean, and especially of the *Ægean*. When such disks are subjected to Microscopic examination, they are found (if uninjured by abrasion) to present the structure represented in Fig. 206; where we see on the surface (by incident light) a number of rounded elevations, arranged in concentric circles around a sort of nucleus (which has been laid open in the figure to show its internal structure); whilst at the margin we observe a row of rounded projections, with a single aperture or pore in each of the intervening depressions. In very thin disks, the structure is often brought into view, by mounting them in Canada balsam, and transmitting light through them, sufficiently well to render any other mode of preparation unnecessary; but in those which are too opaque to be thus seen through, it is suffi-

cient to rub down one of the surfaces upon a stone, and then to mount the specimen in balsam. Each of the superficial elevations will then be found to be the roof or cover of an ovate cavity or cell, which communicates by means of a lateral passage with the cavity on either side of it in the same ring; so that each circular zone of cells might be described as a continuous annular passage, dilated into cavities at intervals. On the other hand, each zone communicates with the zones that are internal and external to it, by means of passages in a radiating direction; and it is curious that these passages run, not from the cells of the inner zone to those of the outer; but from the connecting passages of the former to the cells of the latter; so that the cells of each zone *alternate* in position with those of the zones that are internal and external to it. The radial passages from the outermost annulus make their way at once to the margin, where they terminate, forming the "pores" which (as already mentioned) are to be seen on its exterior. The central nucleus, when rendered sufficiently transparent by the means just adverted to, is found to consist of a central cell (*a*), usually somewhat pear-shaped, that communicates by a narrow passage with a much larger circumambient cell (*b*), which nearly surrounds it, and which sends off a variable number of radiating passages towards the cells of the first zone, which forms a complete ring around the nucleus.¹

288. The idea of the nature of the living occupant of these cavities, which might be suggested by the foregoing account of their arrangement, is fully borne out by the results of the examination of the sarcode body, that may be obtained by the maceration in dilute acid (so as to remove the shelly investment) of specimens of *Orbitolite*, that have been gathered fresh from the sea-weeds to which in the living state they are found adherent, and have been kept in spirit. For this body is found to be composed (Fig. 207) of a multitude of segments of "sarcode," pre-

FIG. 206.




Simple disk of *Orbitolites complanatus*, laid open to show its interior structure:—*a*, central cell; *b*, circumambient cell surrounded by concentric zones of cells, connected with each other by annular and radiating passages.

¹ Although the above may be considered the *typical* form of the *Orbitolite*, yet, in a very large proportion of specimens, the first few zones are not complete circles, the early growth having taken place rather in a *spiral* than in a *radial* direction; between these two plans, there is every variety of gradation; and even where the spiral is most distinctly marked in the first instance, the additions soon come to be made in concentric zones.

senting not the least trace of higher organization in any part, and connected together by "stolons" of the like substance. The "central" pear-shaped segment, *a*, is seen to have budded off its "circumambient" segment, *b*, by a narrow footstalk or stolon; and this circumambient segment, after passing almost entirely round the central one, has budded off three stolons, which swell into new segments from which the first annulus is formed. Scarcely are any two specimens precisely alike, as to the mode

FIG. 207.



Composite Animal of simple type of *Orbitolites complanatus*:—*a*, central mass of sarcod; *b*, circumambient mass, giving off peduncles, in which originate the concentric zones of segments connected by annular bands.

in which the first annulus originates from the "circumambient" segment; for sometimes a score or more of radial passages extend themselves from every part of the margin of the latter (and this, as corresponding with the plan of growth afterwards followed, is probably the typical arrangement), whilst in other cases (as in the example before us) the number of these primary offsets is extremely small. Each zone is seen to consist of an assemblage of ovate segments, whose height (which could not be shown in the figure) corresponds with the thickness of the disk; these segments, which are all exactly


similar and equal to one another, are connected by annular "stolons;" and each zone is connected with that on its exterior, by radial extensions of those stolons, passing off between the segments. Although no opportunity has yet been obtained, for a microscopic examination of these animals in their living state, yet there can be no reasonable doubt that the radial extensions of the outermost zone issue forth as pseudopodia from the marginal pores; and that they search for and draw in alimentary materials, in the same manner as do those of other Foraminifera; the whole of the soft body, which has no communication whatever with the exterior, save through these marginal pores, being nourished by the transmission of the products of digestion from segment to segment and from zone to zone, through similar bands of gelatinous substance. In all cases in which the growth of the disk takes place with normal regularity, it is probable that a complete circular zone is added at once. When the sarcod body has increased beyond the capacity of its enveloping disk, it may be presumed that its pseudopodial extensions, proceeding from the marginal pores, coalesce, so as to form a com-

plete annulus of sarcode round the margin of the outermost zone; and it is probable that it is by a deposit of calcareous matter in the surface-portion of this annulus, that the new zone of shelly substance is formed, which constitutes the walls of the cells and passages occupied by the soft sarcode body. Thus we find this simple type of organization giving origin to fabrics of by no means microscopic dimensions, in which, however, there is no other differentiation of parts than that concerned in the formation of the shell; every segment and every stolon (with the exception of the two forming the "nucleus") being, so far as can be ascertained, a precise repetition of every other, and the segments of the nucleus differing from the rest in nothing else than their form. The equality of the endowments of the segments is shown by the fact, of which accident has repeatedly furnished proof,—that a small portion of a disk, entirely separated from the remainder, will not only continue to live, but will so increase as to form a new disk; the loss of the nucleus not appearing to be of the slightest consequence, from the time that active life is established in the outer zones. In what manner the multiplication and reproduction of the species are accomplished, we can as yet do little more than guess; but from appearances sometimes presented by the sarcode body, it seems reasonable to infer that "gemmales," corresponding with the "zoospores" of Protophytes (§ 197), are occasionally formed by the breaking up of the sarcode into globular masses; and that these, escaping through the marginal pores, are sent forth to develop themselves into new fabrics. Of the mode wherein that sexual operation is performed, however, in which alone true Generation consists, nothing whatever is known.

289. One of the most curious features in the history of this animal, is its capacity for developing itself into a form, which, whilst fundamentally the same as that previously described, is very much more complex. In all the larger specimens of Orbitolite, we observe that the marginal pores, instead of constituting but a single row, form many rows, one above another; and besides this, the cells of the two surfaces, instead of being rounded or ovate in form, are usually oblong and straight-sided, their long diameters lying in a radial direction. When a vertical section is made of such a disk, it is found that these oblong chambers constitute two superficial layers, between which are interposed columnar chambers of a rounded form; and these last are connected together by a complex series of passages, the arrangement of which will be best understood from the examination of a part of the sarcode body that occupies them (Fig. 208). For the oblong superficial chambers are occupied by segments of sarcode, *c c*, *d d*, lying side by side so as to form part of an annulus, but each of them being disconnected from its neighbors, and communicating only by a double footstalk with the two circular stolons *a a'*, *b b'*, which obviously correspond with the single stolon

of the simple type (Fig. 207).

FIG. 208.



Portion of Composite Animal of complex type of *Orbitolites complanatus*:—*a a'*, *b b'*, the upper and lower annular bands of two concentric zones; *c c'*, the upper layer of superficial segments, and *d d'* the lower layer, connected with the annular bands of both zones; *e e'*, *e' e'*, vertical segments of the two zones.

These indirectly connect together, not merely all the superficial cells of each zone, but also the columnar segments of the intermediate layer; for these columns (*e e'*, *e' e'*) terminate above and below in the annular stolons, sometimes passing directly from one to the other, but sometimes going out of the direct course to coalesce with another column. The columns of the successive zones (two sets of which are shown in the figure) communicate with each other by threads of sarcode, in such a manner, that (as in the simple type) each column is thus brought into connection with two columns of the zone next interior, to which it alternates in position. Similar threads, passing off from the outermost zone, through the multiple ranges of marginal pores, would doubtless act as pseudopodia. Now

this plan of growth is so different from that previously described, that there would at first seem ample ground for distinguishing the *simple* and the *complex* types as two species. But the test furnished by the examination of a large number of specimens, which ought never to be passed by when it can possibly be appealed to, furnishes these very singular results:—1st, That the two forms must be considered as specifically identical; since there is not only a gradational passage from one to the other, but they are often combined in the same individual, the inner and first formed portion of a large disk frequently presenting the simple type, whilst the outer and later formed part has developed itself upon the complex:—2d, That although the last mentioned circumstance would naturally suggest that the change from the one plan to another may be simply a feature of advancing age, yet this cannot be the case; since the complex plan sometimes evolves itself even from the very first (the nucleus, though resembling that of the simple form, sending out two or more tiers of radiating threads), whilst, more frequently, the simple prevails for an indefinite number of zones, and then changes itself in the course of a few zones into the complex. The mode in which this change occurs is not a little curious. In the first place, the short segments, threaded (so to speak) upon their annular stolon, undergo elongation, and the annular stolon itself becomes double, being first, as it were, split in two, and its upper and lower halves being separated by the interpolation of a length-

ening piece to each columnar segment. These additional pieces are at first very short; but with the growth of every new zone, they commonly increase in length; and this interpolated portion comes to constitute the principal part of the thickness of the disk. While this change is going on, another is taking place in the position of the superficial portions of the segments which are above and below the annular stolons; for these, whilst at first seeming to be mere continuations of the columns beneath, and being connected (like them) with the stolons of their own zones alone, are so displaced, in the course of two or three zones, as to arch over the space between the zones (as shown in Fig. 208), and to connect themselves with the stolon, not only of their own zone, but of the next. It has been thought desirable to enter into the foregoing detail, since a more striking instance could scarcely be drawn from any department of Natural History, of the wide range of variation that may occur within the limits of one and the same species; and the Microscopist needs to be specially put on his guard as to this point in respect to the lower types of Animal, as to those of Vegetable life, since the determination of form seems to be far less precise among such, than it is in the higher types.¹

290. The type of Foraminiferous structure which has been just described, is in many respects peculiar, and may be considered as verging towards the Sponges. It is obvious, from what has been said of the extreme freedom with which the several segments of the sarcode body communicate with each other, that they form *one whole*, in a far greater degree than they do in the ordinary composite Foraminifera, whose segments are more completely separated, and are very commonly connected only by a few very slender threads of sarcode. Indeed if we were to imagine a discoidal mass of sarcode to be traversed by a reticulated calcareous skeleton, somewhat resembling that open areolar texture which forms the *shell* of the *Echinida* (§ 312), and this network were to possess somewhat of that regularity in the disposition of its successively formed parts, which is presented to us in the *spines* of that group (Fig. 237), we should have no unapt representation of the calcareous skeleton of the Orbitolite, and of its relation to the animal which it envelopes and protects. Now there are certain *Sponges* which have a reticular skeleton composed of mineral matter (§ 296), differing from that of the Orbitolite in little else than the want of the zonular arrangement which marks successive epochs of growth; and we shall see that the constitution of the soft body is essentially the same in one case as in the other. A remarkable connecting link between the Orbitolite and certain Sponges, seems, in fact, to be presented

¹ For a full account of the Organization of the *Orbitolite*, and of the various conditions under which it presents itself, see the Author's memoir upon that genus in "Philos. Transact." 1856.

to us in the curious *Thalassicolla*,¹ discovered by Mr. Huxley, and since observed by Prof. Müller, which seems also to have relations to the *Polycystina* (§ 292).

291. The essential simplicity of the animal, and the absence of anything like structure in the shell, of the Orbitolite, obviously place it much lower in the scale than those Foraminifera, which have the segments not only more completely divided, but enclosed in a shell which is itself distinctly organized. Such is the case with *Nummulites* and their nearest allies among the recent forms. For in these, as will be more fully shown hereafter (Chap. XIX), each segment has its own separate envelope of shell, so that the partition between any two adjacent chambers is double; the chambers are so far cut off from one another, as to communicate only by very narrow passages; but means are afforded, by which even the innermost chambers are brought into tolerably direct relation with the exterior. For in this type of structure, we observe that those parts of their walls which look towards the outer surfaces, are perforated with immense numbers of pores resembling those of Fig. 205, but more numerous, minute, and closely set; and where the walls are thick, these pores are continued as tubes through their entire substance (Figs. 209, 338). However fine they may be, the extraordinary tenuity of the threads into which the sarcode is occasionally seen to divide itself (Fig. 204), shows that this need not be an obstacle to the passage of pseudopodial prolongations through them. But further, in the spaces between the walls of contiguous chambers, we find a system of large tubes, which make their way directly from the central to the peripheral portion of the disk, and which, communicating on the one hand with the innermost chambers, and on the other with the margin (being extended and carried on to it whenever the previous edge is covered by a new growth), serve to bring the former into a very direct relation with the external sources of supply of nutriment and oxygen. A strikingly developed example of this system of "interseptal" canals is presented in the genus *Faujasina* (Fig. 209);² where the large size of the canals, and the extreme simplicity of their arrangement, enable them to be very readily traced out. In some instances the arrangement becomes extremely complex:

¹ See "Annals of Natural History," 2d Ser. vol. viii, p. 433; and "Quart. Journ. of Microsc. Science," vol. iv, p. 72.

² See Prof. Williamson's Memoir on the *Faujasina*, in the "Transact. of Microsc. Soc." 2d Ser. vol. i. As the correctness of the account of the interseptal system of canals, which has been given by the Author (in his Memoir on Nummulite, &c., in "Quart. Geol. Journ." Feb. 1850), and confirmed by the researches of Prof. W. and himself upon numerous recent types, has been called in question by no less an authority than Prof. Schulze, who has, in consequence, altogether ignored this important character in his classification, the Author thinks it right to state, that although the above figure is copied from one of those which illustrate Prof. Williamson's Memoir, yet it is the almost precise counterpart of a section of the same species prepared by himself. The incredulity of Prof. Schulze and others, upon this point, simply depends upon their want of aptitude in the preparation of sufficiently thin sections.

this is especially the case, where, as frequently happens, there is an interstitial calcareous skeleton to be nourished (often extending itself into outgrowths of various forms and sizes), in addition to the immediate investments of the segments; and it becomes obvious from the far greater development of this canal system wherever such is the case, that this interstitial skeleton is chiefly if not entirely maintained through the instrumentality of the "stolons" of sarcode which occupy these canals, and whose remains may often be distinctly traced in the dried shell.¹ Now this, the highest type of Foraminiferous structure, is not only presented by such spiral forms as Nummulite, but is found also in a genus that is



Section of *Favosina* near its base and parallel to it:—showing *a a*, the radiating interseptal canals; *b*, their internal bifurcations; *c*, a transverse branch; *d* tubular wall of the chambers.

conformable, in its concentric plan of growth, to Orbitolite. Hence it is obvious that no arrangement founded, as in that of M. D'Orbigny, upon a character of such secondary importance as the direction of gemmation, is likely to be in accordance with physiological features which a perfect knowledge of the animal might be expected to afford; and as these can be partly judged of from the structure of the shell, it seems obvious that this ought to be made the first consideration. To carry out a classification on such a basis, however, will involve a large amount of patient investigation.²

292. Many of the Foraminifera attach themselves in the living state to Sea-weeds, Zoophytes, &c.; and they should, therefore, be carefully looked for on such bodies, especially when it is desired to observe their internal organization and their habits of life. They are often to be collected in much larger numbers, however, from the sand or mud dredged up from the sea-bottom, or even from that taken from between the tide marks. In a paper containing some valuable hints on this subject,³ Mr. Legg

¹ The Author has been enabled to make out this curious point completely, in the *Calcarina*, a little body resembling a spur-rowel. For he has obtained ample evidence that the *spire* with its regularly added segments, and *interstitial skeleton* extending itself into radiating spines, may grow quite independently of one another. The proof will be submitted in future Memoirs to the Royal Society.

² To this labor, the Author has been for some years devoting a portion of his very limited leisure, in conjunction with his friend Prof. Williamson, of Manchester; and the results of their united labors will appear at the earliest practicable period, in the Ray Society's publications.

³ "Transactions of the Microscopical Society," 2d Series, vol. ii, p. 19.

mentions that, in walking over the Small Mouth Sand, which is situated on the north side of Portland Bay, he observed the sand to be distinctly marked with white ridges, many yards in length, running parallel with the edge of the water; and upon examining portions of these, he found Foraminifera in considerable abundance. One of the most fertile sources of supply that our own coasts afford, is the "ouze" of the Oyster-beds, in which large numbers of living specimens will be found; the variety of specific forms, however, is usually not very great. In separating these bodies from the particles of sand, mud, &c., with which they are mixed, various methods may be adopted, in order to shorten the tedious labor of picking them out, one by one, under the Simple Microscope; and the choice to be made among these will mainly depend upon the condition of the Foraminifera, the importance (or otherwise) of obtaining them alive, and the nature of the substances with which they are mingled. Thus, if it be desired to obtain living specimens from the oyster-ouze, for the examination of their soft parts, or for preservation in a vivarium, much time will be saved by stirring the mud (which should be taken from the surface only of the deposit) in a jar with water, and then allowing it to stand for a few moments; the finer particles will remain diffused through the liquid, while the heavier will subside; and as the Foraminifera (in the present case) belong to the latter category, they will be found almost entirely free from extraneous matter, at the bottom of the vessel, after the operation has been repeated two or three times. It would always be well to examine the first deposit let fall by the water that has been poured away; as this may contain the smaller and lighter forms of Foraminifera. But supposing that it be only desired to obtain the dead shells from a mass of sand brought up by the dredge, a very different method should be adopted. The whole mass should be exposed for some hours to the heat of an oven, and be turned over several times, until it is found to have been thoroughly dried throughout; and then, after being allowed to cool, it should be stirred in a large vessel of water. The chambers of their shells being now occupied by air alone (for the bodies of such as were alive will have shrunk up almost to nothing), the Foraminifera will be the lightest portion of the mass; and they will be found floating on the water, while the particles of sand, &c., subside. Another method, devised by Mr. Legg, consists in taking advantage of the relative sizes of different kinds of Foraminifera and of the substances that accompany them. This, which is especially applicable to the sand and rubbish obtainable from sponges (which may be got in large quantity from the sponge merchants), consists in sifting the whole aggregate through successive sieves of wire-gauze, commencing with one of ten wires to the inch, which will separate large extraneous particles, and proceeding to those of 20, 40, 70, and 100 wires to the inch, each (especially that of 70) retaining

a much larger proportion of Foraminiferous shells than of the accompanying particles; so that, a large portion of the extraneous matters being thus got rid of, the final selection becomes comparatively easy. Certain forms of Foraminifera are found attached to shells, especially bivalves (such as the *Chamaeceras*), with foliated surfaces; and an extensive examination of those of the Indian Seas, when brought home "in the rough," has yielded to Mr. W. K. Parker some most valuable and novel results, which will be made public in due time.

293. The final selection of specimens for mounting, should always be made under some appropriate form of Single Microscope (§§ 27-30); a fine camel-hair pencil, with the point wetted between the lips, being the instrument which may be most conveniently and safely employed, even for the most delicate specimens. In mounting Foraminifera as microscopic objects, the method to be adopted must entirely depend upon whether they are to be viewed by transmitted or by reflected light. In the former case, they should be mounted in Canada balsam; the various precautions to prevent the retention of air bubbles which have been already described (§ 128), being carefully observed. In the latter, no plan is so simple, easy, and effectual, as the attaching them with a little gum to a blackened disk of card, and guarding them by a perforated wooden slide (§ 123). They should be fixed in various positions, so as to present all the different aspects of the shell, particular care being taken that its mouth is clearly displayed; and where, as will often happen, the several individuals differ considerably from one another, special care should be taken to form *series* illustrative of their range of variation, and of the mutual connections of even the most diverse forms. For the display of the internal structure of Foraminifera, it will often be necessary to make extremely thin sections, in the manner already described (§§ 108-110); and much time will be saved, by attaching a number of specimens to the glass at once, and by grinding them down together. For the preparation of sections, however, of the extreme thinness that is often required, those which have been thus reduced should be transferred to separate glasses, and finished off each one by itself.

294. *Polycystina*.—It is probable that we are to refer to the same general type, an extensive group of very interesting microscopic bodies, possessing great beauty and variety of form and structure (Figs. 210-216) of whose essential character extremely little is known. These are minute siliceous shells, which appear, from the recent observations of Prof. Müller, to contain in the living state an olive-brown "sarcode," extending itself into pseudopodial prolongations (resembling those of *Actinophrys*, § 262), that pass through the large apertures by which the shells are perforated. The sarcode body does not seem always to fill the shell; being stated by Prof. Müller to occupy, in the *Encyrtidium*

of Messina, only the upper part or vault of the shell, and to be very regularly divided into four lobes. It is a peculiar feature

FIG. 210.



Fig. 210. *Podocyrtis Schomburgkii*.

FIG. 211.



Fig. 211. *Rhopalocanium ornatum*.

in these Polycystinae, that their shells are often prolonged into spines or other projections, which are sometimes arranged in such a manner as to give them a very singular aspect (Figs. 210, 211). It seems probable that these creatures are almost as widely diffused at the present time as are the Foraminifera; although, from their greater minuteness, they have not been so often recognized. For having been first discovered by Prof. Ehrenberg at Cuxhaven on the North Sea, they were afterwards found by him in collections made

in the Antarctic Seas, and have been recently described by Prof. Bailey as presenting themselves (with Foraminifera and Diatomaceae) in the deposits brought up by the sounding-lead from the bottom of the Atlantic Ocean, at depths of from 1000 to 2000 fathoms. They appear to have been much more abundant, however, during the later geological periods; for not only have certain forms (among them *Haliomma*, Fig. 212) been detected by

FIG. 212.




Fig. 212. *Haliomma Humboldtii*.

FIG. 213.




Fig. 213. *Perichlamyidium proteustum*.

Prof. Ehrenberg in the chalks and marls of Sicily and Greece, and of Oran in Africa, and also in the diatomaceous deposits of Bermuda and Richmond (Virginia); but a large proportion of the rock that prevails through an extensive district in the island of Barbadoes, has been found by him to be composed of Poly-

cystina, mingled with Diatomaceæ, with a few calcareous Foraminifera, and with calcareous earth which was probably derived

FIG. 214.



Fossil *Polycystina*, &c., from Barbadoes:—*a*, *Podocyrtes mitra*; *b*, *Rhabdolithus sceptrum*; *c*, *Lychnocanium fulciferum*; *d*, *Encyrtidium tubulus*; *e*, *Flustrella concentrica*; *f*, *Lychnocanium lucerna*; *g*, *Encyrtidium elegans*; *h*, *Dictyospyris clathrus*; *i*, *Encyrtidium mongolfieri*; *k*, *Stephanolithis spinescens*; *l*, *S. nodosa*; *m*, *Lithocyclia ocellus*; *n*, *Cephalolithis sylvina*; *o*, *Podocyrtes cothurnata*; *p*, *Rhabdolithes pipa*.

from the decomposition of corals, &c., so as to form, according to the relative proportions of these constituents (which differed in different parts of the deposit), a tripoli-like sandstone, whitish and very friable, a compact calcareous sandstone, and strata of a marly character sometimes containing semi-opal. Previously to this last discovery, which was made in the year 1846 (the materials having been furnished by the geological researches of Sir R. H. Schomburgk), 39 species of *Polycystina* had been established by Prof. E.; but in the Barbadoes deposit, he has detected no fewer than 282 forms which he considers to be specifically distinct, besides 25 species of Diatomaceæ and Foraminifera, and 54 forms which he cannot distinctly determine, but which he classes under the provisional designations of *Geolitharia* and *Phytolitharia*, making 361 in all, of which more than 300 were previously unknown. The 282 species of *Polycystina* are arranged by Prof. E. in seven families, which include forty-four genera; but it is obvious that in our present state of almost entire ignorance of the structure and physiology of the animals to which these shells belong, no classification can be otherwise than provisional.—Few Microscopic objects are more beautiful than an assemblage of the most remarkable forms of Barbadian *Polycys-*

tina, especially when seen brightly illuminated upon a black ground; since (for the reason formerly explained, § 62) their

FIG. 215.

FIG. 216.




Fig. 215. *Stylodictya gracilis*.
Fig. 216. *Astromma Aristotelis*.

“solid forms” become much more apparent than they are when these objects are examined by light transmitted through them. And the “black ground illumination,” either by the “spotted lens” or by the “paraboloid” (§ 61), is much to be preferred for this purpose, to the ordinary mode of illuminating opaque objects by incident light from a condenser, although this may be advantageously had recourse to, by the Microscopist who is unprovided with these appurtenances. No class of objects is more suitable than these to the “Binocular Microscope” (§ 40); the stereoscopic projection of which causes them to be presented to the mind’s eye in complete relief, so as to bring out with the most marvelous and beautiful effect all their delicate sculpture, reminding the observer (to compare small things with great) of the finest specimens of the hollow ivory balls carved by the Chinese.¹


295. *Sponges*.—Although this tribe has been bandied from the Animal to the Vegetable kingdom, and back again, several times in succession, yet its claim to a place among the Protozoa may now be considered as pretty certainly determined, by the information derived from Microscopic examination of its minute structure. For in the living Sponge, the skeleton, usually composed of a fibrous network strengthened by spicules of mineral matter, is clothed with a soft flesh; and this flesh has been found by Dujardin and all subsequent observers to consist of an aggregation of Amœba-like bodies (Fig. 217, B), some of which (as Mr. Dobie has shown)² are furnished with one or more long

¹ For a fuller description of this group, see Prof. Ehrenberg’s *Memoirs in the “Transactions of the Berlin Academy”* for 1846, 1847, and his recently published “*Microgeologie*,” also “*Ann. of Nat. Hist.*” 1847.

² *Goodsir’s Annals of Anatomy and Physiology*, No. 2, May, 1852.

cilia, closely resembling those of *Volvox* (Fig. 70, 1), by the agency of which, a current of water is kept up through the passages and canals excavated in the substance of the mass.

FIG. 217.



Structure of *Grantia compressa*.—A, portion moderately magnified, showing general arrangement of triradiate spicules and intervening tissue;—B, small portion highly magnified, showing ciliated cells.

And from the observations of Mr. Carter¹ upon the early development of Sponges, it appears that they begin life as solitary *Amœbæ*, and that it is only in the midst of aggregations formed by the multiplication of these, that the characteristic *Sponge*-structure makes its appearance, the formation of spicules being the first indication of such organization. The ciliated cells seem to form the walls of the canals by which the whole fabric of the Sponge is traversed; these canals, which are very irregular in their distribution, may be said to commence in the small pores of the surface, and to terminate in the large vents; and a current is continually entering at the former, and passing forth from the latter, during the whole life of the Sponge, bringing in alimentary particles and oxygen, and carrying out excrementitious matter.

296. The skeleton which gives shape and substance to the mass of sarcode particles that constitutes the living animal, is composed, in the Sponges with which we are most familiar, of an irregular reticulation of fibres. The arrangement of these may be best made out, by cutting thin slices of a piece of Sponge submitted to firm compression, and viewing these slices, mounted upon a dark ground, with a low magnifying power, under incident light. Such sections, thus illuminated, are not merely striking objects, but serve to show, very characteristically, the general disposition of the larger canals and of the smaller areolæ with which they communicate. In the ordinary Sponge, the fibrous skeleton is almost entirely destitute of spicules, the absence of which, in fact, is one important condition of that

¹ "Annals of Natural History," Ser. 2, vol. iv.

flexibility and compressibility on which its uses depend. When spicules exist in connection with such a skeleton, they are either altogether imbedded in the fibres, or they are implanted into them at their bases, as shown in Fig. 218. In the curious and beautiful *Dictyochalix pumiceus* of Barbadoes, however, the entire network of fibres is composed of silex, and is so transparent that it looks as if composed of spun glass. There are many Sponges in which no fibrous network can be discerned, the spicules lying imbedded in the midst of the sarcode mass; such is the case in *Grantia* (Fig. 217, A), whose triradiate spicules are composed of carbonate of lime. Sponge-spicules are much more frequently siliceous than calcareous; and the variety of forms presented by the siliceous spicules is much greater than that which we find in the comparatively small number of species in which they are composed of carbonate of lime. The long needle-like spicules (Fig. 219) which are extremely abundant in several Sponges, lying close together in bundles, are sometimes straight, sometimes slightly curved; they are sometimes pointed at both ends, sometimes at one only; one or both ends may be furnished with a head like that of a pin, or may carry three or more diverging points, which sometimes curve back so as to form hooks (Fig. 334, H). When the spicules project from the horny framework, they are usually somewhat conical in form, and their surface is often beset with little spines, arranged at regular intervals, giving them a jointed appearance (Fig. 218). Sponge-spicules frequently occur, however, under forms very different from the preceding; some being short and many-branched; and the branches being themselves very commonly stunted into mere

tubercles (some examples of which type are presented in Fig. 334, A, C); whilst others are stellate, having a central body with conical spines projecting from it in all directions (as at D of the same figure). Great varieties

FIG. 218.



Fig. 218. Portion of *Halichondria* (?) from Madagascar, with spicules projecting from the fibrous network.

FIG. 219.



Fig. 219. Siliceous Spicules of *Pachymatisma*.

present themselves in the stellate form, according to the relative predominance of the body and of the rays; in those represented

in Fig. 219, the rays, though very numerous, are extremely short; in other instances the rays are much longer, and scarcely any central nucleus can be said to exist. The varieties in the form of Sponge-spicules are, in fact, almost endless; and a single sponge often presents two or more (as shown in Fig. 219), the *stellate* spicules usually occurring either in the interspaces between the elongated kinds, or in the external crust. In one curious Sponge described by Mr. Bowerbank (the *Dusideia fragilis*), the spicules are for the most part replaced by particles of sand, of very uniform size, which are found imbedded in the horny fibre. The spicules of Sponges cannot be considered, like the "raphides" of Plants (§ 230), simply as deposits of mineral matter in a crystalline state. For the forms of many of them are such as no mere crystallization can produce; many of them possess internal cavities, which contain organic matter; and the calcareous spicules, whose mineral matter can be readily dissolved away by an acid, are found to have a distinct animal basis. Hence it seems probable, that each spicule was originally a cell or segment of sarcode, which has undergone calcification, and by the self-shaping power of which, the form of the spicule is mainly determined.

297. Of the Reproductive process in Sponges, much has yet to be learned. The following is perhaps the most probable account of it:—Multiplication by gemmation is effected by the detachment of minute globular particles of sarcode from the interior of the canals, where they sprout forth as little protuberances, whose footstalks gradually become narrower and narrower until they give way altogether; these *gemmales*, like the zoospores of Alge, possess cilia, and issuing forth from the vents, transport themselves to distant localities, where they may lay the foundation of new fabrics. But according to the recent observations of Mr. Huxley,¹ a true sexual generation also takes place, as might be anticipated; both ova and sperm-cells being found imbedded in the substance of the Sponge. The bodies distinguished as *capsules*, which are larger than the gemmules, and which usually have their investment strengthened with siliceous spicules very regularly disposed, are probably the products of this operation. They contain numerous globular particles of sarcode, every one of which, when set free by the rupture of its envelope, becomes an independent Amœba-like body, and may develop itself into a complete Sponge.

298. With the exception of those that belong to the genus *Spongilla*, all known Sponges are marine; but they differ very much in habit of growth. For whilst some can only be obtained by dredging at considerable depths, others live near the surface, whilst others attach themselves to the surfaces of rocks, shells, &c., between the tide-marks. The various species of *Grantia*, in which alone of all the marine Sponges has ciliary movement

¹ "Ann. of Nat. Hist." Ser. 2, vol. vii.

yet been seen, belong to this last category. They have a peculiarly simple structure, each being a sort of bag whose wall is so thin that no system of canals is required, the water absorbed by the outer surface passing directly towards the inner, and being expelled by the mouth of the bag. The cilia may be plainly distinguished with a 1-8th inch objective, on some of the cells of the gelatinous substance scraped from the interior of the bag; or they may be seen *in situ*, by making very thin transverse sections of the substance of the Sponge.¹ It is by such sections alone, that the internal structure of sponges, and the relation of their spicular and horny skeletons to their fleshy substance, can be demonstrated. In order to obtain the spicules in an isolated condition, however, the animal matter must be got rid of, either by incineration, or by chemical reagents. The latter method is preferable, as it is difficult to free the mineral residue from carbonaceous particles by heat alone. If (as is commonly the case) the spicules are siliceous, the Sponge may be treated with strong nitric or nitro-muriatic acid, until its animal substance is dissolved away; if, on the other hand, they be calcareous, a strong solution of potass must be employed instead of the acid. The operation is more rapidly accomplished by the aid of heat; but if the saving of time be not of importance, it is preferable on several accounts to dispense with it. The spicules, when obtained in a separate state, should be mounted in Canada balsam. Sponge-tissue may often be distinctly recognized in sections of agate, chalcedony, and other siliceous accretions, as will hereafter be stated in more detail (Chap. XIX).

¹ See Dobie, *loc. cit.*; and Bowerbank in "Transactions of Microscopical Society," 1st Ser. vol. iii, p. 137.

CHAPTER XI.


ZOOPHYTES.

299. THE term Zoophyte, although sometimes used in a wider signification, is properly restricted to the class of *Polypifera*, or polype-bearing animals, whose composite skeletons or "poly-paries" have more or less of a plant-like form; even the *Polyzoa* (or Bryozoa) being now excluded, on account of their truly Molluscan structure (Chap. XIII), notwithstanding the zoophytic character of their forms and of their habits of life. The true Zoophytes may be divided into two primary groups, the *Hydrozoa* and the *Anthozoa*; the *Hydra* (or fresh-water polype) standing as the type of the one, and the *Sea-Anemone* as the representative of the other. As the Hydrozoa are essentially microscopic animals, they need to be described with some minuteness; whilst in regard to the Anthozoa, only those points can be dwelt on, which are of special interest to the Microscopist.

300. The *Hydra* is to be searched for in pools and ditches, where it is most commonly to be found attached to the leaves or stems of aquatic plants, floating pieces of stick, &c. Two species are common in this country, the *H. viridis* or green polype, and the *H. vulgaris*, which is usually orange-brown, but sometimes yellowish or red (its color being liable to some variation according to the nature of the food on which it has been subsisting); a third less common species, the *H. fusca*, is distinguished from both the preceding by the length of its tentacula, which in the former are scarcely as long as the body, whilst in the latter they are, when fully extended, many times longer (Fig. 220). The body of the *Hydra* consists of a simple bag or sac, which may be regarded as a stomach, and which is capable of varying its shape and dimensions in a very remarkable degree; sometimes extending itself in a straight line, so as to form a long narrow cylinder, at other times being seen (when empty) as a minute contracted globe, whilst, if distended with food, it may present the form of an inverted flask or bottle, or even of a button. At the upper end of this sac is a central opening, the "mouth," and this is surrounded by a circle of tentacula or "arms," usually from six to ten in number, which are arranged with great regularity around the orifice. The body is prolonged at its lower

end into a narrow base, which is furnished with a suctorial disk; and the Hydra usually attaches itself by this, whilst it allows its

FIG. 220.



Hydra fusca, with a young bud at *b*, and a more advanced bud at *c*.

tendrillike tentacula to float freely in the water, like so many fishing-lines. The wall of the body is composed of cells, imbedded in a kind of sarcode; and it consists of two principal layers, an outer and more compact, of which the cells form a tolerably even surface, and an inner that lines the stomach, into the cavity of which some of the cells project. Between these layers, there is a space chiefly occupied by "sarcode," having many vacuoles or lacunæ (which often seem to communicate with one another) excavated in its substance. The arms are made up of the same materials as the body; but their surface is beset with little wart-like prominences, which, when carefully examined, are found to be composed of clusters of small "thread-cells," having a single large cell with a long spiculum in the centre of each. The structure of these thread-cells or "urticating organs" will be described hereafter (§ 310); at present it will be enough to point out, that this ap-

paratus, repeated many times on each tentacle, is doubtless intended to give to the organ a great prehensile power; the minute filaments forming a rough surface, adapted to prevent the object from readily slipping out of the grasp of the arm, whilst the central spiculum is projected into its substance, and probably conveys into it a poisonous fluid secreted by a vesicle at the base of the dart. The latter inference is founded upon the oft-repeated observation, that if the living prey seized by the tentacles have a body destitute of hard integument, as is the case with the minute aquatic Worms which constitute a large part of its aliment, this speedily dies, even if, instead of being swallowed, it escapes from their grasp; on the other hand, minute Entomostracous Crustacea, Insects, and other animals with hard envelopes, may escape without injury, even after having been detained for some time in the polype's embrace. The contractility of the tentacula (the interior of which is traversed by a canal

which communicates with the cavity of the stomach) is very remarkable, especially in the *Hydra fusca*; whose arms, when extended in search of prey, are not less than seven or eight inches in length; whilst they are sometimes so contracted, when the stomach is filled with food, as to appear only like little tubercles around its entrance. By means of these instruments, the Hydra is enabled to derive its subsistence from animals, whose activity, as compared with its own slight powers of locomotion, might have been supposed to remove them altogether from its reach; for when, in its movements through the water, a minute worm or a water-flea happens to touch one of the tentacula of the polype, spread out as these are in readiness for prey, it is immediately seized by this, other arms are soon coiled around it, and the unfortunate victim is speedily conveyed to the stomach, within which it may frequently be seen to continue moving for some little time. Soon, however, its struggles cease, and its outline is obscured by a turbid film, which gradually thickens, so that at last its form is wholly lost. The soft parts are soon completely dissolved, and the harder indigestible portions are rejected through the mouth. A second orifice has been observed at the lower extremity of the stomach; but this would not seem to be properly regarded as anal, since it is not used for the discharge of such exuviæ; it is probably rather to be considered as representing, in the Hydra, the entrance to that ramifying cavity, which, in the compound Hydroida, brings into connection the lower extremities of the stomach of all the individual polypes (Fig. 223). A striking proof of the simplicity of the structure of the *Hydra*, is the fact that it may be turned inside out like a glove; that which was before its external tegument becoming the lining of its stomach, and *vice versa*.

301. The ordinary mode of multiplication in this animal, is by a gemmation resembling that of Plants. Little bud-like processes (Fig. 220, *b*, *c*) are developed from its external surface, which are soon observed to resemble the parent in character, possessing a digestive sac, mouth, and tentacula; for a long time, however, their cavity is connected with that of the parent, but at last the communication is cut off by the closure of the canal of the footstalk, and the young Polype quits its attachment and goes in quest of its own maintenance. A second generation of buds is sometimes observed on the young Polype, before quitting its parent; and as many as nineteen young Hydræ, in different stages of development, have been seen thus connected with one original stock (Fig. 221). Another very curious endowment seems to depend on the same condition,—the extraordinary power which one portion possesses of reproducing the rest. Into whatever number of parts a *Hydra* may be divided, each may retain its vitality, and give origin to a new and entire fabric; so that thirty or forty individuals may be formed by the section of one. The *Hydra* also propagates itself, however, by a truly

Sexual process; the fecundating apparatus, or vesicle-producing "sperm-cells," and the ovum (containing the "germ-cell," imbedded in a store of nutriment adapted for its early development), being evolved in the substance of the walls of the stomach, the former just beneath the arms, the latter nearer to the lower end of the body. It would appear that sometimes one individual Hydra develops only the male cysts or sperm-cells, while another develops only the female cysts or ovisacs; but the general rule seems to be, that the same individual forms both organs. The fertilization of the ova, however, cannot take place until after the rupture of the spermatocyst and that of the ovisac also; so that the parent has no further participation in it, than has the *Fucus* in the analogous fertilization of its germ-cells after their discharge (§ 205). Although the production, from such an egg, of a new Hydra, similar in all respects to its parent, has not yet been witnessed, there seems no reason to doubt the fact. It would seem that this alternation in the method of reproduction, between the gemmiparous and the sexual, is greatly influenced

FIG. 221.



FIG. 222.




Fig. 221. *Hydra fusca* in gemmation; *a*, mouth; *b*, base; *c*, origin of one of the buds.

Fig. 222. Development of Medusa-buds in *Syncoryna Sarsii*:—*a*, an ordinary polype, with its club-shaped body covered with tentacula:—*b*, a polype putting forth medusan gemme; *a*, a very young bud; *b*, a bud more advanced, the quadrangular form of which, with the four nuclei whence the cirrhi afterwards spring, is shown at *d*; *c*, a bud still more advanced.

by external temperature; the eggs being produced at the approach of winter, and serving to regenerate the species in the

spring, the parents not being able to survive the cold season ; whilst the budding process naturally takes place only during the warmer part of the year, but may be made to continue through the whole winter, by keeping the water inhabited by the polypes at a sufficiently high temperature. The Hydra possesses the power of free locomotion, being able to remove from the spot to which it has attached itself, to any other that may be more suitable to its wants ; its changes of place, however, seem rather to be performed under the influence of *light*, towards which the Hydra seeks to move itself, than with reference to the search after food.

302. Some of the simpler forms of the composite *Hydrozoa* may be likened to a *Hydra*, whose gemmæ, instead of becoming detached, remain permanently connected with the parent ; and as these in their turn may develop gemmæ from their own bodies, a structure of more or less arborescent character may be produced. The form which this will present, and the relation of the component polypes to each other, will depend upon the mode in which the gemmation takes place ; in all instances, however, the entire cluster is produced by continuous growth from a single individual ; and the stomachs of the several polypes are united by tubes, which proceed from the base of each other, along the stalk and branches, to communicate with the cavity of the central stem. This is the case with the family *Corynidae*, which are composite fabrics, sometimes quite arborescent in form, but unpossessed of any firm investment, the external wall being only strengthened by a thin horny cuticle. A very beautiful marine species of this family (the *Coryne pusilla*), is common on seaweeds and stones between the tide-marks ; sometimes clustering parasitically round the stalks of *Tubularia* so as to form a thick beard-like mossiness ; each aggregate structure, however, not being more than an inch in length. The tentacula (as in Fig. 222, A) are short, and arise from the whole surface of the body of the polype, instead of from the margin of the mouth alone ; and at first it seems difficult to understand how they can be of service in bringing food to the mouth, which is situated at the very extremity of the branch. Observation of the living animal, however, soon removes this difficulty ; for the head is so very flexible, that the mouth can bend itself down towards any of the tentacula which may have entrapped prey ; all its movements are performed, however, in a very leisurely manner. The fresh-water genus *Cordylophora* has yet been only found in a few localities ; and the chief interest attaching to it is derived from the fact of its having been made the subject of an admirable Memoir by Prof. Allman,¹ to which every one should refer, who desires to acquaint himself with the minute organization of this group of Zoophytes. The phenomena of the Reproductive process exhibited by these Hydrozoa, are extremely curious. In *Coryne*

¹ "Philos. Transact." 1853.

and its allies, besides the ordinary gemmation which extends the original fabric, certain gemmæ are developed, which gradually come to present an organization altogether comparable to that of the simpler *Medusæ* (Fig. 222, B) and which, when detached, swim freely away. These, there is good reason to believe, stand in the same relation to the ordinary polype-buds, as the flower-buds of a plant do to its leaf-buds; each medusa-bud containing either male or female sexual organs, and performing its part in the sexual act, after it has been set free from the polype structure that bore it, just as the male (or stamiferous) flower of the *Vallisneria spiralis* discharges its pollen upon the female (or pistilline) flower, whilst floating on the surface of the water, after it has broken itself off from the stem. The ova thus fertilized, being deposited by the free swimming Medusa-buds, evolve themselves (it is probable) into single polypes, from every one of which there is gradually produced by continuous gemmation a composite fabric, that in its turn develops Medusa-buds, whose offspring resume the polype form. In *Cordylophora*, instead of detached Medusa-buds, peculiar capsules sprout forth from the stem, some of which contain sperm-cells, and others ova; and the spermatozoa set free from the former enter the ovigerous capsules and fertilize their ova, after the fashion of *Vaucheria* (§ 197). The fertilized ova undergo "segmentation" according to the ordinary type, the whole yolk-mass subdividing successively into 2, 4, 8, 16, 32 or more parts, until a "mulberry mass" is formed; this then begins to elongate itself, the surface becoming smooth, and showing a transparent margin; and this surface becomes covered with cilia, by whose agency these little bodies, now called "gemmales," first move about within the capsule, and then swim forth freely when liberated by the opening of its mouth. At this period, the embryo can be made out to consist of an outer and an inner layer of cells, with a hollow interior; after some little time, the cilia disappear, one extremity becomes expanded into a kind of disk by which it attaches itself to some fixed object; a mouth is formed, and tentacula sprout forth around it; and the body increases in length and thickness, so as gradually to acquire the likeness of one of the parent polypes, after which the plant-like structure characteristic of the genus is gradually evolved, by the successive development of polype-buds from the first formed polype and its subsequent offsets.¹

303. In the family *Tubularidæ*, the long polype-stems are invested by tubular horny sheaths; but these stop short below the polype-heads, which are consequently unprotected; and the reproductive gemmæ bud forth, as in the preceding case, from the base of the tentacula. The most common form of this family is the *Tubularia indivisa*, which receives its specific name from the infrequency with which branches are given off from the stems, these for the most part standing erect and parallel,


¹ Allman (loc. cit.).

like the stalks of wheat, upon the base to which they are attached. This beautiful zoophyte, which sometimes grows between the tide-marks, but is more abundantly obtained by dredging in deep water, often attains a size which renders it scarcely a microscopic object; its stems being sometimes no less than a foot in height, and a line in diameter. Several curious phenomena, however, are brought into view by microscopic examination. The polype-stomach is connected with the cavity of the stem by a circular opening, which is surrounded by a sphincter; and an alternate movement of dilatation and contraction takes place in it, fluid being apparently forced up from below, and then expelled again, after which the sphincter closes, in preparation for a recurrence of the operation; this, as observed by Mr. Lister, being repeated at intervals of eighty seconds. Besides the foregoing movement, a regular flow of fluid, carrying with it solid particles of various sizes, may be observed along the whole length of the stem, passing in a somewhat spiral direction, and a good deal resembling the rotation in *Chara* (§ 201). The Reproductive process in this family seems to be effected in various modes; and the true relation between them has not yet been clearly made out. The polype-stem sometimes puts forth branches, at the termination of which new polypes ultimately make their appearance, as in other composite Hydrozoa; and in the genus *Eudendrium*, which is found on many parts of our coasts, attached to old shells or stones dredged up from deep water, a beautiful tree-like structure, from three to six inches high, is thus formed. But around the polype-heads are evolved gemmæ of a different kind, as in *Coryne*; these being capsules, within which are formed either one or several ovoid bodies, that begin to develop themselves into the polype form even before their escape from their containing cases, and soon fix themselves after their immersion, shooting up into stems like those of the parent. Whether this is a method of producing free gemmæ, or whether it is a process of sexual generation, is not yet certainly known; no spermatozoa have been observed in any of these capsules; and if none should be detected by careful search, the polypes thus evolved may be presumed to be buds. In several Tubularidæ, the evolution of free Medusa-like buds, resembling those of *Syncoryne* (Fig. 222), has been observed; and all analogy would indicate that they act as the sexual organs, the development of spermatozoa in some, and of ova in others, probably not taking place until after they have led an independent life for some time. It is worthy of mention here, that when a Tubularia is kept in confinement, the polype-heads almost always drop off after a few days, but are soon renewed again by a new growth from the stem beneath; and this exuviation and regeneration may take place many times in the same individual.

304. It is in the families *Campanularidæ* and *Sertularidæ*, that the horny polypary attains its completest development; since it

not only affords an investment to the stem, but forms cups or cells for the protection of the polypes, as well as capsules for that of the reproductive bodies. In the *Campanularida*, the polype-cells are campanulate or bell-shaped, and are borne at the extremities of ringed stalks (Fig. 223, *c*); in the *Sertularida*, on the other hand, the polype-cells lie along the stem and branches, attached either to one side only, or to both sides (Fig. 224). In both, the general structure of the individual polypes (Fig. 223, *d*)

FIG. 223.



Campanularia gelatinosa.—A, upper part of the stem and branches, of the natural size; B, a small portion enlarged, showing the structure of the animal; a, terminal branch bearing polypes; b, polype-bud partially developed; c, horny cell, containing the expanded polype; d, ovarian capsule, containing medusiform gemmæ in various stages of development; f, fleshy substance extending through the stem and branches, and connecting the different polype-cells and ovarian capsules; g, annular constrictions at the base of the branches.

closely corresponds with that of the Hydra; and the mode in which they obtain their food is essentially the same. Of the products of digestion, however, a portion finds its way down into the tubular stem, for the nourishment of the general fabric; and very much the same kind of circulatory movement can be seen in *Campanularia*, as in *Tubularia*, the circulation being most vigorous in the neighborhood of growing parts. It is from the soft flesh (*f*) contained in the stem and branches, that new polype-buds (*b*) are evolved; these carry before them (so to speak) a portion of the horny integument, which at first completely invests the bud; but as the latter acquires the organization of a polype, the case thins away at its most prominent part and an opening is formed, through which the young polype protrudes itself. The origin of the bodies commonly but erroneously designated "ovarian capsules" (*e*), is exactly similar; but their destination is very different. Within them

are evolved, by a budding process, the generative organs of the

Zoophyte; and these sometimes develop themselves into the form of independent Medusæ, which completely detach themselves from the stock that bore them, make their way out of the capsule, and swim forth freely, to mature their sexual products (some developing spermatozoa, and others ova) and give origin to a new generation of polypes; whilst in other cases, these flower-buds, whose Medusan structure is less distinctly pronounced, do not completely detach themselves, but expand one after another at the mouth of the capsule, withering and dropping off after they have matured their generative products; and in other cases, again, the Medusan conformation is altogether obscured by want of development, the sexual act being performed by those bodies whilst they are still enclosed within their capsules. There is reason to believe that the male and female Medusoids are always developed within separate capsules, possibly on distinct polypidoms; the males give forth spermatozoa; whilst the females prepare ova, which, when fertilized by the entrance of spermatozoa, develop themselves into ciliated "gemmules," and these, escaping from the capsules, soon evolve themselves into true polypes. This last is the only mode of generation that has been yet witnessed among the *Sertularidæ*; for no free Medusoids have been observed to make their way out of the capsules of any members of this family (Fig. 224), within which may be seen several bodies that are commonly reputed to be eggs, but are really imperfectly developed gemmæ of the Medusan type. It is worthy of notice, that the horny capsule has been shown by Prof. E. Forbes, to be essentially a metamorphosed branch, whose numerous small cells have coalesced (as it were) into a single large one; this is made obvious by a careful comparison of the forms under which it presents itself, in different members of these two families.

305. There are few parts of our coasts, which will not supply some or other of the beautiful and interesting forms of Zoophytic life which have been thus briefly noticed, without any more trouble in searching for them, than that of examining the surfaces of rocks, stones, sea-weeds, and dead shells between the tide-marks. Many of them habitually live in that situation; and others are frequently cast up by the waves from the deeper waters, especially after a storm. Many kinds, however, can only be obtained by means of the dredge. For observing them during their living state, no means is so convenient as the zoophyte-trough (§ 69), invented for that express purpose by Mr. Lister, to whom we owe not only many improvements in the Microscope and its appurtenances, but also some of the earliest and best observations upon this class of Zoophytes which the application of the Achromatic principle permitted. Before mounting them for preservation as microscopic objects, the Author has found it best to keep them for some time in strong spirit; after a prolonged maceration in which, they may be mounted in spirit sufficiently

dilute to be destitute of any injurious action on the cement. Goadby's fluid also may be used; but the preservation of the soft parts is not quite so

FIG. 224.



Sertularia cupressina.—A, natural size; B, portion magnified.

complete with it as with spirit. The size of the cell must of course be proportioned to that of the object; and if it be desired to mount such a specimen as may serve for a characteristic illustration of the mode of growth of the species it represents, the large shallow cells, whose walls are made by cementing four strips of glass to the plate that forms the bottom (§ 136), will generally be found preferable. The horny polyparies of the Sertularidæ, when mounted in Canada balsam, are beautiful objects for the Polariscope; but in order to prepare them successfully, some nicety of management is required. The following are the outlines of the method recommended by Dr. Golding Bird, who

very successfully practised it. The specimens selected, which should not exceed two inches in length, are first to be submitted, while immersed in water of 120° , to the vacuum of an air-pump. The ebullition which will take place within the cavities, will have the effect of freeing the polyparies from dead polypes and other animal matter; and this cleansing process should be repeated several times. The specimens are then to be dried, by first draining them for a few seconds on bibulous paper, and then by submitting them to the vacuum of an air-pump, within a thick earthenware ointment pot fitted with a cover, which has been previously heated to about 200° ; by this means, the specimens are very quickly and completely dried, the water being evaporated so quickly that the cells and tubes hardly collapse or wrinkle. The specimens are then to be placed in camphine, and again subjected to the exhausting process, for the displacement of the air by that liquid; and when they have been thoroughly saturated, they should be mounted in Canada balsam in the usual mode. When thus prepared, they become very beautiful transparent objects for low magnifying powers; and they present a gorgeous display of colors when examined by polarized light, with the interposition of a plate of selenite. These objects are


peculiarly fitted for Mr. Furze's combination of the polarizing plate with the spotted lens (§ 63); as they then exhibit all the richness of coloration which the former developes, with the peculiar solidity or appearance of projection which they derive from the use of the latter.

306. No result of Microscopic research has been more unexpected, than the discovery of the close relationship subsisting between the Hydroid Zoophytes and the Medusoid Acalephæ (or jelly-fish). We have seen that many of the small free-swimming Medusans, belonging to that simple tribe of which *Thaumantias* (Fig. 225) may be taken as a representative, are really to be considered as the detached sexual apparatus of the Zoophytes from which they have been budded off, endowed with independent organs of nutrition and locomotion, whereby they become capable of maintaining their own existence and of developing their generative products. The general conformation of these organs will be understood from the accompanying figure.

Many of this group are very beautiful objects for Microscopic examination, being small enough to be viewed entire in the zoophyte-trough. There are few parts of the coast on which they may not be found, especially on a calm warm day, by skimming the surface of the sea with a fine muslin net attached to a ring, which may either be fixed to the end of a stick held in the hand, or

may be fastened by a string to the stern of the boat as a tow-net. In either case, the net should be taken up from time to time, held so as to allow the water it contains to drain through it, and then turned inside out (so that what was previously its internal surface shall now be the external), and moved about in a bucket of water, so that any minute animals adhering to it may be washed off. When we turn from these small and simple forms, to the large and highly-developed Medusans which are commonly known as "jelly-fish," we find that their history is essentially similar; for their progeny have been ascertained to develop themselves in the first instance under the polype form, and to lead a life which in all essential respects is zoophytic; their development into Medusæ taking place only in the closing phase of their existence, and then rather by gemmation from the original polype, than by a metamorphosis of its own fabric. The embryo emerges from the cavity of its parent, within which the first stages of its development have taken place, in the condition of a ciliated gemmule, of rather oblong form, very closely resembling an Infusory animalcule, but destitute of a mouth. One end soon contracts and attaches itself, however, so


Fig. 225.



Thaumantias pilosella, one of the "naked-eyed" Medusæ:—*a a*, oral tentacula; *b*, stomach; *c*, gastro-vascular canals, having the ovaries, *d d*, on either side, and terminating in the marginal canal, *e e*.

as to form a foot; the other enlarges and opens to form a mouth, four tubercles sprouting around it, which grow into tentacula; whilst the central cells melt down to form the cavity of the stomach. Thus a Hydra-like polype is formed, which soon acquires many additional tentacula; and this, according to the observations of Sir J. G. Dalyell, leads in every important particular the life of a Hydra, propagates like it by repeated gemmation, so that whole colonies are formed as offsets from a single stock, and can be multiplied like it by artificial division, each segment developing itself into a perfect Hydra. There seems to be no definite limit to its continuance in this state, or to its power of giving origin to new polype buds; but under conditions not yet ascertained, the *Strobila* (as it is termed) ceases to propagate by ordinary gemmation, and enters upon an entirely new series of changes. In the first place, the body becomes more cylindrical in form than it previously was; then a constriction or indentation is seen around it, just below the ring which encircles the mouth and gives origin to the tentacula; and similar constrictions are soon repeated around the lower parts of the cylinder, so as to give to the whole body somewhat the appearance of a rouleau of coins; a sort of fleshy bulb, somewhat of the form of the original polype, being still left at the attached extremity (Fig. 226, A). The number of circles is

FIG. 226.




Successive Stages of Development of *Medusa buds* from *Strobila larva*.—a, polype body; b, its original circle of tentacula; c, its secondary circle of tentacula; d, proboscis of most advanced *Medusa disk*; e, polype bud from side of polype body.

indefinite, and all are not formed at once, new constrictions appearing below, after the upper portions have been detached: as many as 30 or even 40 have thus been produced in one specimen. The constrictions then gradually deepen, so as to divide the cylinder into a pile of saucer-like bodies; the division being most complete above, and the upper disks usually presenting some increase in their diameter: and whilst this is taking place, the edges of the disks become divided into lobes (B), each lobe soon presenting the cleft with the supposed rudimentary eye (more probably an auditory organ) at the bottom of it, which is

to be plainly seen in the detached Medusæ (Fig. 227, c). Up to this period, the tentacula of the original polype surmount the highest of the disks; but before the detachment of the topmost disk, this circle disappears, and a new one is developed at the summit of the bulb which remains at the base of the pile (c, c). At last, the topmost and largest disk begins to exhibit a sort of convulsive struggle; it becomes detached and swims freely away; and the same series of changes takes place from above downwards, until the whole pile of disks is detached and converted into free swimming Medusæ. But the original polypoid body still remains; and may return to its polype-like and original mode of gemmation (d, e), becoming the progenitor of a new colony of Strobile, every one of which may in its turn bud off a pile of Medusa disks.

307. The bodies thus detached have all the essential characters of the adult *Medusæ*. Each consists of an umbrella-like disk, divided at its edge into a variable number of lobes, usually eight; and of a stomach, which occupies a considerable proportion of the disk, and projects downwards in the form of a proboscis, in the centre of which is the quadrangular mouth (Fig. 227, A, B). As the animal advances towards maturity, the in-

FIG. 227.




Development of *Medusæ* from the detached gemmæ of *Strobila*:—A, individual viewed sideways, and enlarged, showing the proboscis, a, and b the bifid lobes; B, individual seen from above, showing the bifid lobes of the margin, and the quadrilateral mouth; C, one of the bifid lobes still more enlarged, showing the ocellus (?) at the bottom of the cleft; D, group of young *Medusæ* as seen swimming in the water, of the natural size.

tervals between the segments of the border of the disk gradually fill up, so that the divisions are obliterated; tubular prolongations of the stomach extend themselves over the disk; and from its borders there sprout forth tendril-like filaments, which

hang down like a fringe around its margin. From the four angles of the mouth, which, even in the youngest detached animal, admits of being greatly extended and protruded, prolongations are put forth, which form the four large tentacula of the adult. And finally the generative organs make their appearance in four chambers disposed around the stomach; which are occupied by plaited membranous ribands, containing sperm-cells in the male, and ova in the female; and the embryos evolved from the latter, when they have been fertilized by the agency of the former, repeat the extraordinary cycle of phenomena which has been now described.

308. In connection with the preceding, it will be convenient to mention two curious little marine animals of frequent occurrence, the true place of which in the scale it seems difficult to determine, but which, having the free swimming habits and the soft texture of the Medusæ, have been very commonly ranked as members of the same class. One of these is the *Cydidippe pileus* (Fig. 228, A) very commonly known as the *Beroë*, which

FIG. 228.



A, *Cydidippe pileus* with its tentacles extended; B, *Beroë Forskalii*, showing the tubular prolongations of the stomach.

designation, however, properly appertains to another animal (B) of the same grade of organization. The body of *Cydidippe* is a nearly globular mass of soft jelly, usually about three-eighths of an inch in diameter; and it may be observed, even with the naked eye, to be marked by eight bright bands, which proceed from pole to pole like meridian lines. These bands are seen with the microscope to be formed of rows of large cilia, which are in a state of pretty constant vibration, though sometimes they are at rest; and if the sunlight should fall upon them when they are in activity, they display very beautiful iridescent colors. The mouth of the animal, situated at one of the poles, leads to a stomachal cavity of cylindrical shape, which extends about as far as the centre of the body, and then narrows into an intestinal tube which terminates at the opposite pole; from this stomach

tabular prolongations pass off beneath the ciliated bands, very much as in the true *Beroë* (B). In addition to the rows of cilia, the *Cydippe* is furnished with a pair of locomotive organs of a very peculiar kind; these are long tendril-like filaments, arising from the bottom of a pair of cavities in the posterior part of the body, and furnished with lateral branches (A); within these cavities they are often doubled up, so as not to be visible externally; and when they are ejected, which often happens quite suddenly, the main filaments first come forth, and the lateral tendrils subsequently uncoil themselves, to be drawn in again and packed up within the cavities, with almost equal suddenness. The liveliness of this little creature, which may sometimes be collected in large quantities at once by the muslin net, renders it a most beautiful subject for observation when due scope is given to its movements; but for the sake of microscopic examination, it is of course necessary to confine these. Various species of true *Beroë*, some of them even attaining the size of a small lemon, are occasionally to be met with on our coasts; in all of which the movements of the body are effected by the like agency of cilia, arranged in meridional bands. Very different, however, is the structure of another little globular jelly-like animal, the *Noctiluca miliaris* (Fig. 229), to which the diffused luminosity of the sea, a beautiful phenomenon that is of very frequent occurrence on our shores, is chiefly attributable. This animal is just large enough to be discerned by the naked eye, when the water in which it may be swimming is contained in a glass jar exposed to the light; and a tail-like appendage, marked with transverse rings, which is employed by the animal as an instrument of locomotion, may also be observed with a hand-glass. Near the point of its implantation in the body, is a definite mouth, on one side of which a projecting tooth has been seen by Mr. Huxley; and this mouth leads through a sort of œsophagus, into a large irregular cavity, apparently channelled out in the jelly-like substance of the body, and therefore considered by some in the light of a mere "vacuole," though by Mr. Huxley it is considered to possess regular walls; whilst from its cavity there passes forth a prolongation, which leads, in his belief, to a distinct anal orifice.¹ The external coat is denser than the contained sarcode; and the former sends thread-like prolongations through the latter, so as to divide the entire body into irregular chambers, in some of which "vacuoles" are fre-

FIG. 229.

*Noctiluca miliaris.*

¹ "Quart. Journ. of Microsc. Science," vol. iii, p. 49; see also pp. 102, 199.

quently to be seen. Very little is known about the reproduction of this animal; and until the mode in which it performs that important function shall have been made out, and it shall have also been determined whether it passes through any other phase of existence, we are scarcely in a position to speak positively of its true affinities. The nature of its luminosity is found by microscopic examination to be very peculiar; for what appears to the eye to be a uniform glow, is resolvable under a sufficient magnifying power into a multitude of evanescent scintillations; and these are given forth with increased intensity, whenever the body of the animal receives any mechanical shock, such as that produced by shaking the vessel or pouring out its contents, or is acted on by various chemical stimuli, such as dilute acids, which, however, speedily exhaust the light-producing power, occasioning disorganization of the body.

309. *Anthozoa*.—This group, which constitutes the second great division of the class of Zoophytes, includes all those larger forms, whose polypes, when expanded, present the likeness of “animal flowers:” and it consists of two principal subdivisions,—the *Asteroida*, or Alcyonian zoophytes, whose polypes, having only six or eight broad short tentacula, present a star-like aspect when expanded,—and the *Helianthoida*, whose polypes, having numerous tentacula disposed in several rows, bear a resemblance to sun-flowers or other composite blossoms. Of the first of these orders, which contains no solitary species, a characteristic example is found in the *Alcyonium digitatum* of our coasts, which is commonly known under the name of “dead-man’s toes,” or

FIG. 230.



FIG. 231.



FIG. 232.



230. Spicules of *Alcyonium* and *Gorgonia*. FIG. 231. Spicules of *Gorgonia guttata*. FIG. 232. Spicules of *Muricea elongata*.

“sea-paps.” When a specimen of this is first torn from the rock to which it has attached itself, it contracts into an unshapely mass, whose surface presents nothing but a series of slight depressions arranged with a certain regularity. But after being


immersed for a little time in a jar of sea-water, the mass swells out again, and from every one of these depressions an eight-armed polype is protruded, "which resembles a flower of exquisite beauty and perfect symmetry. In specimens recently taken, each of the petal-like tentacula is seen with a hand-glass to be furnished with a row of delicately slender *pinnæ* or filaments, fringing each margin, and arching outwards; and with a higher power, these *pinnæ* are seen to be roughened, throughout their whole length, with numerous prickly rings. After a day's captivity, however, the petals shrink up into short, thick, unshapely masses, rudely notched at their edges" (Gosse). When a mass of this sort is cut into, it is found to be channelled out, somewhat like a sponge, by ramifying canals; the vents of which open into the stomachal cavities of the polypes, which are thus brought into free communication with each other,—a character that especially distinguishes this order. A movement of fluid is kept up within these canals, as may be distinctly seen through their transparent bodies, by means of cilia lining the internal surfaces of the polypes; but no cilia can be discerned on their external surfaces. The tissue of this spongy polypidom is strengthened throughout, like that of Sponges (§ 296), with mineral spicules (always, however, calcareous), which are remarkable for the elegance of their forms; these are disposed with great regularity around the bases of the polypes, and even extend part of their length upwards on their bodies. The presence of such spicules is, in fact, a very constant character throughout this group. Thus in the *Gorgonia* or Sea-fan, whilst the central part of the polypidom is consolidated into a horny axis, the soft flesh which clothes this axis is so full of tuberculated spicules, especially in its outer layer, that, when this dries up, the spicules form a thick yellowish or reddish incrustation upon the horny stem; this is, however, so friable, that it may be easily rubbed down between the fingers, and, when examined with the Microscope, it is found to consist of spicules of different shapes and sizes, more or less resembling those shown in Figs. 230–232, sometimes colorless, but sometimes of a beautiful crimson, yellow, or purple. These spicules are best seen by the methods of illumination that give a black ground, on which they stand out with great brilliancy. They are, of course, to be separated from the animal substance in the same manner as the calcareous spicules of Sponges (§ 298); and they should be mounted, like them, in Canada balsam. It is interesting to remark that the hard calcareous stem of the *Red Coral*, which takes the place of the horny axis of the Sea-fan, is found, by the examination of very thin sections, to be made up of a solid aggregation of separate spicules, closely resembling those of Alcyonian zoophytes in general. The spicules always possess an organic basis; as is proven by the fact, that when their lime is dissolved by dilute acid, a gelatinous-

looking residuum is left, which preserves the form of the spicule, and is probably to be considered as a cell in an early stage of formation, its wall not yet being differentiated as a distinct membrane.

310. Of the order *Helianthoida*, the common *Actinia* or "Sea-Anemone" may be taken as the type; the individual polypes of all the composite structures included in the group being constructed upon the same model. In by far the larger proportion of these Zoophytes, the bases of the polypes, as well as the soft flesh that connects together the members of aggregate masses, are consolidated by calcareous deposit into stony corals; and the surfaces of these are beset with cells, usually of a nearly circular form, each having numerous lamellæ radiating from its centre towards its circumference, which are formed by the consolidation of the lower portions of the radiating partitions, that divide the space intervening between the stomach and the general integument of the animal into separate chambers. This arrangement is seen on a large scale in the *Fungia* or "mushroom coral" of tropical seas, which is the stony base of a solitary anemone-like polype; on a far smaller scale, it is seen in the little *Caryophyllia*, a like solitary polype of our own coasts, which is scarcely distinguishable from an *Actinia* by any other character than the presence of this disk, and also on the surface of many of those stony corals known as "Madrepores;" whilst in some of these the individual polype-cells are so small, that the lamellated arrangement can only be made out when they are considerably magnified. Portions of the surface of such corals, or sections taken at a small depth, are very beautiful objects for the lower powers of the Compound Microscope, the former being viewed by reflected and the latter by transmitted light. And thin sections of various fossil Corals of this group are very striking objects for the lower powers of the Oxyhydrogen Microscope. The chief point of interest to the Microscopist, however, in the structure of these animals, lies in the extraordinary abundance and high development of those "filiferous capsules," or "thread-cells," the presence of which on the tentacles of the Hydraform polypes has been already noticed (§ 300), and which are also to be found, sometimes sparingly, sometimes very abundantly, in the tentacles surrounding the mouth of the Medusæ, as well as on other parts of their bodies. If a tentacle of any of the Sea-Anemonies, so abundant on our coasts (the smaller and more transparent kinds being selected in preference), be cut off, and be subjected to gentle pressure between the two glasses of the aquatic box or of the compressorium, multitudes of little dart-like organs will be seen to project themselves from its surface near its tip; and if the pressure be gradually augmented, many additional darts will every moment come into view. Not only do these organs present different forms in different species; but

even in one and the same individual very strongly marked diversities are shown, of which a few examples are given in Fig. 233.

At A, B, C, and D, is shown the appearance of the "filiferous capsules," whilst as yet the thread lies coiled up in their interior; whilst at E, F, G, H, are seen a few of the most striking forms which they exhibit, when the thread or dart has started forth. The most probable account of their organization seems to be, that each is a cell, of which one end is extended into the thread-like or dart-like prolongation, but which is doubled in upon itself, in such a manner that the armature appears to be contained in its interior; and that the springing out of the dart is due to the eversion of the portion of the cell which had previously been pressed inwards. These thread-cells are found, however, not merely in the tentacles and other parts of the external integument of Helianthoid Zoophytes, but also in the long filaments which lie in coils within the chambers that surround the stomach, in contact with the sexual organs which are attached to the lamellæ dividing the chambers. It was formerly supposed that the last-named organs were always ovaria, and that the long and slender filaments contain sperm-cells and are consequently the male organs. But since it has been proved that the peculiar "filiferous capsules" which lie side by side in these filaments are really identical in structure with those



Filiferous Capsules of Helianthoid Polypes:—A, B, *Corynactis Allmanni*; C, E, F, *Caryophyllia Smithii*; D, G, *Actinia crassicornis*; H, *Actinia candida*.

which are found in the skin, the idea of their sexual nature has been abandoned; and a more careful examination of the organs attached to the walls of the chambers has shown that these are not always ovaries, but that they sometimes contain sperm-cells, the two sexes being here divided, not united, in the same individual. What can be the office of the filiferous filaments thus contained in the interior of the body, it is difficult to guess at. They are often found to protrude from rents in the external tegument, when any violence has been used in detaching the animal from its base; and when there is no external rupture, they are

often forced through the wall of the stomach into its cavity, and may be seen hanging out of the mouth. The largest of these capsules, in their unprojected state, are about 1-300th of an inch in length; and the thread or dart, in *Corynactis Allmanni*, when fully extended, is not less than 1-8th of an inch, or thirty-seven times the length of the capsule.¹

¹ For the fullest description of these curious bodies, as well as for much other valuable information upon Zoophytes, see Mr. Gosse's "Naturalist's Rambles on the Devonshire Coast." Those who may desire to acquire a more systematic and detailed acquaintance with this group, may be especially referred to the following Treatises and Memoirs:—Dr. Johnston's "History of British Zoophytes," Prof. Owen's "Lectures on the Comparative Anatomy and Physiology of the Invertebrate Animals," Prof. Rymer Jones's "General Outline of the Organization of the Animal Kingdom," Prof. Milne Edwards's "Recherches sur les Polypes," Prof. Van Beneden "Sur les Tubulaires," and "Sur les Campanulaires," in "Mem. de l'Acad. Roy. de Bruxelles," tom. xvii. Sir J. G. Dalyell's "Rare and Remarkable Animals of Scotland," vol. i. Trembley's "Mem. pour servir à l'histoire d'un genre de Polype d'Eau douce," M. Hollard's "Monographie du Genre *Actinia*," in "Ann. des Sci. Nat." Sér. 3, tom. xv, Mr. Mummery, "On the development of *Tubularia indivisa*," in "Transact. of Microsc. Soc." 2d Ser. vol. i, p. 28, and Prof. Max. Schultze, "On the Male Reproductive Organs of *Campanularia geniculata*," in "Quart. Journ. of Microsc. Sci." vol. iii, p. 59.


CHAPTER XII.

OF ECHINODERMATA.

311. As we ascend the scale of Animal life, we meet with such a rapid advance in complexity of structure, that it is no longer possible to acquaint one's self with any organism by microscopic examination of it as a whole; and the dissection or analysis which becomes necessary, in order that each separate part may be studied in detail, belongs rather to the Comparative Anatomist than to the ordinary Microscopist. This is especially the case with the *Echinus* (sea-urchin), *Asterias* (star-fish), and other members of the class Echinodermata; since even a general account of their complex organization would be quite foreign to the purpose of this work; whilst there are certain parts of their structure, which furnish microscopic objects of such beauty and interest that they cannot by any means be passed by; besides which, recent observations on their embryonic forms have revealed a most unexpected order of facts, the extension and verification of which will be of the greatest service to science,—a service that can only be effectually rendered by well-directed Microscopic research in fitting localities.

312. It is in the structure of that calcareous skeleton, which probably exists, under some form or other, in every member of this class, that the Microscopist finds most to interest him. This attains its highest development in the *Echinida*; in which it forms a box-like shell, or "test," composed of numerous polygonal plates jointed to each other with great exactness, and beset on its external surface with "spines," which may have the form of prickles of no great length, or may be stout club-shaped bodies, or, again, may be very long and slender rods. The intimate structure of the shell is everywhere the same; for it is composed of a *network*, which consists of carbonate of lime with a very small quantity of animal matter as a basis, and which extends in every direction (*i. e.* in thickness, as well as in length and breadth), its *areolæ* or interspaces freely communicating with each other (Fig. 234). These "areolæ," and the solid structure which surrounds them, may bear an extremely variable propor-


FIG. 234.



Section of Shell of *Echinus*, showing the calcareous network of which it is composed:—*a*, portions of a deeper layer.

mediate solid structure in the next; and their transparency is such, that when we are examining a section thin enough to contain

FIG. 235.



Transverse Section of the medullary portion of Spine of *Acrocladia*, showing its more open network.

this will be rapidly sucked up into its substance. A very beautiful example of the same kind of calcareous skeleton, having a

FIG. 236.




One of the segments of the calcareous skeleton of an Ambulacral disk of *Echinus*.

tion, one to the other; so that in two masses of equal size, the one or the other may greatly predominate; and the texture may have either a remarkable lightness and porosity, if the network be a very open one, like that of Fig. 235, or may possess a considerable degree of compactness if the solid portion be strengthened. Generally speaking, the different layers of this network, which are connected together by pillars that pass from one to the other in a direction perpendicular to their plane, are so arranged that the perforations in one shall correspond to the intermediate solid structure in the next; and their transparency is such, that when we are examining a section thin enough to contain only two or three such layers, it is easy, by properly "focussing" the Microscope, to bring either one of them into distinct view. From this very simple but very beautiful arrangement, it comes to pass that the plates of which the entire "test" is made up, possess a very considerable degree of strength, notwithstanding that their porousness is such, that if a portion of a fractured edge, or any other part from which the investing membrane has been removed, be laid upon fluid of almost any description, this will be rapidly sucked up into its substance. A very beautiful example of the same kind of calcareous skeleton, having a more regular conformation, is furnished by the disk or rosette which is contained in the tip of every one of the tubular suckers put forth by the living *Echinus* from the ambulacral pores of its shell. If the entire disk be cut off, and be mounted when dry in Canada balsam, the calcareous rosette may be seen sufficiently well; but its beautiful structure is better made out, when the animal membrane that encloses it has been got rid of

by boiling in caustic potass; and the appearance of one of the five segments of which it is composed, when thus prepared, is shown in Fig. 236.

813. The most beautiful display of this reticulated structure, however, is shown in the structure of the "spines" of *Echinus*, *Cidaris*, &c.; in which it is combined with solid ribs or pillars, disposed in such a manner as to increase the strength of these organs; a regular and elaborate pattern being formed by their intermixture, which shows considerable variety in different species. When we make a thin transverse section of almost any spine belonging to the genus *Echinus* (the small spines of our British species, however, being exceptional in this respect), we are at once made aware of the existence of a number of concentric layers, arranged in a manner that strongly reminds us of the concentric rings of an Exogenous tree (Fig. 167). The number of these layers is extremely variable; depending not merely upon the age of the spine, but (as will presently appear) upon the part of its length from which the section happens to be taken. The centre is usually occupied by a very open network (Fig. 285); and this is bounded by a row of transparent spaces (like those at *a a'*, *b b'*, *c c'*, Fig. 237), which, on a cursory inspection, might be supposed to be void spaces, but which on a closer examination are found to be the sections of solid ribs or pillars, which run in the direction of the length of the spine, and form the exterior of every layer. Their solidity becomes very obvious, when we either examine a section of a spine whose substance is pervaded (as often happens) with a coloring matter of some

FIG. 237.


Portion of transverse section of Spine of *Acrocladia mammillata*.

depth, or when we look at a very thin section by the "black-ground" illumination. Around the innermost circle of these solid pillars, there is another layer of the calcareous network, which again is surrounded by another circle of solid pillars; and this arrangement may be repeated many times, as shown in Fig. 237, the outermost row of pillars forming the projecting ribs that are very commonly to be distinguished on the surface of the spine. Around the cup-shaped base of the spine is a membrane

which is continuous with that covering the surface of the shell, and which serves not merely to hold down the cup upon the tubercle over which it works, but also, by its contractility, to move the spine in any required direction. This membrane is probably continued onwards over the whole surface of the spine, although it cannot be clearly traced to any distance from the base; and the new formations may be presumed to take place in its substance. Each new formation completely ensheaths the old; not merely surrounding the part previously formed, but also projecting considerably beyond it; and thus it happens that the number of layers shown in a transverse section, will depend in part upon the place of the section. For if it cross near the base, it will traverse every one of the successive layers from the very commencement; whilst, if it cross near the apex, it will traverse only the single layer of the last growth, notwithstanding that, in the club-shaped spines, this terminal portion may be of considerably larger diameter than the basal; and in any intermediate part of the spine, so many layers will be traversed as have been formed since the spine first attained that length. The basal portion of the spine is enveloped in a reticulation of a very close texture, without concentric layers; forming the cup or socket which works over the tubercle of the shell. The combination of elegance of pattern with richness of coloring, renders well-prepared specimens of these spines among the most beautiful objects that the Microscopist can anywhere meet with. The large spines of the various species of the genus *Acrocladia* furnish sections most remarkable for size and elaborateness as well as for depth of color (in which last point, however, the deep purple spines of *Echinus lividus* are pre-eminent); but for exquisite neatness of pattern, there are no spines that can approach those of *Echinometra heteropora* and *E. lucunter*. The spines of *Heliocidaris variolaris* are also remarkable for their beauty. No succession of concentric layers is seen in the spines of the British Echini, probably because (according to the opinion of the late Sir J. G. Dalyell) these spines are cast off and renewed every year; each new formation thus going to make an entire spine, instead of making an addition to that previously existing. Most curious indications are sometimes afforded by sections of Echinus-spines, of an extraordinary power of reparation inherent in these bodies. For irregularities are often seen in the transverse sections, which can be accounted for in no other way, than by supposing the spines to have received an injury when the irregular part was at the exterior, and to have had its loss of substance supplied by the growth of new tissue, over which the subsequent layers have been formed as usual. And sometimes a peculiar ring may be seen upon the surface of a spine, which indicates the place of a complete fracture, all beyond it being a new growth, whose unconformableness to the older or basal portion is clearly shown by a longitudinal section.

314. The spines of *Cidaris* present a marked departure from the plan of structure exhibited in *Echinus*; for not only are they destitute of concentric layers, but the calcareous network which forms their principal substance, is ensheathed in a solid calcareous cylinder perforated with tubules, which seems to take the place of the separate pillars of the Echini. This is usually found to close in the spine at its tip also; and thus it would appear that the entire spine must be formed at once, since no addition could be made either to its length or to its diameter, save on the outside of this sheath, where it is never to be found. The sheath itself often rises up in prominent points or ridges on the surface of these spines; thus giving them a character by which they may be distinguished from those of Echini. The slender, almost filamentary spines of *Spatangus* (Fig. 238), and the innumerable minute hair-like processes attached to the shell of *Clypeaster*, are composed of the like regularly reticulated sub-

FIG. 238.

Spines of *Spatangus*.

stance; and many of these are very beautiful objects for the lower powers of the Microscope, when examined by reflected light, and laid upon a black ground, without any further preparation. It is interesting also to find that the same structure presents itself in the curious *Pedicellariæ* (forceps-like bodies mounted on long stalks), which are found on the surface of many Echinida, and the nature of which has been a source of much perplexity to Naturalists, some maintaining that they are parasites, whilst others consider them as proper appendages of the *Echinus* itself. The complete conformity which exists between the structure of their skeleton and that of the animal to which they are attached, would seem to remove all reasonable doubt of their being truly appendages to it, as observation of their actions in the living state would indicate. Another example of the same structure is found in the peculiar system of plates which surrounds the interior of the oral orifice of the shell, and which gives support to the five teeth that may often be seen projecting externally through that orifice; the whole forming what is known as the "lantern of Aristotle." The

texture of the plates or jaws resembles that of the shell in every respect, save that the network is more open; but that of the teeth is much more compact. The latter have been described by Mr. Quekett as consisting of a substance not altogether unlike the "dentine" of the teeth of higher animals, save that the tubuli, though sometimes parallel, usually have more of a reticulated arrangement, and sometimes dilate into irregular "lacunæ" or spaces excavated in the hard substance.¹ The Author is not prepared to speak with confidence on this point; but he is disposed to think that the structure of the teeth is essentially the same as that of the shell, save in the interspaces of the network being much narrower; and that the appearance of tubuli (in which Mr. Quekett has not been able to make out distinct walls) is due merely to the elongation of these interspaces.

FIG. 239.

Calcareous plate and claw of *Astrophyton* (Euryale).

315. The calcareous plates which form the less compact skeletons of the *Asteriada* (star-fish and their allies) and of the *Ophiurida* (sand-stars and brittle-stars), have the same texture as those of the shell of Echinus. And this presents itself, too, in the spines or prickles of their surface, when these (as in the large *Goniaster equestris*) are large enough to be furnished with a calcareous framework, and are not mere projections of the horny integument. An example of this kind, furnished by the *Astrophyton* (better known as the *Euryale*), is represented in Fig. 239. The spines with which the arms of the species of *Ophiocoma* (brittle-star) are beset, are often remarkable for their beauty of conformation; that of *O. rosula*, one of the most common kinds, might serve (as Prof. E. Forbes justly remarked) in point of lightness and beauty, as a model for the spire of a cathedral.

316. The calcareous skeleton is very highly developed in the *Crinoidea*; their stems and branches being made up of a calcareous network, closely resembling that of the shell of the Echinus. This is extremely well seen, not only in the recent *Pentacrinus Caput Medusæ*, a somewhat rare animal of the West Indian seas, but also in a large proportion of the fossil Crinoidea, whose remains are so abundant in many of the older geological formations; for, notwithstanding that these bodies have been penetrated in the act of fossilization by a mineral infiltration, which seems to have substituted itself for the original fabric (a regularly crystalline cleavage being commonly found to exist in the fossil stems of *Enerinites*, &c., as in the fossil spines of Echinidans), yet their organic structure is often most perfectly preserved. In the circular stems of *Enerinites*, the texture of

¹ "Lectures on Histology," vol. ii, p. 234.

the calcareous network is uniform, or nearly so, throughout; but in the pentangular Pentacrini, a certain figure or pattern is formed by variations of texture in different parts of the transverse section; and the patterns, though formed upon one general plan, are sufficiently diverse in different species, to enable these to be recognized by the examination of a transverse section of a single joint of the stem.

317. The structure of the shells, spines, and other solid parts of the skeleton of Echinodermata can only be displayed by thin sections, made upon the general plan already described (§§ 109, 110). But their peculiar texture requires that certain precautions should be taken; in the first place, in order to prevent the section from breaking, whilst being reduced to the desirable thinness; and in the second, to prevent the interspaces of the network from being clogged by the particles abraded in the reducing process. A section of the shell, spine, or other portion of the skeleton, should first be cut with a fine saw, and rubbed on a flat file until it is about as thin as an ordinary card, after which it should be smoothed on one side by friction with water on a Water-of-Ayr stone. It should then be carefully dried, first on white blotting-paper, afterwards by exposure for some time to a gentle heat, so that no water may be retained in the interstices of the network, which would oppose the complete penetration of the balsam. Next, it is to be attached to a glass slip by balsam hardened in the usual manner; but particular care should be taken, first, that the balsam be brought to exactly the right degree of hardness, and second, that there be enough, not merely to attach the specimen to the glass, but also to saturate its substance throughout. The right degree of hardness is that at which the cement can be with difficulty indented by the thumb-nail; if it be made harder than this, it is apt to chip off the glass in grinding, so that the specimen also breaks away; and if it be softer, it holds the abraded particles, so that the openings of the network becomes clogged with them. If, when rubbed down nearly to the required thinness, the section appears to be uniform and satisfactory throughout, the reduction may be completed without displacing it; but if (as often happens) some inequality in thickness should be observable, or some minute air-bubbles should present themselves between the glass and the under surface, it is desirable to loosen the specimen by the application of just enough heat to melt the balsam (special care being taken to avoid the production of fresh air-bubbles), and to turn it over so as to attach the side last polished to the glass, taking care to remove or to break with the needle-point any air-bubbles that there may be in the balsam covering the part of the glass on which it is laid. The surface now brought uppermost is then to be very carefully ground down; special care being taken to keep its thickness uniform through every part (which may be even better judged of by the touch than by the eye), and to



carry the reducing process far enough, without carrying it too far. Until practice shall have enabled the operator to judge of this by passing his finger over the specimen, he must have continual recourse to the microscope during the later stages of his work; and he should bear constantly in mind, that, as the specimen will become much more transparent when mounted in balsam and covered with glass, than it is when the ground surface is exposed, he need not carry his reducing process so far as to produce at once the entire transparency he aims at, the attempt to accomplish which would involve the risk of the destruction of the specimen. In "mounting" the specimen, liquid balsam should be employed, and only a very gentle heat (not sufficient to produce air-bubbles, or to loosen the specimen from the glass) should be applied; and if, after it has been mounted, the section should be found too thick, it will be easy to remove the glass cover, and to reduce it further, care being taken to harden the balsam which has been newly laid on, to the proper degree.

§18. If a number of sections are to be prepared at once (and it is often useful to do this for the sake of economy of time, or in order to compare sections taken from different parts of the same spine), this may be most readily accomplished by laying them down, when cut off by the saw, without any preliminary preparation save the blowing the calcareous dust from their surfaces, upon a thick slip of glass well covered with hardened balsam; a large proportion of its surface may thus be occupied by the sections attached to it, the chief precaution required being that all the sections come into equally close contact with it. Their surfaces may then be brought to an exact level, by rubbing them down, first upon a flat piece of grit (which is very suitable for the rough grinding of such sections), and then upon a large Water-of-Ayr stone whose surface is "true." When this level has been attained, the ground surface is to be well washed and dried, and some balsam previously hardened is to be spread over it, so as to be sucked in by the sections, a moderate heat being at the same time applied to the glass slide; and this being increased to a sufficient degree to loosen the sections without overheating the balsam, the sections are to be turned over one by one, so that the ground surfaces are now to be attached to the glass slip, special care being taken to press them all into close contact with it. They are then to be very carefully rubbed down, until they are nearly reduced to the required thinness; and if, on examining them from time to time, their thinness should be found to be uniform throughout, the reduction of the entire set may be completed at once; and when it has been carried sufficiently far, the sections, loosened by warmth, are to be taken up upon a camel-hair brush dipped in turpentine, and transferred to separate slips of glass whereon some liquid balsam has been previously laid, in which they are to be mounted in the usual man-



ner. It more frequently happens, however, that, notwithstanding every care, the sections, when ground in a number together, are not of uniform thickness, owing to some of them being underlain by a thicker stratum of balsam than others are; and it is then necessary to transfer them to separate slips, before the reducing process is completed, attaching them with hardened balsam, and finishing each section separately.

319. It now remains for us to notice the curious and often very beautiful structures, which represent, in the order *Holothurida*, the solid calcareous skeleton of the orders already noticed. All the animals belonging to this order are distinguished by the flexibility and absence of firmness of their envelopes; and excepting in the case of certain species which have a set of calcareous plates, supporting teeth, disposed around the mouth, very much as in the Echinida, we do not find among them any representation that is apparent to the unassisted eye, of that skeleton which constitutes so distinctive a feature of the class generally. But a microscopic examination of their integument at once brings to view the existence of great numbers of minute isolated plates, every one of them presenting the characteristic reticulated structure, which are set with greater or less closeness in the substance of the skin. Various forms of the plates which thus present themselves in *Holothuria* are shown in Fig. 240; and

FIG. 240.

Calcareous plates in skin of *Holothuria*.

at A is seen an oblique view of the kind marked *a*, more highly magnified, showing the very peculiar manner wherein one part is superposed on the other, which is not at all brought into view when it is merely seen through in the ordinary manner. In the *Synapta*, one of the long-bodied forms of this order, which does not occur upon our own coasts, but is abundant in the Adriatic Sea, the calcareous plates of the integument have the regular form shown at *a*; Fig. 241; and each of these carries the curious anchor-like appendage, *c*, which is articulated to it by the notched piece at the foot, in the manner shown (in side view) at *b*. The anchor-like appendages project from the surface of the skin, and may be regarded as representing the spines of Echinida. Nearly allied to *Synapta* is the *Chirodota*, of which one species (the *C. digita*) has been previously accounted a very rare inhabi-

tant of our seas, has lately been found in considerable numbers at Torquay (by Mr. Kingsley), and might probably be met with




Fig. 241. Calcareous skeleton of *Synapta*:—A, plate imbedded in skin; B, the same, with its anchor-like spine attached; C, anchor like spine separated.

more frequently if carefully searched for. Not having had the opportunity of examining a specimen of this animal, the Author

Fig. 242.



Wheel-like plates from skin of *Chirodota violacea*.

is unable to say whether or not its integument possesses the very remarkable wheel-like plates, represented in Fig. 242, which are found in the skin of *Chirodota violacea*, a species inhabiting the Mediterranean. These plates are objects of singular beauty and delicacy, being especially remarkable for the very minute notching (scarcely to be discerned in the figures without


the aid of a magnifying glass) which is traceable round the inner margin of their "tires." There can be scarcely any reasonable doubt, that every member of this order has some kind of calcareous skeleton, disposed in a manner conformable to the examples now cited; and it would be very valuable to determine how far the very marked peculiarities by which they are respectively distinguished, are characteristic of genera and species. The plates may be obtained separately, by the usual method of treating the skin with a solution of potass; and they should be mounted in Canada balsam. But their position in the skin can only be ascertained by making sections of the integument, both vertical and parallel to its surface; and these sections, when dry, are most advantageously mounted in the same medium, by which their transparency is greatly increased. All the objects of this class are most beautifully displayed by the black-ground illumination (§§ 61, 62); and the same method, when applied to very thin sections of Echinus-spines, brings out some effects of marvellous beauty.

320. *Echinoderm Larvæ*.—We have now to notice that most remarkable set of objects, furnished to the microscopic inquirer by the larval forms of this class, for our present knowledge of which, imperfect as it still is, we are already so largely indebted to

the painstaking and widely extended investigations of Prof. Müller. All that our limits permit, is a notice of two of the most curious forms of these larvæ, by way of sample of the wonderful phenomena which his researches have brought to light; so as (it may be hoped) to excite such an interest among those Microscopists in particular who may have the opportunity of pursuing these inquiries, as may induce them to apply themselves perseveringly to them, and thus to supply the numerous links which are at present wanting in the chain of developmental history. The peculiar feature by which the early history of the Echinoderms generally seems to be distinguished, is this,—that the embryonic mass of cells is converted, not into a larva which subsequently attains the adult form by a process of metamorphosis, but into a peculiar *zooid*, which seems to exist for no other purpose than to give origin to the Echinoderm by a kind of internal gemmation, and to carry it to a distance by its active locomotive powers, so as to prevent the spots inhabited by the respective species from being overcrowded by the accumulation of their progeny. The larval zooids are formed upon a type quite different from that which characterizes the adults; for instead of a *radial* symmetry, they exhibit a *bilateral*, the two sides being precisely alike, and each having a ciliated fringe along the greater part of the whole of its length. The two fringes are united by a superior and an inferior transverse ciliated band; and between the two, the mouth of the zooid is always situated. Further, although the adult Star-fish and Sand-stars have neither intestinal tube nor anal orifice, their larval zooids, like those of other Echinoderms, always possess both. The external forms of these larvæ, however, vary in a most remarkable degree, owing to the unequal evolution of their different parts; and there is also a considerable diversity in the several orders, as to the proportion of the fabric of the larva which enters into the composition of the adult form. In the fully developed Star-fish and Sea-urchin, the only part retained is a portion of the stomach and intestine, which is pinched off, so to speak, from that of the Larval zooid.

321. One of the most remarkable forms of Echinoderm

FIG. 243.



Bipinnaria asterigera, or Larva of Star-Fish:—a, mouth; a', oesophagus; b, intestinal tube and anal orifice; c, furrow in which the mouth is situated; d d', bilobed peduncle; 1, 2, 3, 4, 5, 6, 7, ciliated arms.

larvæ is that which has received the name of *Bipinnaria* (Fig. 243), from the symmetrical arrangement of its natatory organs. The mouth (*a*), which opens in the middle of a transverse furrow, leads through an œsophagus *b'* to a large stomach, around which the body of a Star-fish is developing itself; and on one side of this mouth is observed the intestinal tube and anus (*b*). On either side of the anterior portion of the body, are six or more narrow fin-like appendages, which are fringed with cilia; and the posterior part of the body is prolonged into a sort of pedicle, bilobed towards its extremity, which also is covered with cilia. The organization of this larva seems completed, and its movements through the water are very active, before the mass at its anterior extremity presents anything of the aspect of the Star-fish; in this respect corresponding with the movements of the "pluteus" of the Echinida (§ 322). The temporary mouth of the larva does not remain as the permanent mouth of the Star-fish; for the œsophagus of the latter enters on what is to become the dorsal side of its body, and the true mouth is subsequently formed by the thinning away of the integument on its ventral surface. The young Star-fish is separated from the bipinnarian larva, by the forcible contractions of the connecting pedicle, as soon as the calcareous consolidation of its integument has taken place, and its true mouth has been formed, but long before it has attained the adult condition; and as its ulterior development has not hitherto been observed in any instance, it is not yet known what are the species in which this mode of evolution prevails. The larva continues active for several days after its detachment; and it is possible, though perhaps scarcely probable, that it may develop another Asteroid by a repetition of this process of gemmation.¹

322. In the *Bipinnaria*, as in other larva zooids of the Asterozoa, there is no internal calcareous framework; such a framework, however, is found in the larvæ of the Echinida and Ophiurida, of which the form delineated in Fig. 244 is an example.² The embryo issues from the ovum as soon as it has attained, by the repeated segmentation of the yolk, the condition of the "mulberry mass;" and the superficial cells of this are covered with cilia, by whose agency it swims freely through the water. So rapid are the early processes of development, that no more than from twelve to twenty-four hours intervene between fecundation


¹ See the observations of Kœren and Danielsen (of Bergen) in the "Zoologische Bidrag," Bergen, 1847 (translated in the "Ann. des Sci. Nat." 3e Sér. Zool. tom. iii, p. 347); and the Memoir of Prof. Müller, "Ueber die Larven und die Metamorphose der Echinodermen," in "Abhandlungen der Königlichen Akademie der Wissenschaften zu Berlin," 1848.

² See Prof. Müller, "Ueber die Larven und die Metamorphose der Ophiuren und Seeigel," in "Abhandlungen der Königlichen Akademie der Wissenschaften zu Berlin," 1846. See also, for the earlier stages, a Memoir by M. Derbès, in "Ann. des Sci. Nat." 3e Sér. Zool. tom. viii, p. 80; and for the later, Krohn's "Beitrag zur Entwicklungsgeschichte der Seeigellarven," Heidelberg, 1849, and his Memoir in "Müller's Archiv," 1851.

and the emersion of the embryo; the division into two, four, or even eight segments taking place within three hours after impregnation. Within a few hours after its emersion, the embryo changes from the spherical into a sub-pyramidal form with a flattened base; and in the centre of this base is a depression, which gradually deepens, so as to form a mouth that communicates with a cavity in the interior of the body, which is surrounded by a portion of the yolk-mass that has returned to the liquid granular state. Subsequently a short intestinal tube is found, with an anal orifice, opening on one side of the body. The pyramid is at first triangular, but it afterwards becomes quadrangular; and the angles are greatly prolonged round the mouth (or base), whilst the apex of the pyramid is sometimes much extended in the opposite direction, but is sometimes rounded off into a kind of dome (Fig. 244, A). All parts of this curious body, and especially its most projecting portions, are strengthened by a framework of thread-like calcareous rods (e). In this condition, the embryo swims freely through the water, being propelled by the action of cilia, which clothe the four angles of the pyramid and its projecting arms, and which are sometimes thickly set upon two or four projecting lobes (f); and it has received the designation of *Pluteus*. The mouth is usually surrounded by a sort of proboscis, the angles of which are prolonged into four slender processes (g, g, g, g), shorter than the four outer legs, but furnished with a similar calcareous framework.

328. The first indication of the production of the young Echi-

FIG. 244.



Embryonic development of *Echinus*:—A, *Pluteus* larva at the time of the first appearance of the disk; a, mouth in the midst of the four-pronged proboscis; b, stomach; c, echinoid disk; d, d, d, d, four arms of the *Pluteus* body; e, calcareous framework; f, ciliated lobes; g, g, g, g, ciliated processes of the proboscis:—b, disk with the first indication of the cirrhi:—c, disk, with the origin of the spines between the cirrhi:—d, more advanced disk, with the cirrhi and spines projecting considerably from the surface. (N.B. In Figs. b, c, and d, the pluteus is not represented, its parts having undergone no change, save in becoming relatively smaller.)

nus from its "pluteus," is given by the formation of a circular disk (Fig. 244, A, e), on one side of the central stomach (b); and this disk soon presents five prominent tubercles (B), which subsequently become elongated into tubular cirrhi. The disk gradually extends itself over the stomach, and between its cirrhi the rudiments of spines are seen to protrude (c); these, with the cirrhi, increase in length, so as to project against the envelope of the "pluteus," and to push themselves through it; whilst, at the same time, the original angular appendages of the "pluteus" diminish in size, the ciliary movement becomes less active, being superseded by the action of the cirrhi and spines, and the mouth of the "pluteus" closes up. By the time that the disk has grown over half of the gastric sphere, very little of the "pluteus" remains, except some of the slender calcareous rods; and the number of tentacula and spines rapidly increases. The calcareous framework of the shell at first consists, like that of the Starfishes, of a series of isolated networks developed between the cirrhi; and upon these rest the first formed spines (D). But they gradually become more consolidated, and extend themselves over the granular mass, so as to form the series of plates. The mouth of the Echinus (which is altogether distinct from that of the "pluteus") is formed at that side of the granular mass, over which the shell is last extended; and the first indication of it consists in the appearance of five calcareous concretions, which are the summits of the five portions of the framework of jaws and teeth that surround it. All traces of the original "pluteus" are now lost; and the larva, which now presents the general aspect of an Echinoid animal, gradually augments in size, multiplies the number of its plates, cirrhi, and spines, evolves itself into its particular generic and specific type, and undergoes various changes of internal structure, tending to the development of the complete organism. In collecting the free-swimming larvæ of Echinodermata, a fine muslin net should be employed in the manner already described (§ 306); and the search for them is of course most likely to be successful in those localities in which the adult animals of the respective species abound, and on warm calm days, in which they seem to come to the surface in the greatest numbers.

324. It is remarkable that the *Comatula*, one of the most active of all Echinoderms in its adult state, passes a portion of the early period of its life in a fixed condition; being attached by a stem to sea-weeds or zoophytes, precisely after the manner of the *Crinoids* or "lily-stars," which were the most common types of this class in the older epochs of the world's history. In this phase of its life, which was first discovered by Mr. J. V. Thompson, of Cork, in 1823, it is very minute, and forms a most beautiful object for the lower magnifying powers, when viewed in fluid by a strong incident light, as nearly as possible in its natural condition. It has hitherto been found so rarely, however, the Micro-

scopists have been able to become possessed of it; but the Author has been fortunate enough to discover a locality (Lamlash Bay, in the Isle of Arran) in which it is so abundant, that it may hereafter find its way into almost every cabinet. It has been made next to certain, by the observations of Busch, that this fixed stage is preceded by a free-swimming larval condition; and the passage from one phase to the other is a problem of the greatest interest, which the Author hopes to have it in his power to work out.


CHAPTER XIII.

POLYZOA, AND COMPOUND TUNICATA.

At the lower extremity of the great series of Molluscan animals we find two very remarkable groups, whose mode of life has much in common with Zoophytes, whilst their type of structure is conformable in all essential particulars to that of the true Mollusks. These animals are for the most part microscopic in their dimensions; and as some members of both these groups are found on almost every coast, and are most interesting objects for anatomical examination, as well as for observation in the living state, a brief general account of them will be here appropriate.

325. *Polyzoa*.—The group which is known under this name to British naturalists, corresponds with that which by Continental

FIG. 245.



Cells of *Lepadidæ*:—A, *L. Hyndmanni*; B, *L. Agularis*; C, *L. verrucosa*.

zoologists is designated *Bryozoa*: the former name (though first used in the singular instead of the plural number), having been introduced by Mr. J. V. Thompson in a memoir published in 1830, seems to have precedence in point of time over the latter, which was conferred by Prof. Ehrenberg in 1831 on a most heterogeneous group, wherein the *Bryozoa*, as now limited, were combined with the *Foraminifera*. As the history of the researches by which the *Polyzoa* have been raised from the class of *Zoophytes* (in which they

were formerly ranked, for the most part in apposition with the *Hydrozoa*), to the Molluscan sub-kingdom, has already been sketched (p. 49), we may now proceed, without further preface, to a survey of the leading features of their organization. The

animals of the Polyzoa, in consequence of their universal tendency to multiplication by gemmation, are seldom or never found solitary, but form clusters or colonies of various kinds; and as each is enclosed in either a horny or calcareous sheath, or "cell," a composite structure is formed, closely corresponding with the polypidom of a Zoophyte, which has been appropriately designated the "polyzoary." The individual cells of the "polyzoary" are sometimes only connected with each other by their common relation to a creeping stem or "stolon," as in *Laguncula* (Fig. 246); but more frequently they bud forth directly, one from another, and extend themselves in different directions over plane surfaces, as is the case with *Flustrea*, *Lepralia*, &c. (Fig. 245); whilst not unfrequently the Polyzoary develops itself into an arborescent structure (Fig. 247) which may even present somewhat of the density and massiveness of the stony Corals. Each individual is composed externally of a sort of sac, of which the outer or tegumentary layer is either simply membranous, or is horny, or in some instances calcified, so as to form the cell; this investing sac is lined by a more delicate mem-

FIG. 246.



Laguncula repens, as seen in its expanded state at A, and in its contracted state, in two different aspects, at B and C. The same references answer for each figure:—*a a*, tentacula clothed with vibratile cilia; *b*, pharyngeal cavity; *c*, valve separating this cavity from *d* the œsophagus; *e*, the stomach, with *f* its pyloric valve, and *g* the circle of cilia surrounding that orifice; *h*, wall of the stomach with biliary follicles; *i*, the intestine, containing *k* excrementitious matter, and terminating at *l* the anus; *m*, the testicle; *n*, the ovary; *o*, an ovum set free from the ovary; *p*, openings for the escape of the ova; *q*, spermatozoa freely moving in the cavity that surrounds the viscera; *r*, retractor muscle of the angle of the aperture of the sheath; *s*, retractor of the sheath; *t*, retractor of the tentacular circle; *u*, retractor of the œsophagus; *v*, retractor of the stomach; *w*, principal extensor muscle; *x*, transverse wrinkles of the sheath; *y*, fibres of the sheath, themselves probably muscular; *z*, muscles of the tentacula; *a* (at the base of the tentacular circle in A), nervous or œsophageal ganglion; *β*, stem, —*n*, a portion of the tentacular circle shown separately on a larger scale; *a a*, the tentacula clothed with cilia; *b b*, their internal canals; *c*, muscles of the tentacula; *d*, transverse muscles forming a ring at the base of the tentacula; *e*, muscles of the tentacular circle.

brane, which closes its orifice, and which then becomes continuous with the wall of the alimentary canal; this lies freely in the visceral sac, floating (as it were) in the liquid which it contains. The further details of the anatomy will be best understood from the examination of a characteristic example, such as the *Laguncula repens*; which is shown in the state of expansion at *A*, Fig. 246, and in the state of contraction at *B* and *C*. The mouth is surrounded by a circle of tubular tentacula, which are clothed with vibratile cilia; these tentacula, in the species we are considering, vary from ten to twelve in number; but in some other instances they are more numerous. By the ciliary investment of their tentacula, the Polyzoa are at once distinguishable from those Hydraform polypes to which they bear a superficial resemblance, and with which they were at one time confounded; and accordingly, whilst still ranked among the Zoophytes, they were characterized as *Ciliobrachiata*. The tentacula are seated upon an annular disk, which is termed the "lophophore," and which forms the roof of the visceral or perigastric cavity; and this cavity extends itself into the interior of the tentacula, through perforations in the "lophophore." The mouth, situated in the centre of the "lophophore," leads to a funnel-shaped cavity, or pharynx, *b*, which is separated from the œsophagus, *d*, by a valve at *c*; and this œsophagus opens into the stomach, *e*, which occupies a considerable part of the visceral cavity. In the *Bowerbankia*, and some other Polyzoa, a muscular stomach or gizzard, for the trituration of the food, intervenes between the œsophagus and the true digestive stomach. The walls of the stomach, *h*, have considerable thickness; and they are beset with minute follicles, which seem to have the character of a rudimentary liver. This, however, is more obvious in some other members of the group. The stomach is lined, especially at its upper part, with vibratile cilia, as seen at *c*, *g*; and by the action of these, the food is kept in a state of constant agitation during the digestive process. From the upper part of the stomach, which is (as it were) doubled upon itself, the intestine *i* opens, by a pyloric orifice *f*, which is furnished with a regular valve; within the intestine are seen at *k* particles of excrementitious matter; which are discharged by the anal orifice at *l*. No circulating apparatus here exists; but the liquid which fills the cavity that surrounds the viscera, contains the nutritive matter which has been prepared by the digestive operation, and which has transuded through the walls of the alimentary canal; a few corpuscles of irregular size are seen to float in it. The visceral sacs of the different individuals put forth from the same stem, appear to communicate with each other. No other respiratory organs exist than the tentacula; into whose cavity the nutritive fluid is probably sent from the visceral cavity, for aeration by the current of water that is continually flowing over them.

326. The production of gemmæ may take place either from

the bodies of the animals themselves, which is what always happens when the cells are in mutual apposition; or from the connecting stem or stolon, where the cells are detached from each other, as in *Laguncula*. There is first seen a bud-like protuberance of the horny external integument, into which the soft membranous lining prolongs itself; the cavity thus formed, however, is not to become (as in Hydra and its allies) the stomach of the new zooid; but it constitutes the chamber surrounding the digestive viscera, which organs have their origin in a thickening of the lining membrane, that projects from one side of the cavity into its interior, and gradually shapes itself into the alimentary canal with its tentacular appendages. Of the production of gemmæ from the zooids themselves, the best examples are furnished by the *Flustra* and their allies. From a single cell of a *Flustra*, five such buds may be sent off, which develop themselves into new zooids around it; and these, in their turn, produce buds from their unattached margins, so as rapidly to augment the number of cells to a very large amount. To this extension there seems no definite limit; and it often happens that the cells in the central portion of the leaf-like expansion of a *Flustra* are devoid of contents and have lost their vitality, whilst the edges are in a state of active growth. Independently of their propagation by gemmation, the Polyzoa have a true sexual generation; the sexes, however, being usually, if not invariably, united in the same individuals. The sperm-cells are developed in a glandular body, the testicle *m*, which lies beneath the base of the stomach; when mature, they rupture, and set free the spermatozoa *q*, which swim freely in the liquid of the visceral cavity. The ova, on the other hand, are formed in an ovarium *n*, which is lodged in the membrane lining the tegumentary sheath, near its outlet; the ova, having escaped from this into the visceral cavity, as at *o*, are fertilized by the spermatozoa which they there meet with; and are finally discharged by an outlet at *p*, beneath the tentacular circle.

327. These creatures possess a considerable number of muscles, by which their bodies may be projected from their sheaths or drawn within them; of these muscles, *r, s, t, u, v, w, x*, the direction and points of attachment sufficiently indicate the uses; they are for the most part *retractors*, serving to draw in and double up the body, to fold together the circle of tentacula, and to close the aperture of the sheath, when the animal has been completely withdrawn into its interior. The projection and expansion of the animal, on the contrary, appear to be chiefly accomplished by a general pressure upon the sheath, which will tend to force out all that can be expelled from it. The tentacula themselves are furnished with distinct muscular fibres, by which their separate movements seem to be governed; the arrangement of these is seen at *D*. At the base of the tentacular circle, just above the anal orifice, is a small body (seen at *A, a*), which is a nervous

ganglion; as yet no branches have been distinctly seen to be connected with it in this species; but its character is less doubtful in some other Polyzoa.

328. If we scrutinize the foregoing characters, we shall find that the most important of them are Molluscan, rather than Zoophytic. In the first place, all true Polypes use their tentacula to grasp their food and convey it to the mouth; and these tentacula are destitute of cilia; whilst, on the other hand, in all the Acephalous Mollusca, the nutritive matter is drawn in by a ciliary current, which also serves to aerate the fluids. Now the latter, as we have just seen, is the case with the Polyzoa; and thus, although their arms very commonly present a circular disposition around the mouth, they may be considered as representing, in their relation to the economy of the animal, the ciliated branchial sac of the Ascidians (§ 331). But they do not by any means constantly present this radial symmetry; thus, in the *Plumatella*, a beautiful fresh-water genus of Polyzoa, the ciliated arms are set upon two lobes or projections, one on either side of the mouth. The structure of the alimentary canal, again, removes the Polyzoa from the zoophytic series. In no true polype is there a separate intestine and anal orifice, nor does the whole apparatus hang freely in the visceral cavity; and the existence of a gizzard-like organ, and of a rudimentary liver (closely resembling that found in the lowest Tunicata), are also characters of elevation. The most important of all the single characters furnished by the anatomy of these animals, is their nervous system; which, as already pointed out, is distinctly Molluscan in its type. The absence of a heart and a distinct circulating system is, it is true, a Zoophytic character; but we shall presently find that even in the Tunicata, which are true Mollusks, the character of the circulating apparatus is extremely degraded. The propagation by gemmation, although formerly supposed to be a character exclusively Zoophytic, is known to belong also to the greater part of the "tunicated" Mollusks; and from this, therefore, no argument can be drawn in favor of the zoophytic nature of the Polyzoa. And although many of their composite fabrics have a stony density, and closely resemble the solid polypidoms of the helianthoid and asteroid Polypes, yet in others, especially amongst the fresh-water species, we find a very close resemblance to the gelatinous bed or leathery crust in which the Compound Ascidians are lodged; and if we imagine calcareous matter to be deposited in this bed or crust, we should have a fabric closely resembling that of many stony polyzoaries.

329. Of all the Polyzoa of our own coasts, the *Flustræ* or "sea-mats" are the most common; these present flat expanded surfaces, resembling in form those of many sea-weeds (for which they are often mistaken), but exhibiting, when viewed, even with a low magnifying power, a most beautiful network, which at once indicates their real character. The cells are arranged on

both sides; and it has been calculated by Dr. Grant, that as a single square inch of an ordinary Flustra contains 1800 such cells, and as an average specimen presents about 10 square inches of surface, it will consist of no fewer than 18,000 zooids. The want of transparency in the cell-wall, however, and the infrequency with which the animal projects its body far beyond the mouth of the cell, renders the Polyzoa of this genus less favorable subjects for microscopic examination, than are those of the *Bowerbankia*, a Polyzoan with a trailing stem and separated cells like those of *Laguncula*, which is very commonly found clustering around the bases of Flustræ. It was in this, that many of the details of the organization of the interesting group we are considering, were first studied by Dr. A. Farre, who discovered it in 1837, and subjected it to a far more minute examination than any Polyzoan had previously received;¹ and it is one of the best adapted of all the marine forms yet known, for the display of the beauties and wonders of this type of organization. The *Halodactylus* (formerly called *Alcyonidium*), however, is among the most remarkable of all the marine forms, for the comparatively large size of the tentacular crowns; these, when expanded, being very distinctly visible to the naked eye, and presenting a spectacle of the greatest beauty when viewed under a sufficient magnifying power. The polyzoary of this genus has a spongy aspect and texture, very much resembling that of the Alcyonian Zoophytes, for which it might readily be mistaken when its contained animals are all withdrawn into their cells; when these are expanded, however, the aspect of the two is altogether different, as the minute plumose tufts which then issue from the surface of the *Halodactylus*, making it look as if it were covered with the most delicate downy film, are in striking contrast with the larger, solid-looking polypes of the Alcyonium. The opacity of the polyzoary of the *Halodactylus* renders it quite unsuitable for the examination of anything more than the tentacular crown and the œsophagus which it surmounts; the stomach and the remainder of the visceral apparatus being always retained within the cell. Several of the fresh-water Polyzoa are peculiarly interesting subjects for microscopic examination; alike on account of the remarkable distinctness with which the various parts of their organization may be seen, and the very beautiful manner in which their ciliated tentacula are arranged upon a deeply crescentic or horse-shoe-shaped "lophophore." By this peculiarity, the fresh-water Polyzoa are separated as a distinct sub-class from the marine; the former being designated as *Hippocrepiæ* (horseshoe-like), while the latter are termed *Infundibulata* (funnel-like).

330. The *Infundibulata* or Marine Polyzoa, constituting by far the most numerous division of the class, are divided into four

¹ See his Memoir in the "Philosophical Transactions," for that year.

orders, as follows;—I. *Cheilostomata*, in which the mouth of the cell is sub-terminal, or not quite at its extremity (Fig. 245), is somewhat crescentic in form, and is furnished with a movable (generally membranous) lip, which closes it when the animal retreats. This includes a large part of the species that most abound on our own coasts, notwithstanding their wide differences in form and habit. Thus the polyzoaries of some (as *Flustra*) are horny and flexible, whilst those of others (as *Eschara* and *Retepora*) are so penetrated with calcareous matter as to be quite rigid; some grow as independent plant-like structures (as *Bugula* and *Gemellaria*), whilst others, having a like arborescent form, creep over the surfaces of rocks or stone (as *Hippothoa*), and others, again, have their cells in close apposition, and form crusts which possess no definite figure (as is the case with *Lepralia* and *Membranipora*). A large proportion of the Polyzoa of this order are furnished with very peculiar motile appendages, which are of two kinds, *avicularia* and *vibracula*. The “avicularia,” or “bird’s-head processes,” are so named from the striking resemblance they present to the head and jaws of a bird (Fig. 247, B). They are generally “sessile” upon the angles or margins of the

FIG. 247.



A, Portion of *Cellularia ciliata*, enlarged; B, one of the “bird’s-head processes” of *Bugula avicularia*, more highly magnified, and seen in the act of grasping another.

cells, that is, are attached at once to them, without the intervention of a stalk, as in Fig. 247, A, being either “projecting” or “immersed;” but in the genera *Bugula* and *Bicellaria*, where they are present at all, they are “pedunculate” or mounted on footstalks (B). Under one form or the other, they are wanting in but few of the genera belonging to this order; and their presence or absence furnishes valuable characters for the discrimination of species. Each avicularium has two “mandibles,” of which one is fixed, like the upper jaw of a bird, the other movable like its lower jaw; the latter is opened and closed by two sets of muscles which are seen in the interior of the “head;” and

between them is a peculiar body, furnished with a pencil of

bristles, which is probably a tactile organ, being brought forwards when the mouth is open, so that the bristles project beyond it, and being drawn back when the mandible closes. The "avicularia" keep up a continual snapping action, during the life of the polyzoary; and they may often be observed to lay hold of minute worms or other bodies, sometimes even closing upon the beaks of adjacent organs of the same kind, as shown in Fig. 247, B. In the pedunculate forms, besides the snapping action, there is a continual rhythmical nodding of the head upon the stalk; and few spectacles are more curious than a portion of the polyzoary of *Bugula avicularia* (a very common British species) in a state of active vitality, when viewed under a power sufficiently low to allow a number of these bodies to be in sight at once. It is still very doubtful what is their precise function in the economy of the animal; whether it is to retain bodies that may serve as food within the reach of the ciliary current, or whether it is, like the "pedicellaria" of Echini (§ 314), to remove extraneous particles that may be in contact with the surface of the polyzoary. The latter would seem to be the function of the *vibracula*, which are long bristle-shaped organs, each one springing at its base out of a sort of cup (Fig. 245, A), that contains muscles by which it is kept in almost constant motion, sweeping slowly and carefully over the surface of the polyzoary, and removing what might be injurious to the delicate inhabitants of the cells when their tentacula are protruded. Out of 191 species of Cheilostomatous Polyzoa described by Mr. Busk, no fewer than 126 are furnished either with "avicularia," or with "vibracula," or with both of these organs.¹ II. The second order, *Cyclostomata*, consists of those Polyzoa which have the mouth at the termination of tubular calcareous cells, without any movable appendage or lip. This includes a comparatively small number of genera, of which *Crinia* and *Tubulipora* contain the largest proportion of the species that occur on our own coasts. III. The distinguishing character of the third order, *Ctenosomata*, is derived from the presence of a comb-like circular fringe of bristles, connected by a delicate membrane, around the mouth of the cell, when the animal is projected from it; this fringe being drawn in when the animal is retracted. The polyzoaries of this group are very various in character, the cells being sometimes horny and separate (as in *Laguncula* and *Bowerbankia*), sometimes fleshy and coalescent (as in *Halodactylus*). IV. In the fourth order, *Pedicellineæ*, which includes only a single genus, *Pedicellina*, the lophophore is produced upwards on the back of the tentacles, uniting them at their base in a sort of muscular calyx, and giving to the animal when expanded somewhat the form of an inverted bell, like that of

¹ See Mr. G. Busk's "Remarks on the Structure and Function of the Avicularian and Vibracular Organs of Polyzoa," in "Transact. of Microscop. Soc." Ser. II, vol. ii, p. 26.

Vorticella (Fig. 196). Among the *Hippocrepia* may be noticed, as exceptional forms, the *Cristatella*, whose polyzoary is unattached, so as to be capable of moving freely through the water, and the *Fredericella*, the lophophore of which is rather circular than crescentic, the prolongation being so slight as only to be discernible on a careful examination. Generally speaking, the cells are lodged in a sort of gelatinous substratum, which spreads over the leaves of aquatic plants, sometimes forming masses of considerable size. As the animals of this group altogether resemble the true Zoophytes in their habits, and are found in the same localities, it is not requisite to add anything to what has already been said (§ 305) respecting the collection, examination, and mounting, of this very interesting class of objects.¹


331. *Compound Tunicata*.—The Tunicated Mollusca are so named from the enclosure of their bodies in a "tunic" which is sometimes leathery or even cartilaginous in its texture, and which very commonly includes calcareous spicules, whose forms are often very beautiful. They present a strong resemblance to the Polyzoa, not merely in their general plan of conformation, but also in their tendency to produce composite structures by gemmation; they are differentiated from them, however, by the absence of the ciliated tentacula, which form so conspicuous a feature in the external aspect of the Polyzoa, by the presence of a distinct circulating apparatus, and by their peculiar respiratory apparatus, which may be regarded as a dilatation of their pharynx. In their habits, too, they are more inactive, exhibiting scarcely anything comparable to those rapid movements of expansion and retraction, which it is so interesting to watch among the Polyzoa; whilst, with the exception of the *Salpidae* and other floating species which are chiefly found in seas warmer than those that surround our coast, they are rooted to one spot during all but the earliest period of their lives. The larger forms of the Ascidian group, which constitutes the bulk of the class, are always solitary; either not propagating by gemmation at all, or, if this process does take place, the gemmæ being detached before they have advanced far in their development. Since these cannot be considered as Microscopic objects (although no part of their organization can be properly made out without the assistance of that instrument), our attention will be confined to those "Compound Ascidiæ," the small size and transparency of whose bodies, when detached from the mass in which they are imbedded, not only enables their structure to be clearly

¹ For a more detailed account of the Structure and Classification of this group, see Prof. Allman's "Report on the Fresh-water Polyzoa" in the "Transactions of the British Association" for 1850; Prof. Van Beneden's "Recherches sur les Bryozoaires de la Côte d'Ostende," in "Mem. de l'Acad. Roy. de Bruxelles," tom. xvii; Mr. G. Busk's "Catalogue of the Marine Polyzoa in the Collection of the British Museum;" and Dr. G. Johnston's "History of British Zoophytes."

discerned without dissection, but allows many of their living actions to be watched. Of these we have a characteristic example in *Amaroucium proliferum*; of which the form of the composite mass and the anatomy of a single individual are displayed in Fig. 248. The clusters of these animals present themselves on rocks, sea-weeds, &c., very commonly between the tide-marks. They appear almost completely inanimate, exhibiting no very obvious movements when irritated; but if they be placed when fresh in sea water, a slight pouting of the orifices will soon be perceptible, and a constant and energetic series of currents will be found to enter by one set and to be ejected by the other, indicating that all the machinery of active life is going on within these apathetic bodies. In the tribe of *Polyclinians*, to which this genus belongs, the body is elongated, and may be divided into three regions, the thorax (A), which is chiefly occupied by the respiratory sac, the abdomen (B), which contains the digestive apparatus, and the post-abdomen (c), in which the heart and generative organs are lodged. At the summit of the thorax is seen the oral orifice, *c*, which leads to the branchial sac, *e*; this is perforated by an immense number of slits, which allow part of the water to pass into the space between the branchial sac and the muscular mantle, where it is especially collected in the thoracic sinus, *f*. At *k* is seen the œsophagus, which is continuous with the lower part of the pharyngeal cavity; this leads to the stomach, *l*, which is surrounded by biliary tubuli; and from this passes off the intestine, *m*, which terminates at *n* in the cloaca. The long post-abdomen is principally occupied by the large ovarium, *p*, which contains ova

in various stages of development. These, when matured and set free, find their way into the cloaca; where two large ova are seen (one marked *p'* and the other immediately below

FIG. 248.




Compound mass of *Amaroucium proliferum*, with the anatomy of a single zooid:—A, thorax; B, abdomen; c, post-abdomen:—c, oral orifice; e, branchial sac; f, thoracic sinus; k, anal orifice; *s'*, projection overhanging it; j, nervous ganglion; k, œsophagus; l, stomach surrounded by biliary tubuli; m, intestine; n, termination of intestine in cloaca; o, heart; *o'*, pericardium; p, ovarium; *p'*, egg ready to escape; q, testis; r, spermatic canal; *r'*, termination of this canal in the cloaca.

it), waiting for expulsion. In this position they receive the fertilizing influence from the testis, *q*, which discharges its products by the long spermatic canal, *r*, that opens into the cloaca at *r'*. At the very bottom of the post-abdomen, we find the heart, *o*, enclosed in its pericardium, *o'*. In the tribe we are now considering, a number of such animals are imbedded together in a sort of gelatinous mass, and covered with an integument common to them all; the composition of this gelatinous substance is remarkable as including "cellulose," which generally ranks as a purely vegetable product. The mode in which new individuals are developed in this mass, is by the extension of "stolons" or creeping stems from the bases of those previously existing; and from each of these stolons several buds may be put forth, every one of which may evolve itself into the likeness of the stock from which it proceeded, and may in its turn increase and multiply after the same fashion. A communication between the circulating systems of the different individuals is kept up, through their connecting stems, during the whole of life; and thus their relationship to each other is somewhat like that of the several polypes on the polypidom of a *Campanularia* (§ 304).

332. In the family of *Didemnians*, the post-abdomen is absent,

the heart and generative apparatus being placed by the side of the intestine in the abdominal portion of the body. The zooids are frequently arranged in star-shaped clusters, their anal orifices being all directed towards a common vent which occupies the centre. This shortening is still more remarkable, however, in the family of *Botryllians*, whose beautiful stellate gelatinous incrustations




Botryllus violaceus.—A. cluster on the surface of a *Fucus*
B. portion of the same enlarged.

are extremely common upon sea-weeds and submerged rocks (Fig. 249). The anatomy of these animals is very similar to that of the *Amaroucium* already described; with this exception, that the body exhibits no distinction of cavities, all the organs being brought together in one, which must be considered as thoracic. In this respect, there is an evident approximation towards the solitary species.

333. This approximation is still closer, however in the "social" Ascidians, or *Clavellinidæ*; in which the general plan of structure is nearly the same, but the zooids are simply connected by their

stolons, instead of being included in a common investment (Fig. 250); so that their relation to each other is very nearly the same as that of the zooids of *Laguncula* (§ 322), the chief difference

FIG. 250.



A, Group of *Perophora* (enlarged), growing from a common stalk:—B, single *Perophora*; a, test; b, inner sac; c, branchial sac, attached to the inner sac along the line c' c'; e, e, finger-like processes projecting inwards; f, cavity between test and internal coat; f', anal orifice or funnel; g, oral orifice; g', oral tentacula; h, downward stream of food; h', œsophagus; i, stomach; k, vent; l, ovary (?); n, vessels connecting the circulation in the body with that in the stalk.

being that a regular circulation takes place through the stolon in the one case, such as has no existence in the other. A better opportunity of studying the living actions of the Ascidians can scarcely be found, than that which is afforded by the genus *Perophora*, first discovered by Mr. Lister, which occurs not unfrequently on the south coast of England and in the Irish Sea, living attached to sea-weeds, and looking like an assemblage of minute globules of jelly, dotted with orange and brown, and linked by a silvery winding thread. The isolation of the body of each zooid from that of its fellows, and the extreme transparence of its tunics, not only enable the movements of fluid within the body to be distinctly discerned, but also allow the action of the cilia that border the slits of the respiratory sac to be clearly made out. This sac is perforated with four rows of narrow oval openings, through which a portion of the water that enters its branchial orifice (g) escapes into the space between the sac and the mantle, and is thus discharged immediately by the funnel (f'). Whatever little particles, animate or inanimate, the current of water brings, flow into the sac, unless stopped by the tentacula (g') at its entrance, which do not appear fastidious. The particles which are admitted usually lodge somewhere on the sides of the sac, and then travel horizontally until they arrive at that part of it down which the current proceeds to the entrance of the sto-

mach (*i*), which is situated at the bottom of the sac. Minute animals are often swallowed alive, and have been observed darting about in the cavity for some days, without any apparent injury either to themselves or to the creature which encloses them. In general, however, particles which are unsuited for reception into the stomach, are ejected by the sudden contraction of the mantle (or muscular tunic), the vent being at the same time closed, so that they are forced out by a powerful current through the branchial orifice.

334. The circulation of blood, throughout the entire class, is remarkable for the *alternation* which it presents, from time to time, in the direction of its flow; and this curious phenomenon may be particularly well studied in the *Perophora*. The creeping-stalk (Fig. 250) that connects the individuals of any group, contains two distinct canals, which send off branches into each peduncle. One of these branches terminates in the heart, which is nothing more than a contractile dilatation of the principal trunk; this trunk subdivides into vessels (or rather *sinuses*, which are mere channels not having proper walls of their own), of which some ramify over the respiratory sac, branching off at each of the passages between the oval slits, whilst others are first distributed to the stomach and intestines, and to the soft surface of the mantle. All these reunite, and form a trunk, which passes to the peduncle, and constitutes the returning branch. Now whilst at some periods, the heart may be seen vigorously contracting from behind forwards, so as to propel the blood along the course just described, the observer, if he continue to watch, will see its pulsations becoming fainter for a few beats, and the flow in the vessels becoming slower; and then, after a slight pause, the whole current in all its windings is reversed. The heart gives the opposite impulse, receiving the blood from the body, and sending it back into the peduncle through the tube that previously conveyed it thence; while the tube that had previously served to carry the returning stream, now brings the blood from the stem, and distributes it to the branchial sac, mantle, and visceral apparatus, whence it finds its way back to the heart. After the circulation has continued for a certain time in this new direction, the intermission is repeated; and then a reversal takes place to the first course. The average time during which the circulation persists in each direction, seems to be about the same for the one as for the other; but the period between the changes varies as much as from thirty seconds to two minutes. Although the circulation in the different bodies is brought into connection by the common stem, yet that of each is independent of the rest, continuing when the current through its own footstalk is interrupted by a ligature; and the stream which returns from the branchial sac and the viscera is then poured into the posterior part of the heart, instead of entering the peduncle. This is the course which it takes in

the "solitary" Ascidians; and also in those composite forms, whose connecting stolons do not contain bloodvessels.

335. The development of the Ascidians presents some phenomena of much interest to the Microscopist; the early stages of which are observable, whilst the ova are still within the cloaca of the parent. After the ordinary repeated segmentation of the yolk, whereby a "mulberry mass," is produced, a sort of ring is seen, encircling its central portion; but this soon shows itself as a tapering tail-like prolongation from one side of the yolk, which gradually becomes more and more detached from it, save at the part from which it springs. Either whilst the egg is still within the cloaca, or soon after it has escaped from the vent, its envelope bursts, and the larva escapes; and in this condition it presents very much the appearance of a tadpole, the tail being straightened out, and propelling the body freely through the water by its lateral strokes. The centre of the body is occupied by a mass of liquid yolk; and this is continued into the interior of three prolongations which extend themselves from the opposite extremity, each terminating in a sort of sucker. After swimming about for some hours with an active wriggling movement, the larva attaches itself to some solid body by means of one of these suckers; if disturbed from its position, it at first swims about as before; but it soon completely loses its activity, and becomes permanently attached; and important changes manifest themselves in its interior. The prolongations of the central yolk-substance into the anterior processes and tail are gradually drawn back, so that the whole of it is concentrated into one mass; and the tail, now consisting only of the gelatinous envelope, is either detached entire from the body by the contraction of the connecting portion, or withers and is thrown off gradually in shreds. The shaping of the internal organs out of the yolk-mass, takes place very rapidly; so that by the end of the second day of the sedentary state, the outlines of the branchial sac and of the stomach and intestine may be traced; no external orifices, however, being as yet visible. The pulsation of the heart is first seen on the third day, and the formation of the branchial and anal orifices takes place on the fourth; after which the ciliary currents are immediately established through the branchial sac and alimentary canal. The embryonic development of other Ascidians, solitary as well as composite, takes place on a plan essentially the same as the foregoing, a free tadpole-like larva being always produced in the first instance; and in the curious *Appendicularia*, which occasionally presents itself on our own coasts, this larval form is retained through life.¹

¹ For more special information respecting the Compound Ascidians, see especially the admirable Monograph of Prof. Milne Edwards on that group. Mr. Lister's Memoir "On the Structure and Functions of Tubular and Cellular Polypi, and of Ascidiæ," in the "Philos. Transact." 1834, and Mr. Huxley's Memoir "On Doliolum and Appendicularia," in "Philos. Transact." 1851; also the Art. *Tunicata* in the "Cyclopaedia of Anatomy and Physiology."

CHAPTER XIV.

MOLLUSCOUS ANIMALS GENERALLY.

THE various forms of "Shell-fish," with their "naked" or shell-less allies, furnish a great abundance of objects of interest to the Microscopist; of which, however, the greater part may be grouped under three heads;—namely (1), the structure of the *Shell*, which is most interesting in the *Conchifera* or "Bivalves;" (2) the structure of the *Tongue* of the *Gasteropoda*, most of which have "Univalve" shells, others, however, being "naked;" and (3) the *Developmental History* of the embryo, for the study of which certain of the *Gasteropods* present the greatest facilities. These three subjects, therefore, will be first treated of systematically; and a few miscellaneous facts of interest will be subjoined.

336. *Shells of Mollusca*.—These investments were formerly regarded as mere inorganic exudations, composed of calcareous particles cemented together by animal glue; Microscopic examination, however, has shown that they possess a distinctly organic structure; and this structure presents certain very remarkable variations, in some of the natural groups of which the Molluscous series is composed. We shall first describe that which may be regarded as the characteristic structure of the ordinary Bivalves; taking as a type the group of *Margaritaceæ*, which includes the "Pearl-oyster" and its allies, the common *Pinna* ranking amongst the latter. In all these shells, we readily distinguish the existence of two distinct layers; an external, of a brownish-yellow color; and an internal, which has a pearly or "nacreous" aspect, and is commonly of a lighter hue. The structure of the *outer* layer may be conveniently studied in the shell of *Pinna*, in which it commonly projects beyond the inner, and there often forms laminæ sufficiently thin and transparent to exhibit the general nature of its organization without any artificial reduction. If a small portion of such a lamina be examined with a low magnifying power, even without any preparation by transmitted light, each of its surfaces will present very much the appearance of a honeycomb; whilst its broken edge exhibits an aspect which is evidently fibrous to the eye, but which, when examined under the microscope with reflected light, resembles that of an assemblage of basaltic columns (Fig. 334, P).

The shell is thus seen to be composed of a vast number of prisms, having a tolerably uniform size, and usually presenting an approach to the hexagonal shape. These are arranged per-

FIG. 251.




FIG. 252.





Fig. 251. Section of Shell of *Pinna*, transversely to the direction of its prisms. Fig. 252. Membranous basis of the same.

pendicularly (or nearly so) to the surface of the lamina of the shell; so that its thickness is formed by their length, and its two surfaces by their extremities. A more satisfactory view of these prisms is obtained by grinding down a lamina until it possesses a high degree of transparency; and it is then seen (Fig. 251) that the prisms themselves appear to be composed of a very homogeneous substance, but they are separated by definite and strongly marked lines of division. When such a lamina is submitted to the action of dilute acid, so as to dissolve away the carbonate of lime, a tolerably firm and consistent membrane is left, which exhibits the prismatic structure just as perfectly as did the original shell (Fig. 252); the hexagonal divisions being apparently the walls of cells resembling those of the pith or bark of a plant, in which the cells are frequently hexagonal prisms. In very thin natural laminae, the nuclei of the cells can often be plainly distinguished. By making a section of the shell perpendicularly to its surface, we obtain a view of the prisms cut in the direction of their length (Fig. 253); and they are frequently seen to be marked by delicate transverse striæ (Fig. 254), closely resembling those observable on the prisms of the enamel of teeth, to which this kind of shell-structure may be considered as bearing a very close resemblance, except as regards the mineralizing ingredient. If a similar section be decalcified by dilute acid, the membranous residuum will exhibit the walls of the prismatic cells viewed longitudinally; and these will be seen to be more or less regularly marked by the transverse striæ just alluded to. It sometimes


FIG. 253.



Section of the Shell of *Pinna*, in the direction of its prisms.

happens in recent, but still more commonly in fossil shells, that the decay of the animal membrane leaves the contained prisms without any connecting medium; as they are then quite isolated, they can be readily detached one from another; and each one may be observed to be marked by the like striations, which, when a sufficiently high magnifying power is used, are seen to

FIG. 251.



Oblique Section of Prismatic Shell-substance.

be minute grooves, apparently resulting from a thickening of the intermediate wall in those situations. This thickening seems best accounted for, by supposing (as first suggested by Prof. Owen) that each long prismatic cell is made up by the coalescence of a pile of flat epidermic cells, the transverse striation marking their lines of junction; and this view corresponds well with the fact, that the shell-membrane not unfrequently shows a tendency to split into thin laminae along the lines of striation; whilst we occasionally meet with an excessively thin natural lamina, composed of flat pavement-like cells, lying between the thicker prismatic layers, with one of which it would have probably coalesced, but for some accidental cause which preserved its distinctness. That the prism is not formed in its entire length at once, but that it is progressively lengthened and consolidated at its lower extremity, would appear also from the fact, that where the shell presents a deep color (as in *Pinna nigrina*), this color is usually disposed in distinct strata, the outer portion of each layer being the part most deeply tinged, whilst the inner extremities of the prisms are almost colorless. This prismatic arrangement of the carbonate of lime in the shells of *Pinna* and its allies, has been long familiar to conchologists; but it has been usually regarded as the result of crystallization.

337. It is only in the shells of a few families of Bivalves, that the combination of organic with mineral components is seen in this very distinct form; and these families are for the most part nearly allied to *Pinna*. In all the genera of the *Margaritaceæ*, we find the external layer of the shell formed upon this plan, and of considerable thickness; the internal layer being nacreous. In the *Unionidæ* (fresh-water Mussels), on the contrary, nearly the whole thickness of the shell is made up of the internal or nacreous layer; but a uniform stratum of prismatic cellular substance is always found between the nacre and the periostracum.¹ In

¹ The periostracum is the yellowish-brown membrane covering the surface of many shells, which is often (but erroneously) termed the epidermis.

the *Ostraceæ* (or Oyster tribe), the greater part of the shell is composed of a sub-nacreous substance, the successively formed laminae of which have very little adhesion to each other; but every one of these laminae is bordered at its free edge by a layer of the prismatic cellular substance, distinguished by its brownish-yellow color. In these and some other cases, a distinct organic structure is left after the decalcification of the prismatic layer by dilute acid; and this is most tenacious and substantial, where (as in the *Margaritaceæ*) there is no proper periostracum; as if the horny matter which would have otherwise gone to form this investment, had been diffused as an intercellular substance between the proper cell-walls. Generally speaking, a cellular layer may be detected upon the external surface of bivalve shells, where this has been protected by a periostracum, or has been prevented in any other manner from undergoing abrasion; thus it is found pretty generally in *Chama*, *Trigonia*, and *Solen*, and occasionally in *Anomia* and *Pecten*.

338. In many other instances, however, although a cellular arrangement is apparent in sections of the shell, no corresponding structure can be distinctly seen in the delicate membrane left after decalcification. In all such cases, the animal basis bears but a very small proportion to the calcareous deposit, and the shell is usually extremely hard. But there are numerous other cases, in which no distinct traces of cellular structure can be detected in the fully-formed shell; and in which it would seem as if the consolidation of the animal basis by calcareous deposit had taken place whilst the former was as yet in the condition of a uniform gelatinous sarcode, before the commencement of any differentiation into cell-wall and cell-contents. A very curious appearance is presented by a section of the large hinge-tooth of *Mya arenaria* (Fig. 255), in which the carbonate of lime seems to be deposited in nodules, not distinctly separated from each other by intervening membrane, and possessing a crystalline structure resembling that of the mineral termed Wavellite. Approaches to this curious arrangement are seen in many other shells.


339. The *internal* layer of Bivalve shells rarely presents any distinct structure, when examined in a thin section; and the residuum left after decalcification is usually a structureless "basement-membrane." This form of shell-substance may consequently be distinguished as *membranous*. In the *Margaritaceæ* and many other families, this internal layer has a nacreous or iride-

FIG. 255.

Section of hinge-tooth of *Mya arenaria*.

scent lustre, which depends (as Sir D. Brewster has shown¹) upon the striation of its surface with a series of grooved lines, which usually run nearly parallel to each other (Fig. 256).

FIG. 256.



Section of Nacreous lining of Shell of *Arctica margaritacea*
(Pearl-oyster).

As these lines are not obliterated by any amount of polishing, it is obvious that their presence depends upon something peculiar in the texture of this substance, and not upon any mere superficial arrangement. When a piece of nacre is carefully examined, it becomes evident that the lines are produced by the cropping out of laminae of shell, situated more or less obliquely to the plane of the surface. The greater the *dip* of these laminae, the closer will their edges

be; whilst the less the angle which they make with the surface, the wider will be the interval between the lines. When the section passes for any distance in the plane of a lamina, no lines will present themselves on that space. And thus the appearance of a section of nacre is such, as to have been aptly compared by Sir J. Herschel² to the surface of a smoothed deal board, in which the woody layers are cut perpendicularly to their surface in one part, and nearly in their plane in another. Sir D. Brewster (loc. cit.) appears to suppose that nacre consists of a multitude of layers of carbonate of lime alternating with animal membrane; and that the presence of the grooved lines on the most highly polished surface, is due to the wearing away of the edges of the animal laminae, whilst those of the hard calcareous laminae stand out. If each line upon the nacreous surface, however, indicates a distinct layer of shell-substance, a very thin section of mother-of-pearl ought to contain many thousand laminae, in accordance with the number of lines upon its surface; these being frequently no more than 1-7500th of an inch apart. But when the nacre is treated with dilute acid, so as to dissolve its calcareous portion, no such repetition of membranous layers is to be found; on the contrary, if the piece of nacre be the product of one act of shell-formation, there is but a single layer of membrane. The membrane is usually found to present a more or less *folded* or *plaited* arrangement; but this has generally been obviously disturbed by the disengagement of carbonic acid in the act of decalcification, which tends to unfold the plaits. There is one shell, however,—


¹ "Philos. Transact." 1814.

² "Edinb. Philos. Journal," vol. ii.

the well-known *Haliotis splendens*,—which affords us the opportunity of examining the plaits without any disturbance of their arrangement, and thus presents a clear demonstration of the real structure of nacre. This shell is for the most part made up of a series of plates of animal matter, resembling tortoise-shell in its aspect, alternating with thin layers of nacre; and if a piece of it be submitted to the action of dilute acid, the calcareous portion of the nacreous layers being dissolved away, the plates of animal matter fall apart, each one carrying with it the membranous residuum of the layer of nacre that was applied to its inner surface. It will usually be found that the nacre-membrane covering some of these horny plates will remain in an undisturbed condition; and *their surfaces then exhibit their iridescent lustre, although all the calcareous matter has been removed from their structure.* On looking at the surface with reflected light under a magnifying power of 75 diameters, it is seen to present a series of folds or plaits more or less regular; and the iridescent hues which these exhibit, are often of the most gorgeous description. If the membrane be extended, however, with a pair of needles, these plaits are unfolded, and it covers a much larger surface than before; but its iridescence is then completely destroyed. This experiment, then, demonstrates that the peculiar lineation of the surface of nacre (on which its iridescence undoubtedly depends, as originally shown by Sir D. Brewster) is due, not to the outcropping of alternate layers of membranous and calcareous matter, but to the disposition of a single membranous layer in folds or plaits, which lie more or less obliquely to the general surface. There are several bivalve shells which present what may be termed a *sub-nacreous* structure, their polished surfaces being covered with lines indicative of folds in the basement-membrane; but these folds are destitute of that regularity of arrangement, which is necessary to produce the iridescent lustre. This is the case, for example, with most of the *Pectinidæ* (or Scallop tribe), also with some of the *Mytilaceæ* (or Mussel tribe), and with the common *Oyster*. Where there is no indication of a regular corrugation of the shell-membrane, there is not the least approach to the nacreous aspect; and this is the case with the internal layer of by far the greater number of shells, the presence of true nacre being exceptional, save in a small number of families. It is of the inner layer that those rounded concretions are usually formed, which are often found in the interior of shells, and which, when composed of nacreous substance resembling that of the lining of the shell of *Avicula*, are known as *pearls*. Such concretions are found in many other shells; but they are usually less remarkable for their pearly lustre; and when formed at the edge of the valves, they may be partly or even entirely made up of the prismatic substance of the external layer, and may be consequently altogether destitute of the pearly aspect. The “membranous” shell-substance, some form of which constitutes the internal layer

of most bivalve shells, is occasionally traversed by *tubes*, which seem to commence from the inner surface of the shell, and to pass towards the exterior. These tubes vary in size from about the 1-20,000th of an inch, or even less, to about the 1-2000th; but their general diameter, in the shells in which they most


FIG. 257.

Tubular Shell-structure of *Anomia*.

abound, is about 1-4000th of an inch. The direction and distribution of these tubes are extremely various in different genera: in *Anomia ephippium* they are scantily distributed in the internal nacreous lamina, whilst in the yellow outer layer they are very abundant, forming an irregular network (Fig. 257), which spreads out in a plane parallel to the surface.

340. The ordinary account of the mode of growth of the shells of Bivalve Mollusca,—that they are progressively enlarged by the deposition of new laminæ, each of which is in contact with the internal surface of the preceding, and extends beyond it,—does not express the whole truth; for it takes no account of the fact that most shells are composed of two layers of very different texture, and does not specify whether *both* these layers are thus formed by the entire surface of the mantle whenever the shell has to be extended, or whether only *one* is produced. An examination of Fig. 258 will clearly show the mode in which the

FIG. 258.

Vertical section of the lip of one of the valves of the shell of *Unio*.—*a, b, c*, successive formations of the outer layer, *a' b' c'*, the same of the inner layer.

operation is effected. This figure represents a section of one of the valves of *Unio occidentis*, taken perpendicularly to its surface, and passing from the margin or lip (at the left hand of the figure) towards the hinge (which would be at some distance beyond the right). This section brings into view the two substances of which the shell is composed; traversing the outer or prismatic layer in the direction of the length of its cells, and passing through the nacreous lining, in such a manner as to bring into view its numerous laminæ, separated by the lines *a a', b b', c c'*,

&c. These lines evidently indicate the successive formations of this layer; and it may be easily shown, by tracing them towards the hinge on the one side, and towards the margin on the other, that at every enlargement of the shell, its whole interior is lined by a new nacreous lamina, in immediate contact with that which preceded it. The number of such laminæ, therefore, in the oldest part of the shell, indicates the number of enlargements which it has undergone. The outer or prismatic layer of the growing shell, on the other hand, is only formed where the new structure projects beyond the margin of the old; and thus we do not find one layer of it overlapping another, except at the lines of junction of two distinct formations. When the shell has attained its full dimensions, however, new laminæ of both layers still continue to be added; and thus the lip becomes thickened by successive formations of prismatic structure, each being applied to the inner surface of the preceding, instead of to its free margin. A like arrangement may be well seen in the *Oyster*; with this difference, that the successive layers have but a comparatively slight adhesion to each other.

341. The shells of *Terebratulæ*, and of several other genera of *Brachiopoda*, are distinguished by peculiarities of structure, which serve to distinguish them from all others. When thin sections of them are microscopically examined, they exhibit the appearance of long flattened prisms (Fig. 259, *b*), which are arranged with such obliquity, that their rounded extremities crop out upon the inner surface of the shell in an imbricated (tile-like) manner (*a*). All true *Terebratulidæ*, both recent and fossil, exhibit another very remarkable peculiarity; namely, the presence of a large number of perforations in the shell, generally passing nearly perpendicularly from one surface to the other (as is shown in vertical sections, Fig. 261), and terminating internally by open orifices




Fig. 259. Internal surface (*a*), and oblique section (*b*), of Shell of *Terebratula* (*Waldheimia*) *australis*.
Fig. 260. External surface of same.

(Fig. 259), whilst externally they are covered by the periostracum (Fig. 260). Their diameter is greatest towards their external surface, where they sometimes expand suddenly, so as to become

trumpet-shaped; and it is usually narrowed rather suddenly,

FIG. 261.



Vertical sections of Shell of *Terebratula* (*Waldheimia*) *australis*:—showing at A the canals opening by large trumpet-shaped orifices on the outer surface, and contracting at *d, d*, into narrow tubes; and presenting at *b* a bifurcation of the canals.

when, as sometimes happens, a new internal layer is formed as a lining to the preceding (Fig. 261, A, *d d*). Hence the diameter of these canals, as shown in different transverse sections of one and the same shell, will vary according to the part of its thickness which the section happens to traverse. The different species of *Terebratulidæ*, however, present very striking diversities in the size and closeness of the canals, as shown by sections taken in corresponding parts of their shells; three examples of this kind are given for the sake of comparison in Figs. 262–264. These canals

are occupied, in the living state, by tubular prolongations of the mantle, the interior of which is filled with a fluid containing minute cells and granules, which, from its corresponding in appearance with the fluid contained in the great sinuses of the mantle, may be considered to be the animal's blood. Hence these cæcal tubes may be inferred to

FIG. 262.



FIG. 263.




FIG. 264.



Fig. 262. Horizontal section of Shell of *Terebratula bullata* (fossil, oolite).
 Fig. 263. " " " of *Megerlia lima* (fossil, chalk).
 Fig. 264. " " " of *Spiriferina rostrata* (triassic).

possess a respiratory function; and seem to be analogous to tubes of a very similar nature, which extend into the "test" of many Tunicata from their sinus system (§ 334). In the family *Rhynchonellidæ*, which is represented by only two recent species (the *Rh. psittacea* and *Rh. nigricans*, both of which formerly ranked as *Terebratulæ*), but which contains a very large proportion of fossil Brachiopods, these canals are entirely absent; so that the uniformity of their presence in the *Terebratulidæ*, and of their absence in the *Rhynchonellidæ*, supplies a character of great value in the

discrimination of the fossil shells belonging to these two groups respectively. Great caution is necessary, however, in applying this test; mere surface-markings cannot be relied on; and no statement on this point is worthy of reliance, which is not based on a microscopic examination of thin sections of the shell. In the families *Spiriferidæ* and *Strophonemidæ*, on the other hand, some species possess the perforations, whilst others are destitute of them; so that their presence or absence *there* only serves to mark out subordinate groups. This, however, is what holds good in regard to characters of almost every description, in other departments of Natural History, as well as in this; a character which is of fundamental importance from its close relation to the general plan of organization in one group, being, from its want of constancy, of far less account in another.¹

342. There is not by any means the same amount of diversity in the structure of the shell in the class of *Gasteropoda*, as that which exists among the several tribes of *Conchifera*; a certain typical plan of construction being common to by far the greater number of them. The small proportion of animal matter contained in most of these shells, is a very marked feature in their character; and it serves to render other features indistinct, since the residuum left after the removal of the calcareous matter is usually so imperfect, as to give no clue whatever to the explanation of the appearances shown by sections. Nevertheless, the structure of these shells is by no means homogeneous, but always exhibits indications, more or less clear, of an original organic arrangement. The "porcellanous" shells are composed of three layers, all presenting the same kind of structure, but each differing from the others in the mode in which this is disposed. For each layer is made up of an assemblage of thin laminæ placed side by side, which separate one from another, apparently in the planes of rhomboidal cleavage, when the shell is fractured; and, as was first pointed out by Mr. Bowerbank, each of these laminæ consists of a series of elongated spicules (considered by him as prismatic cells filled with carbonate of lime) lying side by side in close apposition; and these series are disposed alternately in contrary directions, so as to intersect each other nearly at right angles, though still lying in parallel planes. The direction of the planes is different, however, in the three layers of the shell, bearing the same relation to each other as have those three sides of a cube which meet each other at the same angle; and by this arrangement, which is better seen in the fractured edge of *Cypræa* or any similar shell, than in thin sections, the strength of the shell is greatly augmented. A similar arrangement, obviously designed with the same purpose, has been shown by Mr. Tomes to exist in the enamel of the

¹ For a particular account of the Author's researches on this group, see his memoir on the subject, forming part of the Introduction of Mr. Davidson's "Monograph of the British Fossil Brachiopoda," published by the Palæontographical Society.

teeth of Rodentia. The principal departures from this plan of structure are seen in *Patella*, *Chiton*, *Haliotis*, *Turbo*, and its allies, and in the "naked" Gasteropods, many of which last, both terrestrial and marine, have some rudiment of a shell. Thus in the common Slug, *Limax rufus*, a thin oval plate, of calcareous texture, is found imbedded in the shield-like fold of the mantle covering the fore-part of its back; and if this be examined in an early stage of its growth, it is found to consist of an aggregation of cell-like bodies, generally somewhat hexagonal in form, and consolidated by a deposit of calcareous matter, which is sometimes so arranged as to be quite transparent, whilst in other instances it presents an appearance closely resembling that delineated in Fig. 255. In the epidermis of the mantle of some species of *Doris*, on the other hand, we find long calcareous spicules, generally lying in parallel directions, but not in contact with each other, giving firmness to the whole of its dorsal portion; and these are sometimes covered with small tubercles, like the spicules of *Gorgonia* (§ 309). They may be separated from the soft tissue in which they are imbedded, by means of caustic potash; and when treated with dilute acid, whereby the calcareous matter is dissolved away, an organic basis is left, retaining in some degree the form of the original spicule. This basis cannot be said to be a true cell; but it seems to be rather a cell in the earliest stage of its formation, being an isolated particle of "sarcode" without wall or cavity; and the close correspondence between appearances presented by thin sections of various "univalve" shells, and the forms of the spicules of *Doris*, seems to justify the conclusion, that even the most compact shells of this group are constructed out of the like elements, in a state of closer aggregation and more definite arrangement, with the occasional occurrence of a layer of more spheroidal bodies of the same kind, like those forming the rudimentary shell of *Limax*.

343. The animals composing the class of *Cephalopoda* (Cuttlefish and *Nautilus* tribe), are for the most part unpossessed of shells; and the structure of the few that we meet with in the genera *Nautilus*, *Argonauta* (Paper-nautilus) and *Spirula*, does not present any peculiarities that need here detain us. The rudimentary shell or *sepiostaire* of the common Cuttlefish, however, which is frequently spoken of as the "cuttlefish bone," exhibits a very beautiful and remarkable structure, such as causes sections of it to be very interesting microscopic objects. The outer shelly portion of this body consists of horny layers, alternating with calcified layers, in which last may be seen an hexagonal arrangement somewhat corresponding with that in Fig. 255. The soft friable substance that occupies the hollow of this boat-shaped shell, is formed of a number of delicate plates, running across it from one side to the other in parallel directions, but separated by intervals several times wider than

the thickness of the plates; and these intervals are in great part filled up by what appear to be fibres or slender pillars, passing from one plate or floor to another. A more careful examination shows, however, that instead of a large number of detached pillars, there exists a comparatively small number of very thin sinuous laminæ, which pass from one surface to the other, winding and doubling upon themselves, so that each lamina occupies a considerable space. Their precise arrangement is best seen by examining the parallel plates, after the sinuous laminæ have been detached from them; the lines of junction being distinctly indicated upon these. By this arrangement, each layer is most effectually supported by those with which it is connected above and below; and the sinuosity of the thin intervening laminæ, answering exactly the same purpose as the "corrugation" given to iron plates for the sake of diminishing their flexibility, adds greatly to the strength of this curious texture; which is at the same time lightened by the large amount of space between the parallel plates that intervenes between the sinuosities of the laminæ. The best method of examining this structure, is to make sections of it with a sharp knife in various directions, taking care that the sections are no thicker than is requisite for holding together; and these may be mounted on a black ground as opaque objects, or in Canada balsam as transparent objects.

344. The structure of Shells generally is best examined by making sections in different planes, as nearly parallel as may be possible to the surfaces of the shell; and other sections at right angles to these; the former may be designated as *horizontal*, the latter as *vertical*. Nothing need here be added to the full directions for making such sections, which have already been given (§§ 108-110). Much valuable information may also be derived, however, from the examination of the surfaces presented by fracture. The membranous residua left after the decalcification of the shell by dilute acid, may be mounted in weak spirit or in Goadby's solution.¹

345. *Tongue of Gasteropod Mollusks.*—The organ which is commonly known under this designation, is one of a very singular nature; and we should be altogether wrong in conceiving of it as having any likeness to that on which our ordinary ideas of such an organ are founded. For instead of being a projecting body, lying in the cavity of the mouth, it is a tube that passes backwards and downwards beneath the mouth; its hinder end being closed, whilst in front it opens obliquely upon the floor of the mouth, being (as it were) slit up and spread out, so as to form a nearly flat surface. On the interior of the tube, as well as on the flat expansion of it, we find numerous transverse

¹ For fuller details on the minute structure of the shells of the Mollusca, see the Author's memoirs on that subject in the "Reports of the British Association" for 1844 and 1847; also Mr. Bowerbank's memoir on the same subject in "Transact. of Microscopical Society." Ser. 1, vol. i; and Mr. Quekett's "Lectures on Histology," vol. ii, Chaps. xvi-xxii.

rows of minute teeth, which are set upon flattened plates; each principal tooth sometimes having a basal plate of its own, whilst in other instances one plate carries several teeth. Of the former arrangement we have an example in the tongue of many

FIG. 265.



FIG. 266.



Fig. 265. Portion of the left half of the Palate of the *Helix hortensis*; the rows of teeth near the edge separated from each other to show their form.

Fig. 266. Palate of *Zonites cellarius*.

terrestrial Gasteropods, such as the Snail (*Helix*) and Slug (*Limax*), in which the number of plates in each row is very considerable (Figs. 265, 266), amounting to 180 in the large garden Slug (*Limax maximus*); whilst the latter prevails in many marine Gasteropods, such as the common Whelk (*Buccinum undatum*), the tongue of which has only three plates in each row, one bearing the small central teeth, and the two others the large lateral teeth (Fig. 269). The length of the tongue, and the number of rows of teeth, vary greatly in different species. Generally speaking, the tongue of the terrestrial Gasteropods is short, and is contained entirely within the nearly globular head; but the rows of teeth being closely set together, they are usually very numerous, there being frequently more than 100, and in some species as many as 160 or 170; so that the total number of teeth may mount up, as in *Helix pomatia*, to 21,000, and in *Limax maximus*, to 26,800. The transverse rows are usually more or less curved, as shown in Fig. 266, whilst the longitudinal rows are quite straight; and the curvature takes its departure on each side from a central longitudinal row, the teeth of which are symmetrical, whilst those of the lateral portions of each transverse row present a modification of that symmetry, the prominences on the *inner* side of each tooth being suppressed, whilst those on the *outer* side are increased; this modification being observed to augment in degree, as we pass from the central line towards the edges. The tongue of the marine Gasteropods is generally longer, and its teeth larger; and in many instances it extends far beyond the head, which may, indeed, contain but a small part of it. Thus in the common Limpet (*Patella*), we find the principal part of the tongue to lie folded up, but perfectly free, in the abdominal cavity, between the intestines and the

muscular foot; and in some species its length is twice or even three times as great as that of the entire animal. In a large proportion of cases, these tongues exhibit a very marked separation between the central and the lateral portions (Figs. 267, 269); the teeth of the central band being frequently small and smooth at their edges, whilst those of the lateral are large and serrated. The tongue of *Trochus zizyphinus*, represented in Fig. 267, is one of the most beautiful examples of this form; not only the large teeth of the lateral bands, but the delicate leaf-like teeth of the central portion, having their edges minutely serrated. A yet more complex type, however, is found in the tongue of *Haliotis*; in which there is a central band of teeth having nearly straight edges instead of points; then, on each side, a lateral band consisting of large teeth shaped like those of the shark; and beyond this, again, another lateral band on either side, composed of several rows of smaller teeth. Very curious differences also present themselves among the different species of the same genus. Thus in *Doris pilosa*, the central band is almost entirely wanting, and each lateral band is formed of a single row of very large hooked teeth, set obliquely, like those of the lateral bands in Fig. 267; whilst in *Doris tuberculata*, the central band is the

FIG. 267.

Palate of *Trochus zizyphinus*.

FIG. 268.

Palate of *Doris tuberculata*.

part most developed, and contains a number of rows of conical teeth, standing almost perpendicularly, like those of a harrow (Fig. 268).

346. Many other varieties might be described, did space permit; but we must be content with adding, that the form and arrangement of the teeth afford characters of great value in classification, as was first pointed out by Prof. Lovèn (of Stockholm) in 1847, and has been since very strongly urged by Dr. J. E. Gray, who considers that the structure of the tongue is one

of the best guides to the natural affinities of the species, genera, and families of this group, since any important alteration in the form or position of the teeth must be accompanied by some corresponding peculiarity in the habits and manners of the animal.¹ Hence a systematic examination and delineation of the structure and arrangement of these organs, by the aid of the Microscope and Camera Lucida, would be of the greatest service to this department of Natural History. The short thick tube of the *Limax* and other terrestrial Gasteropods, appears adapted for the trituration of the food previously to its passing into the cesophagus; for in these animals we find the roof of the mouth furnished with a large strong horny plate, against which the flat end of the tongue can work. On the other hand, the flattened portion of the tongue of *Buccinum* and its allies is used by these animals as a file, with which they bore holes through the shells of the mollusks that serve as their prey; this they are enabled to effect, by everting that part of the proboscis-shaped mouth whose floor is formed by the flattened part of the tongue, which is thus brought to the exterior, and by giving a kind of sawing motion to the organ, by means of the alternate action of two pairs of muscles,—a protractor, and a retractor,—which put forth and draw back a pair of cartilages whereon the tongue is supported, and also elevate and depress its teeth.² Of the use of the long blind tubular part of the tongue in these Gasteropods, however, scarcely any probable guess can be made; unless it be a sort of "cavity of reserve," from which a new toothed surface may be continually supplied, as the old one is worn away, somewhat as the front teeth of the Rodents are constantly being regenerated from the surface of the pulps which occupy their hollow conical bases, as fast as they are rubbed down at their edges.

347. The preparation of these tongues for the Microscope, can, of course, be only accomplished by carefully dissecting them from their attachments within the head; and it will be also necessary to remove the membrane that forms the sheath of the tube, when this is thick enough to interfere with its transparency. The tube itself should be slit up with a pair of fine scissors, through its entire length; and should be so opened out, that its expanded surface may be a continuation of that which forms the floor of the mouth. The mode of mounting it will depend upon the manner in which it is to be viewed. For the ordinary purposes of Microscopic examination, no method is so good as mounting in fluid; either weak spirit or Goadby's solution answering very well. But many of these tongues, especially those of the marine Gasteropods, become most beauti-

¹ "Annals of Natural History," Ser. 2, vol. x, p. 413.

² For additional details on the organization of the tongue and teeth of the Gasteropod Mollusks, see Mr. W. Thomson, in "Cyclop. of Anat. and Physiol." vol. iv, pp. 1142, 1143; and in "Ann. of Nat. Hist." Ser. 2, vol. vii, p. 86.

ful objects for the Polariscopes, when they are mounted in Canada balsam; the form and arrangement of the teeth being very strongly brought out by it (Fig. 269), and a gorgeous play of colors being exhibited when a selenite plate is placed behind the object, and the analyzing prism is made to rotate.

348. The stomachs, also, of many Gasteropod Mollusks are furnished with teeth, which are implanted on their walls for the further reduction of the food; such teeth, very numerous but of small size, and bearing a strong resemblance to those of its tongue, are found in the stomach of the common *Slug*. In several marine Gasteropods, however, especially *Bulla*, *Scyllæa*, and *Aplysia*, the gastric teeth are individually much larger, though less numerous, and constitute a very efficient reducing apparatus, especially when combined, as they frequently are, with a horny or calcareous deposit in the walls of the stomach, which converts it into a "gizzard" for the trituration of the substances that have been divided by the teeth.

349. *Development of Gasteropod Mollusks.*—The history of embryonic development may be studied with peculiar facility in certain members of this class, and presents numerous phenomena of great interest. The eggs (save among the terrestrial species) are usually deposited in aggregate masses, each enclosed in a common protective envelope. The nature of this envelope, however, varies greatly: thus in the common *Lymnæus stagnalis*, or "water-snail," of our ponds and ditches, it is nothing else than a mass of soft jelly, about the size of a sixpence, in which from 50 to 60 eggs are imbedded, and which is attached to the leaves or stems of aquatic plants; in the *Buccinum undatum*, or common Whelk, it is a membranous case, connected with a considerable number of similar cases by short stalks, so as to form large globular masses, which may often be picked up on our shores, especially between April and June; in the *Purpura lapillus*, or Rock-whelp, it is a little flask-shaped capsule, having a firm horny wall, which is attached by a sort of foot to the surface of rocks between the tide-marks, great numbers being often thus found standing erect side by side; whilst in the *Nudibranchiate* order generally (consisting of the *Doris*, *Eolis*, and other "sea-slugs") it forms a long tube with a membranous wall, in which immense numbers of eggs (even half a million or more) are packed closely together in the midst of a jelly-like substance, this tube being disposed in coils of various forms,

FIG. 269.



Palate of *Buccinum undatum*, as seen under polarized light.

which are usually attached to sea-weeds or zoophytes. The course of development, in the first and last of these instances, may be readily observed from the very earliest period, down to that of the emersion of the embryo; owing to the extreme transparency of the "nidamentum," and of the egg-membranes themselves. The first change which will be noticed by the ordinary observer, is the "segmentation" of the yolk-mass, which divides itself (after the manner of a cell undergoing duplicative subdivision) into two parts, each of these two into two others, and so on, until a mulberry-like mass of minute yolk segments is evolved. Generally speaking, however, there may be noticed at a very early stage of this process, as performed by Gasteropod Mollusks, an inequality in the size of the segments (Fig. 271, c); one set, derived from the larger of the two divisions (D) into which the yolk-sphere first separates itself, being destined to form the internal organs, whilst the other set of segments, of much inferior dimensions, and formed by the subdivision of the smaller half of the original sphere, furnishes the material for the superficial parts. Soon after the "mulberry mass" has been formed, it commonly begins to exhibit a very curious alternating movement within the egg, two or three turns being made in one direction, and the same number in a reverse direction: this movement, which is due to ciliary action, is often extremely transitory in its duration; but in the *Lymnaeus* it continues almost up to the escape of the embryo, and, when several ova are brought into view at once under a low magnifying power, the spectacle is a very curious one.

350. A separation is usually seen at an early period, between the anterior or cephalic portion, and the posterior or visceral portion, of the embryonic mass; and the development of the former advances with the greater activity. One of the first changes which is seen in it, consists in its extension into a sort of fin-like membrane on either side, the edges of which are fringed with long cilia (Fig. 270), whose movements may be clearly distinguished whilst the embryo is still shut up within

FIG. 270.



Embryoes of Nudibranchiate Gasteropods.

the egg; at a very early period may also be discerned the "auditory vesicles" or rudimentary organs of hearing (§ 353), which scarcely attain any higher development in these creatures during the whole of life; and from the immediate neighborhood of these is put forth a projection, which is afterwards to be evolved

into the "foot" or muscular disk of the animal. While these organs are making their appearance, the shell is being formed

on the surface of the posterior portion, appearing first as a thin covering over its hinder part, and gradually extending itself until it becomes large enough to enclose the embryo completely, when this contracts itself. The ciliated lobes are best seen in the embryos of *Nudibranchs*, in which they are much larger than in *Lymnæus*; and the fact of the universal presence of a shell in the embryos of the former group, is of peculiar interest, as it is destined to be cast off very soon after they enter upon active life. These embryos may be seen to move about as freely as the narrowness of their prison permits, for some time previous to their emersion; and when set free by the rupture of the egg-cases, they swim forth with great activity by the action of their ciliated lobes,—these, like the wheels of *Rotifera*, serving also to bring food to the mouth, which is at that time unprovided with the reducing apparatus subsequently found in it. The same is true of the embryo of *Lymnæus*, save that its swimming movements are less active, in consequence of the inferior development of the ciliated lobes; and the currents produced by these seem to have reference chiefly to the provision of supplies of food, and of aerated water for respiration. The disappearance of the cilia has been observed by Mr. Hogg to be coincident with the development of the teeth to a degree sufficient to enable the young water-snail to crop its vegetable food; and he has further ascertained, that if the growing animal be kept in fresh water alone for some time, with vegetable matter of any kind, the gastric teeth are very imperfectly developed, and the cilia are still retained.¹

351. A very curious modification of the ordinary plan of development, is presented in the *Purpura lapillus*; and it is probable that something of the same kind exists also in *Buccinum*, as also in other Gasteropods of the same extensive order (*Pectinibranchiata*). Each of the capsules already described (§ 349) contains from 500 to 600 egg-like bodies (Fig. 271, A), imbedded in a viscid gelatinous substance; but only from 12 to 30 embryos usually attain complete development; and it is obvious from the large comparative size which these attain (Fig. 272, B), that each of them must include an amount of substance equal to that of a great number of the bodies originally found within the capsule. The explanation of this fact (long since noticed by Dr. J. E. Gray in regard to *Bucci-*

FIG. 271.




Early stages of embryonic development of *Purpura lapillus*:—A, egg-like spherule; B, C, E, F, G, successive stages of segmentation of yolk-spherules; D, H, I, J, K, successive stages of development of early embryos.

¹ "Transactions of Microscopical Society," 2d Ser. vol. ii, p. 93.

num) seems to be as follows:—Of those 500 or 600 egg-like bodies, only a small part are true *ova*, the remainder being merely *yolk-spherules*, which are destined to serve for the nutrition of the embryos. The distinction between them manifests itself at a very early period, even in the first segmentation; for while the yolk-spherules divide into two equal hemispheres (Fig. 271, *β*), the real *ova* divide into a larger and a smaller segment (*δ*); in the cleft between these are seen the minute “directive vesicles,” which appear to be always double or even triple, although, from being seen “end on,” only one may be visible; and near these is generally to be seen a clear space in each segment. The difference is still more strongly marked in the subsequent divisions; for whilst the cleavage of the yolk-spherules goes on irregularly, so as to divide each into from 14 to 20 segments having no definiteness of arrangement (*c, e, f, g*), that of the *ova* takes place in such a manner as to mark out the distinction already alluded to between the cephalic and the visceral portions of the mass (*η*); and the evolution of the former into distinct organs very speedily commences. In the first instance, a narrow transparent border is seen around the whole embryonic mass, which is broader at the cephalic portion (*ι*); next, this border is fringed with a short cilia, and the cephalic extension into two lobes begins to show itself; and then between the lobes a large mouth is formed, opening through a short, wide œsophagus, the interior of which is ciliated, into the visceral cavity, occupied as yet only by the yolk particles originally belonging to the ovum (*κ*). Whilst these developmental changes are taking place in the embryo, the whole aggregate of segments formed by the subdivision of the yolk-spherules coalesces into one mass, as shown at *λ*, Fig. 272; and the embryos are often, in the first instance, so completely buried within this, as only to be discoverable by tearing its portions asunder; but some of them may commonly be found upon its exterior; and those contained in one capsule very commonly exhibit the different stages of development represented in Fig. 271, *η-κ*. After a short time, however, it becomes apparent that the most advanced embryos are beginning to *swallow* the yolk-segments of the conglomerate mass; and capsules will not unfrequently be met with, in which embryos of various sizes, as *a, b, c, d, e* (Fig. 272, *λ*), are projecting from their surface, their difference in size not being accompanied by advance in development, but merely depending upon the amount of this “supplemental” yolk which the individuals have respectively gulped down. For during the time in which they are engaged in appropriating this additional supply of nutriment, although they increase in size, yet they scarcely exhibit any other change; so that the large embryo, Fig. 272, *e*, is not apparently more advanced as regards the formation of its organs, than the small embryo, Fig. 271, *κ*. So soon as this operation has been completed, however, and the embryo has attained its full bulk, the

evolution of its organs takes place very rapidly; the ciliated lobes are much more highly developed, being extended in a long sinuous margin, so as almost to remind the observer of the "wheels" of Rotifera (§ 277), and being furnished with very long cilia (Fig. 272, B); the auditory vesicles, the tentacula, the eyes, and the foot, successively make their appearance; a curious rhythmically contractile vesicle is seen, just beneath the edge of the shell in the region of the neck, which may perhaps, serve as a temporary heart; a little later, the real heart may be seen pulsating beneath the dorsal part of the shell; and the mass of yolk-segments of which the body is made up, gradually shapes itself into the various organs of digestion, respiration, &c., during the evolution of which (and while they are as yet far from complete) the capsule thins away at its summit, and the embryos make their escape from it.



Later stages of embryonic development of *Purpura lapillus*:—A, conglomerate mass of vitelline segments, to which were attached the embryos, a, b, c, d, e:—B, full-sized embryo, in more advanced stage of development.

352. It happens not unfrequently, that one of the embryos which a capsule contains, does not acquire its supplemental yolk in the manner now described, and can only proceed in its development as far as its original yolk will afford it material; and thus, at the time when the other embryos have attained their full size and maturity, a strange-looking creature, consisting of two large ciliated lobes with scarcely the rudiment of a body, may be seen in active motion among them. This may happen, indeed, not only to one but to several embryos within the same capsule, especially if their number should be considerable; for it sometimes appears as if there were not food enough for all, so that whilst some attain their full dimensions and complete development, others remain of unusually small size, without being deficient in any of their organs, and others again are more or less completely abortive,—the supply of supplemental yolk which they have obtained having been too small for the development of their viscera, although it may have afforded what was needed for that of the ciliated lobes, eyes, tentacles, auditory vesicles, and even the foot,—or, on the other hand, no additional supply whatever having been acquired by them, so that their development has been arrested at a still earlier stage.

These phenomena are of so novel and remarkable a character, that they furnish an abundant source of interest to any Microscopist who may happen to be spending the months of August and September in a locality in which the *Purpura* abounds; since, by opening a sufficient number of capsules, no difficulty need be experienced in arriving at all the facts which have been noticed in this brief summary.¹ It is much to be desired that such Microscopists as possess the requisite opportunity, would apply themselves to the study of the corresponding history in other Pectinibranchiate Gasteropods, with a view of determining how far the plan now described prevails through the order. And now that these Mollusks have been brought not only to live, but to breed, in artificial "vivaria," it may be anticipated that a great addition to our knowledge of this part of their life-history will ere long be made.²

353. *Ciliary Motion on Gills*.—There is no object that is better suited to exhibit the general phenomena of ciliary motion (§ 276), than a portion of the gill of some "bivalve" Mollusk. The *Oyster* will answer the purpose sufficiently well; but the cilia are much larger on the gills of the *Mussel*,³ as they are also on those of the *Anodon* or common "fresh-water mussel" of our ponds and streams. Nothing more is necessary, than to detach a small portion of one of the riband-like bands, which will be seen running parallel with the edge of each of the valves when the shell is opened; and to place this, with a little of the liquor contained within the shell, upon a slip of glass,—taking care to spread it out sufficiently with needles to separate the *bars* of which it is composed, since it is on the edges of these, and round their knobbed extremities, that the ciliary movement presents itself,—and then covering it with a thin glass disk. Or it will be convenient to place the object in the animalcule cage, which will enable the observer to subject it to any degree of pressure that he may find convenient. A magnifying power of about 120 diameters is amply sufficient to afford a general view of this spectacle; but a much greater amplification is needed, to bring into view the peculiar mode in which the stroke of each cilium is made. Few spectacles are more striking to the unpre-

¹ Fuller details on this subject will be found in the Author's account of his researches, in the Second Series of the "Transactions of the Microscopical Society," vol. iii, p. 17; and he would refer such of his readers as may be desirous of knowing what has been done by other observers in regard to the development of different species of Gasteropods, to the various Memoirs there cited.

² The Author thinks it worth his while to mention the method which he has found most convenient for examining the contents of the capsules of *Purpura*; as he believes that it may be advantageously adopted in many other cases. This consists in cutting off the two ends of the capsule (taking care not to cut far into its cavity), and in then forcing a jet of water through it, by inserting the end of a fine-pointed syringe (§ 132) into one of the orifices thus made, so as to drive the contents of the capsule before it through the other. These should be received into a shallow cell, and first examined under a single microscope.

³ This shell-fish may be obtained, not merely at the sea-side, but likewise at the shops of the Fishmongers who supply the humbler classes, even in midland towns.

pared mind, than the exhibition of such wonderful activity as will then become apparent, in a body which to all ordinary observation is so inert. This activity serves a double purpose; for it not only drives a continual current of water over the surface of the gills themselves, so as to effect the aeration of the blood, but it also directs a portion of this current (as in the *Tunicata*, § 333) to the mouth, so as to supply the digestive apparatus with the aliment afforded by the *Diatomaceæ*, *Infusoria*, &c., which it carries in with it.

354. *Organs of Sense of Mollusks.*—Some of the minuter and more rudimentary forms of the special organs of sight, hearing, and touch, which the Molluscous series presents, are very interesting objects of Microscopic examination. Thus just within the margin of each valve of *Pecten*, we see (when we observe the animal in its living state, under water) a row of minute circular points of great brilliancy, each surrounded by a dark ring; these are the eyes, with which this creature is provided for the purpose (it can scarcely be doubted) of directing its peculiarly active movements. Each of them, when their structure is carefully examined, is found to be protected by a sclerotic coat with a transparent cornea in front, and to possess a colored iris (having a pupil) that is continuous with a layer of pigment lining the sclerotic, a crystalline lens and vitreous body, and a retinal expansion proceeding from an optic nerve which passes to each eye from the trunk that runs along the margin of the mantle. Eyes of still higher organization are borne upon the head of most Gasteropod Mollusks, generally at the base of one of the pairs of tentacles, but sometimes, as in the *Snail* and *Slug*, at the points of these organs. In the latter case, the tentacles are furnished with a very peculiar provision for the protection of the eyes; for when the extremity of either of them is touched, it is drawn back into the basal part of the organ, much as the finger of a glove may be pushed back into the palm. The retraction of the tentacle is accomplished by a long muscular slip, which arises within the head, and proceeds to the extremity of the tentacle; whilst its protrusion is effected by the agency of the circular bands with which the tubular wall of the tentacle is itself furnished, the inverted portion being (as it were) squeezed out by the contraction of the lower part into which it has been drawn back. The structure of the eyes, and the curious provision just described, may easily be examined by snipping off one of the eye-bearing tentacles with a pair of scissors. None but the Cephalopod Mollusks have distinct organs of hearing; but rudiments of such organs may be found in most Gasteropods, attached to some part of the nervous collar that surrounds the œsophagus; and even in many Bivalves, in connection with the nervous ganglion imbedded in the base of the foot. These "auditory vesicles," as they are termed, are minute sacculi, each of which contains a fluid, wherein are suspended a

number of minute calcareous particles (named *otolithes* or ear-stones), which are kept in a state of continual movement by the action of cilia lining the vesicles. This "wonderful spectacle," as it was truly designated by its discoverer Siebold, may be brought into view without any dissection, by submitting the head of any small and not very thick-skinned Gasteropod, or the young of the larger forms, to gentle compression under the Microscope, and transmitting a strong light through it. The very early appearance of the auditory vesicles in the embryo Gasteropod has been already alluded to (§ 350). Those who have the opportunity of examining young specimens of the common *Pecten*, will find it extremely interesting to watch the action of the very delicate tentacles which they have the power of putting forth from the margin of their mantle, the animal being confined in a shallow cell, or in the zoophyte-trough; and if the observer should be fortunate enough to obtain a specimen so young that the valves are quite transparent, he will find the spectacle presented by the ciliary movement of the gills, as well as the active play of the foot (of which the adult animal can make no such use), to be worthy of more than a cursory glance.

355. *Chromatophores of Cephalopods.*—Almost any species of Cuttle-fish (*Sepia*) or Squid (*Loligo*) will afford the opportunity of examining the very curious provision which their skin contains for changing its hue. This consists in the presence of numerous large "pigment-cells," containing coloring matter of various tints; the prevailing color, however, being that of the fluid of the ink-bag. These pigment-cells may present very different forms, being sometimes nearly globular, whilst at other times they are flattened and extended into radiating prolongations; and, by the peculiar contractility with which they are endowed, they can pass from one to the other of these conditions, so as to spread their colored contents over a comparatively large surface, or to limit them within a comparatively small area. Very commonly there are different layers of these pigment-cells, their contents having different hues in each layer; and thus a great variety of coloration may be given, by the alteration of the form of the cells of which one or another layer is made up. It is curious that the changes in the hue of the skin appear to be influenced, as in the case of the Chameleon, by the color of the surface with which it may be in proximity. The alternate contractions and extensions of these pigment-cells or "chromatophores" may be easily observed in a piece of skin detached from the living animal, and viewed as a transparent object; since they will continue for some time, if the skin be placed in sea-water. And they may also be well seen in the embryo Cuttle-fish, which will sometimes be found in a state of sufficient advancement, in the grape-like eggs of these animals attached to sea-weeds, zoophytes, &c.

CHAPTER XV.

ANNULOSA, OR WORMS.

856. UNDER the general designation of "annulose" animals, or Worms, may be grouped together all that lower portion of the *Articulated* sub-kingdom, in which the division of the body into longitudinally arranged segments is not distinctly marked out, and in which there is an absence of those "articulated" or jointed limbs, that constitute so distinct a feature of Insects and their allies. This group includes the classes of *Entozoa* or Intestinal Worms, *Rotifera* or Wheel-Animalcules, *Turbellaria*, and *Annelida*; each of which furnishes many objects for Microscopic examination, that are of the highest scientific interest. As our business, however, is less with the professed Physiologist, than with the general inquirer into the minute wonders and beauties of Nature, we shall pass over these classes (the *Rotifera* having been already treated of in detail, Chap. IX), with only a notice of such points as are likely to be specially deserving the attention of observers of the latter order.

857. *Entozoa*.—This class consists almost entirely of animals of a very peculiar plan of organization, which are parasitic within the bodies of other animals, and which obtain their nutriment by the absorption of the juices of these,—thus bearing a striking analogy to the parasitic Fungi (§§ 209–212). The most remarkable feature in their structure, consists in the entire absence, or the extremely low development, of their nutritive system, and the extraordinary development of their reproductive apparatus. Thus, in the common *Tenia* (tape-worm), which may be taken as the type of the "cestoid" group, there is neither mouth nor stomach, the so-called "head" being merely an organ for attachment, whilst the segments of the "body" contain repetitions of a complex generative apparatus, the male and female sexual organs being so united in each, as to enable it to fertilize and bring to maturity its own very numerous eggs; and the chief connection between these segments is established by two pairs of longitudinal canals, which, though regarded by some as representing a digestive apparatus, and by others as a circulating system, appear really to represent the "water-vascular system," whose simplest condition has been noticed in the Wheel-animalcule (§ 278). Few among the recent results of microscopic inquiry have been more curious, than the elucidation of the real

nature of the bodies formerly denominated *cystic* Entozoa, which have always ranked until recently as a distinct group. These are not found, like the preceding, in the cavity of the alimentary canal of the animals they infest; but always occur in the substance of solid organs, such as the glands, muscles, &c. They present themselves to the eye as bags or vesicles of various sizes, sometimes occurring singly, sometimes in groups; but upon careful examination, each vesicle is found to bear upon some part a "head" furnished with hooklets and suckers; and thus may be either single, as in *Cysticercus* (the entozoon whose presence gives to pork what is known as the "measly" disorder), or multiple, as in *Cœnurus*, which is developed in the brain, chiefly of sheep, giving rise to the disorder known as "the staggers." Now in none of these "cystic" forms has any generative apparatus ever been discovered; and hence they are obviously to be considered as imperfect animals. The close resemblance between the "head" of certain *Cysticerci* and that of certain *Tæniæ*, first suggested that the two might be different states of the same animal; and experiments recently made by those who have devoted themselves to the working out of this curious subject, have led to the assured conclusion, that the "cystic" Entozoa are nothing else than "cestoid" worms, whose development has been modified by the peculiarity of their position,—the large bag being formed by a sort of dropsical accumulation of fluid when the young are evolved in the midst of solid tissues, whilst the very same bodies, conveyed into the alimentary canal of some carnivorous animal which has fed upon the flesh infested with them, begin to bud forth the generative segments, the long succession of which, united end to end, gives to the entire series a worm-like aspect.

358. In the intestinal canal of Insects, Centipedes, &c., a very curious kind of animal parasite is often to be met with, the simplicity of whose structure seems to carry us back to the Protozoa (Chap. IX). It is not yet by any means certain, however, that we know the entire life-history of this parasite, the *Gregarina*; and it may be only a phase in the existence of some higher kind of Entozoon. Each individual essentially consists of a single cell, usually more or less ovate in form, and sometimes considerably elongated; a sort of beak or proboscis frequently projects from one extremity; and in some instances this is furnished with a circular row of hooklets, closely resembling that which is seen on the head of *Tænia*. Within the cavity of the cell, whose contents are usually milk white and minutely granular, there is generally seen a pellucid nucleus; and this becomes first constricted and then cleft, when, as often happens, the cell subdivides into two, by a process exactly analogous to that which takes place in the simplest Protophytes (§ 150). The membrane and its contents, except the nucleus, are soluble in acetic acid. Cilia have been detected both upon the outer and the inner sur-

face ; but these would seem destined, not so much to give motion to the body, as to renew the stratum of fluid in contact with it ; for such change of place as the animal does exhibit, is effected by the contractions and extensions of the body generally, as in the *Amœba* (§ 261). A sort of "conjugation" has been seen to take place between two individuals, whose bodies, coming into contact with each other by corresponding points, first become more globular in shape, and are then encysted by the formation of a capsule around them both ; and the partition walls between their cavities disappear ; and the substance of the two bodies becomes completely fused together. As the product of this conjugation, there are first seen a number of globules or cell-like bodies ; and these gradually assume a form so like that of *Naviculæ* (§ 184) as to have been mistaken for them ; their walls, however, are destitute of silex, and there is no further resemblance between the two kinds of bodies, than that of figure. These "pseudo-naviculæ" are set free, in time, by the bursting of the capsule that encloses them ; and they develop themselves into a new generation of Gregarinæ, first passing through an *Amœbo*-like form. It appears, however, that the "pseudo-naviculæ" or "psorosperms" may be also formed by the simple division of the granular matter of a single Gregarina body, without any conjugation.¹

359. The higher forms of Entozoa, belonging to the "nematoid" or thread-like order—of which the common *Ascaris* may be taken as a type, one species of it (the *A. lumbricoidea*, or "round worm") being a common parasite in the small intestine of man, while another (the *A. vermicularis*, or "thread worm") is found rather in the lower bowel,—approach more closely to the ordinary type of conformation of Worms ; having a distinct alimentary canal, which commences with a mouth at the anterior extremity of the body, and which terminates by an anal orifice near the other extremity ; and also possessing a regular arrangement of circular and longitudinal muscular fibres, by which the body can be shortened, elongated, or bent in any direction. The smaller species of *Ascaris*, by some or other of which almost every Vertebrated animal is infested, are so transparent, that every part of their internal organization may be made out, especially with the assistance of the "compressor," without any dissection ; and the study of the structure and actions of their generative apparatus has yielded many very interesting results, especially in regard to the first formation of the ova, the mode of their fertilization, and the history of their subsequent development. Some of the worms belonging to this group are not parasitic in the bodies of other animals, but live in the midst of dead or decomposing vegetable matter. The *Gordius* or hair-worm, which is peculiar in not having any perceptible anal

¹ For the most recent information on this point, see a memoir by M. Nat. Lieberkühn, in *Mem. de l'Acad. Roy. de Belgique*, tom. xvi.

orifice, seems to be properly a parasite in the intestines of water insects; but it is frequently found in large knot-like masses (whence its name) in the water or mud of the pools inhabited by such insects, and may apparently be developed in these situations. The *Anguillulæ* are little eel-like worms, of which one species, *A. fluviatilis*, is very often found in fresh water amongst *Desmidiæ*, *Confervæ*, &c., also in wet moss and moist earth, and sometimes also in the alimentary canal of snails, frogs, fishes, insects, and larger worms; whilst another species, *A. tritici*, is met with in the ears of wheat affected with the blight termed the "cockle;" another, the *A. glutinis*, is found in sour paste; and another, the *A. aceti*, was often found in stale vinegar, until the more complete removal of mucilage and the addition of sulphuric acid, in the course of the manufacture, rendered this liquid a less favorable "habitat" for these little creatures. A writhing mass of any of these species of "eels," is one of the most curious spectacles which the Microscopist can exhibit to the unscientific observer; and the capability which they all possess (in common with the Rotifera and Tardigrada) of revival after desiccation, at however remote an interval, enables him to command this spectacle at any time. A grain of wheat within which these worms (often called *vibriones*) are being developed, gradually assumes the appearance of a black pepper corn; and if it be divided in two, the interior will be found almost completely filled with a dense white cottony mass, occupying the place of the flour, and leaving merely a small space for a little glutinous matter. The cottony substance seems to the eye to consist of bundles of fine fibres closely packed together; but on taking out a small portion, and putting it under the microscope, with a little water under a thin glass cover, it will be found after a short time (if not immediately) to be a wriggling mass of life, the apparent fibres being really *Anguillulæ*, or the "eels" of the Microscopist. If the seeds be soaked in water for a couple of hours before they are laid open, the eels will be found in a state of activity from the first; their movements, however, are by no means so energetic as those of the *A. glutinis* or "paste-eel." This last frequently makes its appearance spontaneously in the midst of paste that is turning sour; but the best means of securing a supply for any occasion, consists in allowing any portion of a mass of paste in which they may present themselves, to dry up, and then, laying this by so long as it may not be wanted, to introduce it into a mass of fresh paste, which, if it be kept warm and moist, will be found after a few days to swarm with these curious little creatures.

360. Besides the foregoing orders of Entozoa, the "trematode" group must be named; of which the *Distoma hepaticum*, or "fluke," found in the livers of sheep affected with the "rot," is a typical example. Into the details of the structure of this animal, which has the general form of a sole, there is no occasion for us


here to enter: it is remarkable, however, for the branching form of its digestive cavity, which extends throughout almost the entire body, very much as in the Planariæ (Fig. 273); and also for the curious phenomena of its development, several distinct forms being passed through between one sexual generation and another. These have been especially studied in the Distoma which infests the *Lymnæus*; the ova of which are not developed into the likeness of their parents, but into minute worm-like bodies, which seem to be little else than masses of cells enclosed in a contractile integument, no formed organs being found in them; these cells, in their turn, are developed into independent "zooids," which escape from their containing cyst in the condition of free ciliated animalcules; in this condition they remain for some time, and then imbed themselves in the mucus that covers the tail of the Mollusk, in which they undergo a gradual development into true Distomata; and having thus acquired their perfect form, they penetrate the soft integument, and take up their habitation in the interior of the body. Thus a considerable number of Distomata may be produced from a single ovum, by a process of cell-multiplication in an early stage of its development. In some instances, the free ciliated larva possesses distinct eyes; although they are wanting in the fully developed Distoma, the peculiar "habitat" of which would render them useless.

361. *Turbellaria*.—This group of animals, which is distinguished by the presence of cilia over the entire surface of the body, seems intermediate in some respects between the "trematode" Entozoa and the Leech-tribe among Annelida. It deserves special notice here, chiefly on account of the frequency with which the worms of the *Planarian* tribe present themselves among collections both of marine and fresh-water animals (particular species inhabiting either locality), and on account of the curious organization which many of these present. Most of the members of this tribe have elongated flattened bodies, and move by a sort of gliding or crawling action over the surfaces of aquatic plants and animals. Some of the smaller kinds are sufficiently transparent to allow of their internal structure being seen by transmitted light, especially when they are slightly compressed; and the accompanying figure (Fig. 273) displays the general conformation of their principal organs, as thus seen. The body has the flattened sole-like shape of the "trematode" Entozoa; its mouth, which is situated at a considerable distance from the anterior extremity of the body, is surrounded by a circular sucker, that is applied to the living surface from which the animal draws its nutriment; and the buccal cavity (*b*) opens into a short œsophagus (*c*), which leads at once to the cavity of the stomach. In the true *Planariæ*, the mouth is furnished with a sort of long funnel-shaped proboscis; and this, even when detached from the body, continues to swallow anything presented

to it. The cavity of the stomach does not give origin to any intestinal tube, nor is it provided with any second orifice; but a large number of ramifying canals are prolonged from it, which carry its contents into every part of the body. This seems to render unnecessary any system of vessels for the circulation of nutritive fluids; and the two principal trunks, with connecting and ramifying branches, which may be observed in them, are probably to be regarded in the light of a "water-vascular" system, the function of which is essentially respiratory. Both sets of sexual organs are combined in the same individuals; though the congress of two, each impregnating the ova of the other, seems to be generally necessary. The ovaria, as in the

Entozoa, extend through a large part of their body, their ramifications proceeding from the two oviducts (*k, k*), which have a dilatation (*l*) at their point of junction. There is much obscurity about the history of the embryonic development of these animals; and the facts observed by Siebold seem to be best explained upon the hypothesis, that what has been usually considered as an egg is really an egg-capsule containing several embryos with a store of supplemental yolk, as in *Purpura* (§ 351), which yolk is swallowed by the embryos at a very early period of their development within the capsule. After their emersion from the capsule, the embryos bear so strong a resemblance to certain Infusoria, as to have led Prof. Agassiz to the conclusion, that the genera *Paramecium* and *Kolpoda* are nothing else than Planarian larvæ (§ 266, note). This point, however, is still a matter for investigation.¹ The Planariæ, however, do not multiply by eggs alone; for they occasionally undergo spontaneous fission in a transverse direction, each segment becoming a perfect animal; and an artificial division into two or even more parts may be practised with a like

FIG. 273.



Structure of *Polycelis levigatus* (a Planarian worm):—*a*, mouth, surrounded by its circular sucker; *b*, buccal cavity; *c*, œsophageal orifice; *d*, stomach; *e*, ramifications of gastric canals; *f*, cephalic ganglia and their nervous filaments; *g, g*, testes; *h*, vesicula seminalis; *i*, male genital canal; *k, k*, oviducts; *l*, dilatation at their point of junction; *m*, female genital orifice.


result. In fact, the power of the Planariæ to reproduce portions

¹ See § 129 of Siebold and Stannius's "Vergleichende Anatomie;" also "Muller's Archiv," 1850, p. 485.

which have been removed, seems but little inferior to that of the *Hydra* (§ 301); a circumstance which is peculiarly remarkable, when the much higher character of their organization is borne in mind. They possess a distinct pair of nervous ganglia (*ff*), from which branches proceed to various parts of the body; and in the neighborhood of these are usually to be observed a number (varying from 2 to 40) of *ocelli*, or rudimentary eyes, each having its refracting body or crystalline lens, its pigment layer, its nerve-bulb, and its cornea-like bulging of the skin. The integument of many of these animals is furnished with "thread-cells" or "filiferous capsules," very much resembling those of Zoophytes (§ 310).

362. *Annelida*.—This class includes all the higher kinds of worm-like animals, the greater part of which are marine, though there are several species which inhabit fresh water, and some which live on land. The body in this class is usually very long, and nearly always presents a well-marked segmental division, the segments being for the most part similar and equal to each other, except at the two extremities; but in the lower forms, such as the Leech and its allies, the segmental division is very indistinctly seen, on account of the general softness of the integument. A large proportion of the marine Annelids have special respiratory appendages, into which the fluids of the body are sent for aeration; and these are situated upon the head (Fig. 274), in those species which (like the *Serpula*, *Terebella*, *Sabellaria*, &c.) have their bodies enclosed by tubes, either formed of a shelly substance produced from their own surface, or built up by the agglutination of grains of sand, fragments of shell, &c.; whilst they are distributed along the two sides of the body, in such as swim freely through the water, or crawl over the surfaces of rocks, as is the case with the *Nereidæ*, or simply bury themselves in the sand, as the *Arenicola* or "lob-worm." In these respiratory appendages, the circulation of the fluids may be distinctly seen by microscopic examination; and these fluids are of two

FIG. 274.



Circulating apparatus of *Terebella conchilega*:—*a*, labial ring; *b, b*, tentacula; *c*, first segment of the trunk; *d*, skin of the back; *e*, pharynx; *f*, intestine; *g*, longitudinal muscles of the inferior surface of the body; *h*, glandular organ (liver?); *i*, organs of generation; *j*, feet; *k, k*, branchiæ; *l*, dorsal vessel acting as a respiratory heart; *m*, dorso-intestinal vessel; *n*, venous sinus surrounding œsophagus; *n'*, inferior intestinal vessel; *o, o*, ventral trunk; *p*, lateral vascular branches.

kinds,—first, a colorless fluid, containing numerous cell-like corpuscles, which can be seen in the smaller and more transparent species to occupy the space that intervenes between the outer surface of the alimentary canal and the inner wall of the body, and to pass from this into canals which often ramify extensively in the respiratory organs, but are never furnished with a returning series of passages,—and second, a fluid which is usually red, contains few floating particles, and is enclosed in a system of proper vessels, that communicates with a central propelling organ, and not only carries the fluid away from this, but also brings it back again. In *Terebella*, we find a distinct provision for the aeration of both fluids; for the first is transmitted to the tendril-like tentacula which surround the mouth (Fig. 274, *b*, *b*), whilst the second circulates through the beautiful arborescent branchiæ (*k*, *k*) situated just behind the head. The former are covered with cilia, the action of which continually renews the stratum of water in contact with them, whilst the latter are destitute of these organs; and this seems to be the general fact as to the several appendages to which these two fluids are respectively sent for aeration, the nature of their distribution varying greatly in the different members of the class. The red fluid is commonly considered as blood, and the tubes through which it circulates as bloodvessels; but the Author has elsewhere given his reasons¹ for coinciding in the opinion of Mr. Huxley, that the colorless corpusculated fluid which moves in the general cavity of the body and in its extensions, is that which really represents the blood of other Articulated Animals; and that the system of vessels carrying the red fluid is to be likened on the one hand to the “water-vascular system” of the inferior Worms, and on the other to the tracheal apparatus of Insects (§ 391).

363. In the observation of the beautiful spectacle presented by the respiratory circulation of the various kinds of Annelids which swarm on most of our shores, and in the examination of what is going on in the interior of their bodies (where this is rendered possible by their transparency), the Microscopist will find a most fertile source of interesting occupation, and may easily, with care and patience, make many valuable additions to our present stock of knowledge on these points. There are many of these marine Annelids, in which the appendages of various kinds put forth from the sides of their bodies, furnish very beautiful microscopic objects; as do also the different forms of teeth, jaws, &c., with which the mouth is commonly armed in the free or non-tubicolar species, these being eminently carnivorous. The early history of their development, too, is extremely curious; for they come forth from the egg in a condition very little more advanced than the ciliated gemmules of polypes, consisting of a globular mass of untransformed cells, certain parts of whose surface are covered with cilia; in a few hours, however, this embryonic mass elon-

¹ See his “Principles of Comparative Physiology,” 4th Edit. §§ 218, 219, 222.

gates, and indications of a segmental division become apparent, the head being (as it were) marked off in front, whilst behind this is a large segment thickly covered with cilia, then a narrower and non-ciliated segment, and lastly the caudal or tail-segment which is furnished with cilia. A little later, a new segment is seen to be interposed in front of the caudal; and the dark internal granular mass shapes itself into the outline of an alimentary canal. The number of segments progressively increases by the interposition of new ones between the caudal and its preceding segments; the various internal organs become more and more distinct, eye-spots make their appearance, little bristly appendages are put forth from the segments, and the animal gradually assumes the likeness of its parent; a few days being passed by the tubicolar kinds, however, in the actively moving condition, before they settle down to the formation of a tube. In places where Annelida abound, free swimming larvæ are often to be obtained at the same time and in the same manner as those of the Echinoderms (§ 323); but to carry out any systematic observations on their embryonic development, the eggs should be searched for in the situations which these animals haunt. To one other phenomenon of the greatest interest, presented by various small marine Annelida, the attention of the Microscopist should be specially directed; this is their *luminosity*, which is not a steady glow like that of the glow-worm or fire-fly, but a series of vivid scintillations (strongly resembling those produced by an electric discharge through a tube spotted with tin foil), that pass along a considerable number of segments, lasting for an instant only, but capable of being repeatedly excited by any irritation applied to the body of the animal. These scintillations may be discerned under the microscope, even in separated segments, when they are subjected to the irritation of a needle-point or to a gentle pressure; and it has been ascertained by the careful observations of M. de Quatrefages, that they are given out by the muscular fibres in the act of contraction.¹

364. Among the fresh-water *Annelida*, those most interesting to the Microscopist are the worms of the *Nais* tribe, which are common in our rivers and ponds, living chiefly amidst the mud at the bottom, and especially among the roots of aquatic plants. Being blood-red in color, they give to the surface of the mud, when they protrude themselves from it in large numbers and keep the protruded portion of their bodies in constant undulation, a very peculiar appearance; but if disturbed, they withdraw themselves suddenly and completely. These worms, from the extreme transparency of their bodies, present peculiar facilities for microscopic examination, and especially for the study of the internal circulation of the red liquid commonly considered as blood. There are here no external respiratory organs; and the

¹ See his Memoirs on the Annelida of La Mancha, in "Ann. of Nat. Sci." Sér. 2, tom. xix, and Sér. 3, tom. xiv.

thinness of the general integument appears to supply all needful facility for the aeration of the fluids. One large vascular trunk (dorsal) may be seen lying above the intestinal canal, and another (ventral) beneath it; and each of these enters a contractile dilatation, or heart-like organ, situated just behind the head. The fluid moves forwards in the dorsal trunk as far as the heart, which it enters and dilates; and when this contracts, it propels the fluid partly to the head, and partly to the ventral heart, which is distended by it. The ventral heart, contracting in its turn, sends the blood backwards along the ventral trunk to the tail, whence it passes towards the head as before. In this circulation, it branches off from each of the principal trunks into numerous vessels proceeding to different parts of the body, which then return into the other trunk; and there is a peculiar set of vascular coils, hanging down in the perigastric space that contains the corpusculated liquid representing the true blood, which seem specially destined to convey to it the aerating influence received by the red fluid in its circuit, thus acting (so to speak) like internal gills. The *Naid*-worms have been observed to undergo spontaneous division during the summer months, a new head and its organs being formed for the posterior segment behind the line of constriction, before its separation from the interior. It has been generally believed that each segment continues to live as an entire worm; but Dr. T. Williams has lately asserted, that from the time when the division occurs, neither half takes in any more food, and that the two segments only retain vitality enough to enable them to be (as it were) the "nurses" of the eggs which both include. In the *Leech* tribe, the apparatus of teeth with which the mouth is furnished, is one of the most curious among their points of minute structure; and the common "medicinal" leech affords one of the most interesting examples of it. What is commonly termed the "bite" of the leech, is really a saw cut, or rather a combination of three saw cuts, radiating from a common centre. If the mouth of the leech be examined with a hand-magnifier, or even with the naked eye, it will be seen to be a triangular aperture in the midst of a sucking disk; and on turning back the lips of that aperture, three little white ridges are brought into view. Each of these is the convex edge of a horny semicircle, which is bordered by a row of eighty or ninety minute hard and sharp teeth; whilst the straight border of the semicircle is imbedded in the muscular substance of the disk, by the action of which it is made to move backwards and forwards in a saw-like manner, so that the teeth are enabled to cut into the skin to which the suctorial disk has affixed itself.

CHAPTER XVI.


CRUSTACEA.

PASSING from the lower division of the Articulated series to the higher—that in which the body is furnished with distinctly articulated or jointed limbs,—we come first to the class of *Crustacea*, which includes (when used in its most comprehensive sense) all those animals belonging to this group, which are fitted for aquatic respiration. It thus comprehends a very extensive range of forms; for although we are accustomed to think of the Crab, Lobster, Cray-fish, and other well-known species of the order *Decapoda* (ten-footed), as its typical examples, yet all these belong to the highest of its many orders; and among the lower are many of a far simpler structure, and not a few which would not be recognized as belonging to the class at all, were it not for the information derived from the study of their development as to their real nature, which is far more apparent in their early than it is in their adult condition. Many of the inferior kinds of *Crustacea* are so minute and transparent, that their whole structure may be made out by the aid of the Microscope without any preparation; this is the case, indeed, with nearly the whole group of *Entomostraca* (§ 366), and with the larval forms even of the *Crab* and its allies (§ 375); and we shall give our first attention to these, afterwards noticing such points in the structure of the larger kinds, as are likely to be of general interest.

865. One of the most curious examples of the reduction of an elevated type to its very simplest form, which the Animal Kingdom affords, is presented by the group of *Pycnogonidæ*; some members of which may be found by attentive search in almost every locality where sea-weeds abound, it being their habit to crawl (or rather to sprawl) over the surfaces of these, and probably to imbibe as food the gelatinous substance with which they are invested. The general form of their bodies (Fig. 275) usually reminds us of that of some of the long-legged Crabs; the abdomen being almost or altogether deficient, whilst the head is very small, and fused (as it were) into the thorax; so that the last named region, with the members attached to it, constitutes nearly the whole bulk of the animal. The head is extended in front into a proboscis-like projection, at the extremity of which is the

narrow orifice of the mouth ; which seems to be furnished with vibratile cilia, that serve to draw into it the semi-fluid aliment. Instead of being furnished (as in the higher Crustaceans) with two pairs of antennæ and numerous pairs of "feet-jaws," it has but a single pair of either; it also bears four minute *ocelli*, or

FIG. 275.



Ammotea pycnogonoides:—*a*, narrow œsophagus; *b*, stomach; *c*, intestine; *d*, digestive cœca of the feet-jaws; *e*, *e*, digestive cœca of the legs.

rudimentary eyes, set at a little distance from each other on a sort of tubercle. From the thorax proceed four pairs of legs, each composed of several joints, and terminated by a hooked claw; and by these members the animal drags itself slowly along, instead of walking actively upon them like a crab. The mouth leads to a very narrow œsophagus (*a*), which passes back to the central stomach (*b*), situated in the midst of the thorax, from the hinder end of which a narrow intestine (*c*) passes off, to terminate at the posterior extremity of the body. From the central stomach, five pairs of cœcal prolongations radiate; one pair (*d*) entering the feet-jaws, the other four (*e*, *e*) penetrating the legs, and passing along them as far as the last joint but one; and these extensions are covered with a layer of brownish-yellow granules, which are probably to be regarded as a diffused and rudimentary condition of the liver. The stomach and its cœcal prolongations are continually executing peristaltic movements of a very curious kind; for they contract and dilate with an irregular alternation, so that a flux and reflux of their contents is constantly taking place between the central portion and its radiating extensions, and between one of these extensions and another. The space

between the widely-extended stomach and the walls of the body and limbs is occupied by a transparent liquid, in which are seen floating a number of minute transparent corpuscles of irregular size; and this fluid, which represents the blood, is kept in continual motion, not only by the general movements of the body and limbs, but also by the actions of the digestive apparatus; since, whenever the cæcum of any one of the legs undergoes dilatation, a part of the circumambient liquid will be pressed out from the cavity of that limb, either into the thorax, or into some other limb whose stomach is contracting. The fluid must obtain its aeration through the general surface of the body, as there are no special organs of respiration. The nervous system consists of a single ganglion in the head (formed by the coalescence of a pair), and of another in the thorax (formed by the coalescence of four pairs), with which the cephalic ganglion is connected in the usual mode, namely, by two nervous cords which diverge from each other to embrace the œsophagus. Of the reproduction of this animal, nothing is yet known. In the study of the very curious phenomena exhibited by the digestive apparatus, as well as of the various points of internal conformation which have been described, the achromatic condenser will be found useful, even with the 1 inch, 2-3ds inch, or $\frac{1}{2}$ inch objectives; for the imperfect transparence of the bodies of these animals renders it of importance to drive a large quantity of light through them, and to give to this light such a quality, as shall define the internal organs as sharply as possible.


366. The *Entomostracous* group of Crustaceans, nearly all the existing members of which are of such minute size as to be only just visible to the naked eye, is distinguished by the enclosure of the entire body within a horny or shelly casing, which sometimes closely resembles a bivalve shell in form and in the mode of junction of its parts, whilst in other instances it is formed of only a single piece, like the hard envelope of certain Rotifera (§ 282, III). The segments into which the body is divided, are frequently very numerous, and are for the most part similar to each other; but there is a marked difference in regard to the appendages which they bear, and to the mode in which these minister to the locomotion of the animals. For in the *Lophyropoda*, or bristly-footed tribe, the number of legs is small, not exceeding five pairs, and their function is limited to locomotion, the respiratory organs being attached to the parts in the neighborhood of the mouth; whilst in the *Branchiopoda*, or gill-footed tribe, the same members serve both for locomotion and for respiration, and the number of these is commonly large, being in *Apus* not less than sixty pairs. The character of their movements differs accordingly; for whilst all the members of the first-named tribe dart through the water in a succession of jerks, so as to have acquired the common name of "water-fleas," those among the

latter which possess a great number of "fin-feet," swim with an easy gliding movement, sometimes on their back alone (as is the case with *Branchipus*), sometimes with equal facility on the back, belly, or sides (as is done by *Artemia salina*, the "brine shrimp"). Some of the most common forms of both tribes will now be briefly noticed.

367. The tribe of *Lophyropoda* is divided into two orders; of which the first, *Ostracoda*, is distinguished by the complete enclosure of the body in a bivalve shell, by the small number of legs, and by the absence of an external ovary. One of the best-known examples is the little *Cypris*, which is a common inhabitant of pools and streams; this may be recognized by its possession of two pairs of antennæ, the first having numerous joints with a pencil-like tuft of filaments, and projecting forwards from the front of the head, whilst the second has more the shape of legs, and is directed downwards; and by the limitation of its legs to two pairs, of which the posterior does not make its appearance outside the shell, being bent upwards to give support to the ovaries. The valves are generally opened sufficiently wide, to allow the greater part of both pairs of antennæ, and of the front pair of legs, to pass out between them; but when the animals are alarmed, they draw these members within the shell, and close the valves firmly. They are very lively creatures, being almost constantly seen in motion, either swimming by the united action of their foot-like antennæ and legs, or walking upon plants and other solid bodies floating in the water. Nearly allied to the preceding is the *Cythere*, whose body is furnished with three pairs of legs, all projecting out of the shell, and whose superior antennæ are destitute of the filamentous brush; this genus is almost entirely marine, and some species of it may almost invariably be met with in little pools among the rocks between the tide-marks, creeping about (but not swimming) amongst *Confervæ* and *Corallines*. There is abundant evidence of the former existence of Crustacea of this group, of larger size than any now existing, to an enormous extent; for in certain fresh-water strata, both of the Secondary and Tertiary series, we find layers, sometimes of great extent and thickness, which are almost entirely composed of the fossilized shells of *Cyprides*; whilst in certain parts of the Chalk, which was a marine deposit, the remains of bivalve shells resembling those of *Cythere*, present themselves in such abundance as to form a considerable part of its composition. In the order *Copepoda*, there is a jointed shell forming a kind of buckler that almost entirely encloses the head and thorax, an opening being left beneath, through which the members project; and there are five pairs of legs, mostly adapted for swimming, the fifth pair, however, being rudimentary in the genus *Cyclops*, the commonest example of the group. This genus receives its name from possessing only a single eye,

or rather a single cluster of ocelli; which character, however, it has in common with the two genera already named, as well as with *Daphnia* (§ 368), and with many other Entomostraca. It contains numerous species, some of which belong to fresh water, whilst others are marine. The fresh-water species often abound in the muddiest and most stagnant pools, as well as in the clearest springs; the ordinary water with which London is supplied, frequently contains large numbers of them. Of the marine species, some are to be found in the localities in which the *Cythere* is most abundant, whilst others inhabit the open ocean, and must be collected by a fine muslin net. The body of the Cyclops is soft and gelatinous, and it is composed of two distinct parts, a thorax (Fig. 276, *a*) and an abdomen (*b*), of which the latter, being comparatively slender, is commonly considered as a tail, though traversed by the intestine which terminates near its extremity. The head, which coalesces with the thorax, bears one very large pair of antennæ (*c*) possessing numerous articulations, and furnished with bristly appendages, and another small pair (*d*); it is also furnished with a pair of "mandibles" or true jaws, and with two pairs of "feet-jaws," of which the hinder pair is the longer, and most abundantly supplied with bristles. The legs (*e*) are all beset with plumose tufts, as is also the tail (*f, f*) which is borne at the extremity of the abdomen. On either side of the abdomen of the female, there is often to be seen an egg-capsule or external ovarium (*b*), within which the ova, after being fertilized, undergo the earlier stages of their development. The Cyclops is a very active creature, and strikes the water in swimming, not merely with its legs and tail, but also with its antennæ. The rapidly repeated movements of its feet-jaws serve to create a whirlpool in the surrounding water, by which minute animals of various kinds, and even its own young, are brought to its mouth to be devoured.

FIG. 276.



A, female of *Cyclops quadricornis*:—*a*, body; *b*, tail; *c*, antenna; *d*, antennule; *e*, feet; *f*, plumose setæ of tail:—*B*, tail, with external egg-sac; *C, D, E, F, G*, successive stages of development of young.

slender, is commonly considered as a tail, though traversed by the intestine which terminates near its extremity. The head, which coalesces with the thorax, bears one very large pair of antennæ (*c*) possessing numerous articulations, and furnished with bristly appendages, and another small pair (*d*); it is also furnished with a pair of "mandibles" or true jaws, and with two pairs of "feet-jaws," of which the hinder pair is the longer, and most abundantly supplied with bristles. The legs (*e*) are all beset with plumose tufts, as is also the tail (*f, f*) which is borne at the extremity of the abdomen. On either side of the abdomen of the female, there is often to be seen an egg-capsule or external ovarium (*b*), within which the ova, after being fertilized, undergo the earlier stages of their development. The Cyclops is a very active creature, and strikes the water in swimming, not merely with its legs and tail, but also with its antennæ. The rapidly repeated movements of its feet-jaws serve to create a whirlpool in the surrounding water, by which minute animals of various kinds, and even its own young, are brought to its mouth to be devoured.

368. The tribe of Branchiopoda also is divided into two orders;

of which the *Cladocera* present the nearest approach to the preceding, having a bivalve carapace, no more than from four to six pairs of legs, two pairs of antennæ, of which one is large and branched and adapted for swimming, and a single eye. The commonest form of this is the *Daphnia pulex*, sometimes called the "arborescent water-flea" from the branching form of its antennæ. It is very abundant in many ponds and ditches, coming to the surface in the mornings and evenings and in cloudy weather, but seeking the depths of the water during the heat of the day. It swims by taking short springs; and feeds on minute particles of vegetable substances, not, however, rejecting animal matter when offered. Some of the peculiar phenomena of its reproduction will be presently described (§ 370). The order *Phyllopoda* includes those Branchiopoda whose body is divided into a great number of segments, nearly all of which are furnished with leaf-like members, or "fin-feet." The two families which this order includes, however, differ considerably in their conformation; for in that of which the genera *Apus* and *Nebalis* are representatives, the body is enclosed in a shell, either shield-like or bivalve, and the feet are generally very numerous; whilst in that which contains *Branchipus* and *Artemia*, the body is entirely unprotected, and the number of pairs of feet does not exceed eleven. The *Apus canceriformis*, which is an animal of comparatively large size, its entire length being about $2\frac{1}{2}$ inches, is an inhabitant of stagnant waters; but although occasionally very abundant in particular pools or ditches, it is not to be met with nearly so commonly as the Entomostraca already noticed. It is recognized by its large oval carapace, which covers the head and body like a shield; by the nearly cylindrical form of its body, which is composed of thirty articulations; and by the multiplication of its legs, which amount to about sixty pairs. The number of joints in these and in the other appendages is so great, that in a single individual they may be safely estimated at not less than two millions. These organs, however, are for the most part small; and the instruments chiefly used by the animal for locomotion are the first pair of feet, which are very much elongated (bearing such a resemblance to the principal antennæ of other Entomostraca, as to be commonly ranked in the same light), and are distinguished as *rami* or oars. With these they can swim freely in any position; but when the "rami" are at rest and the animal floats idly on the water, its fin-feet may be seen to be in incessant motion, causing a sort of whirlpool in the water, and bringing to the mouth the minute animals (chiefly the smaller Entomostraca inhabiting the same localities) that serve them as food. The *Branchipus stagnalis* has a slender, cylindrical, and very transparent body of nearly an inch in length, furnished with eleven pairs of fin-feet, but is destitute of any protecting envelope; its head is furnished with a pair of very curious prehensile organs (which are really modified an-

tennæ), whence it has received the name of *Cheirocephalus*; but these are not used by it for the seizure of prey, the food of this animal being vegetable, and their function is to clasp the female in the act of copulation. The Branchipus or Cheirocephalus is certainly the most beautiful and elegant of all the Entomostraca, being rendered extremely attractive to the view by the "uninterrupted undulatory wavy motion of its graceful branchial feet, slightly tinged as they are with a light reddish hue, the brilliant mixture of transparent bluish green and bright red of its prehensile antennæ, and its bright red tail with the beautiful plumose setæ springing from it:" unfortunately, however, it is a comparatively rare animal in this country. The *Artemia salina* or "brine shrimp" is an animal of very similar organization, and almost equally beautiful in its appearance and movements, but of smaller size, its body being about half an inch in length. Its "habitat" is very peculiar; for it is only found in the salt-pans or brine-pits in which sea-water is undergoing concentration (as at Lymington); and in these situations it is sometimes so abundant, as to communicate a red tinge to the liquid.

369. Some of the most interesting points in the history of the Entomostraca lie in the peculiar modes in which their Generative function is performed, and in their tenacity of life when desiccated, in which last respect they correspond with many Rotifera (§ 280). This provision is obviously intended to prevent them from being completely exterminated, as they might otherwise soon be, by the drying up of the pools, ditches, and other small collections of water which constitute their usual "habitats." It does not appear, however, that the adult animals can bear a *complete* desiccation, although they will preserve their vitality in mud that holds the smallest quantity of moisture; but their eggs are more tenacious of life, and there is ample evidence that these will become fertile on being moistened, after having continued for a long time in the condition of fine dust. Most Entomostraca, too, are killed by severe cold, and thus the whole race of adults perishes every winter; but their eggs seem unaffected by the lowest temperature, and thus continue the species which would otherwise be exterminated. Again, we frequently meet in this group with that reproduction by gemmation, which we have seen to prevail so extensively among the lower Radiata and Mollusca. In many species there is a double mode of multiplication, the sexual and the non-sexual. The former takes place at certain seasons only, the males (which are often so different in conformation from the females, that they would not be supposed to belong to the same species, if they were not seen in actual congress) disappearing entirely at other times; whilst the latter continues at all periods of the year, so long as warmth and food are supplied, and is repeated many times (as in the Hydra), so as to give origin to as many successive "broods." Further, a single act of impregnation serves to

fertilize not merely the ova which are then mature or nearly so, but all those subsequently produced by the same female, which are deposited at considerable intervals. In these two modes, the multiplication of these little creatures is carried on with great rapidity, the young animal speedily coming to maturity and beginning to propagate; so that, according to the computation of Jurine, founded upon data ascertained by actual observation, a single fertilized female of the common *Cyclops quadricornis* may be the progenitor in one year of 4,442,189,120 young.

370. The eggs of some Entomostraca are deposited freely in the water, or are carefully attached in clusters to aquatic plants; but they are more frequently carried for some time by the parent in special receptacles developed from the posterior part of the body; and in many cases they are retained there until the young are ready to come forth, so that these animals may be said to be ovo-viviparous. In the *Daphnia*, the eggs are received into a large cavity between the back of the animal and its shell, and there the young undergo almost their whole development, so as to come forth in a form nearly resembling that of their parent. Soon after their birth, a moult or exuviation of the shell takes place; and the egg coverings are cast off with it. In a very short time afterwards, another brood of eggs is seen in the cavity, and the same process is repeated, the shell being again exuviated after the young have been brought to maturity. At certain times, however, the *Daphnia* may be seen with a dark opaque substance within the back of the shell, which has been called the *ephippium* from its resemblance to a saddle. This, when carefully examined, is found to be of dense texture, and to be composed of a mass of hexagonal cells; and it contains two oval bodies, each consisting of an ovum covered with a horny casing, enveloped in a capsule which opens like a bivalve shell. The first traces of the "ephippium" are seen after the third moult, as a green matter in the ovaries, which differs both in color and appearance from that of the eggs; after the fourth moult, this green matter passes from the ovaries into the matrix or open space on the back, and there becomes developed into the "ephippium;" and at the fifth moult this is thrown off, and the ephippium, with the two eggs enclosed, floats on the water until the next spring, when the young are hatched with the returning warmth of the season. This curious provision is obviously destined to afford protection to the eggs which are to endure the severity of winter cold; and some approach to it may be seen in the remarkable firmness of the envelopes of the "winter eggs" of some of the Rotifera (§ 279). It has been ascertained by Dr. Baird, that the young produced from the ephippial eggs have the same power of continuing the race by non-sexual reproduction, as the young developed under ordinary circumstances.

371. In most Entomostraca, the young at the time of their

emersion from the egg differ considerably from the parent, especially in having only the thoracic portion of the body as yet evolved, and in possessing but a small number of locomotive appendages; the visual organs, too, are frequently wanting at first. (See Fig. 276, c-g.) The process of development, however, takes place with great rapidity; the animal at each successive moult (which process is very commonly repeated at intervals of a day or two) presenting some new parts, and becoming more and more like its parent, which it very early resembles in its power of multiplication, the female laying eggs before she has attained her own full size. Even when the Entomostraca have attained their full growth, they continue to exuviate their shell at short intervals during the whole of life; and the purpose which seems to be answered by this repeated moulting, is the preventing the animal from being injured, or its movements obstructed, by the overgrowth of parasitic Animalcules and Confevæ; weak and sickly individuals being frequently seen to be so covered with such parasites, that their motion and life are soon arrested, apparently because they have not strength to cast off and renew their envelopes. The process of development appears to depend in some degree upon the influence of light, being retarded when the animals are secluded from it; but its rate is still more influenced by heat; and this appears also to be the chief agent that regulates the time which elapses between the moultings of the adult, these, as in the *Daphnia*, taking place at intervals of two days in warm summer weather, whilst several days intervene between them when the weather is colder. The cast shell carries with it the sheaths not only of the limbs and plumes, but of the most delicate hairs and setæ which are attached to them. If the animal have previously sustained the loss of a member, it is generally renewed at the next moult, as in higher Crustacea.¹

372. Closely connected with the Entomostracous group is the tribe of *Suctorial* Crustacea; which for the most part live as parasites upon the exterior of other animals (especially Fish), whose juices they imbibe by means of the peculiar proboscis-like organ which takes in them the place of the jaws of other Crustaceans; whilst other appendages, representing the feet-jaws, are furnished with hooks, by which these parasites attach themselves to the animals from whose juices they derive their nutriment. Many of the Suctorial Crustacea bear a strong resemblance, even in their adult condition, to certain Entomostraca; but more commonly it is between the earlier forms of the two groups that the resemblance is the closest, most of the *Suctorial* undergoing such extraordinary changes in their progress towards the adult condition, that if their complete forms were alone attended to, they might be excluded from the class alto-

¹ For a complete and detailed account of this group, see Dr. Baird's "Natural History of the British Entomostraca," published by the Ray Society.

gether, as has (in fact) been done by many Zoologists. Among those Suctorial Crustacea which present the nearest approach to the ordinary Entomostracous type, may be specially mentioned the *Argulus foliaceus*, which attaches itself to the surface of the bodies of fresh-water fish, and is commonly known under the name of the "fish-louse." This animal has his body covered with a large firm oval shield, which does not extend, however, over the posterior part of the abdomen. The mouth is armed with a pair of styliform mandibles; and on each side of the proboscis there is a large short cylindrical appendage, terminated by a curious sort of sucking disk, with another pair of longer jointed members, terminated by prehensile hooks. These two pairs of appendages, which are probably to be considered as representing the feet-jaws, are followed by four pairs of legs, which, like those of the Branchiopoda, are chiefly adapted for swimming; and the tail, also, is a kind of swimmeret. This little animal can leave the fish on which it feeds, and then swims freely in the water, usually in a straight line, but frequently and suddenly changing its direction, and sometimes turning over and over several times in succession. The stomach is remarkable for the large cæcal prolongations which it sends out on either side, immediately beneath the shell; for these subdivide and ramify in such a manner, that they are distributed almost as minutely as the cæcal prolongations of the stomach of the *Planaria* (Fig. 273). The proper alimentary canal, however, is continued backwards from the central cavity of the stomach, as an intestinal tube, which terminates in an anal orifice at the extremity of the abdomen.¹


373. From the parasitic suctorial Crustacea, the transition is not really so abrupt as it might at first sight appear, to the class of *Cirrhipeda*, consisting of the *Barnacles* and their allies; which, like many of the Suctoria, are fixed to one spot during the adult portion of their lives, but come into the world in a condition that bears a strong resemblance to the early state of many of the Crustacea. The departure from the ordinary Crustacean type in the adult, is, in fact, so great, that it is not surprising that Zoologists in general should have separated them; their superficial resemblance to the Mollusca, indeed, having caused most systematists to rank them in that series, until due weight was given to those structural features which mark their Articulated character. We must limit ourselves in our notice of this group, to that very remarkable part of their history, the Microscopic study of which has contributed most essentially to the elucidation of their real nature. The observations of Mr. J. V. Thompson,² with the extensions and rectifications which they have subsequently re-

¹ As this group is rather interesting to the professed Naturalist than to the amateur Microscopist, even an outline view of it would be unsuitable to the present treatise: and the Author would refer such of his readers as may desire to study it, to the admirable treatise by Dr. Baird, already referred to.

² "Zoological Researches," No. III, 1830.

ceived from others, show that there is no essential difference between the early forms of the *sessile* (Balanidæ, or "acorn-shells") and of the *pedunculated* Cirrhipeds (Lepadidæ or "barnacles"); for that both are active little animals (Fig. 277, A), possessing three pairs of legs and a pair of compound eyes, and having the body covered with an expanded shield, like that of many Entomostracous Crustaceans, so as in no essential particular to differ from the larva of *Cyclops* (Fig. 276, c). After going through a series of metamorphoses, one stage of which is represented in Fig. 277, B, c, these larvæ come to present a form D, which reminds us strongly of that of *Daphnia*; the body being enclosed

FIG. 277.



Development of *Balanus balanoides*:—A, earliest form;—B, larva after second moult;—C, side view of the same;—D, stage immediately preceding the loss of activity; a, stomach (?); b, nucleus of future attachment (?).

in a shell composed of two valves, which are united along the back, whilst they are free along their lower margin, where they separate for the protrusion of a large and strong anterior pair of prehensile limbs provided with an adhesive sucker and hooks, and of six pairs of posterior legs adapted for swimming. This bivalve shell, with the members of both kinds, is subsequently thrown off; the animal then attaches itself by its head, a portion of which becomes excessively elongated into the "peduncle" of the Barnacle, whilst in *Balanus* it expands into a broad disk of adhesion; the first thoracic segment sends backwards a prolongation, which arches over the rest of the body so as completely to enclose it, and of which the exterior layer is consolidated into the "multivalve" shell; whilst from the other thoracic segments are evolved the six pairs of *cirri*,—which are long, slender, many-jointed, tendril-like appendages, fringed with delicate filaments covered with cilia, whose action serves both to bring food


to the mouth, and to maintain aerating currents in the water,—from whose peculiar character the name of the group is derived.

374. The chief points of interest to the Microscopist in the more highly organized forms of Crustacea, are furnished by the structure of the shell, and by the metamorphosis of the larvæ, both of which may be best studied in the commonest kinds. The shell of the Decapods in its most complete form consists of three strata; namely, 1, a horny structureless layer covering the exterior; 2, a cellular stratum; and 3, a laminated tubular substance. The innermost and even the middle layers, however, may be altogether wanting; thus in the *Phyllosomæ* or “glass-crabs,” the envelope is formed by the transparent horny layer alone; and in many of the small Crabs belonging to the genus *Portuna*, the whole substance of the carapace beneath the horny investment is made up of hexagonal thick-walled cells. (It may be here noticed, that the carapace of *Daphnia*, *Branchipus*, and some other Entomostraca, exhibits the hexagonal division; whilst in many species this is not distinguishable.) It is in the large thick-shelled Crabs that we find the three layers most differentiated. Thus in the common *Cancer pagurus*, we may easily separate the structureless horny covering after a short maceration in dilute acid; the cellular layer, in which the pigmentary matter of the colored parts of the shell is contained, may be easily brought into view by grinding away as flat a piece as can be selected, from the *inner* side, having first cemented the outer surface to the glass slide, and by examining this with a magnifying power of 250 diameters, driving a strong light through it with the achromatic condenser; whilst the tubular structure of the thick inner layer may be readily demonstrated, by means of sections parallel and perpendicular to its surface. This structure, which very strongly resembles that of *dentine* (§ 406), save that the tubuli do not branch, but remain of the same size through their whole course, may be particularly well seen in the black extremity of the claw, which (apparently from some difference in the molecular arrangement of the mineral particles, the organic structure being precisely the same) is much denser than the rest of the shell, the former having almost the semi-transparency of ivory, whilst the latter has a chalky opacity. In a transverse section of the claw, the tubuli may be seen to radiate from the central cavity towards the surface, so as very strongly to resemble their arrangement in a tooth; and the resemblance is still further increased by the presence, at tolerably regular intervals, of minute sinuosities corresponding with the laminations of the shell, which seem, like the “secondary curvatures” of the dentinal tubuli, to indicate successive stages in the calcification of the animal basis. This inner layer rises up through the pigmentary layer of the Crab’s shell, in little papillary elevations; and it is from the deficiency of the pigmentary layer at these parts, that the colored portion of the shell derives its

minutely speckled appearance. Many departures from this type are presented by the different species of Decapods; thus in the *Prawns* there are large stellate pigment-cells (resembling those of Fig. 327, c), the colors of which are often in remarkable conformity with those of the bottom of the rock-pools frequented by these creatures; whilst in the *Shrimps* there is seldom any distinct trace of the cellular layer, and the calcareous portion of the skeleton is disposed in the form of concentric rings, an approach to which arrangement is seen in the papillæ of the surface of the deepest layer of the Crab's shell.

375. It is a very curious circumstance, that a strongly marked difference exists between Crustaceans that are otherwise very closely allied, in regard to the degree of change to which their young are subject in their progress towards the adult condition. For whilst the common *Crab*, *Lobster*, *Spiny Lobster*, *Prawn*, and *Shrimp*, undergo a regular metamorphosis, the young of the *Land Crab* and the *Cray-fish* come forth from the egg in a form which corresponds in all essential particulars with that of their parents. Generally speaking, a strong resemblance exists among the young of all the species of Decapods which undergo a metamorphosis, whether they are afterwards to belong to the *brachy-ourous* (short-tailed) or to the *macrourous* (long-tailed) division of the group; and the forms of these larvæ are so peculiar, and so entirely different from any of those into which they are ultimately to be developed, that they were considered as belonging to a distinct genus, *Zoea*, until their real nature was first ascertained by Mr. J. V. Thompson. Thus in the earliest state of *Carcinus menas* (small edible crab), we see the head and thorax, which form the principal bulk of the body, included within a large carapace or shield (Fig. 278, A) furnished with a long projecting spine, beneath which the fin-feet are put forth; whilst the abdominal

FIG. 278.



Metamorphosis of *Carcinus menas*:—A, first stage; B, second stage; C, third stage, in which it begins to assume the adult form; D, perfect form.

segments, narrowed and prolonged, carry at the end a flattened tail-fin, by the strokes of which upon the water, the propulsion of the animal is chiefly effected. Its condition is hence compa-

rable, in almost all essential particulars, to that of *Cyclops* (§ 367). In the case of the Lobster, Prawn, and other "macrourous" species, the metamorphosis chiefly consists in the separation of the locomotive and respiratory functions, true legs being developed from the thoracic segments for the former, and true gills (concealed within a special chamber formed by an extension of the carapace beneath the body) for the latter; and the abdominal segments increase in size, and become furnished with appendages (false feet) of their own. In the Crabs, or "brachyurous" species, on the other hand, the alteration is much greater; for besides the change first noticed in the thoracic members and respiratory organs, the thoracic region becomes much more developed at the expense of the abdominal, the latter remaining in an almost rudimentary condition, and being bent under the body; the thoracic limbs are more completely adapted for walking, save the first pair, which are developed into *chela* or pincers; and the little creature entirely loses the active swimming habits which it originally possessed, and takes on the mode of life peculiar to the adult. We have, in this history, a most characteristic example of Von Bär's great law of "progress from the general to the special" in organic development; for the *Entomostracous* form is thus seen to be common to the highest and the lowest Crustaceans in the earliest phase of their lives; but whilst the latter remain and go on to completion upon that type, the former entirely diverge from it; and whilst diverging from it, they also become differentiated from each other, the distinctive characters of their families, genera, and species, evolving themselves, as the individuals advance towards their mature forms.

376. In collecting minute Crustacea, whether fresh-water or marine, the use of the ring-net, as for minute Acalephæ or Echinoderm larvæ, will be found the most efficient instrument; and in favorable localities, the same "gathering" will often contain multitudes of various species of Entomostraca, accompanied, perhaps, by the larvæ of higher Crustacea, by Echinoderm larvæ, by Annelid larvæ, and by the smaller Medusæ. The water containing these should be put into a large glass jar freely exposed to the light; and after a little practice, the eye will become so far habituated to the general appearance and modes of movement of these different forms of Animal life, as to be able to distinguish them, one from the other. In selecting any specimen for microscopic examination, the dipping tube (§ 71) will be found invaluable. The study of the metamorphoses of the larvæ will be best prosecuted, by obtaining the fertilized eggs which are carried about by the females, and watching the history of their products.

CHAPTER XVII.

INSECTS AND ARACHNIDA.

THERE is no class in the whole Animal Kingdom which affords to the Microscopist such a wonderful variety of interesting objects, and such facilities for obtaining an almost endless succession of novelties, as that of Insects. For, in the first place, the number of different kinds that may be brought together (at the proper time) with extremely little trouble, far surpasses that which any other group of Animals can supply to the most painstaking collector; then, again, each specimen will afford, to him who knows how to employ his materials, a considerable number of microscopic objects of very different kinds; and, thirdly, although some of these objects require much care and dexterity in their preparation, a large proportion may be got out, examined, and mounted, with very little skill or trouble. Take, for example, the common House-Fly:—its *eyes* may be easily mounted, one as a transparent, the other as an opaque object (§ 383); its *antennæ*, although not such beautiful objects as those of many other Diptera, are still well worth examination (§ 385); its *tongue* or “proboscis” is a peculiarly interesting object (§ 386), though requiring some care in its preparation; its *spiracles*, which may be easily cut out from the sides of its body, have a very curious structure (§ 392); its alimentary canal affords a very good example of the minute distribution of the “trachææ” (§ 391); its *wings*, examined on a living specimen, newly come forth from the pupa state, exhibit the circulation of the blood in the “nervures” (§ 390); the wing of the insect when dead, moreover, exhibits a most beautiful play of iridescent colors, and shows a remarkable areolation of surface, when it is examined by light reflected from its surface at a particular angle (§ 395); its *foot* has a very peculiar conformation, which is doubtless connected with its singular power of walking over smooth surfaces in direct opposition to the force of gravity, although the mode in which it serves this purpose is not yet certainly ascertained (§ 397); and the structure and physiology of its sexual apparatus, with the history of its development and metamorphoses, would of itself suffice to occupy the whole time of an

observer who should desire thoroughly to work it out, not only for months but for years. Hence in treating of this department in such a work as the present, the author labors under the *embarras des richesses*; for to enter into such a description of the parts of the structure of Insects most interesting to the Microscopist, as should be at all comparable in fulness with the accounts which it has been thought desirable to give of other classes, would swell out the volume to an inconvenient bulk; and no course seems open, but to limit the treatment of the subject to a notice of the *kinds* of objects which are likely to prove most generally interesting, with a few illustrations that may serve to make the descriptions more clear, and with an enumeration of some of the sources whence a variety of specimens of each class may be most readily obtained. And thus limitation is the less to be regretted, since there already exist in our language numerous elementary treatises on Entomology, wherein the general structure of Insects is fully explained, and the conformation of their minute parts as seen with the Microscope is adequately illustrated.

377. A considerable number of the smaller Insects,—especially those belonging to the orders *Coleoptera* (beetles), *Neuroptera* (dragon fly, May fly, &c.), *Hymenoptera* (bee, wasp, &c.), and *Diptera* (two-winged flies), may be mounted entire as opaque objects for low magnifying powers; care being taken to spread out their legs, wings, &c., so as adequately to display them, which may be accomplished even after they have dried in other positions, by softening them by steeping them in hot water, or, where this is objectionable, by exposing them to steam. Full directions on this point, applicable to small and large Insects alike, will be found in all text-books of Entomology. There are some, however, whose translucency allows them to be viewed as transparent objects; and these are either to be mounted in Canada balsam, or in weak spirit or glycerine, according to the degree in which the horny opacity of their integument requires the assistance of the former to facilitate the transmission of light through it, or the softness and delicacy of their textures renders a preservative liquid more desirable. Thus an ordinary *Flea* or *Bug* will best be mounted in the former medium; but the various parasites of the *Louse* kind, with some or other of which almost every kind of animal is affected, should be set up in the latter. Some of the aquatic larvæ of the *Diptera* and *Neuroptera*, which are so transparent that their whole internal organization can be made out without dissection, are very beautiful and interesting objects, when examined in the living state, especially because they allow the circulation of the blood and the action of the dorsal vessel to be discerned (§ 389). Among these, there is none preferable to the larva of the *Ephémère marginata* (day fly), which is distinguished by the possession of a number of beautiful appendages on its body and

tail, and is, moreover, an extremely common inhabitant of our ponds and streams. This insect passes two or even three years in its larva state, and during this time it repeatedly throws off its skin; the cast-skin, when perfect, is an object of extreme beauty, since, as it formed a complete sheath to the various appendages of the body and tail, it continues to exhibit their outlines with the utmost delicacy; and by keeping these larvæ in a Vivarium, and by mounting the entire series of their cast skins, a record is preserved of the successive changes they undergo. Much care is necessary, however, to extend them upon their slides, in consequence of their extreme fragility; and the best plan is to place the slip of glass under the skin whilst it is floating on water, and to lift the object out upon the slide.

378. *Structure of the Integument.*—In treating of the separate parts of the organization of Insects, which furnish the most interesting objects of Microscopic study, we may most appropriately commence with their integument and its appendages (scales, hairs, &c.) The body and members are closely invested by a hardened skin, which acts as their skeleton, and affords points of attachment to the muscles by which their several parts are moved; being soft and flexible, however, at the joints. The skin is usually more or less horny in its texture, and is consolidated by the animal substance termed *chitine*, as well as, in some cases, by a small quantity of mineral matter. It is in the Coleoptera that it attains its greatest development; the dermo-skeleton of many beetles being so firm, as not only to confer upon them an extraordinary power of passive resistance, but also to enable them to put forth enormous force, by the action of the powerful muscles which are attached to it. It may be stated as a general rule, that the external layer of this dermo-skeleton is always cellular, taking the place of an epidermis; and that the cells are straight-sided and closely fitted together, so as to be polygonal (usually hexagonal) in form. Of this we have a very good example in the *superficial* layers (Fig. 286, B) of the thin horny lamellæ or blades, which constitute the terminal portion of the antenna of the *Cockchaffer* (Fig. 285); this layer being easily distinguished from the intermediate portion of the lamina (A), by careful focussing. In many beetles, the hexagonal areolation of the surface is often distinguishable when the light is reflected from it at a particular angle, even when not discernible in transparent sections. The integument of the common *Red Ant* exhibits the hexagonal cellular arrangement very distinctly throughout; and the broad flat expansion on the leg of the *Crabro* (sand-wasp) affords another beautiful example of a distinctly cellular structure in the outer layer of the integument. The inner layer, however, which constitutes the principal part of the thickness of the horny casing of the Beetle tribe, seldom exhibits any distinct organization; though it may be usually separated into several laminæ, which are sometimes traversed by tubes that pass into

them from the inner surface, and extend towards the outer without reaching it. Occasionally, however, even this exhibits very clear indications of cellular structure; of which a good example is afforded by the *middle* layer of the lamellæ of the antenna of the *Cockchaffer* (Fig. 286, A), wherein is plainly to be seen an assemblage of rounded cells with large nuclei, lying in the midst of a homogeneous intercellular substance, and thus closely resembling Cartilage (Fig. 324) in structure, though differing from it in chemical composition.

379. *Tegumentary Appendages*.—The surface of many *Insects* is beset, and is sometimes completely covered with *appendages*, having sometimes the form of broad flat scales, sometimes that of hairs more or less approaching the cylindrical shape, and sometimes being intermediate between the two. The *scaly* investment is most complete among the *Lepidoptera* (butterfly and moth tribe); the distinguishing character of the insects of this order being derived from the presence of a regular layer of scales, upon each side of their large membranous wings. It is to the peculiar coloration of the scales, that the various hues and figures are due, by which these wings are so commonly distinguished; all the scales of one patch (for example) being green, those of another red, and so on; for the subjacent membrane remains perfectly transparent and colorless, when the scales have been brushed off from its surface. Each scale seems to be composed of two superficial colored laminæ, enclosing a central lamina of structureless membrane, the surface of which is highly polished, and which acts as a "foil" to increase their brilliancy by reflecting back the light that passes through them,—an arrangement which may often be discerned in scales that have lost a portion of their superficial layer by some accidental injury (Fig. 281, c). The color of the superficial laminæ seems to be generally inherent in their substance, especially in the *Lepidoptera*; but it sometimes appears to be (like the prismatic hues of a soap-bubble) a purely optical effect of their extreme thinness, this being especially the case among those beetles, as the *Curculio imperialis* (diamond beetle), the scales of which have a metallic lustre, and exhibit colors that vary with the mode in which the light glances from them. Each scale is furnished with a sort of handle at one end (Figs. 279–281), by which it is fitted into a minute socket attached to the surface of the insect; and on the wings of *Lepidoptera* these sockets are so arranged, that the scales lie in very regular rows, each row overlapping a portion of the next, so as to give to their surface, when sufficiently magnified, very much the appearance of being *tiled* like the roof of a house. Such an arrangement is said to be "imbricated." The forms of these scales are often very curious, and frequently differ a good deal on the several parts of the wings and of the body of the same individual; being usually more expanded on the former, and narrower and more hair-like on the latter. The


peculiar markings which many of these scales exhibit, very early attracted the attention of those engaged in the improvement of the Microscope by the application of the principle of achromatic correction (p. 41); since these markings are entirely invisible, however great may be the magnifying power employed, under microscopes of the older construction, owing to the necessary limitation of their angular aperture; whilst, as they are brought into view with a clearness and strength that are proportionate to the extension of the angular aperture and the perfection with which the aberrations are corrected, they serve as "test objects" of the goodness of an achromatic combination. At first, the scale of the *Podura* (Fig. 281) was the most difficult test known for the *highest* powers; and a microscope which could only exhibit an alternation of dark and light bands or striæ upon its surface, was considered a good one. But even the complete "resolution" of these striæ into their component markings, is now considered as but a very ordinary "test" for the *medium* powers of the Microscope; and tests of much greater difficulty, and therefore more suitable for the higher, are afforded (as we have seen, § 102, III) by the valves of the Diatomaceæ. Still, the test scales of Insects have their use, in enabling us to appreciate the performance of achromatics of *medium* power (§ 102, III); and it will therefore be advantageous here to

FIG. 279.



Scale of *Morpho Menelaus*.

FIG. 280.




Battledoor Scale of *Polyommatus argus*
(azure blue).

notice a few of those which are most commonly employed for this purpose.

380. Among the most beautiful of all these scales, both for color and for regularity of marking, are those of the butterfly termed *Morpho Menelaus* (Fig. 279). These are of a rich blue

tint, and exhibit strong longitudinal striæ, which seem due to ribbed elevations of the superficial colored layer. There is also an appearance of transverse striation, which cannot be seen at all with an inferior objective, becomes very decided with a good objective of medium focus, but is found, when submitted to the test of a high power and achromatic condenser, to depend upon a sort of beaded subdivision of the longitudinal ribs,—the transverse striæ that may be seen *between* these ribs, being apparently produced by the beading of the ribs on the other surface of the scale. The large scales of the *Polyommatus argus* (azure-blue butterfly) resemble those of the Menelaus in form and structure, but are more delicately marked. The same insect, however, furnishes small scales, which are commonly termed the “battledoor” scales, the resemblance which their form presents to that instrument being usually much greater than in the specimen represented in Fig. 280; these scales, also, are marked by narrow longitudinal ribbings, which at intervals expand into rounded or oval elevations, that give to the scale a dotted appearance; at the lower part of the scale, however, these dots are wanting; and in the interval between the two portions, we observe a sort of crescent, formed of minute pigment-granules, crossing the scale transversely. The scales of the *Pontia brassica* (cabbage butterfly) and of the *Hipparchia janira* (meadow-brown butterfly), have longitudinal markings of a somewhat similar nature, but less sharply defined; these are further noticeable for the brush-like appendage which each scale bears at the end furthest from its implantation. The *Podura plumbea* or “spring-tail” is a little wingless insect that is found amidst the sawdust of many wine-cellars, especially such as are damp, leaping about like a flea, by means of that peculiar power of using its tail, from which its name is derived. Its scales are of different sizes and of different degrees of strength of marking (Fig. 281, A, B), and are therefore by no means of uniform value as tests. The general appearance of their surface, under a power not sufficient to resolve their marking, is that of watered silk, light and dark bands passing across with wavy irregularity; but a well-corrected lens

FIG. 281.



Scales of *Podura plumbea*:—A. large strongly marked scale; B. small scale, more faintly marked; C. portion of an injured scale, showing the nature of the markings.

of very moderate angular aperture, now suffices to resolve every dark band into a row of short lines, each of these being thick at

of very moderate angular aperture, now suffices to resolve every dark band into a row of short lines, each of these being thick at

one end and coming to a point at the other, so that the impression conveyed is that of a set of spines projecting obliquely from the flat surface of the scale, like the teeth of a "hackle." A more careful examination of scales, however, of which the superficial layers have been partly removed (c), serves to show that these dark lines are but the spaces between the minute wedge-like particles, arranged side by side, and end to end, of which those layers are made up; so that the structure of the scale does not in reality differ essentially from the ordinary type.¹ Although scarcely useful as a "test-object," since its structure is too easily resolved, the scale of the *Lepisma saccharina* or "sugar-louse" deserves notice; the longitudinal ribbings being so strongly marked and so regular, as to give them an appearance resembling that of many bivalve shells. The long narrow scale of the common *Gnat*, also, exhibits a few very prominent ribbings; and, from its small size, it serves as a good test-object for the medium powers.

381. The *Hairs* of many Insects, and still more of their larvæ, are very interesting objects for the microscope, on account of their branched or tufted conformation; this being particularly remarkable in those with which the common hairy Caterpillars are so abundantly beset. Some of these afford very good tests for the perfect correction of objectives. Thus, the hair of the *Bee* is pretty sure to exhibit strong prismatic colors, if the chromatic aberration should not have been exactly neutralized; and that of the larva of the *Dermestes*, or "bacon-beetle," was once thought a very good test of defining power, and is still useful for this purpose. It has a cylindrical shaft (Fig. 282, B) with closely set whorls of spiny protuberances, four or five in each whorl; the highest of these whorls is composed of more knobby spines; and the hair is surmounted by a curious circle of six or seven large filaments, attached by their pointed ends to its shaft, whilst at their free extremities they dilate into knobs. An approach to this structure is seen in the hairs of certain *Myriapods* (centipedes, gally-worms, &c.), of which an example is shown in Fig. 282, A.

382. In examining the integument of Insects, and its appendages, parts of the surface may be viewed either by reflected or transmitted light, according to their degree of transparency and the nature of their covering. The Beetle and Butterfly tribes furnish the greater number of objects suitable to be viewed in

¹ *Podura* may be obtained by sprinkling a little oatmeal on a piece of black paper near their haunts; and after leaving it there for a few hours, removing it carefully to a large glazed basin, so that, when they leap from the paper (as they will when brought to the light) they may fall into the basin, and may thus separate themselves from the meal. The best way of obtaining their scales, is to confine several of them together beneath a wine-glass inverted upon a piece of fine smooth paper; for the scales will be detached by their leaps against the glass, and will fall upon the paper; and if they be left thus confined for some time, they will be very likely, by treading upon some of the scales, to bring them into the condition represented at c, Fig. 281, which best illustrates their true nature.

the former of these modes; and nothing is easier than to mount portions of the *elytra* of the former (which are usually the

FIG. 282.



A, Hair of *Myriapod*.
B, Hair of *Dermestes*.

most showy portions of their bodies), or of the wings of the latter, in the manner described in § 123. The tribe of *Curculionidæ*, in which the surface of the body is beset with scales having the most varied and lustrous hues, is distinguished among all other Coleoptera for the brilliancy of the objects it affords; the most remarkable in this respect being the well-known *Curculio imperialis*, or "diamond beetle" of South America, parts of whose *elytra*, when properly illuminated and looked at with a low power, show like clusters of jewels flashing against a dark velvet ground. In many of the British *Curculionidæ*, which are smaller and far less brilliant, the scales lie at the bottom of little depressions of the surface; and if the *elytra* of the "diamond beetle" be carefully examined, it will be found that each of the clusters of scales which are arranged upon it in rows, seems to rise out of a deep pit which sinks in by its side. The transition from scales to hairs is extremely well seen, by comparing the different parts of the surface of the "diamond beetle" with each other.

The beauty and brilliancy of many objects of this kind are increased by mounting them in cells in Canada balsam, even though they are to be viewed with reflected light; other objects, however, are rendered less attractive by this treatment; and in order to ascertain whether it is likely to improve or to deteriorate the specimen, it is a good plan first to test some other portion of the body, having scales of the same kind, by touching it with turpentine, and then to mount the part selected as an object, either in balsam, or dry, according as the turpentine increases or diminishes the brilliancy of the scales on the spot to which it was applied. Portions of the wings of *Lepidoptera* are best mounted as opaque objects, without any other preparation than gumming them flat down to the card-board surface of the slide (§ 123); care being taken to avoid disturbing the arrangement of the scales, and to keep the objects, when mounted, as secluded as possible from dust. In selecting such portions, it is well to choose those which have the brightest and the most contrasted colors, foreign butterflies being in this respect usually preferable to British; and before attaching them to their slides, care should be taken to ascertain in what position, with the arrangement of light ordinarily used, they are seen to the best advantage, and to fix them there accordingly. Whenever portions of the integument of Insects are to be viewed as transparent


objects, for the display of their intimate structure, they should be mounted in Canada balsam, after soaking for some time in turpentine; since this substance has a peculiar effect in increasing the translucency. Not only the horny casings of perfect Insects of various orders, but also those of their pupæ, are worthy of this kind of study; and objects of great beauty (such as the chrysalis case of the Emperor-moth), as well as of scientific interest, are sure to reward any who may prosecute it with any assiduity. Further information may often be gained, by softening such parts in potash, and viewing them in fluid. The scales of the wings of Lepidoptera, &c., are best transferred to the slide, by simply pressing a portion of the wing either upon the slip of glass or upon the cover; if none should adhere, the glass may first be gently breathed on. Some of them are best seen when examined "dry," whilst others are more clear when mounted in fluid; and for the determination of their exact structure, it is well to have recourse to both these methods. If these scales are to be used as "test-objects," it is preferable to place them between two pieces of thin glass, in the manner specified in § 122. Hairs, on the other hand, are best mounted in balsam.

383. *Parts of the Head.*—The *Eyes* of Insects, situated upon the upper and outer part of the head, are usually very conspicuous organs, and are frequently so large as to touch each other in front (Fig. 283). We find in their structure a remarkable example of that multiplication of similar parts, which seems to be the predominating idea in the conformation of Articulated animals; for each of the large protuberant bodies which we designate as an *eye*, is really an aggregate of many hundred, or even many thousand minute eyes, which are designated *ocelli*.

Approaches to this structure are seen in the Annelida and Entomostraca; but the number of "ocelli" thus grouped together is usually small. In the higher Crustacea, however, the ocelli are very numerous; their compound eyes being constructed upon the same general plan as those of Insects, although their shape and position are often very peculiar (Fig. 348). The individual ocelli are at once recognized, when the composite eyes are examined under even a low magnifying power, by the "faceted" appearance of the surface (Fig. 283), which is marked

out by very regular divisions either into hexagons or into squares: each facet is the cornea of a separate ocellus, and


FIG. 283.



Head and Compound Eyes of the *Bee*, showing the ocelli *in situ* on one side (A), and displaced on the other (B); a, a, a, stemmata; b, b, antennæ.

has a convexity of its own; hence by counting the facets, we can ascertain the number of ocelli in each composite eye. In the two eyes of the common *Fly*, there are as many as 4000; in those of the *Cabbage Butterfly*, there are about 17,000; in the *Dragon fly*, 24,000; and in the *Mordella beetle*, 25,000. Behind each "corneule" is a layer of dark pigment, which takes the place, and serves the purpose, of the "iris" in the eyes of Vertebrate animals; and this is perforated by a central aperture or "pupil," through which the rays of light that have traversed the cornea gain access to the interior of the eye. The further structure of these bodies is best examined by vertical sections; and these show that the shape of each ocellus is conical, or rather pyramidal (Fig. 284), the cornea forming its base (*a*), whilst its apex abuts upon a bulbous expansion of the optic

FIG. 284.



A, Section of the eye of *Melolontha vulgaris* (Cockchafer):—*a*, a portion more highly magnified:—*a*, facets of the cornea; *b*, transparent pyramids surrounded with pigment; *c*, fibres of the optic nerve; *d*, trunk of the optic nerve.

nerve. Each "corneule" acts as a distinct lens; as may be shown by detaching the entire assemblage by maceration, and then drying it (flattened out) upon a slip of glass; for when this is placed under the microscope, if the point of a knife, scissors, or any similar object, be interposed between the mirror and the stage, the image of this point will be seen, by a proper adjustment of the focus of the microscope, in every one of the lenses. The focus of each "corneule" has been ascertained by experiment to be equivalent to the length of the pyramid behind it; so that the image which it produces will fall upon the extremity of the filament of the optic nerve which passes to its point. This pyramid consists of a transparent substance (*B, b*), which may be considered as representing the "vitreous humor;" and the pyramids are separated from each other by a layer of dark pigment, which completely encloses them, save at the pupillary apertures which admit the rays that have passed through the "corneules," and at their smaller ends, where the pigment is perforated by a set of apertures that give passage to the fibres of

the optic nerve (*c*), of which one proceeds to each ocellus. Thus the rays which have passed through the several "corneules" are prevented from mixing with each other; and no rays save those which pass in the axis of the pyramids, can reach the fibres of the optic nerve. Hence it is evident that, as no two ocelli on the same side have exactly the same axis, no two can receive their rays from the same point of an object; and thus, as each composite eye is immovably fixed upon the head, the combined action of the entire aggregate will probably only afford but a single image, resembling that which we obtain by means of our single eyes. Although the foregoing may be considered as the typical structure of the eyes of Insects, yet there are various departures from it (most of them slight) in the different members of the class. Thus in some cases the posterior surface of each "corneule" is concave; and a space is left between it and the iris-like diaphragm, which seems to be occupied by a watery fluid or "aqueous humor;" in other instances, again, this space is occupied by a double convex body, which seems to represent the "crystalline lens;" and this body is sometimes found behind the iris, the number of ocelli being reduced, and each one being larger, so that the cluster presents more resemblance to that of Spiders, &c. Besides their composite eyes, Insects usually possess a small number of rudimentary single eyes, resembling those of the Arachnida; these are seated upon the top of the head, and are termed *stemmata* (Fig. 283, *a, a, a*). It is remarkable that the *larvæ* of Insects which undergo a complete metamorphosis, only possess single eyes; the composite eyes being developed, at the same time with the wings and other parts which are characteristic of the Imago state, during the latter part of Pupal life.

384. Various modes of preparing and mounting the Eyes of Insects may be adopted, according to the manner wherein they are to be viewed. For the observation of their external faceted surface by reflected light, it is better to lay down the entire head, so as to present a front-face or a side-face, according to the position of the eyes; the former giving a view of *both* eyes, when they approach each other so as nearly or quite to meet (as in Fig. 283); whilst the latter will best display *one*, when the eyes are situated more at the sides of the head. For the minuter examination of the "corneules," however, these must be separated from the hemispheroidal mass whose exterior they form, by prolonged maceration; and the pigment must be carefully washed away by means of a fine camel-hair brush, from their inner or posterior surface. In flattening them out upon the glass slide, one of two things must necessarily happen; either the margin must tear when the central portion is pressed down to a level; or, the margin remaining entire, the central portion must be thrown into plaits, so that its corneules overlap one another. As the latter condition interferes with the examination of the

structure much more than the former does, it should be avoided by making a number of slits in the margin of the convex membrane before it is flattened out. Such preparations may be mounted either in liquid, or in Canada balsam; the latter being preferable when (as sometimes happens) the membrane is so horny as to be but imperfectly transparent. Vertical sections, adapted to demonstrate the structure of the ocelli and their relations to the optic nerve, can of course be only made when the body of the insect is fresh; and these should be mounted in fluid. The following are some of the Insects, whose eyes are best adapted for Microscopic preparations:—*Coleoptera*, *Cicindela*, *Dytiscus*, *Melolontha* (cockchafer), *Lucanus* (stag-beetle); *Orthoptera*, *Acheta* (house and field crickets), *Locusta*; *Hemiptera*, *Notonecta* (boat-fly); *Neuroptera*, *Libellula* (dragon-fly), *Agrion*; *Hymenoptera*, *Vespidæ* (wasps) and *Apidæ* (bees) of all kinds; *Lepidoptera*, *Vanessa* (various species of butterflies), *Sphinx ligustri* (privet hawk-moth), *Bombyx* (silkworm-moth, and its allies); *Diptera*, *Tabanus* (gad-fly), *Asilus*, *Eristalis* (drone-fly), *Tipula* (crane-fly), *Musca* (house-fly), and many others.

385. The *Antennæ*, which are the two jointed appendages arising from the upper part of the head of Insects (Fig. 283, *b, b*), present a most wonderful variety of conformation in the several tribes of Insects; often differing considerably in the several species of one genus, and even in the two sexes of the same species. Hence the characters which they afford are extremely useful in classification; especially since their structure must almost necessarily be in some way related to the habits and general economy of the creatures to which they belong (although our imperfect acquaintance with their function prevents us from clearly discerning this relation), so that their resemblances and differences will generally be found to coincide with those resemblances and differences in general conformation, on which every "natural" arrangement must be founded. Thus, in the Coleopterous order, we find one large family, including the glow-worm, fire-fly, skip-jack, &c., distinguished by the toothed or serrated form of the antennæ, and hence called *Serricornes*; in another, of which the "burying-beetle" is the type, the antennæ are terminated by a club-shaped enlargement, so that these beetles are termed *Clavicornes*; in another, again, of which the *Hydrophilus* or "large water-beetle" is an example, the antennæ are never longer and are commonly shorter than one of the pairs of palpi, whence the name of *Palpicornes* is given to this group; in the very large family that includes the *Lucani* or "stag-beetles," with the *Scarabæi* of which the "cockchafer" is the commonest example, the antennæ terminate in a set of leaf-like appendages, which are sometimes arranged like a fan or the leaves of an open book (Fig. 285), are sometimes parallel to each other like the teeth of a comb, and sometimes fold one over the other, thence giving the name of *Lamellicornes*; whilst another large family is distinguished by the appellation *Longi-*


cornes, from the great length of the antennæ, which are at least as long as the body, and often longer. Among the Lepidoptera, again, the conformation of the antennæ frequently enables us at once to distinguish the group to which any specimen belongs. As every treatise on Entomology contains figures and descriptions of the principal types of conformation of these organs, there is no occasion here to dwell upon them longer than to specify such as are most interesting to the Microscopist; *Coleoptera*, *Brachinus*, *Calathus*, *Harpalus*, *Dytiscus*, *Staphylinus*, *Philonthus*, *Elater*, *Lampyrus*, *Silpha*, *Hydrophilus*, *Aphodius*, *Melolontha*, *Cetonia*, *Curculio*; *Orthoptera*, *Forficula* (carwig), *Blatta* (cockroach); *Lepidoptera*, *Sphinxes* (hawk-moths) and *Nocturna* (moths) of various kinds, the large "plumed" antennæ of the latter being peculiarly beautiful objects under a low magnifying power; *Diptera*, *Culicidæ* (gnats of various kinds), *Tipulidæ* (crane-flies and midges), *Tabanus*, *Eristalis*, and *Muscidæ* (flies of various kinds). All the larger antennæ should be put up in balsam, after being soaked for some time in turpentine; but the small feathery antennæ of gnats and midges are so liable to distortion when thus mounted, that it is better to set them up in fluid, the head with its pair of antennæ being thus preserved together when not too large.

386. The next point in the organization of Insects, to which the attention of the Microscopist may be directed, is the structure of the *Mouth*. Here, again, we find almost infinite varieties in the details of conformation; but these may be for the most part reduced to a small number of types or plans, which are characteristic of the different orders of Insects. It is among the *Coleoptera*, or beetles, that we find the several parts of which the mouth is composed, in their most distinct form; for although

FIG. 285.


Antenna of *Melolontha* (Cockchafer).

FIG. 286.

Minute structure of Leaf-like expansions of Antenna of *Melolontha*:—A, their internal layer; B, their superficial layer.

some of these parts are much more highly developed in other insects, other parts may be so much altered or so little developed, as to be scarcely recognizable. The Coleoptera present the typical conformation of the *mandibulate* mouth, which is adapted for the prehension and division of solid substances; and this consists of the following parts:—1. A pair of jaws, termed *mandibles*, frequently furnished with powerful teeth, opening laterally on either side of the mouth, and serving as the chief instruments of manducation; 2. A second pair of jaws, termed *maxillæ*, smaller and weaker than the preceding, beneath which they are placed, and serving to hold the food, and to convey it to the back part of the mouth; 3, an upper lip, or *labrum*; 4, a lower lip, or *labium*; 5, one or two pairs of small jointed appendages termed palpi, attached to the maxillæ, and hence called *maxillary palpi*; 6, a pair of *labial palpi*. The labium is often composed of several distinct parts; its basal portion being distinguished as the *mentum* or chin, and its anterior portion being sometimes considerably prolonged forwards, so as to form an organ which is properly designated the *ligula*, but which is more commonly known as the “tongue,” though not really entitled to

FIG. 287.



Tongue of common Fly:—*a*, lobes of ligula; *b*, portion enclosing the lancets formed by the metamorphosis of the maxilla; *c*, maxillary palpi:—*A*, portion of one of the metamorphosed tracheæ enlarged.


that designation, the real *tongue* being a soft and projecting organ that forms the floor of the mouth, and being only found as a distinct part in a comparatively small number of insects, as

the Cricket. The ligula is extremely developed in the *Fly* kind, in which it forms the chief part of what is commonly called the "proboscis" (Fig. 287); and it also forms the "tongue" of the *Bee* and its allies (Fig. 288). In the *Diptera* or two-winged flies generally, the labrum, maxillæ, mandibles, and the internal tongue (where it exists) are converted into delicate lancet-shaped organs termed *setæ*, which, when closed together, are received into a hollow on the upper side of the labium (Fig. 287, *b*), but which are capable of being used to make punctures in the skin of animals or the epidermis of plants, whence the juices may be drawn forth by the proboscis. Frequently, however, two or more of these organs may be wanting, so that their number is reduced from six to four, three, or two. In the *Hymenoptera* (bee and wasp tribe), however, the labrum and the mandibles (Fig. 288, *b*), much resemble those of mandibulate insects, and are used for corresponding

purposes; the maxillæ (*c*) are greatly elongated, and form, when closed, a tubular sheath for the *ligula* or "tongue," through which the honey is drawn up; the labial palpi (*d*) also are greatly developed, and fold together like the maxillæ, so as to form an inner sheath for the "tongue;" while the "ligula" itself (*e*) is a long tapering muscular organ, marked by an immense number of short annular divisions, and densely covered over its whole length with long hairs (*b*). It is not tubular, as some have stated, but is solid; when actively employed in taking food, it is extended to a great distance beyond the other parts of the mouth; but when at rest, it is closely packed up and concealed between the maxillæ. "The manner," says Mr. Newport, "in which the honey is obtained when the organ is plunged into it at the bottom of a flower, is by 'lapping,' or a constant succession of short and quick extensions and contractions of the organ, which occasion the fluid to accumulate upon it and to ascend along its upper surface, until it reaches the orifice of the tube formed by the approximation of the maxillæ above, and of the labial palpi and this part of the ligula below."

387. By the plan of conformation just described, we are led to that which prevails among the *Lepidoptera* or butterfly tribe,


FIG. 288.



A. Parts of the mouth of *Apis mellifica* (Honey-bee):—*a*, mentum; *b*, mandibles; *c*, maxilla; *d*, labial palpi; *e*, ligula, or prolonged labium. commonly termed the tongue:—*b*, portion of the surface of the ligula, more highly magnified.

and which, being pre-eminently adapted for suction, is termed the *haustellate* mouth. In these insects, the labrum and mandibles are reduced to three minute triangular plates; whilst the maxillæ are immensely elongated, and are united together along the median line, to form the *haustellium* or proboscis, which contains a tube formed by the junction of the two grooves that are channelled out along their mutually applied surfaces, and which serves to pump up the juices of deep cup-shaped flowers, into which the size of their wings prevents these insects from entering. The length of this haustellium varies greatly; thus in such Lepidoptera as take no food in their perfect state, it is a very insignificant organ; in some of the white Hawk-moths, which hover over blossoms without alighting, it is nearly two inches in length; and in most Butterflies and Moths it is about as long as the body itself. This haustellium, which, when not in use, is coiled up in a spiral beneath the mouth, is an extremely beautiful microscopic object, owing to the peculiar banded arrangement it exhibits (Fig. 289), which is probably due to the disposition of

FIG. 289.

Haustellium (proboscis) of *Vanessa*.

its muscles. In many instances, the two halves may be seen to be locked together by a set of hooked teeth, which are inserted into little depressions between the teeth of the opposite side. Each half, moreover, may be ascertained to contain a trachea or air-tube (§ 391); and it is probable, from the observations of Mr. Newport,¹ that the sucking up of the juices of a flower through the haustellium

(which is accomplished with great rapidity) is effected by the agency of the respiratory apparatus. The proboscis of many Butterflies is furnished, for some distance from its extremity, with a double row of small projecting barrel-shaped bodies (shown in Fig. 289), which are surmised by Mr. Newport to be organs of taste. Numerous other modifications of the structure of the mouth, existing in the different tribes of Insects, are well worthy of the careful study of the Microscopist; but as detailed descriptions of most of these will be found in every systematic treatise on Entomology, the foregoing general account of the principal types must suffice.

388. *Parts of the Body.*—The conformation of the several divisions of the *Alimentary Canal* presents such a multitude of

¹ "Cyclopædia of Anatomy and Physiology," vol. ii, p. 902.

diversities, not only in different tribes of Insects, but in different states of the same individual, that it would be utterly vain to attempt here to give even a general idea of it; more especially as it is a subject of far less interest to the ordinary Microscopist, than it is to the professed Anatomist. Hence we shall only stop to mention, that the muscular gizzard in which the œsophagus very commonly terminates, is often lined by several rows of strong horny teeth for the reduction of the food, which furnish very beautiful microscopic objects. These are particularly developed among the Grasshoppers, Crickets, and Locusts, the nature of whose food causes them to require powerful instruments for its reduction.

389. The *Circulation of Blood* may be distinctly watched in many of the more transparent *Larvæ*, and may sometimes be observed in the perfect Insect. It is kept up, not by an ordinary heart, but by a "dorsal vessel," which really consists of a succession of muscular hearts or contractile cavities, one for each segment, opening one into another from behind forwards, so as to form a continuous trunk, divided by valvular partitions. In many larvæ, however, these partitions are very indistinct, and the walls of the "dorsal vessel" (so named from the position it always occupies along the middle of the back) are so thin and transparent that it can with difficulty be made out, a limitation of the light by the diaphragm being often necessary. The blood which moves through this trunk, and which is distributed by it to the body, is a transparent and nearly colorless fluid, carrying with it a number of "oat-shaped" corpuscles, by the motion of which its flow can be followed. The current enters the dorsal vessel at its posterior extremity, and is propelled by the contractions of the successive chambers towards the head, being prevented from moving in the opposite direction by the valves between the chambers, which only open forwards. Arrived at the anterior extremity of the dorsal vessel, the blood is distributed into three principal channels; a central one, namely, passing to the head, and a lateral one to either side, descending so as to approach the lower surface of the body. It is from the two lateral currents that the secondary streams diverge, which pass into the legs and wings, and then return back to the main stream; and it is from these also, that, in the larva of the *Ephemera marginata* (day-fly), the extreme transparency of which renders it one of the best of all subjects for the observation of Insect circulation, the smaller currents diverge into the gill-like appendages with which the body is furnished (§ 393). The blood-currents seem rather to pass through channels excavated among the tissues, than through vessels with distinct walls; but it is not improbable that in the perfect Insect the case may be different. In many aquatic larvæ, especially those of the *Culicidæ* (gnat tribe), the body is almost entirely occupied by the visceral cavity; and the blood may be seen to move backwards


in the space that surrounds the alimentary canal, which here serves the purpose of the channels usually excavated through the solid tissues, and which freely communicates at each end with the dorsal vessel. This condition strongly resembles that found in many Annelida. In some larvæ, whose development is yet less advanced, even the dorsal vessel appears to be wanting, although the fluid of the visceral cavity (in which corpuscles abound) is in a state of continual oscillatory movement.

390. The Circulation may be easily seen in the wings of many insects in their *Pupa* state, especially in those of the Neuropterous order (such as dragon-flies and day-flies) which pass this part of their lives in water, in a condition of activity; the pupæ of *Agrion puella*, one of the smaller dragon-flies, is a particularly favorable subject for such observations. Each of the "nerves" of the wings contains a "trachea" or air-tube (§ 391), which branches off from the tracheal system of the body; and it is in a space around the trachea that the blood may be seen to move, when the hard framework of the nerve itself is not too opaque. The same may be seen, however, in the wings of the pupæ of Bees, Butterflies, &c., which remain shut up motionless in their cases; for this condition of apparent torpor is one of great activity of their nutritive system, those organs, especially, which are peculiar to the perfect insect, being then in a state of rapid growth, and having a vigorous circulation of blood through them. In certain Insects of nearly every order, a movement of fluid has been seen in the wings for some little time after their last metamorphosis; but this movement soon ceases, and the wings dry up. The common *Fly* is as good a subject for this observation, as can be easily found; it must be caught within a few hours or days of its first appearance; and the circulation may be most conveniently brought into view, by enclosing it (without water) in the animalcule cage, and pressing down the cover sufficiently to keep the body at rest, without doing it any injury.

391. The *Respiratory Apparatus* of Insects affords a very interesting series of microscopic objects; for, with great uniformity in its general plan, there is almost infinite variety in its details. The aeration of the blood in this class is provided for, not by the transmission of the fluid to any special organ representing the *lung* of a Vertebrated animal (§ 438) or the *gill* of a Mollusk (§ 353), but by the introduction of air into every part of the body, through a system of minutely distributed *tracheæ* or air-tubes, which penetrate even the smallest and most delicate organs. Thus, as we have seen, they pass into the *haustellium* or "proboscis" of the Butterfly (§ 386), and they are minutely distributed in the elongated *labium* or "tongue" of the fly (Fig. 287). Their general distribution is shown in Fig. 290; where we see two long trunks (*f*) passing from one end of the body to the other, and connected with each other by a transverse canal in every segment; these trunks communicate, on the one hand, by

short wide passages, with the "stigmata," "spiracles," or breathing-pores (*g*), through which the air enters and is discharged; whilst they give off branches to the different segments, which divide again and again into ramifications of extreme minuteness. They usually communicate also with a pair of air-sacs (*h*) which are situated in the thorax; but the size of these (which are only found in the perfect insect, no trace of them existing in the larvæ) varies greatly in different tribes, being usually greatest in those insects which (like the bee) can sustain the longest and most powerful flight, and least in such as habitually live upon the ground or upon the surface of the water. The structure of the air-tubes reminds us of that of the "spiral vessels" of Plants, which seem destined (in part at least) to perform a similar office (§ 232); for within the membrane that forms their outer wall, an elastic fibre winds round and round, so as to form a spiral, closely resembling in its position and functions the spiral wire-spring of flexible gas-pipes; within this again, however, there is another membranous wall to the air-tubes, so that the spire winds between their inner and outer coats. The tongue of the Fly presents a curious modification of this structure, the purpose of which is not apparent; for instead of its tracheæ being kept pervious after the usual fashion, by the

FIG. 290.



Tracheal system of *Nepa* (Water-scorpion):—*a*, head; *b*, first pair of legs; *c*, first segment of the thorax; *d*, second pair of wings; *e*, second pair of legs; *f*, tracheal trunk; *g*, one of the stigmata; *h*, air-sac.

FIG. 291.



Portion of a large Trachea of *Dytiscus*, with some of its principal branches.

winding of a continuous spiral fibre through their interior, the fibre is broken into rings, and these rings do not surround the whole tube, but are terminated by a set of arches that pass from one to another (Fig. 287, A). When a portion of one of the great trunks with some of the principal branches of the tracheal system has been dissected out, and so pressed in mounting that the sides of the tubes are flattened against each other (as has happened in the specimen represented in Fig. 291), the spire forms two layers which are brought into close apposition; and a very beautiful appearance, resembling that of "watered silk," is produced by the crossing of the two sets of fibres, of which one overlies the other. That this appearance, however, is altogether an optical illusion, may be easily demonstrated by carefully following the course of any one of the fibres, which will be found to be perfectly regular.

392. The "stigmata" or "spiracles" through which the air enters the tracheal system, are generally visible on the exterior of the body of the Insect (especially on the abdominal segments) as a series of pores along each margin of the under surface. In most larvæ, nearly every segment is provided with a pair; but in the perfect insect, several of them remain closed, especially in the thoracic region, so that their number is often considerably reduced. The structure of the spiracles varies greatly in regard to complexity in different Insects; and even where the general plan is the same, the details of conformation are peculiar, so that perhaps in scarcely any two species are they alike. Generally speaking, they are furnished with some kind of sieve at their entrance, by which particles of dust, soot, &c., which would otherwise enter the air-passages, are filtered out; and this sieve may be formed by the interlacement of the branches of minute arborescent growths from the borders of the spiracle, as in the common *Fly* (Fig. 292), or in the *Dytiscus*; or it may be a membrane perforated with minute holes, and supported upon a frame-

FIG. 292.


Spiracle of common *Fly*.

FIG. 293.

Spiracle of Larva of *Cockchafer*.

work of bars that is prolonged in like manner from the thickened margin of the aperture (Fig. 293), as in the larva of the *Melo-*

lontha (cockchafer). Not unfrequently, the centre of the aperture is occupied by an impervious disk, from which radii proceed to its margin, as is well seen in the spiracle of *Tipula* (crane-fly). In those aquatic larvæ which breathe air, we often find one of the spiracles of the last segment of the abdomen prolonged into a tube, the mouth of which remains at the surface, while the body is immersed; the larvæ of the *Gnat* tribe may frequently be observed in this position.

393. There are many aquatic Larvæ, however, which have an entirely different provision for respiration; being furnished with external leaf-like or brush-like appendages, into which the tracheæ are prolonged, so that, by absorbing air from the water that bathes them, they may convey this into the interior of the body. We cannot have a better example of this than is afforded by the larva of the common *Ephemera* (day-fly), the body of which is furnished with a set of branchial appendages resembling the "fin-feet" of Branchiopods (§ 368), whilst the three-pronged tail also is fringed with clusters of delicate hairs which appear to minister to the same function. In the larva of the *Libellula* (dragon-fly), the extension of the surface for aquatic respiration takes place within the termination of the intestine; the lining membrane of which is folded into an immense number of plaits, each containing a minutely ramified system of tracheæ; the water, slowly drawn in through the anus for bathing this surface, is ejected with such violence that the body is impelled in the opposite direction; and the air taken up by its tracheæ is carried, through the system of air-tubes of which they form a part, into the remotest organs. This apparatus is a peculiarly interesting object for the Microscope, on account of the extraordinary copiousness of the distribution of the tracheæ in the intestinal folds.

394. The main trunks of the Tracheal system, with their principal ramifications, may generally be got out with little difficulty, by laying open the body of an insect or larva, under water, in a dissecting-trough (§ 104), and removing the whole visceral mass, taking care to leave as many as possible of the branches which will be seen proceeding to this from the two great longitudinal tracheæ, to whose position these branches will serve as a guide. Mr. Quekett recommends the following as the most simple method of obtaining a perfect system of tracheal tubes from a larva:—a small opening having been made in its body, this is to be placed in strong acetic acid, which will soften or decompose all the viscera; and the tracheæ may then be well washed with the syringe, and removed from the body with the greatest facility, by cutting away the connections of the main tubes with the spiracles by means of fine-pointed scissors. In order to mount them, they should be floated upon the slide, on which they should then be laid out in the position best adapted for displaying them. If they are to be mounted in Canada balsam, they should be

allowed to dry upon the slide, and should then be treated in the usual way; but their natural appearance is best preserved by mounting them in fluid (weak spirit or Goadby's solution), using a shallow cell to prevent pressure. The finer ramifications of the tracheal system may generally be seen particularly well in the membranous wall of the stomach or intestine; and this, having been laid out and dried upon the glass, may be mounted in balsam so as to keep the tracheæ full of air (whereby they are much better displayed), if care be taken to use balsam that has been previously thickened, to drop this on the object without liquefying it more than is absolutely necessary, and to heat the slide and the cover (the heat may be advantageously applied directly to the cover, after it has been put on, by turning over the slide so that its upper face shall look downwards) only to such a degree as to allow the balsam to spread and the cover to be pressed down. The spiracles are easily dissected out by means of a pointed knife or a pair of fine scissors; they should be mounted in fluid, when their texture is soft; and in balsam, when the integument is hard and horny.

395. *Wings*.—These organs are essentially composed of an extension of the external membranous layer of the integument, over a framework formed by prolongations of the inner horny layer; within which prolongations, tracheæ are nearly always to be found, whilst they also contain channels through which blood circulates during the growth of the wing and for a short time after its completion. This is the simple structure presented to us in the wings of *Neuroptera* (dragon-flies, &c.), *Hymenoptera* (bees and wasps), *Diptera* (two-winged flies), and also of many *Homoptera* (cicadæ and aphides); and the principal interest of these wings as microscopic objects, lies in the distribution of their "veins" or "nerves" (for by both names are the ramifications of their skeleton known), and in certain points of accessory structure. The venation of the wings is most beautiful in the smaller *Neuroptera*; since it is the distinguishing feature of this order, that the veins, after subdividing, reunite again, so as to form a close network; whilst in the *Hymenoptera* and *Diptera* such reunions are rare, especially towards the margin of the wings, and the areolæ are much larger. Although the membrane of which these wings are composed, appears perfectly homogeneous when viewed by transmitted light, even with a high magnifying power, yet, when viewed by light reflected obliquely from their surfaces, an appearance of cellular areolation is often discernible; this is well seen in the common *Fly*, in which each of these areolæ has a hair in its centre. In order to make this observation, as well as to bring out the very beautiful iridescent hues which the wings of many minute insects (as the *Aphides*) exhibit when thus viewed, it is convenient to hold the wing in the stage-forceps, for the sake of giving it every variety of inclination; and when that position has been found, which

best displays its most interesting features, it should be set up as nearly as possible in the same. For this purpose it should be mounted on an opaque slide; but instead of being laid down upon its surface, the wing should be raised a little above it, its "stalk" being held in the proper position by a little cone of soft wax, in the apex of which it may be imbedded. The wings of most Hymenoptera are remarkable for the peculiar apparatus by which those of the same side are connected together, so as to constitute in flight but one large wing; this consists of a row of curved hooks on the anterior margin of the posterior wing, which lay hold of the thickened and doubled-down posterior edge of the anterior wing. These hooks are sufficiently apparent in the wings of the common *Bee*, when examined with even a low magnifying power; but they are seen better in the *Wasp*, and better still in the *Hornet*. The peculiar scaly covering of the wings of the Lepidoptera has already been noticed (§ 381); but it may here be added that the entire wings of many of the smaller and commoner insects of this order, such as the *Tineidæ* or "clothes' moths," form very beautiful opaque objects for low powers; the most beautiful of all being the divided wings of the *Fissipennes* or "plumed moths," especially those of the genus *Pterophorus*.


396. There are many Insects, however, in which the wings are more or less consolidated by the interposition of a layer of horny substance between the two layers of membrane. This plan of structure is most fully carried out in the Coleoptera (beetles), in which the anterior wings are so much thickened and are so little extended, that they are useless in flight, and serve merely as cases or covers for the posterior, which lie folded up beneath them when not in use; hence these are distinguished as *elytra*. These *elytra*, when the insect is at rest, meet along the median line of the back, and cover nearly the whole upper surface of the body; and it is upon them that the brilliant hues, by which the integument of many of these insects is distinguished, are most strikingly displayed. In the anterior wings of the *Forficulidæ* or earwig tribe (which form the connecting link between this order and the Orthoptera), the cellular structure may often be readily distinguished when they are viewed by transmitted light, especially after having been mounted in Canada balsam. The anterior wings of the Orthoptera (grasshoppers, crickets, &c.) although not by any means so solidified as those of Coleoptera, contain a great deal of horny matter; they are usually rendered sufficiently transparent, however, by Canada balsam, to be viewed with transmitted light; and many of them are so colored as to be very showy objects (as are also the posterior fan-like wings) for the solar or gas-microscope, although their large size, and the absence of any minute structure, prevent them from affording much interest to the ordinary Microscopist. We must not omit to mention, however, the curious sound-producing apparatus which

is possessed by most insects of this order, and especially by the common *House-Cricket*; this consists of the "tympanum" or drum, which is a space on each of the upper wings, scarcely crossed by veins, but bounded externally by a large dark vein provided with three or four longitudinal ridges, and of the "file" or "bow," which is a transverse horny ridge in front of the tympanum, furnished with numerous teeth; and it is believed that the sound is produced by the rubbing of the two bows across each other, while its intensity is increased by the sounding-board action of the tympanum. The wings of the *Fulgoridæ* (lantern-flies) have much the same texture with those of the Orthoptera, and possess about the same value as microscopic objects; differing considerably from the purely membranous wings of the Cicade and Aphides, which are associated with them in the order Homoptera. In the order Hemiptera, to which belong various kinds of land and water insects that have a suctorial mouth resembling that of the common *Bug*, the wings of the anterior pair are usually of parchmenty consistence, though membranous near their tips, and are often so richly colored as to become very beautiful objects, when mounted in balsam and viewed by transmitted light; this is the case especially with the terrestrial vegetable-feeding kinds, such as the *Pentatoma* and its allies, some of the tropical forms of which rival the most brilliant of the Beetles. The British species are by no means so interesting; and the aquatic kinds, which, next to the bed-bugs, are the most common, always have a dull brown or almost black hue; even among these last, however,—of which the *Notonecta* (water-boatman) and the *Nepa* (water-scorpion) are well-known forms,—the wings are beautifully variegated by differences in the depth of that hue.

397. *Feet*.—Although the feet of Insects are formed pretty much on one general plan, yet that plan is subject to considerable modifications, in accordance with the habits of life of different species. The entire limb usually consists of five divisions, namely, the *coxa* or hip, the *trochanter*, the *femur* or thigh, the *tibia* or shank, and the *tarsus* or foot; and this last portion is made up of several successive joints. The typical number of these joints seems to be five; but that number is subject to reduction; and the vast order Coleoptera is subdivided into primary groups, according as the tarsus consists of five, four, or three segments. The last joint of the tarsus is usually furnished with a pair of strong hooks or claws (Figs. 294, 295); and these are often serrated (that is, furnished with saw-like teeth), especially near the base. The under surface of the other joints is frequently beset with tufts of hairs, which are arranged in various modes, sometimes forming a complete "sole;" this is especially the case in the family *Curculionidæ*; so that a pair of the feet of the "diamond-beetle," mounted so that one shows the upper surface made resplendent by its jewel-like scales, and the other the hairy cushion beneath, is a very interesting object. In many Insects,


especially of the *Fly* kind, the foot is furnished with a pair of membranous expansions, termed *pulvilli* (Fig. 294); and these are beset with numerous hairs, each of which has a minute disk at its extremity. This structure is evidently connected with the power which these insects possess, of walking over smooth surfaces in opposition to the force of gravity; yet there is still considerable uncertainty as to the precise mode in which it ministers to this faculty. Some believe that the "pulvilli" act as suckers, the insect being held up by the pressure of the air against their upper surface, when a vacuum is formed beneath; whilst others maintain that the adhesion is the result of the secretion of a viscid liquid from the under side of the foot. The careful observations of Mr. Hepworth have led him to a conclusion which seems in harmony with all the facts of the case; namely, that the minute disks at

FIG. 294.



Foot of *Fly*.

FIG. 295.



A, Foot of *Dytiscus*, showing its apparatus of suckers; a, b, large suckers; c, ordinary suckers:—
 a, one of the ordinary suckers more highly magnified.

the extremity of the individual hairs act as suckers, and that each of them secretes a liquid, which, though not viscid, serves to make its adhesion perfect.¹ And this view of the case derives

¹ See Mr. Hepworth's communications to the "Quart. Journ. of Microsc. Science," vol. ii, p. 158, and vol. iii, p. 312.

information from the presence of a similar apparatus, on a far larger scale, in the feet of the *Hyacinth*. Fig. 256, A. The first joint of the tarsi of the insect are widely expanded, so as to form a nearly circular base: and this is provided with a very remarkable apparatus of suckers, of which one disk (a) is extremely large, and is furnished with strong radiating fibres, a second (b) is a smaller one formed on the same plan (a third, of the like kind, being often present, whilst the greater number are comparatively small tubular sub-shaped bodies, each having a very delicate membranous sucker at its extremity, as seen on a larger scale at c. These last seem to resemble the hairs of the Fly: but in every particular but dimension: and an intermediate use is possessed by the hairs of many beetles, especially Curculionidae. The feet of Caterpillars differ considerably from those of perfect Insects. Those of the first three segments, which are afterwards to be replaced by true legs, are furnished with strong horny claws; but each of those of the other segments, which are termed "pro-legs," is composed of a circular series of comparatively slender curved hirsutiæ, by which the caterpillar is enabled to cling to the minute roughnesses of the surface of the leaves, &c., on which it feeds. This structure is well seen in the pro-legs of the common Silk-worm.

257. *Stings and Ovipositors*.—The Insects of the order *Hymenoptera* are all distinguished by the prolongation of the last segment of the abdomen into a peculiar organ, which, in one division of the order, is a "sting," and in the other is an "ovipositor,"—an instrument for the deposition of the eggs, which is usually also provided with the means of boring a hole for their reception. The former group consists of the Bees, Wasps, Ants, &c.; the latter of the Saw-flies, Gall-flies, Ichneumon-flies, &c. These two sets of instruments are not so unlike in structure, as they are in function. The "sting" is usually formed of a pair of darts, beset with barbed teeth at their points, and furnished at their roots with powerful muscles whereby they can be caused to project from their sheath, which is a horny case formed by the prolongation of the integument of the last segment, slit into two halves, which separate to allow the protrusion of the sting; whilst the peculiar "venom" of the sting is due to the ejection, by the same muscular action, of a poisonous liquid, from a bag situated near the root of the sting, which passes down a canal excavated between the darts, so as to be inserted into the puncture which they make. The stings of the common Bee, Wasp, and Hornet, may all be made to display this structure without much difficulty in the dissection. The "ovipositor" of such insects as deposit their eggs in holes ready made, or in soft animal or vegetable substances (as is the case with the *Ichneumonidae*), is simply a long tube, which is enclosed, like the sting, in a cleft sheath. In the Gall-flies (*Cynipidæ*), the extremity of the ovipositor has a toothed edge, so as to act as a kind of saw,

whereby harder substances may be penetrated; and thus an aperture is made in the leaf, stalk, or bud of the plant or tree infested by the particular species, in which the egg is deposited, together with a drop of fluid that has a peculiarly irritating effect upon the vegetable tissues, occasioning the production of the "galls," which are new growths that serve not only to protect the larvæ, but also to afford them nutriment. The Oak is infested by several species of these insects, which deposit their eggs in different parts of its fabric; and some of the small "galls" which are often found upon the surface of oak leaves, are extremely beautiful objects for the lower powers of the Microscope. It is in the *Tenthredinidæ*, or Saw-flies, and in their allies the *Siricidæ*, that the ovipositor is furnished with the most powerful apparatus for penetration; and some of these insects can bore by its means into hard timber. Their "saws" are not unlike the "stings" of Bees, &c., but are broader, are toothed for a greater length, and are made to slide along a firm piece that supports each blade, like the "back" of a carpenter's "tenon-saw;" they are worked alternately (one being protruded while the other is drawn back) with great rapidity; and when the perforation has been made, the two blades are separated enough to allow the passage of the eggs between them. Many other Insects, especially of the order *Diptera*, have very prolonged ovipositors, by means of which they can insert their eggs into the integuments of animals, or into other situations in which the larvæ will obtain appropriate nutriment; a remarkable example of this is furnished by the Gad-fly (*Tabanus*), whose ovipositor is composed of several joints, capable of being drawn together or extended like those of a telescope, and is terminated by boring instruments; and the egg being conveyed by its means, not only *into* but *through* the integument of the Ox, so as to be imbedded in the tissue beneath, a peculiar kind of inflammation is set up there, which (as in the analogous case of the gall-fly) forms a nidus appropriate both to the protection and to the nutrition of the larva. Other Insects which deposit their eggs in the ground, such as the *Locusts*, have their ovipositors so shaped as to answer for digging holes for their reception. The preparations which serve to display the foregoing parts, are best seen when mounted in balsam; save in the case of the muscles and poison-apparatus of the sting, which are better preserved in weak spirit or Goadby's solution.

399. The sexual organs of Insects furnish numerous objects of extreme interest to the Anatomist and Physiologist; but as an account of them would be unsuitable to the present work, a reference to a copious source of information respecting one of their most curious features, and to a list of the species that afford good illustrations, must here suffice.¹ The *Eggs* of many Insects are

¹ See the Memoirs of M. Lacaze-Duthiers "Sur l'armure génitale des Insectes," in "Ann. des Sci. Nat." 3ième Sér. tom. xii, xiv, xvii, xviii, xix; and M. Ch. Robin's

objects of great beauty, on account of the regularity of their form, and the symmetry of the markings on their surface (Fig. 296). The most interesting belong for the most part to the Lepidopterous order; and there are few among these that are not worth examination, some of the commonest (such as those of the

FIG. 296.



Eggs of Insects magnified:—A, *Pontia napi*; B, *Vanessa urticae*; C, *Hipparchia tithonus*; D, *Argynnis Lathonia*.

Cabbage-butterfly, which are found covering large patches of the leaves of that plant) being as remarkable as any. Those of the puss-moth (*Cerura vinula*), the privet hawk-moth (*Sphinx ligustri*), the small tortoise-shell butterfly (*Vanessa urticae*), the meadow-brown butterfly (*Hipparchia janira*), the brimstone-moth (*Rumia crataegata*), and the silk-worm (*Bombyx mori*), may be particularly specified; and from other orders, those of the cockroach (*Blatta orientalis*), field cricket (*Acheta campestris*), water scorpion (*Nepa ranatra*), bug (*Cimex lectularius*), cow-dung fly (*Scatophaga stercoraria*), and blow-fly (*Musca vomitoria*). In order to preserve these eggs, they must be mounted in fluid in a cell; since they will otherwise dry up and become misshapen. The remarkable mode of reproduction that exists among the *Aphides* must not pass unnoticed here, from its curious connection with the non-sexual reproduction of *Entomostraca* (§ 369) and *Rotifera* (§ 279), as also of *Hydra* (§ 301) and *Zoophytes* generally, all of which fall specially, most of them exclusively, under the observation of the Microscopist. The wingless *Aphides* which may be seen in the spring and early summer, may be considered as larvæ or pupæ (the earlier states of this insect not being distinguishable from its perfect form, except by their want of wings); and these larvæ, which, though commonly designated as females, are really of no sex, give origin to a brood of similar wingless *Aphides*, which come into the world alive, and which, before long, go through a like process of multiplication. As many as from seven to ten successive broods may thus be produced in the course of a single season; so that from a single *Aphis*, it has been calculated that no fewer than ten thousand million millions may be evolved within that period. In the latter part of the year, however, some of these larval *Aphides* attain their full development into winged males and females, by which the true generative process is performed, whose products are eggs, which, when

"Mémoire sur les Objets qui peuvent être conservés en Préparations Microscopiques" (Paris, 1856), which is peculiarly full in the enumeration of the objects of interest afforded by the class of Insects.

hatched in the succeeding spring, give origin to a new brood of larvæ that repeat the curious life-history of their predecessors.¹ The non-sexual multiplication of the larvæ is obviously a process of *gemmation*, analogous to the multiplication of cells by subdivision; whilst the true Generative process, analogous to the *conjugation* of cells, is only performed when perfect Aphides of distinct sexes have been evolved.

400. *Arachnida*.—The general remarks which have been made in regard to Insects, are equally applicable to this class; which includes, along with the *Spiders* and *Scorpions*, the tribe of *Acarida*, which consists of the *Mites* and *Ticks*. Many of these last are parasitic, and are popularly associated with the wingless parasitic Insects, to which they bear a strong general resemblance, save in having eight legs instead of six. The true "mites" (*Acarinæ*) generally have the legs adapted for walking, and some of them are of active habits. The common *cheese-mite*, as seen by the naked eye, is familiar to every one; yet few who have not seen it under a microscope, have any idea of its real conformation and movements; and a cluster of them, cut out of the cheese they infest, and placed under a magnifying power sufficiently low to enable a large number to be seen at once, is one of the most amusing objects that can be shown to the young. There are many other species, which closely resemble the cheese-mite in structure and habits, but which feed upon different substances; and some of these are extremely destructive. To this group belongs a small species, the *Sarcoptes scabiei*, whose presence appears to be the occasion of one of the most disgusting diseases of the skin—the itch,—and which is hence commonly termed the "itch insect." It is not found in the pustule itself, but in a burrow which passes off from one side of it, and which is marked by a red line on the surface; and if this burrow be carefully examined, the creature will very commonly, but not always, be met with. It is scarcely visible to the naked eye; but when examined under the microscope, it is found to have an oval body, a mouth of conical form, and eight feet, of which the four anterior are terminated by small suckers, whilst the four posterior end in very prolonged bristles. The male is only about half the size of the female. The *Ricinæ* or "ticks" are usually destitute of eyes, but have the mouth provided with lancets, that enable them to penetrate more readily the skins of animals whose blood they suck. They are usually of a flattened, round, or oval form; but they often acquire a very large size by suction, and become distended like a blown bladder. Different species are parasitic upon different animals; and they bury their suckers (which are often furnished with minute recurved hooks) so firmly in the skins of these, that they can hardly be detached without pulling away the skin with them. It is probably the

¹ For a careful examination and philosophical appreciation of the real nature of this process, see Dr. Waldo J. Burnett's Memoir in the "Transactions of the American Academy of Arts and Sciences," 1853.

young of a species of this group, which is commonly known as the "harvest bug," and which is usually designated as the *Acarus autumnalis*; this is very common in the autumn upon grass or other herbage, and insinuates itself into the skin at the roots of the hair, producing a painful irritation; like other Acarida, for some time after their emersion from the egg, it possesses only six legs (the other pair being only acquired after the first moult), so that its resemblance to parasitic insects becomes still stronger. It is probable that to this group also belongs the *Demodex folliculorum*, a creature which is very commonly found parasitic in the sebaceous follicles of the human skin, especially in those of the nose. In order to obtain it, pressure should be made upon any one of these that appears enlarged and whitish, with a terminal black spot; the matter forced out will consist principally of the accumulated sebaceous secretion, having the parasites with their eggs and young mingled with it. These are to be separated by the addition of oil, which will probably soften the sebaceous matter sufficiently to set free the animals, which may be then removed with a pointed brush; but if this mode should not be effectual, the fatty matter may be dissolved away by digestion in a mixture of alcohol and ether. The pustules in the skin of a dog affected with the "mange" have been found by Mr. Topping to contain a *Demodex*, which seems only to differ from that of the human sebaceous follicles in its somewhat smaller size; and M. Gruby is said to have given to a dog a disease resembling the mange, if not identical with it, by inoculating it with the human parasite. The *Acarida* are best preserved as microscopic objects, by mounting in glycerine.

401. The number of objects of general interest, furnished to the Microscopist by the *Spider* tribe, is by no means considerable. Their eyes exhibit a condition intermediate between that of Insects and Crustaceans, and that of Vertebrata; for they are single like the "stemmata" of the former, usually number from six to eight, are sometimes clustered together in one mass, but are sometimes disposed separately, while they present a decided approach in internal structure to the type characteristic of the visual organs of the latter. The structure of the Mouth is always mandibulate, and is less complicated than that of the mandibulate insects. The Respiratory apparatus, which, where developed at all among the Acarida, is tracheary like that of Insects, is here constructed upon a very different plan; for the "stigmata," which are usually four in number on each side, open into a like number of respiratory sacculi, each of which contains a series of leaf-like folds of its lining membrane, upon which the blood is distributed so as to afford a large surface to the air. In the structure of the limbs, the principal point worthy of notice is the peculiar appendage with which they usually terminate; for the strong claws, with a pair of which

the last joint of the foot is furnished, have their edges cut into comb-like teeth (Fig. 297), which seem to be used by the animal as cleansing instruments. One of the most curious parts of the organization of the Spiders is the "spinning apparatus," by means of which they fabricate their elaborately constructed webs. This consists of the "spinnerets," and of the glandular organs in which the fluid that hardens into the thread is elaborated. The



Foot, with comb-like claws, of the common Spider (*Epeira*).

usual number of the spinnerets, which are situated at the posterior extremity of the body, is six; they are little teat-like prominences, beset with hairy appendages; and it is through a certain set of these appendages, which are tubular and terminate in fine-drawn points, that the glutinous secretion is forced out in a multitude of streams of extreme minuteness. These streams harden into fibrils, immediately on coming into contact with the air; and the fibrils proceeding from all the apertures of each spinneret, coalesce into a single thread. It is doubtful, however, whether all the spinnerets are in action at once, or whether those of different pairs may not have dissimilar functions; for whilst the radiating threads of a spider's web are simple (Fig. 298, A), those which lie across these, forming its concentric circles or rather polygons, are studded at intervals with viscid globules (B), which appear to give to these threads their peculiarly adhesive character; and it does not seem by any means unlikely, that each kind of thread should be produced by its own pair of spinnerets. The total number of spinning tubes varies greatly, according to the species of the spider, and the sex and age of the individual; being more than 1000 in some cases, and less than 100 in others. The size and complexity of the secreting glandulæ vary in like manner: thus in the Spiders which are most remarkable for the large dimensions and regular construction of their webs, they occupy a large portion of the abdominal cavity, and are composed of slender branching tubes, whose length is increased by numerous convolutions; whilst in those which have only occasional use for their threads, the secreting organs are either short and simple follicles, or are undivided tubes of moderate length.

FIG. 298.



Ordinary thread (A), and glutinous thread (B), of the common Spider.

CHAPTER XVIII.

VERTEBRATED ANIMALS.

402. WE are now arrived at that highest division of the Animal Kingdom, in which the bodily fabric attains its greatest development, not only as to completeness, but also as to size; and it is in most striking contrast with the class we have been last considering. Since not only the entire bodies of Vertebrated animals, but, generally speaking, the smallest of their integral parts, are far too large to be viewed as Microscopic objects, we can study their structure only by a separate examination of their component elements; and it seems, therefore, to be a most appropriate course, to give, under this head, a sketch of the microscopic characters of those *primary tissues*, of which their fabric is made up, and which, although they may be traced with more or less distinctness in the lower tribes of Animals, attain their most complete development in this group. Since the time when Schwann first made public the remarkable results of his researches (p. 56), it has been very generally believed that all the Animal tissues are formed, like those of Plants, by a metamorphosis of *Cells*; an exception being taken, however, by some Physiologists, in regard to the simple fibrous tissues (§ 417). The tendency of many recent investigations, however, has been to throw further doubt on the generality of this doctrine; since they appear to indicate that many other tissues than the fibrous may be formed (like these) by the consolidation of the *plasma* or formative fluid, without passing through the intermediate condition of cells. Hence no attempt will here be made to do more than describe the most important of those distinctive characters, which the principal tissues present, when subjected to microscopic examination; and as it is of no essential consequence what order is adopted, we may conveniently begin with the structure of the *skeleton*,¹ which gives support and protection to the softer parts of the fabric.

403. *Bone*.—The Microscopic characters of osseous tissue may

¹ This term is used in its most general sense, as including not only the proper *vertebral* or internal skeleton, but also the hard parts protecting the exterior of the body, which forms the *dermal* skeleton.

sometimes be seen in very thin natural plates of bone, such as in that forming the scapula (shoulder-blade) of a Mouse; but they are displayed more perfectly by artificial sections, the details of the arrangement being dependent upon the nature of the specimen selected, and the direction in which the section is made. Thus when the shaft of a "long" bone of a Bird or Mammal is cut across in the middle of its length, we find it to consist of a hollow cylinder of dense bone, surrounding a cavity which is occupied by an oily marrow; but if the section be made nearer its extremity, we find the outside wall gradually becoming thinner, whilst the interior, instead of forming one large cavity, is divided into a vast number of small chambers or *cancelli*, which com-

municate with each other and with the cavity of the shaft, and are filled, like it, with marrow. In the bones of Reptiles and Fishes, on the other hand, this "cancellated" structure usually extends throughout the shaft, which is not so completely differentiated into solid bone and medullary cavity, as it is in the higher Vertebrata. In the most developed kinds of "flat" bones, again, such as those of the head, we find the two surfaces to be composed of dense plates of bone, with a "cancellated" structure between them; whilst in the less perfect type presented to us in the lower Vertebrata, the whole thickness is usually more or less "cancellated," that is, burrowed out by medullary cavities.

When we examine, under a low magnifying power, a longitudinal section of a "long" bone, or a section of a "flat" bone parallel to its surface, we find it traversed by numerous canals, termed *Haversian* after their discoverer Havers, which are in connection with the central cavity, and are filled, like it, with marrow: in the shafts of long bones, these canals usually run in the direction of their length, but are connected here and there by cross branches; whilst in the flat bones, they form an irregular network. On applying a higher magnifying power to a thin transverse section of a long bone, we observe that each of the canals whose orifices present themselves in the field of view (Fig. 299), is the centre of a rod of bony tissue (1), usually more or less circular in its form, which is arranged around it in concentric

FIG. 299.



Minute structure of Bone, as seen in transverse section:—1, an ossicle surrounding an Haversian canal, 3, showing the concentric arrangement of the lamellæ; 2, the same, with the lacunæ and canaliculi; 4, portions of the lamellæ parallel with the external surface.

rings, resembling those of an Exogenous Stem. These rings are marked out and divided by circles of little dark spots; which, when closely examined (2), are seen to be minute flattened cavities excavated in the solid substance of the bone, from the two flattened sides of which pass forth a number of extremely minute tubules, one set extending inwards, or in the direction of the centre of the system of rings, and the other outwards, or in the direction of its circumference; and by the inosculation of the tubules (which are termed *canaliculi*) of the different rings with each other, a continuous communication is established between the central Haversian canal and the outermost part of the bony rod that surrounds it, which doubtless ministers to the nutrition of the texture. Bloodvessels are traceable into the Haversian canals; but the "canaliculi," being far too minute to carry blood-corpuscles, can only convey a nutrient fluid that is separated from the blood for the special service of the bone.

FIG. 300.



Lacuna of Osseous substance—*a*, central cavity,
b, its ramifications.


404. The minute cavities, or *lacunæ* (sometimes, but erroneously termed "bone-corpuscles," as if they were solid bodies), from which the canaliculi proceed, are highly characteristic of the true osseous structure; being never deficient in the minutest parts of the bones of the higher Vertebrata, although those of Fishes are occasionally destitute of them. The dark appearance which they present is not due to opacity, but is simply an optical effect, dependent (like the blackness of air-bubbles in liquids) upon the dispersion of the rays by the highly-refracting substance that surrounds them (§ 98). The size and form of the lacunæ differ considerably in the several Classes of Vertebrata, and even in some instances in the Orders; so as to allow of the determination of the tribe to which a bone belonged, by the microscopic examination of even a minute fragment of it (§ 453). The following are the average dimensions of the lacunæ, in characteristic examples drawn from the four principal classes, expressed in fractions of an inch:—

	Long Diameter.	Short Diameter.
Man,	1-1440 to 1-2400	1-4000 to 1-8000
Ostrich,	1-1333 to 1-2250	1-5425 to 1-9650
Turtle,	1-375 to 1-1150	1-4500 to 1-5840
Conger-eel,	1-550 to 1-1135	1-4500 to 1-8000

The lacunæ of *Birds* are thus distinguished from those of *Mammals* by their somewhat greater length and smaller breadth; but they differ still more in the remarkable tortuosity of the canaliculi, which wind backwards and forwards in a very irregular manner. There is an extraordinary increase in length in the

lacunæ of *Reptiles*, without a corresponding increase in breadth; and this is also seen in some *Fishes*, though in general the lacunæ of the latter are remarkable for their angularity of form, and the fewness of their radiations,—as shown in Fig. 301, which represents the lacunæ and canaliculi in the bony scale of the *Lepidosteus* (“bony pike” of the North American lakes and rivers), with which the bones of its internal skeleton perfectly agree in structure. The dimensions of the lacunæ in any bone do not bear any relation to the size of the animal to which it belonged; thus there is little or no perceptible difference between their size in the enormous extinct *Iguanodon*, and in the smallest *Lizard* now inhabiting the earth. But they bear a close relation to the size of the blood-corpuscles in the several classes; and this relation is particularly obvious in the “perennibranchiate” *Batrachia*, the extraordinary size of whose blood-corpuscles will be presently noticed (§ 414):—

FIG. 301.



Section of the bony scale of *Lepidosteus*:—a, showing the regular distribution of the lacunæ and of the connecting canaliculi; b, small portion more highly magnified.

	Long Diameter.	Short Diameter.
Proteus,	1-570 to 1-980	1-885 to 1-1200
Siren,	1-290 to 1-480	1-540 to 1-975
Menopoma,	1-450 to 1-700	1-1300 to 1-2100
Lepidosiren,	1-375 to 1-494	1-980 to 1-2200
Pterodactyle,	1-445 to 1-1185	1-4000 to 1-5225 ¹

405. In preparing sections of bone, it is important to avoid the penetration of the Canada balsam into the interior of the lacunæ and canaliculi; since, when these are filled by it, they become almost invisible. Hence it is preferable not to employ this cement at all, except, it may be, in the first instance; but to rub down the section beneath the finger, guarding its surface with a slice of cork or a slip of gutta serena (§ 111); and to give it such a polish, that it may be seen to advantage even when mounted dry. As the polishing, however, occupies much time, the benefit which is derived from covering the surfaces of the specimen with Canada balsam may be obtained, without the injury resulting from the penetration of the balsam into its interior, by adopting the following method. A quantity of balsam proportioned to the size of the specimen is to be spread upon a glass slip, and to be

¹ See Prof. J. Quekett's Memoir on this subject, in the "Transact. of the Microsc. Soc." Ser. 1, vol. ii; and his more ample illustration of it in the "Illustrated Catalogue of the Histological Collection in the Museum of the Roy. Coll. of Surgeons," vol. ii.

rendered stiffer by boiling, until it becomes nearly solid when cold; the same is to be done to the thin glass cover; next, the specimen being placed on the balsamed surface of the slide, and being overlaid by the balsamed cover, such a degree of warmth is to be applied, as will suffice to liquefy the balsam, without causing it to flow freely; and the glass cover is then to be quickly pressed down, and the slide to be rapidly cooled, so as to give as little time as possible for the penetration of the liquefied balsam into the lacunar system. The same method may be employed in making sections of Teeth.

406. *Teeth*.—The intimate structure of the Teeth in the several classes and orders of Vertebrata presents differences which are no less remarkable than those of their external form, arrangement and succession. It will obviously be impossible here to do more than sketch some of the most important of these varieties. The principal part of the substance of all teeth is made up of a solid tissue that has been appropriately termed *Dentine*. In the Shark tribe, as in many other Fishes, the general structure of this “dentine” is extremely analogous to that of bone; the tooth being traversed by numerous canals, which are continuous with the Haversian canals of the subjacent bone, and receive bloodvessels from them (Fig. 302); and each of these canals being surrounded by a system of tubuli (Fig.

FIG. 302.



FIG. 303.



Fig. 302. Perpendicular section of tooth of *Lamna*, moderately enlarged, showing network of medullary canals.

Fig. 303. Transverse section of portion of tooth of *Pristis*, more highly magnified, showing orifices of medullary canals, with systems of radiating and inosculating tubuli.

303), which radiate into the surrounding solid substance. These tubuli, however, do not enter lacunæ, nor is there any concentric annular arrangement around the medullary canals; but each system of tubuli is continued onwards through its own division

of the tooth, the individual tubes sometimes giving off lateral branches, whilst in other instances their trunks bifurcate.

This arrangement is peculiarly well displayed, when sections of teeth constructed upon this type are viewed as opaque objects (Fig. 304). In the teeth of the higher Vertebrata, however, we usually find the centre excavated into a single cavity (Fig. 305), and the remainder destitute of vascular canals; but there are intermediate cases (as in the teeth of the great fossil Sloths) in which the inner portion of the dentine is traversed by




FIG. 304.
Transverse Section of Tooth of *Myliobates* (Eagle Ruy) viewed as an opaque object.

prolongations of this cavity, conveying bloodvessels, which do not pass into the exterior layers. The tubuli of the "non-vascular" dentine, which exists by itself in the teeth of nearly all Mammalia, and which in the Elephant is known as "ivory," all radiate from the central cavity, and pass towards the surface of the tooth in a nearly parallel course. Their diameter at their largest part averages 1-10,000th of an inch; their smallest branches are immeasurably fine. It is impossible that even the largest of them can receive blood, as their diameter is far less than that of the blood-discs; but it is probable that, like the canaliculi of bone, they may absorb nutrient matter from the vascular surface upon which their inner extremities open. The tubuli in their course present greater and lesser undulations; the former are few in number; but the latter are numerous, and as they occur at the same part of the course of several contiguous tubes, they give rise to the appearance of lines concentric with the centre of radiation. These secondary curvatures probably indicate, in dentine, as in the crab's shell (§ 374) successive stages of calcification.

407. In the teeth of Man and most other Mammals, and in those of many Reptiles and some Fishes, we find two other substances, one of them harder, and the other softer, than dentine; the former is termed *Enamel*; and the latter *Cementum* or *Crusta Petrosa*. The *Enamel* is composed of long prismatic cells, closely resembling those of the prismatic shell-substance formerly described (§ 336), but on a far more minute scale; the diameter of the cells not being more, in Man, than 1-5600th of an inch. The length of the prisms corresponds with the thickness of the layer of enamel; and the two surfaces of this layer present the ends of the prisms, the form of which usually approaches the hexagonal. The course of the enamel-prisms is

more or less wavy; and they are marked by numerous trans-

FIG. 305.



Vertical section of *Human Molar Tooth*:—1, enamel; 2, cementum or *crusta petrosa*; 3, dentine or ivory; 4, osseous excrescence, arising from hypertrophy of cementum; 5, pulp-cavity; 6, osseous lacunæ at outer part of dentine.

verse striæ, resembling those of the prismatic shell-substance, and probably originating in the same cause,—the coalescence of a series of shorter cells, to form the lengthened prism. In Man, and in Carnivorous animals, the enamel covers the crown of the tooth only, with a simple cap or superficial layer of tolerably uniform thickness (Fig. 305, *a*), which follows the surface of the dentine in all its inequalities; and its component prisms are directed at right angles to that surface, their inner extremities resting in slight but regular depressions on the exterior of the dentine. In the teeth of many of the Herbivorous animals, however, the Enamel forms (with the Cementum) a series of vertical plates, which dip down into the substance of the dentine, and present their edges alternately with it, at the grinding surface of the tooth; and there is in such teeth no continuous layer of enamel over the crown. The purpose of this arrangement is evidently to provide, by the unequal *wear* of these

three substances,—of which the enamel is the hardest, and the cementum the softest,—for the constant maintenance of a rough surface, adapted to triturate the tough vegetable substances on which these animals feed. The enamel is the least constant of the dental tissues. It is more frequently absent than present in the teeth of the class of Fishes; it is wanting in the entire order of Ophidia (serpents) among existing Reptiles; and it forms no part of the teeth of the Edentata (sloths, &c.) and Cetacea (whales) amongst Mammals. The *Cementum*, or *Crusta Petrosa*, has the characters of true bone; possessing its distinctive stellate lacunæ and radiating canaliculi. Where it exists in small amount, we do not find it traversed by medullary canals; but, like dentine, it is occasionally furnished with them, and thus resembles bone in every particular. These medullary canals enter its substance from the exterior of the tooth, and consequently pass towards those which radiate from the central cavity in the direction of the surface of the dentine, where this possesses a similar vascularity,—as was remarkably the case in the teeth of the extinct *Megatherium*. In the Human tooth, however, the cementum has no such vascularity; but forms a thin layer, which envelopes the root of the tooth, commencing near the termination of the capping of enamel (Fig. 305, *b*). In the teeth of many Herbivorous Mammals, it dips down with the enamel to form the vertical plates of the interior of the tooth;

and in the teeth of the Edentata, as well as of many Reptiles and Fishes, it forms a thick continuous envelope over the whole surface, until worn away at the crown.

408. *Dermal Skeleton*.—The skin of Fishes, of most Reptiles, and of a few Mammals, is strengthened by plates of a horny, cartilaginous, bony, or even enamel-like texture, which are sometimes fitted together at their edges, so as to form a continuous box-like envelope, whilst more commonly they are so arranged as partially to overlie one another, like the tiles on a roof; and it is in this latter case that they are usually known as *scales*. Although we are accustomed to associate in our minds the “scales” of Fishes with those of Reptiles, yet they are essentially different structures; the former being developed in the substance of the true skin, with a layer of which, in addition to the epidermis, they are always covered; and bearing a resemblance to cartilage and bone in their texture and composition; whilst the latter are formed upon the surface of the true skin, and are to be considered as analogous to nails, hoofs, &c., and other “epidermic appendages.” In nearly all the existing Fishes, the scales are flexible, being but little consolidated by calcareous deposit; and in some species they are so thin and transparent, that, as they do not project obliquely from the surface of the skin, they can only be detected by raising the superficial layer of the skin, and searching beneath it, or by tearing off the entire thickness of the skin, and looking for them near its under surface. This is the case, for example, with the common *Eel*, and with the *Viviparous Blenny*; of either of which fish, the skin is a very interesting object when dried and mounted in Canada balsam, the scales being seen imbedded in its substance, whilst its outer surface is studded with pigment-cells. Generally speaking, however, the posterior extremity of each scale projects obliquely from the general surface, carrying before it the thin membrane that encloses it, which is studded with pigment-cells; and a portion of the skin of almost any Fish, but especially of such as have scales of the *ctenoid* kind (that is, furnished at their posterior extremities with comb-like teeth, Fig. 307), when dried with its scales *in situ*, is a very beautiful opaque object for the low powers of the Microscope (Fig. 306). Care must be taken, however, that the light is made to glance upon it in the most advantageous manner; since the brilliancy with which it is reflected from the comb-like projections, entirely depends upon the angle at which it falls upon them. The only appearance of structure exhibited by the thin flat scale of the *Eel*, when examined microscopically, is the presence of a layer of isolated spheroidal transparent bodies, imbedded in a plate of like transparency; these, from the researches of Prof. Williamson upon other scales, appear not to be cells (as they might readily be supposed to be), but to be concretions of carbonate of lime. When the scale of the *Eel* is examined by polarized light, its surface exhibits a

beautiful St. Andrew's cross; and if a plate of selenite be placed behind it, and the analyzing prism be made to revolve, a remarkable play of colors is presented.

409. In studying the structure of the more highly-developed scales, we may take as an illustration that of the *Carp*; in which two very distinct layers can be made out by a vertical section, with a third but incomplete layer interposed between them. The outer layer is composed of several concentric laminae of a structureless transparent substance, like that of cartilage; the outermost of these laminae is the smallest, and the size of the plates increases progressively from without inwards, so

FIG. 306.


Portion of Skin of *Sole*, viewed as an opaque object.

FIG. 307.

Scale of *Sole*, viewed as a transparent object.

that their margins appear on the surface as a series of concentric lines; and their surfaces are thrown into ridges and furrows, which commonly have a radiating direction. The inner layer is composed of numerous laminae of a fibrous structure, the fibres of each lamina being inclined at various angles to those of the lamina above and below it. Between these two layers is interposed a stratum of calcareous concretions, resembling those of the scale of the Eel; these are sometimes globular or spheroidal, but more commonly "lenticular," that is, having the form of a double convex lens. The scales which resemble those of the *Carp* in having a form more or less circular, and in being destitute of comb-like prolongations, are called *cycloid*; and such are the characters of those of the salmon, herring, roach, &c. The structure of the "ctenoid" scales (Fig. 307), which we find in the sole, perch, pike, &c., does not differ essentially from that of the "cycloid," save as to the projection of the comb-like teeth from the posterior margin; and it does not appear that the strongly-marked division which Prof. Agassiz has attempted to establish between the "cycloid" and the "ctenoid" orders of fishes, on the basis of this difference, is in harmony with their general organization. Scales of either kind may become consolidated to a considerable extent, by the calcification of their soft substance; but still they never present any approach to the

true bony structure, such as is shown in the two orders to be next adverted to. In the *ganoid* scales, on the other hand, the whole substance of the scale is composed of a substance which is essentially bony in its nature; its intimate structure being almost always comparable to that of one or other of the varieties which present themselves in the bones of the Vertebrate skeleton; and being very frequently identical with that of the bones of the same fish, as is the case with the *Lepidosteus* (Fig. 301), one of the few existing representatives of this order, which, in former ages of the Earth's history, comprehended a large number of important families. Their name (from γάνος splendor) is bestowed on account of the smoothness, hardness, and high polish of the outer surface of the scales; which is due to the presence of a peculiar layer that has been likened (though erroneously) to the enamel of teeth, and is now distinguished as *ganoin*. The scales of this order are for the most part angular in their form; and are arranged in regular rows, the posterior edges of each slightly overlapping the anterior ones of the next, so as to form a very complete defensive armor to the body. The scales of the *placoid* type, which characterizes the existing Sharks and Rays, with their fossil analogues, are irregular in their shape, and very commonly do not come into mutual contact; but are separately imbedded in the skin, projecting from its surface under various forms. In the Rays, each scale usually consists of a flattened plate of a rounded shape, with a hard spine projecting from its centre; in the Sharks (to which tribe belongs the "dog-fish" of our own coast) the scales have more of the shape of teeth. This resemblance is not confined to external form; for their intimate structure strongly resembles that of dentine, their dense substance being traversed by tubuli, which extend from their centre to their circumference in minute ramifications, without any trace of osseous lacunæ. These tooth-like scales are often so small, as to be invisible to the naked eye; but they are well seen by drying a piece of the skin to which they are attached, and mounting it in Canada balsam; and they are most brilliantly shown by the assistance of polarized light. A like structure is found to exist in the spiny rays of the dorsal fin, which, also, are parts of the dermal skeleton; and these rays usually have a central cavity filled with medulla, from which the tubuli radiate towards the circumference. This structure is very well seen in thin sections of the fossil spiny rays, which, with the teeth and scales, are often the sole relics of the vast multitudes of Sharks that must have swarmed in the ancient seas, their cartilaginous internal skeletons having entirely decayed away. In making sections of bony scales, spiny rays, &c., the method must be followed which has been already detailed under the head of bone (§ 405).

410. The *scales* of Reptiles, the *feathers* of Birds, and the *hairs*, *hoofs*, *nails*, *claws*, and *horns* (when not bony) of Mammals, are

all *Epidermic* appendages; that is, they are produced upon the surface, not within the substance, of the true skin, and are allied in structure to the Epidermis (§ 419); being essentially composed of aggregations of cells, filled with horny matter; and frequently much altered in form. This structure may generally be made out in horns, nails, &c., with little difficulty, by treating thin sections of them with a dilute solution of soda; which after a short time causes the cells that had been flattened into scales, to resume their globular form. The most interesting modifications of this structure are presented to us in Hairs and in Feathers; which forms of clothing are very similar to each other in their essential structure, and are developed in the same manner,—namely, by an increased production of epidermic cells at the bottom of a flask-shaped follicle, which is formed in the substance of the true skin, and which is supplied with abundance of blood by a special distribution of vessels to its walls. When a hair is pulled out “by its root,” its base exhibits a bulbous enlargement, of which the exterior is tolerably firm, whilst its interior is occupied by a softer substance, which is known as the “pulp;” and it is to the continual augmentation of this pulp in the deeper part of the follicle, and to its conversion into the peculiar substance of the hair when it has been pushed upwards to its narrow neck, that the growth of the hair is due. The same is true of feathers, the stems of which are but hairs on a larger scale; for the “quill” is the part contained within the follicle, answering to the “bulb” of the hair; and whilst the outer part of this is converted into the peculiarly solid horny substance forming the “barrel” of the quill, its interior is occupied, during the whole period of the growth of the feather, with the soft pulp, only the shrivelled remains of which, however, are found within it after the quill has ceased to grow.

411. Although the *Hairs* of different Mammals differ greatly in the appearances they present, we may generally distinguish in them two elementary parts; namely, a *cortical* or investing substance, of a dense horny texture, and a *medullary* or pith-like substance, usually of a much softer texture, occupying the interior. The former can sometimes be distinctly made out to consist of flattened scales arranged in an imbricated manner, as in some of the hairs of the *Sable* (Fig. 308); whilst in the same hairs, the medullary substance is composed of large spheroidal cells. In the *Musk-deer*, on the other hand, the cortical substance is nearly undistinguishable; and almost the entire hair seems made up of thin-walled polygonal cells (Fig. 309). The hair of the *Reindeer*, though much larger, has a very similar structure; and its cells, except near the root, are occupied with air alone, so as to seem black by transmitted light, except when penetrated by the fluid in which they are mounted. In the hair of the *Mouse*, *Squirrel*, and other small Rodents (Fig. 310, A, B), the cortical substance forms a tube, which we see crossed at

intervals by partitions that are sometimes complete, sometimes only partial; these are the walls of the single or double line of cells, of which the medullary substance is made up. The hairs of the *Bat* tribe are commonly distinguished by the projections on their surface, which are formed by extensions of the component scales of the cortical substance; these are particularly well seen in the hairs of one of the Indian species, which has a set of whorls of long narrow leaflets (so to speak) arranged at regular intervals on the stem (c). In the hair of *Pecari* (Fig. 311), the cortical envelope sends inwards a set of radial prolongations, the interspaces of which are occupied by the polygonal cells of the medullary substance; and this, on a larger scale, is the structure of the "quills" of the *Porcupine*; the radiating partitions of which, when seen through the more transparent parts of the cortical sheath, give to the surface of the latter a fluted appearance. The hair of the *Ornithorhynchus* is a very curious object; for whilst the lower part of it resembles the fine hair of the mouse or squirrel, this thins away and then dilates again into a very

thick fibre, having a central portion composed of polygonal cells, enclosed in a flattened sheath of a brown fibrous substance. The structure of the *Human* hair is in certain respects peculiar. When its outer surface is examined, it is seen to be traversed by irregular lines (Fig. 312, A), which are most strongly marked in foetal hairs; and these are the indications of the imbricated arrangement of the flattened cells or scales which form the cortical layer. This layer, as is shown by transverse sections (c, d), is a very thin and transparent cylinder; and it incloses the peculiar fibrous substance, that constitutes the principal part of the shaft of the hair. The constituent fibres of this substance, which

FIG. 308.



FIG. 309.



FIG. 310.





Fig. 308. Hair of *Sable*, showing large rounded cells in its interior, covered by imbricated scales or flattened cells.

Fig. 309. Hair of *Musk-deer*, consisting almost entirely of polygonal cells.

Fig. 310. A, Small Hair of *Squirrel*; B, Large Hair of *Squirrel*; C, Hair of *Indian Bat*.

thick fibre, having a central portion composed of polygonal cells, enclosed in a flattened sheath of a brown fibrous substance. The structure of the *Human* hair is in certain respects peculiar.

When its outer surface is examined, it is seen to be traversed by irregular lines (Fig. 312, A), which are most strongly marked in foetal hairs; and these are the indications of the imbricated arrangement of the flattened cells or scales which form the cortical layer. This layer, as is shown by transverse sections (c, d), is a very thin and transparent cylinder; and it incloses the peculiar



Transverse Section of Hair of *Pecari*.

fibrous substance, that constitutes the principal part of the shaft of the hair. The constituent fibres of this substance, which

are marked out by the delicate striæ that may be traced in longitudinal sections of the hair (B), may be separated from each other by crushing the hair, especially after it has been macerated for some time in sulphuric acid; and each of them, when completely separated from its fellows, is found to be a long spindle-shaped



Structure of *Human Hair*:—A, external surface of the shaft, showing the transverse striæ and jagged boundary caused by the imbrications of the cortical substance; B, longitudinal section of the shaft, showing the fibrous character of the medullary substance, and the arrangement of the pigmentary matter; C, transverse section, showing the distinction between the transparent envelope, the cylinder of medullary substance, and the cellular centre; D, another transverse section showing deficiency of central cellular substance.

cell. In the axis of this fibrous cylinder, there is very commonly a band which is formed of spheroidal cells; but this is usually deficient in the fine hairs scattered over the general surface of the body, and is not always present in those of the head.¹ The hue of the hair is due, partly to the presence of pigmentary granules, either collected into patches, or diffused through its substance; but partly also to the existence of a multitude of minute air-spaces, which cause it to appear as dark by transmitted and white by reflected light. The cells of the axis-band, in particular, are very commonly found to contain air, giving it the black appearance shown at c. The difference between the blackness of pigment and that of air-spaces, may be readily determined by attending to the characters of the latter as already laid down (§§ 98, 99); and by watching the effects of the penetration of oil of turpentine or other liquids, which do not alter the appearance of pigment spots, but obliterate all the markings produced by air-spaces, these returning again as the hair dries. In mounting hairs as microscopic preparations, they should in the first instance be cleansed of all their fatty matter by maceration in ether; and they may then be put up, either in weak spirit or in

¹ Several writers regard this band of polygonal cells as the "medullary" substance, and the fibrous structure which forms the principal body of the hair, as the "cortical" substance; the transparent sheath receiving some separate designation. To the Author, however, it appears perfectly clear that the transparent horny sheath, with its lines of imbrication, is the representative of the cortical substance of other hairs; and that its entire contents, whether polygonal cells or cells elongated into fusiform fibres, must be considered as equivalent to their medullary substance.

Canada balsam, as may be thought preferable, the former menstruum being well adapted to display the characters of the finer and more transparent hairs, while the latter allows the light to penetrate more readily through the coarser and more opaque. Transverse sections of hairs are best made by gluing or gumming several together, and then putting them into the section-instrument; those of human hair may be easily obtained, however, by shaving a second time, very closely, a part of the surface over which the razor has already passed more lightly, and by picking out from the lather, and carefully washing, the sections thus taken off.

412. The stems of *Feathers* exhibit the same kind of structure as hairs; their cortical portion being the horny sheath that envelopes the shaft, and their medullary portion being their pith-like substance which that sheath includes. In small feathers, this may usually be made very plain by mounting them in Canada balsam; in large feathers, however, the texture is sometimes so altered by the drying up of the pith (the cells of which are always found to be occupied by air alone), that the cellular structure cannot be demonstrated, save by boiling thin slices in a dilute solution of potass, and not always even then. In small feathers, especially such as have a downy character, the cellular structure is very distinctly seen in the *laminae* or "barbs," which are sometimes found to be composed of single files or pear-shaped cells, laid end to end; but in larger feathers, it is usually necessary to increase the transparency of the barbs, especially when these are thick and little pervious to light, either by soaking them in turpentine, mounting them in Canada balsam, or boiling them in a weak solution of potass. In the feathers which are destined to strike the air with great force in the act of flight, we find the barbs fringed on each side with hair-like filaments or *pinnae*; on one side of each barb these filaments are toothed on one edge, whilst on the other side they are furnished with curved hooks; and as the two sets of *pinnae* which spring from two adjacent barbs, cross one another at an angle, and each hooked *pinna* on one locks into the teeth of several of the toothed *pinnae* arising from the other, the barbs are connected together very firmly by this apparatus of "hooks and eyes," which reminds us of that already mentioned as to be observed on the wings of Hymenopterous Insects (§ 395). Feathers or portions of feathers of birds distinguished by the splendor of their plumage, are very good objects for low magnifying powers, when illuminated on an opaque ground; but care must be taken that the light falls upon them at the angle necessary to produce their most brilliant reflection into the axis of the microscope; since feathers which exhibit the most brilliant metallic lustre to an observer at one point, may seem very dull to the eye of another in a different position. The small feathers of Humming birds, portions of the feathers of the Peacock, and others of a like kind, are well worthy of an examination; and the scientific


Microscopist who is but little attracted by mere gorgeousness, may well apply himself to the discovery of the peculiar structure, which imparts to these objects their most remarkable character.

413. Sections of *Horns, Hoofs, Claws*, and other like modifications of Epidermic structure,—which may be made by the Section instrument (§ 107), the substance to be cut having been softened, if necessary, by soaking in warm water,—do not in general afford any very interesting features, when viewed in the ordinary mode; but there are no objects on which Polarized light produces more remarkable effects, or which display a more beautiful variety of colors, when a plate of selenite is placed behind them, and the analyzing prism is made to rotate. A curious modification of the ordinary structure of horn, is presented in the appendage borne by the *Rhinoceros* upon its snout, which in many points resembles a bundle of hairs, its substance being arranged in minute cylinders around a number of separate centres, which have probably been formed by independent papillæ (Fig. 313). When transverse sections of these cylinders are viewed by polarized light, each of them is seen to be marked by a cross, somewhat resembling that of starch-grains; and the lights and shadows of this cross are replaced by contrasted colors, when the selenite-plate is interposed. The substance commonly but erroneously termed *Whalebone*, which is formed from the surface of the membrane that lines the mouth of the whale, and has no relation to its true bony skeleton, is almost identical in structure with *Rhinoceros-horn*, and is similarly affected by polarized light. The central portion of each of its component fibres, like the medullary substance of hairs, contains cells that have been so little altered as to be easily recognized; and the

outer or cortical portion also may be shown to have a like structure, by macerating it in a solution of potass, and then in water. Sections of any of the horny tissues are best mounted in Canada balsam.

414. *Blood*.—Carrying our Microscopic survey, now, to the elementary parts of which those softer tissues are made up, that are subservient to the active life of the body, rather than to its merely mechanical requirements, we shall in the first place notice the isolated floating cells contained in the Blood, and

FIG. 313.



Transverse Section of Horn of *Rhinoceros*, viewed by Polarized Light.

known as the "blood-corpuscles." These are of two kinds; the

“red,” and the “white” or “colorless.” The former present, in every instance, the form of a flattened disk, which is circular in Man and in most Mammalia (Fig. 315), but which is oval in Birds, Reptiles (Fig. 314), and Fishes, and in a few Mammals (all belonging to the *Camel* tribe); in the one form as in the other, this disk is a flattened cell, whose walls are pellucid and colorless, but whose contents are colored. They may be caused to swell up and burst, however, by the imbibition of water; and the perfect transparency and the homogeneous character of their walls then become evident. The “red corpuscles” in the blood of Oviparous Vertebrata are distinguished by the presence of a distinct central spot or *nucleus*, which appears to be composed of an aggregation of minute granules; this is most distinctly brought into view by treating the blood-disks with acetic acid, which renders the remaining portion extremely transparent, while it increases the opacity of the nucleus (Fig. 314, *d*). It is remarkable, however, that the “red corpuscles” of the blood of Mammals should possess no obvious nucleus; the dark spot which is seen in their centre (Fig. 315, *b*), being merely an effect of refrac-

FIG. 314.



FIG. 315.




Fig. 314. Red Corpuscles of *Frog's* Blood:—1, 1, their flattened face; 2, particle turned nearly edgewise; 3, colorless corpuscle; 4, red corpuscles altered by dilute acetic acid.

Fig. 315. Red Corpuscles of *Human* Blood; represented at *a*, as they are seen when rather within the focus of the microscope, and at *b*, as they appear when precisely in the focus.

tion, consequent upon the double concave form of the disk. When the corpuscles are treated with water, so that their form becomes first flat, and then double convex, the dark spot disappears; whilst, on the other hand, it is made more evident when the concavity is increased by the partial emptying of the cell, which may be accomplished by treating the blood-corpuscles with fluids of greater density than their own contents. The size of the “red corpuscles” is not altogether uniform in the same blood; thus it varies in that of Man, from about 1-4000th to the 1-2800th of an inch. But we generally find that there is an average size, which is pretty constantly maintained among the different individuals of the same species; that of Man may be stated at about 1-3200th of an inch. The following Table¹ exhibits the average dimen-

¹ These measurements are chiefly selected from those given by Mr. Gulliver in his edition of *Hewson's Works*, p. 236, *et seq.*

sions of some of the most interesting examples of the Red blood-corpuscles, in the four classes of Vertebrated Animals, expressed in fractions of an inch. Where two measurements are given, they are the long and the short diameters of the same corpuscle.

MAMMALS.

Man,	1-3200	Camel,	1-3254, 1-5921
Dog,	1-3542	Llama,	1-3361, 1-6294
Whale,	1-3099	Java Musk-Deer,	1-12325
Elephant,	1-2745	Caucasian Goat,	1-7045
Mouse,	1-3814	Two-toed Sloth,	1-2865

BIRDS.

Golden Eagle,	1-1812, 1-3832	Ostrich,	1-1649, 1-3000
Owl,	1-1830, 1-3400	Cassowary,	1-1455, 1-2800
Crow,	1-1961, 1-4000	Heron,	1-1913, 1-3491
Blue-Tit,	1-2313, 1-4128	Fowl,	1-2102, 1-3466
Parrot,	1-1898, 1-4000	Gull,	1-2097, 1-4000

REPTILES.

Turtle,	1-1231, 1-1882	Frog,	1-1108, 1-1821
Crocodile,	1-1231, 1-2286	Water-Newt,	1-814, 1-1246
Green Lizard,	1-1555, 1-2743	Siren,	1-420, 1-760
Slow-worm,	1-1178, 1-2666	Proteus,	1-337
Viper,	1-1274, 1-1800	Lepidosiren,	1-570, 1-941

FISHES.

Perch,	1-2099, 1-2824	Pike,	1-2000, 1-3555
Carp,	1-2142, 1-3429	Eel,	1-1745, 1-2842
Gold-Fish,	1-1777, 1-2824	Gymnotus,	1-1745, 1-2599

Thus it appears that the *smallest* red corpuscles known are those of the *Musk-Deer*; whilst the *largest* are those of that curious group of Batrachian (frog-like) Reptiles which retain their gills through the whole of life; and the oval blood-disks of the *Proteus*, being above 36 times as long as those of the Musk-Deer, and probably at least 20 times as broad, would cover no fewer than 720 of them.

415. The "colorless" corpuscles are more readily distinguished in the blood of Reptiles, than in that of Man; being, in the former case, of much smaller size, as well as having a circular outline (Fig. 314, *c*); whilst in the latter, their size and contour are nearly the same, so that, as the red corpuscles themselves, when seen in a single layer, have but a very pale hue, the deficiency of color does not sensibly mark their difference of nature. It is remarkable that, notwithstanding the great variations in the sizes of the red corpuscles in different species of Vertebrated animals, the size of the "colorless" is extremely constant throughout, their diameter being seldom much greater or less than 1-3000th of an inch in the warm-blooded classes, and 1-2500th in Reptiles. Their ordinary form is globular; but their aspect is subject to considerable variations, which seem to depend in great part upon their phase of development. Thus in their early state, in which they seem to be identical with the corpuscles found floating in *Chyle* and *Lymph*, the cell-wall can scarcely be dis-

tinguished from the large nuclear mass which it incloses; by treating the cell with water or acetic acid, however, the membrane is distended, and the nucleus very commonly breaks up into fragments in its interior. This last appearance seems natural to the corpuscles in a more advanced condition; and the isolated particles are often to be seen executing an active molecular movement within the cell, which continues when they are discharged by the bursting of the cell, consequent upon the addition of a solution of potass. These corpuscles are occasionally seen to exhibit very curious changes of form, which remind us of those of the *Amœba* (§ 261); a protrusion taking place from the same portion of the cell-wall, the form of which seems quite indeterminate; and this being soon succeeded by another, from some different part of the cell, the first being either drawn in again, or remaining as it was. These changes have been observed, not only in the "colorless corpuscles" of the blood of various Vertebrated animals, but also in the corpuscles floating in the circulating fluid of the higher Invertebrata, such as the Crab, which resemble the "colorless" corpuscles of Vertebrated blood rather than its "red" corpuscles,—these last, in fact, being altogether peculiar to the circulating fluid of Vertebrated animals.


416. In examining the Blood microscopically, it is, of course, of importance to obtain as thin a stratum of it as possible, so that the corpuscles may not overlies one another. This is best accomplished by selecting a piece of thin glass of perfect flatness, and then, having received a small drop of blood upon a glass slide, to lay the thin glass, not upon this, but with its edge just touching the edge of the drop; for the blood will then be drawn in by capillary attraction, so as to spread in a uniformly thin layer between the two glasses. The inexperienced observer will be surprised at the very pale hue which the red corpuscles exhibit beneath the microscope, when seen in a single stratum; but this surprise need no longer be felt, when it is borne in mind that the thickness of the film of coloring fluid which they contain, is probably not more than 1-20,000th of an inch; and if a drop of ink, or of almost any colored liquid, however dark, be pressed out between two glasses into an equally thin film, its hue will be lightened in the same degree. The red hue of the corpuscles, however, becomes obvious enough, when two or more layers of them are seen through at once. The "colorless corpuscles" in Human blood are usually not more than 1-350 of the "red," so that no more than one or two are likely to be in the field at once; and these may generally be recognized most readily, by their standing apart from the rest; for whilst the "red" corpuscles have a tendency to adhere to each other by their discoidal surfaces, the "colorless" show no such disposition. Thin films of blood may be preserved in the liquid state, with little change, by applying gold size or asphalt round the edge of the thin glass cover before evaporation has had time to take

place; but it is in some respects preferable to dilute the liquid with a small quantity of Goadby's solution, its strength being so adjusted as not to produce any endosmotic change of form in the corpuscles. But it is far simpler to allow such films to dry, without any cover, and then merely to cover them for protection; and in this condition the general characters of the corpuscles can be very well made out, notwithstanding that they have in some degree shrivelled by the desiccation they have undergone. And this method is particularly serviceable, as affording a fair means of comparison, when the assistance of the Microscopist is sought in determining, for Medico-legal purposes, the source of suspicious blood stains; the average dimensions of the dried blood-corpuscles of the several domestic animals, being sufficiently different from each other and from those of Man, to allow the nature of any specimen to be pronounced upon with a high degree of probability.

417. *Simple Fibrous Tissues*.—A large proportion of every animal fabric is made up of *simple fibres*, whose function is to hold other parts together, or to serve as cords for the communication of movement. A very beautiful example of a fibrous tissue of this kind, is furnished by the membrane of the common Fowl's egg, which (as may be seen by examining an egg whose shell remains soft for want of consolidation by calcareous particles) consists of two principal layers, one serving as the basis of the shell itself, and the other forming that lining to it, which is known as the *membrana putaminis*. The latter may be separated, by careful tearing with needles and forceps, after prolonged maceration in water, into several matted lamellæ resembling that represented in Fig. 316; and similar lamellæ may be readily obtained from the shell itself, by dissolving away its lime by dilute acid. The simply fibrous structures of the body generally, however, belong to one of two very definite kinds of tissue; the "white" and the "yellow," whose appearance, composition, and properties are very different. The *white* fibrous tissue, though sometimes apparently composed of distinct fibres, more commonly presents the aspect of bands, usually of a flattened form, and attaining the breadth of 1-500th of an inch, which are marked by numerous longitudinal streaks, but can seldom be torn up into minute fibres of determinate size. The fibres and bands are occasionally somewhat wavy in their direction; and they have a peculiar tendency to fall into undulations, when it is attempted to tear them apart from each other (Fig. 317). This tissue is easily distinguished from the other, by the effect of acetic acid, which swells it up and renders it transparent, at the same time bringing into view certain oval nuclear corpuscles. It is perfectly inelastic; and we find it in such parts as tendons, ordinary ligaments, fibrous capsules, &c., whose function it is to resist tension without yielding to it. The *yellow* fibrous tissue exists in the form of long, single, elastic, branching fila-

ments, with a dark decided border; which are disposed to curl when not put on the stretch (Fig. 318). They are for the most part between 1-5000th and 1-10,000th of an inch in diameter; but they are often met with both larger and smaller. They frequently anastomose, so as to form a network. This tissue does not undergo any change, when treated with acetic

FIG. 316.



Fibrous membrane from Egg-shell.


FIG. 317.



White Fibrous Tissue from Ligament.

acid. It exists alone (that is, without any mixture of the white) in parts which require a peculiar elasticity, such as the middle coat of the Arteries, the Vocal Cords, the "ligamentum nuchæ" of Quadrupeds, the elastic ligament which holds together the valves of a Bivalve shell, and that by which the claws of the Feline tribe are retracted when not in use; and it enters largely into the composition of *Areolar* tissue. This consists of a network of minute fibres and bands, which are interwoven in every direction, so as to leave innumerable *areolæ* or little spaces, which communicate freely with one another. Of these fibres, some

FIG. 318.



Yellow Fibrous Tissue from Ligamentum Nuchæ of Calf.

are of the yellow or elastic kind; but the majority are composed of the white fibrous tissue; and, as in that form of elementary structure, they frequently present the form of broad flattened bands, or membranous shreds, in which no distinct fibrous arrangement is visible. The proportion of the two forms varies, according to the amount of elasticity, or of simple resisting power, which the endowments of the part may require. We find this tissue in a very large proportion of the bodies of higher Animals; thus it binds together the ultimate fibres of the muscles and nerves into minute fasciculi, unites these fasciculi into larger ones, these again into still larger ones which are

obvious to the eye, and these into the entire muscle; whilst it also forms the membranous divisions between distinct muscles. In like manner it unites the elements of nerves, glands, &c., binds together the fat-cells into minute masses, these into larger ones, and so on; and in this way it penetrates and forms part of all the softer organs of the body. For the display of the characters of these tissues, small and thin shreds may be cut with the curved scissors from any part that affords them; and these must be torn asunder with needles under the simple microscope, until the fibres are separated to a degree sufficient to enable them to be examined to advantage under a higher magnifying power.

418. *Skin, Mucous and Serous Membranes.*—The Skin which forms the external envelope of the body, is divisible into two principal layers; the “true skin,” which usually makes up by far the larger part of its thickness, and the “cuticle,” “scarf skin,” or Epidermis, which covers it. At the mouth, nostrils, and other orifices of the open cavities and canals of the body, the skin passes into the membrane that lines these, which is distinguished as the Mucous membrane, from the peculiar glairy secretion of mucus by which its surface is protected. But the great closed cavities of the body, which surround the heart, lungs, intestines, &c., are lined by membranes of a different kind; which, as they secrete only a thin serous fluid from their surfaces, are known as Serous membranes. Both Mucous and Serous membranes consist, like the skin, of a proper membranous basis, and of a thin cuticular layer, which, as it differs in many points from the epidermis, is distinguished as the Epithelium (§ 421). The substance of the “true skin” and of the “mucous” and “serous” membranes, is principally composed of the fibrous tissues last described; but the skin and the mucous membranes are very copiously supplied with bloodvessels and with glandulæ of various kinds; and in the skin we also find abundance of nerves and lymphatic vessels, as well as, in some parts, of hair-follicles. The distribution of the vessels in the skin and mucous membranes, which is one of the most interesting features in their structure, and which will come under our notice hereafter (Figs. 328, 329), is intimately connected with their several functions. In serous membranes, on the other hand, the supply of bloodvessels is more scanty, their function being simply protective.

419. *Epidermic and Epithelial Cell-layers.*—The Epidermis or “cuticle” covers the exterior surfaces of the body, as a thin semi-transparent pellicle, which is shown by microscopic examination to consist of a series of layers of cells, which are continually wearing off at the external surface, and are being renewed at the surface of the true skin; so that the newest and deepest layers gradually become the oldest and most superficial, and are at last thrown off by slow desquamation. In their progress from

the internal to the external surface of the Epidermis, the cells undergo a series of well-marked changes. When we examine the innermost layer, we find it soft and granular; consisting of *nuclei*, in various stages of development into cells, held together by a tenacious semi-fluid substance. This was formerly considered as a distinct tissue, and was supposed to be the peculiar seat of the color of the skin; it received the designation of *rete mucosum*. Passing outwards, we find the cells more completely formed; at first nearly spherical in shape, but becoming polygonal where they are flattened one against another. As we proceed further towards the surface, we perceive that the cells are gradually more and more flattened, until they become mere horny scales, their cavity being obliterated; their origin is indicated, however, by the nucleus in the centre of each. This change in form is accompanied by a change in the chemical composition of the tissue, which seems to be due to the metamorphosis of the contents of the cells, into a horny substance identical with that of which hair, horn, nails, hoofs, &c., are composed. Mingled with the epidermic cells, we find others which secrete coloring-matter instead of horn; these are termed "pigment-cells." The most remarkable development of "pigment-cells" in the higher animals, is on the inner surface of the Choroid coat of the eye, where they have a very regular arrangement, and form several layers, known as the *Pigmentum nigrum*. When examined separately, these cells are found to have a polygonal form (Fig. 319, *a*), and to have a distinct nucleus (*b*) in their interior. The black color is given by the accumulation, within the cell, of a number of flat, rounded, or oval granules, of extreme minuteness, which exhibit an active movement when set free from the cell, and even whilst enclosed within it. The pigment-cells are not always, however, of this simply-rounded or polygonal form; they sometimes present remarkable stellate prolongations, under which form they are well seen in the skin of the Frog (Fig. 327, *c, c*). The gradual formation of these prolongations may be traced in the pigment-cells of the Tadpole during its metamorphosis (Fig. 320). Similar varieties of form are to be met with in the pigmentary cells of Fishes and small Crustacea, which also present a great variety of hues; and these seem to have the power of likening their color to that of the bottom over which the animal may live, so as to serve for its concealment.

420. The structure of the Epidermis may be examined in a variety of ways. If it be removed by maceration from the true skin, the cellular nature of its under-surface is at once recognized,

FIG. 319.



Cells from *Pigmentum Nigrum*:—*a*, pigmentary granules concealing the nucleus; *b*, the nucleus distinct.

when it is subjected to a magnifying power of 200 or 300 diameters, by light transmitted through it, with this surface uppermost; and if the epidermis be that of a negro or any other dark-skinned race, the pigment-cells will be very distinctly seen. This under-surface of the epidermis is not flat, but is excavated into pits and channels for the reception of the papillary elevations of the true skin; an arrangement which is shown on a large scale in the thick cuticular covering of the Dog's foot, the subjacent papillæ being large enough to be distinctly seen (when injected) with the naked eye. The cellular nature of the newly-formed layers is best seen, by examining a little of the soft film that is found upon the surface of the true skin, after the more consistent layers of the cuticle have been raised by a blister. The alteration which the cells of the external layers have undergone, tends to obscure their character; but if any fragment of epidermis be macerated for a little time in a weak solution of soda or potass, its dry scales become softened, and are filled out by imbibition into rounded or polygonal cells. The same mode of treatment enables




Pigment-cells from tail of Tadpole:—*a*, simple forms of recent origin; *b*, more complex forms subsequently assumed.

us to make out the cellular structure in warts and corns, which are epidermic growths from the surface of papillæ enlarged by hypertrophy.

421. The Epithelium may be designated as a delicate cuticle, covering all the free *internal* surfaces of the body, and thus lining all its cavities, canals, &c. Save in the mouth and other parts in which it approximates to the ordinary cuticle both in locality and in nature, its cells usually form but a single layer; and are so deficient in tenacity of mutual adhesion, that they cannot be detached in the form of a continuous membrane. Their shape varies greatly; for sometimes they are broad, flat, and scale-like, and their edges approximate closely to each other, so as to form what is termed a "pavement" or "tessellated" epithelium; such cells are observable on the web of a frog's foot, or on the tail of the tadpole; for, though covering an external surface, the soft moist cuticle of these parts has all the characters of an epithelium. In other cases, the cells have more of the form of cylinders, standing erect side by side, one extremity of each cylinder forming part of the free surface, whilst the other rests upon the membrane to which it serves as a covering. If the cylinders be closely pressed together, their form is changed into prisms; and such epithelium is often known as "prismatic." On the other hand, if the surface on which it rests be convex, the bases or lower ends of the cylinders become smaller than their free extremities; and thus each has the form of a truncated cone rather than of a cylin-

der, and such epithelium (of which that covering the *villi* of the intestine, Fig. 328, is a peculiarly-good example) is termed "conical." But between these primary forms of epithelial cells, there are several intermediate gradations; and one often passes almost insensibly into the other. Any of these forms of Epithelium may be furnished with *cilia*; but these appendages are more commonly found attached to the elongated, than to the flattened forms of epithelium-cells (Fig. 321). "Ciliated epithelium" is found upon the lining membrane of the air-passages in all air-breathing Vertebrata; and it also presents itself in many other situations, in which a propulsive power is needed to prevent an accumulation of mucous or other secretions. Owing to the very slight attachment that usually exists between the epithelium and the membranous surface whereon it lies, there is usually no difficulty whatever in examining it; nothing more being necessary than to scrape the surface of the membrane with a knife, and to add a little water to what has been thus removed. The ciliary action will generally be found to persist for some hours or even days after death, if the animal has been previously in full vigor;¹ and the cells that bear the cilia, when detached from each other, will swim freely about in water. If the thin fluid that is copiously discharged from the nose in the first stage of an ordinary "cold in the head," be subjected to microscopic examination, it will commonly be found to contain a great number of ciliated epithelium-cells that have been thrown off from the lining membrane of the nasal passages.




Ciliated Epithelium; a, nucleated cells, resting on their smaller extremities; b, cilia.

422. *Fat*.—One of the best examples which the bodies of higher animals afford, of a tissue composed of an aggregation of cells, is presented by the Adipose tissue; the cells of which are distinguished by their power of drawing into themselves oleaginous matter from the blood. Fat-cells are sometimes dispersed in the interspaces of Areolar tissue; whilst in other cases they are aggregated in distinct masses,—constituting the proper Adipose tissue. The individual fat-cells always present a nearly spherical or spheroidal form; sometimes, however, when they are closely pressed together, they become somewhat polyhedral, from the flattening of their walls against each other (Fig. 322). Their intervals are traversed by a minute network of blood-vessels, from which they derive their secretion; and it is probably by the constant moistening of their walls with a *watery* fluid, that their contents are retained without the least transudation,

¹ Thus it has been observed in the lining of the windpipe of a decapitated criminal, as much as seven days after death; and in that of the river-tortoise, it has been seen fifteen days after death, even though putrefaction had already far advanced.

though these are quite fluid at the temperature of the living animal, but when filled with their characteristic contents,



have the peculiar appearance which has been already described as appertaining to oil-globules (§ 99), being very bright in their centre, and very dark towards their margin, in consequence of their high refractive power; but if, as often happens in preparations that have been long mounted, the oily contents should have escaped, they then look like any other cells of the same form. Although the fatty matter which fills these cells (consisting of a mixture of stearine or of margarine with oleine) is liquid at the ordinary temperature of the body of a warm-blooded animal, yet its harder portion sometimes crystallizes on cooling; the crystals shooting from a centre, so as to form a star-shaped cluster. In examining the structure

of Adipose tissue, it is desirable, where possible, to have recourse to some specimen in which the fat is deposited in vessels, and in which they can be observed without the necessity of opening them open: such a condition is found in the mesentery of the mouse, and it is also occasionally seen in the fat deposits which present themselves at the junctions of the connective tissues of the muscles, joints, &c. In such cases, the fat cells are found in the deeper layers of the mesentery, and are brought into view by vertical sections of the mesentery. In large masses of fat may be examined in a similar manner, being placed under water in thin cells, so as to prevent the escape of the thin glass from their surface, and to prevent the escape of the oil particles. No method (as far as the Author is aware) is successful in preparing such specimens permanently to retain their contents.

Cartilage.—In the ordinary forms of Cartilage, also, we find a tissue essentially composed of cells; but these are not so separated from each other by an intercellular substance, the thickness of which differs greatly in different species of cartilage, and even in different stages of the growth of the same species. In the cartilage of the external ear of a *Bat* or *Blow-bee* (Fig. 29), the cells are packed as closely together as are those of ordinary vegetable parenchyma (Fig. 150, A); and they therefore present the early condition of most cartilages that are known to us. In the ordinary cartilages, however, that cover the extremities of the bones, so as to present smooth surfaces for the working of the joints, the amount of intercellular substance is usually considerable; and the cartilage is commonly found imbedded in this, in clusters of cells (see, for example, Fig. 324), which are evidently formed by

a process of "duplicative subdivision" analogous to that by which the multiplication of cells takes place in the Vegetable Kingdom (Fig. 67). The substance of these *cellular Cartilages* is entirely destitute of blood-vessels; being nourished solely by imbibition from

FIG. 323.




FIG. 324.




Fig. 323. Cellular Cartilage of Mouse's ear.

Fig. 324. Section of the branchial Cartilage of Tadpole:—a, group of four cells, separating from each other; b, pair of cells in apposition; c, c', nuclei of cartilage-cells; d, cavity containing three cells.

the blood brought to the membrane covering their surface. Hence they may be compared, in regard to their grade of organization, with the larger Algæ; which consist, like them, of aggregations of cells held together by intercellular substance, without vessels of any kind, and are nourished by imbibition through their whole surface. There are many cases, however, in which the structureless intercellular substance is replaced by bundles of fibres, sometimes elastic, but more commonly non-elastic; such combinations, which are termed *fibro-cartilages*, are interposed in certain joints, wherein tension as well as pressure has to be resisted, as, for example, between the vertebræ of the spinal column, and the bones of the pelvis. In examining the structure of Cartilage, nothing more is necessary than to make very thin sections with a sharp razor or scalpel, or with a Valentin's knife (§ 106), or, if the specimen be large and dense (as the cartilage of the ribs), with the section-instrument (§ 107). These sections may be mounted in weak spirit, in Goadby's solution, or in glycerine; but in whatever way they are mounted, they undergo a gradual change by the lapse of time, which renders them less fit to display the characteristic features of their structure.

424. *Structure of Glands.*—The various secretions of the body (as the saliva, bile, urine, &c.) are formed by the instrumentality of organs termed Glands; which are, for the most part, formed on one fundamental type, whatever be the nature of their product. The simplest idea of a gland is that which we gain from an examination of the "follicles" or little bags imbedded in the wall of the stomach; some of which secrete mucus for the protection of its surface, and others gastric juice. These little bags are filled with cells of a spheroidal form, which may be consi-

dered as constituting their epithelial lining; these cells, in the progress of their development, draw into themselves from the blood the constituents of the particular product they are to secrete; and they then seem to deliver it up, either by the bursting or by the melting away of their walls, so that this product may be poured forth from the mouth of the bag, into the cavity in which it is wanted. The liver itself, in the lowest animals wherein it is found, presents this condition. Some of the cells that form the lining of the stomach in the Hydra and Actinia, seem to be distinguished from the rest by their power of secreting bile, which gives them a brownish-yellow tinge; in many Polyzoa, Compound Tunicata, and Annelida, these biliary cells can be seen to occupy follicles in the walls of the stomach; in Insects, these follicles are few in number, but are immensely elongated so as to form biliary tubes, which lie loosely within the abdominal cavity, frequently making many convolutions within it, and discharge their contents into the commencement of the intestinal canal; whilst in the higher Mollusca, and in Crustacea, the follicles are vastly multiplied in number, and are connected with the ramifications of gland ducts, like grapes upon the stalks of their bunch, so as to form a distinct mass, which now becomes known as the liver. The examination of the biliary tubes of the Insect, or of the biliary follicles of the Crab, which may be accomplished with the utmost facility, is well adapted to give an idea of the essential nature of glandular structure. Among Vertebrated animals, the salivary glands, the pancreas (sweetbread), and the mammary glands, are well adapted to display the follicular structure; nothing more being necessary than to make sections of these organs, thin enough to be viewed as transparent objects. The liver of Vertebrata, however, presents certain peculiarities of structure, which are not yet fully understood; for although it is essentially composed, like other glands, of secreting cells, yet it has not yet been determined beyond doubt, whether these cells are contained within any kind of membranous investment. The kidneys of Vertebrated animals are made up of elongated tubes, which are straight, and lined with a pavement-epithelium, in the inner or "medullary" portion of the kidney, whilst they are convoluted, and filled with a spheroidal epithelium, in the outer or "cortical." Certain flask-shaped dilatations of these tubes include curious little knots of bloodvessels, which are known as the "Malpighian bodies" of the kidney; these are well displayed in injected preparations. For such a full and complete investigation of the structure of these organs as the Anatomist and Physiologist require, various methods must be put in practice, which this is not the place to detail. It is perfectly easy to demonstrate the cellular nature of the substance of the liver, by simply scraping a portion of its cut surface; since a number of its cells will be then detached. The general arrangement of the cells in the lobules, may be shown by means of sections thin enough to be

transparent; whilst the arrangement of the bloodvessels can only be shown by means of injections (§ 433). Fragments of the tubules of the kidney, sometimes having the Malpighian capsules in connection with them, may also be detached by scraping its cut surface; but the true relations of these parts can only be shown by thin transparent sections, and by injections of the bloodvessels and tubuli. The simple follicles contained in the walls of the stomach are brought into view by vertical sections; but they may be still better examined by leaving small portions of the lining membrane for a few days in dilute nitric acid (one part to four of water), whereby the fibrous tissue will be so softened, that the clusters of glandular epithelium lining the cells (which are but very little altered) will be readily separated.

425. *Muscular Tissue*.—Although we are accustomed to speak of this tissue as consisting of “fibres,” yet the ultimate structure of the “muscular fibre” is very different from that of the simple fibrous tissues already described. When we examine an ordinary Muscle (or piece of “flesh”) with the naked eye, we observe that it is made up of a number of *fasciculi* or bundles of fibres; which are arranged side by side with great regularity, in the direction in which the muscle is to act; and which are united by areolar tissue. These fasciculi may be separated into smaller parts, which appear like simple fibres; but when these are examined by the microscope, they are found to be themselves fasciculi, composed of minuter fibres bound together by delicate filaments of areolar tissue. By carefully separating these, we may obtain the ultimate “muscular fibre.” This fibre exists under two forms, the *striated* and the *non-striated*. The former is chiefly distinguished by the transversely-striated appearance which it presents, and which is due to an alteration of light and dark spaces along its whole extent; the breadth and distance of those striæ vary, however, in different fibres, and even in different parts of the same fibre, according to its state of contraction or relaxation. Longitudinal striæ are also frequently visible, which are due to a partial separation between the component fibrillæ into which the fibre may be broken up. When a fibre of this kind is more closely examined, it is seen to consist of a delicate tubular sheath, quite distinct on the one hand from the areolar tissue which binds the fibres into fasciculi, and equally distinct from the internal substance of the fibre. This membranous tube, which has been termed the *myolemma*, is not perforated either by nerves or capillary vessels; and forms, in fact, a complete barrier between the real elements of Muscular structure, and the surrounding parts. These elements appear to be very minute cylindrical particles with flattened faces of nearly uniform size, and adherent to each other both by their flat surfaces and by their edges. The former adhesion is usually the most powerful; and causes the substance of the fibre, when it is broken up, to present itself in the form of delicate *fibrillæ*, each

of which is composed of a single row of the primitive particle (Fig. 325). The diameter of the fibres varies greatly in different

Fig. 325.



Striated Muscular fibre, separating into fibrillae.

kinds of Vertebrated animals. Its average is greater in Reptiles and Fishes than in Birds and Mammals, and its extremes also are wider; thus its dimensions vary in the Frog from 1-100th to 1-1000th of an inch, and in the Skate from 1-65th to 1-300th; whilst in the Human subject, the average is about 1-400th of an inch,

and the extremes about 1-200th and 1-600th.

426. When the fibrillae are separately examined, under a magnifying power of from 250 to 400 diameters, they are seen

Fig. 326.



Structure of the ultimate Fibrillae of Striated Muscular fibre:—*a*, a fibril in a state of ordinary relaxation; *b*, a fibril in a state of partial contraction.

to present a cylindrical or slightly beaded form; and their linearly aggregated particles then appear to be minute *cells*. We observe the same alternation of light and dark spaces, as when the fibrillae

are united into fibres or into small bundles; but it may be distinctly seen, that each light space is divided by a transverse line; and that there is a pellucid border at the *sides* of the dark spaces, as well as between their contiguous extremities (Fig. 326). This pellucid border seems to be the cell-wall; the dark space enclosed by it (which is usually bright in the centre) being the cavity of the cell, which is filled with a highly refracting substance. When the fibril is in a state of relaxation, as seen at *a*, the diameter of the cells is greatest in the longitudinal direction: but when it is contracted, the fibril increases in diameter as it diminishes in length; so that the transverse diameter of each cell becomes equal to the longitudinal diameter, as seen at *b*; or even exceeds it.

Thus the act of Muscular contraction seems to consist in a change of form in the cells of the ultimate fibrillae, consequent upon a contraction between the walls of their two extremities; and it is interesting to observe, how very closely it thus corresponds with the contraction of certain Vegetable tissues, of which the component cells are

capable of producing movements, when they are irritated, by means of a similar change of form. The diameter of the ultimate fibrillae will of course be subject to variations, in accordance with their contracted or relaxed condition; but it seems to be otherwise tolerably uniform in different animals, being for

the most part about 1-10,000th of an inch. It has been observed, however, as high as 1-5000th of an inch, and as low as 1-20,000th, even when the fibre was not put upon the stretch.

427. The "smooth" or *non-striated* form of Muscular fibre, which is especially found in the walls of the stomach, intestines, bladder, and other similar parts, is composed of flattened bands whose diameter is usually between 2-2000th and 1-3000th of an inch; and these bands are collected into fasciculi, which do not lie parallel with each other, but which cross and interlace. By macerating a portion of such muscular substance, however, in dilute nitric acid (about one part of ordinary acid to three parts of water) for two or three days, it is found that the bands just mentioned may be easily separated into elongated fusiform cells, not unlike woody fibre in shape; each distinguished for the most part by the presence of a long staff-shaped nucleus, brought into view by the action of acetic acid. These cells, in which the distinction between cell-wall and cell-contents can by no means be clearly seen, are composed of a soft yellow substance, often containing small pale granules, and sometimes yellow globules of fatty matter. In the coats of the bloodvessels are found cells having the same general characters, but shorter and wider in form; and although some of these approach very closely in their general appearance to epithelium-cells, yet they seem to have quite a different nature, being distinguished by their contractile endowments.

428. In the examination of Muscular Tissue, a small portion may be cut out with the curved scissors; and this should be torn up into its component fibres, and these, if possible, should be separated into their fibrillæ, by dissection with a pair of needles, under the simple microscope. The general character of the striated fibre are admirably shown in the large fibres of the Frog; and by selecting a portion in which these fibres spread themselves out to unite with an aponeurotic expansion, they may often be found so well displayed in a single layer, as not only to exhibit all their characters without any dissection, but also to show their mode of connection with the simple fibrous tissue of which the aponeurosis is formed. As the ordinary characters of the fibre are but little altered by boiling, this process may be had recourse to for their more ready separation, especially in the case of the tongue. The separation of the fibres into their fibrillæ is only likely to be accomplished, in the higher Vertebrata, by repeated attempts, of which the greater number are likely to be unsuccessful; but it may be accomplished with much greater facility in the Eel and other fish, the tenacity of whose muscular tissue is much less. The characters of the fibrillæ are not nearly so well pronounced, however, in the Fish, as in the warm-blooded Vertebrata; and among the latter, the Pig has been found by Mr. Lealand (who has been peculiarly successful in this class of preparations) to yield the best examples.

He lays great stress on the freshness of the specimen, which should be taken from the body as soon as possible after death; and when a successful preparation has been made, it should be preserved in Goadby's solution. The shape of the fibres can only be properly seen in cross sections; and these are best made by drying a piece of muscle, so that very thin slices can be cut with a sharp instrument, which on being moistened again, will resume in great part their original characters. Striated muscular fibres are readily obtainable from the limbs of Crustacea and of Insects; and their presence is also readily distinguishable in the bodies of Worms, even of very low organization; so that it may be regarded as characteristic of the Articulated series generally. On the other hand, the Molluscous classes are for the most part distinguished by the non-striation of their fibre; there are, however, two remarkable exceptions, strongly striated fibre having been found in the *Terebratula* and other *Brachiopods*, and also in many *Polyzoa*. Its presence seems always related to energy and rapidity of movement; whilst the non-striated presents itself, where the movements are slower and feebler in their character.

429. *Nerve-substance*.—Whenever a distinct Nervous system can be made out, it is found to consist of two very different forms of tissue; namely, the *vesicular*, which are the essential components of the ganglionic centres, and the *tubular*, of which the connecting trunks consist. The "nerve-vesicles" or "ganglion-globules" are cells, whose typical form may be regarded as globular; but they often present an extension into one or more long processes, which give them a "caudate" or a "stellate" aspect. These processes have been traced into continuity, in some instances, with the axis-cylinders of nerve-tubes; whilst in other cases they seem to inosculate with those of other vesicles. The vesicles are filled with a finely-granular substance, which extends into the prolongations; and they also usually contain pigment-granules, which give them a reddish or yellowish-brown color; but these are commonly absent among the lower animals. It is the presence of this pigment, however, which gives to collections of ganglion-globules in the warm-blooded Vertebrata that peculiar hue, which causes it to be known as the *cineritious* or *gray* matter. Each of the nerve-tubes, on the other hand, of which the trunks are composed, consists, in its most completely developed form, of a delicate membranous sheath, within which is a hollow cylinder of a material known as the "white substance of Schwann," whose outer and inner boundaries are marked out by two distinct lines, giving to each margin of the nerve-tube what is described as a "double contour." The centre or axis of the tube is occupied by a transparent substance, which is known as the "axis-cylinder;" and there is reason to believe that this last is the essential component of the nervous fibre, and that the hollow cylinder that surrounds it, serves, like the

tubular sheath, for its complete isolation. The contents of the membranous envelope are very soft, yielding to slight pressure; and they are so quickly altered by the contact of water or of any liquids that are foreign to their nature, that their characters can only be properly judged of when they are quite fresh. Besides the proper tubular fibres, however, there are others, known as "gelatinous," which are considerably smaller than the preceding, and do not exhibit any differentiation of parts. They are flattened, soft, and homogeneous in their appearance, and contain numerous nuclear particles, which are brought into view by acetic acid. They can sometimes be seen to be continuous with the axis-cylinders of the ordinary fibres, and also with the radiating prolongations of the vesicles; so that their nervous character, which has been doubted by some anatomists, seems established beyond doubt. The ultimate distribution of the nerve-fibres may be readily traced in thin vertical sections of the skin, treated with solution of soda. It was formerly supposed that all its papillæ are furnished with nerve-fibres, and minister to sensation; but it is now known that a large proportion (at any rate) of those furnished with loops of bloodvessels (Fig. 329, D), being destitute of nerve-fibres, must have for their special office the production of the epidermis; whilst those which, possessing nerve-fibres, have sensory functions, are usually destitute of bloodvessels. The greater part of the interior of each sensory papilla of the skin, is occupied by a peculiar "axile body," which seems to be merely a bundle of ordinary fibrous tissue, whereon the nerve-fibre appears to terminate. The nerve-fibres are more readily seen, however, in the "fungiform" papillæ of the tongue, to each of which several of them proceed; these bodies, which are very transparent, may be well seen by snipping off minute portions of the tongue of the Frog; or by snipping off the papillæ themselves from the surface of the living Human tongue, which can be readily done by a dexterous use of the curved scissors, with no more pain than the prick of a pin would give. The transparency of any of these papillæ is increased, by treating them with a solution of soda.

430. For the sake of obtaining a general acquaintance with the microscopic characters of these principal forms of Nerve-substance, it is best to have recourse to minute nerves and ganglia. The small nerves which are found between the skin and the muscles of the back of the Frog, and which become apparent when the former is being stripped off, are extremely suitable for this purpose; and if they be treated with strong acetic acid, a contraction of their tubes takes place, by which the axis cylinder is forced out from their cut extremities, so as to be made more apparent than it can be in any other way. The "gelatinous" fibres are found in the greatest abundance in the Sympathetic nerves; and their characters may be best studied in the smaller branches of that system. So, for the examination of

the ganglionic vesicles, and of their relation to the nerve-tubes, it is better to take some minute ganglion as a whole (such as one of the Sympathetic ganglia of the frog, mouse, or other small animal), than to dissect the larger ganglionic masses, whose structure can only be successfully studied by such as are proficient in this kind of investigation. The nerves of the orbit of the eye of Fish, with the ophthalmic ganglion and its branches, which may be very readily got at in the Skate, and of which the components may be separated without much difficulty, form one of the most convenient objects for the demonstration of the principal forms of nerve-tissue, and especially for the connection of nerve-fibres and ganglionic corpuscles. No method of preserving the nerve-tissue has yet been devised, which makes it worth while to attempt to mount preparations for the sake of displaying its minute characters; but the general course of the nerve-tubes, and the disposition of the ganglionic vesicles, may be demonstrated in preparations preserved in weak spirit; and when the skin has been injected, the passage of the nerve-fibres to the papillæ can sometimes be traced in vertical sections, mounted as opaque objects, and viewed by reflected light. The following method, recommended by Mr. J. Lockhart Clarke, for the examination of the structure of the Spinal Cord,¹ would be equally applicable to that of other large ganglionic masses:—A perfectly fresh cord is to be hardened in strong spirit, so that extremely thin sections can be made with a very sharp knife; and such sections, placed on slips of glass, are to be treated with a mixture of one part of acetic acid and three of spirit, which not only makes the fibrous portion more distinct, but also renders the vesicular portion more transparent. If it be desired to preserve such a section, it should be transferred, after maceration for an hour or two in the mixture of acetic acid and spirit, into pure spirit, in which it should be allowed to remain for about the same space of time; from the spirit it should be transferred to oil of turpentine, which soon expels the spirit, and renders the section perfectly transparent, so that it can be examined with high magnifying powers; and it may then be mounted in Canada balsam in the usual manner.

431. *Circulation of the Blood.*—One of the most interesting spectacles that the microscopist can enjoy, is that which is furnished by the circulation of the blood in the "capillary" blood-vessels, which distribute the fluid through the tissues it nourishes. This, of course, can only be observed in such parts of animal bodies, as are sufficiently thin and transparent to allow of the transmission of light through them without any disturbance of their ordinary structure; and the number of these is very limited. The web of the Frog's foot is perhaps the most suitable for ordinary purposes, more especially since this animal is to be easily obtained in almost every locality; and the following is

¹ See his Memoir on that subject, in "Philos. Transact.," 1851.

the arrangement which the author has found most convenient for the purpose. A piece of thin cork is to be obtained, about 9 inches long and 3 inches wide (such pieces are prepared by the cork-cutters, as soles), and a hole about 3-8ths of an inch in diameter is to be cut at about the middle of its length, in such a position that, when the cork is secured upon the stage, this aperture may correspond with the axis of the microscope. The body of the frog is then to be folded in a piece of wet calico, one leg being left free, in such a manner as to confine its movements, but not to press too tightly upon its body; and being then laid down near one end of the cork plate, the free leg is to be extended, so that the foot can be laid over the central aperture. The spreading out of the foot over the aperture is to be accomplished, either by passing pins through the edge of the web into the cork beneath, or by tying the ends of the toes by threads to pins stuck into the cork at a small distance from the aperture; the former method is by far the least troublesome, and it may be doubted whether it is really the source of more suffering to the animal than the latter is, the confinement being obviously that which is most felt. A few turns of tape, carried *loosely* around the calico bag, the projecting leg, and the cork, serve to prevent any sudden start; and when all is secure, the cork plate is to be laid down upon the stage of the microscope, where a few more turns of the tape will serve to keep it in place. The web being moistened with water (a precaution which should be repeated as often as the membrane exhibits the least appearance of dryness), and an adequate light being reflected through the web from the mirror, this wonderful spectacle is brought into view on the adjustment of the focus (a power of from 75 to 100 diameters being the most suitable for ordinary purposes), provided that no obstacle to the movement of the blood be produced by undue pressure upon the body or leg of the animal. It will not unfrequently be found, however, that the current of blood is nearly or altogether stagnant for a time; this seems occasionally due to the animal's alarm at its new position, which weakens or suspends the action of its heart, the movement recommencing again after the lapse of a few minutes, although no change has been made in any of the external conditions. But if the movement should not renew itself, the tape which passes over the body should be slackened; and if this does not produce the desired effect, the calico envelope must also be loosened. When everything has once been properly adjusted, the animal will often lie for hours without moving, or will only give an occasional twitch. The movement of the blood will be distinctly seen by that of the corpuscles, which course after one another through the network of capillaries that intervenes between the smallest arteries and the smallest veins; in those tubes that pass most directly from the veins to the arteries, the current is always in the same direction; but in those which pass

across between these, it may not unfrequently be seen that the direction of the movement changes from time to time. The larger vessels (Fig. 327), with which the capillaries are seen to

FIG. 327.



Capillary circulation in a portion of the web of a *Frog's* foot: 1, trunk of vein; 2, 2, its branches; 3, 3, pigment-cells.

be connected, are almost always *veins*, as may be known from the direction of the flow of blood in them from the branches (2, 2) towards their trunks (1); the *arteries*, whose ultimate subdivisions discharge themselves into the capillary network, are for the most part restricted to the immediate borders of the toes. When a power of 200 or 250 diameters is employed, the visible area is of course greatly reduced; but the individual vessels and their contents are much more plainly seen; and it may then be observed, that whilst the red corpuscles flow at a very rapid rate along the centre of each tube, the colorless corpuscles which are occasionally discernible, move slowly in the clear stream near its margin.

432. The circulation may also be displayed in the *tongue* of the *Frog*, by laying the animal down on its back, with its head close to the hole in the cork-plate, and, after securing the body in this position, drawing out the tongue with the forceps, and fixing it on the other side of the hole with pins. This method, however, is so much more distressing to the animal, that its employment seems scarcely justifiable for the mere purpose of display; and nothing but some anticipated benefit to science, can justify the laying open of the body of the living animal, for the purpose of

examining the circulation of its lungs or mesentery. The *tadpole* of the Frog, when sufficiently young, furnishes a good display of the circulation in its tail; and the difficulty of keeping it quiet during the observation may be overcome, by gradually mixing some hot water with that in which it is swimming, until it becomes motionless; this usually happens when it has been raised to a temperature between 100 and 110°; and notwithstanding that the muscles of the body are thrown into a state of spasmodic rigidity by this treatment, the heart continues to pulsate, and the circulation is maintained: The *larva of the Water-Newt*, when it can be obtained, furnishes a most beautiful display of the circulation, both in its external gills, and in its delicate feet. It may be enclosed in a large aquatic box or in a shallow cell, gentle pressure being made upon its body, so as to impede its movements, without stopping the heart's action. The circulation may also be seen in the tails of small fish, such as the *Minnow* or *Stickle-back*, by confining these animals in tubes, or in shallow cells, or in a large aquatic box; but although the extreme transparency of these parts adapts them well for this purpose in one respect, yet the comparative scantiness of their bloodvessels prevents them from being as suitable as the Frog's web in another not less important particular. One of the most beautiful of all displays of the circulation, however, is that which may be seen upon the yolk-bag of young Fish (such as the trout) soon after they have been hatched; and as it is their habit to remain almost entirely motionless at this stage of their existence, the observation can be made with the greatest facility by means of the zoophyte-trough, provided that the subject of it can be obtained. Now that the artificial breeding of these fish is largely practised for the sake of stocking fish-ponds, there can seldom be much difficulty in procuring specimens at the proper period. The store of yolk which the yolk-bag supplies for the nutrition of the embryo, not being exhausted in the Fish (as it is in the bird) previously to the hatching of the egg, this bag hangs down from the belly of the little creature on its emersion; and continues to do so until its contents have been absorbed into the body, which does not happen for some little time afterwards. And the blood is distributed over it in copious streams, partly that it may draw into itself fresh nutritive material, and partly that it may be subjected to the aerating influence of the surrounding water.

433. *Injected Preparations.*—Next to the circulation of the blood in the living body, the varied distribution of the Capillaries in its several organs, as shown by means of "injections" of coloring matter thrown into their principal vessels, is one of the most interesting subjects of microscopic examination. The art of making successful preparations of this kind, is one in which perfection can usually be attained only by long practice, and by attention to a great number of minute particulars; and better specimens may be obtained, therefore, from those who have made it a business to prepare them, than are likely to be prepared by amateurs

for themselves. For this reason, no more than a general account of the process will be here offered; the minute details which need to be attended to, in order to attain successful results, being readily accessible elsewhere to such as desire to put it in practice.¹ The coloring matter which is altogether most suitable when only one set of vessels is to be injected, is Chinese vermilion. This, however, as commonly sold, contains numerous particles of far too large a size; and it is necessary first to reduce it to a greater fineness by continued trituration in a mortar (an agate or a steel mortar is the best) with a small quantity of water, and then to get rid of the larger particles by a process of "levigation," exactly corresponding to that by which the particles of coarse sand, &c., are separated from the Diatomaceæ (p. 305). The fine powder thus obtained, ought not, when examined under a magnifying power of 200 diameters, to exhibit particles of any appreciable dimensions. The "size" or "gelatine" should be of a fine and pure quality, and should be of sufficient strength to form a tolerably firm jelly when cold, whilst quite limpid when warm. It should be strained, whilst hot, through a piece of new flannel; and great care should be taken to preserve it free from dust, which may best be done by putting it into clean jars, covering its surface with a thin layer of alcohol. The proportion of levigated vermilion to be mixed with it for injection, is about 2 oz. to a pint; and this is to be stirred in the melted size, until the two are thoroughly incorporated, after which the mixture should be strained through muslin. The injection is thrown into the vessels by means of a brass syringe expressly constructed for the purpose, which has several jet-pipes of different sizes, adapted to the different dimensions of the vessels to be injected; and these should either be furnished with a stop-cock to prevent the return of the injection when the syringe is withdrawn, or a set of small corks of different sizes should be kept in readiness, with which they may be plugged. The pipe should be inserted into the cut end of the trunk which is to be injected, and should be tied therein by a silk thread. In injecting the vessels of fish, mollusks, &c., the softness of the vessels renders them liable to break in the attempt to tie them; and it is therefore better for the operator to satisfy himself with introducing a pipe as large as he can insert, and with passing it into the vessel as far as he can without violence. All the vessels from which the injection might escape, should be tied, and sometimes it is better to put a ligature round a part of the organ or tissue itself; thus, for example, when a portion of the intestinal tube is to be injected through its branch of the mesenteric artery, not only should ligatures be put round any divided vessels of the mesentery, but the cut ends of the intestinal tube should be firmly tied. The operation should either be performed when the body or organ is

¹ See especially the article "Injection," in the "Micrographic Dictionary;" Dr. Beale's treatise on "The Microscope, and its application to Clinical Medicine," Chap. viii; and M. Robin's work, "Du Microscope et des Injections." (Paris, 1849.) (See also Appendix.)

as fresh as possible, or after the expiry of sufficient time to allow the *rigor-mortis* to pass off, the presence of this being very inimical to the success of the injection. The part should be thoroughly warmed, by soaking in warm water for a time proportionate to its bulk; and the injection, the syringe, and the pipes should also have been subjected to a temperature sufficiently high to insure the free flow of the liquid. The force used in pressing down the piston should be very moderate at first; but should be gradually increased as the vessels become filled; and it is better to keep up a steady pressure for some time, than to attempt to distend them by a more powerful pressure, which will be certain to cause extravasation. This pressure should be maintained¹ until the injection begins to flow from the large veins, and the tissue is thoroughly reddened; and if one syringe-ful of injection after another be required for this purpose, the return of the injection should be prevented by stopping the nozzle of the jet-pipe when the syringe is removed for refilling. When the injection has been completed, any openings by which it can escape should be secured, and the preparation should then be placed for some hours in cold water, for the sake of causing the size to "set."²

434. Although no injections look so well by reflected light, as those which are made with vermilion, yet other coloring substances may be advantageously employed for particular purposes. Thus the yellow chromate of lead, which is precipitated when a solution of acetate of lead is mixed with a solution of chromate of potass, is an extremely fine powder, which "runs" with great facility in an injection, and has the advantage of being very cheaply prepared. The best method of obtaining it, is to dissolve 200 grains of acetate of lead and 105 grains of chromate of potass in separate quantities of water, to mix these, and then, after the subsidence of the precipitate, to pour off the supernatant fluid, so as to get rid of the acetate of potash which it contains, since this is apt to corrode the walls of the vessels, if the preparation be kept moist. The solutions should be mixed cold, and the precipitate should not be allowed to dry before being incorporated with the size, four ounces of which will be the proportion appropriate to the quantity of the coloring-substance produced by the above process. The same materials may be used in such a manner that the decomposition takes place within the vessels themselves, one of the solutions being thrown in first, and then the other; and this process involves so little trouble or expense, that it may be considered the best for those who are

¹ A simple mechanical arrangement for this purpose, by which the fatigue of maintaining this pressure with his hand is saved to the operator, is described in the "Micrographic Dictionary," p. 354.

² The kidney of a sheep or pig is a very advantageous organ for the learner to practise on; and he should first master the filling of the vessels from the arterial trunk alone, and then, when he has succeeded in this, he should fill the tubuli uriniferi with white injection, before sending colored injection into the renal artery. The entire systemic circulation of small animals, as mice, rats, frogs, &c., may be injected from the aorta; and the pulmonary vessels from the pulmonary artery.

novices in the operation, and who are desirous of perfecting themselves in the practice of the easier methods, before attempting the more costly. By M. Doyère, who first devised this method, it was simply recommended to throw in saturated solutions of the two salts, one after the other; but Dr. Goadby, who has had much experience in the use of it, advises that gelatine should be employed, in the proportion of 2 oz. dissolved in 8 oz. of water, to 8 oz. of the saturated solutions of each salt. This method answers very well for preparations that are to be mounted dry; but for such as are to be preserved in fluid, it is subject to the disadvantage of retaining in the vessels the solution of acetate of potash, which exerts a gradual corrosive action upon them. Dr. Goadby has met this objection, however, by suggesting the substitution of nitrate for acetate of lead; the resulting nitrate of potash having rather a preservative than a corrosive action on the vessels. When it is desired to inject two or more sets of vessels (as the arteries, veins, and gland-ducts) of the same preparation, different coloring substances should be employed. For a *white* injection, the carbonate of lead (prepared by mixing solutions of acetate of lead and carbonate of soda, and pouring off the supernatant liquid when the precipitate has fallen) is the best material. No *blue* injections can be much recommended, as they do not reflect light well, so that the vessels filled with them seem almost black; the best is freshly-precipitated Prussian blue (formed by mixing solutions of persulphate of iron and ferrocyanide of potassium), which, to avoid the alteration of its color by the free alkali of the blood, should be triturated with its own weight of oxalic acid and a little water, and the mixture should then be combined with size, in the proportion of 146 grains of the former to 4 oz. of the latter.

435. Injected preparations may be preserved either dry or in fluid. The former method is well suited to sections of many solid organs, in which the disposition of the vessels does not sustain much alteration by drying; for the colors of the vessels are displayed with greater brilliancy than by any other method, when such slices, after being well dried, are moistened with turpentine and mounted in Canada balsam. But for such an injection as that shown in Fig. 328, in which the form and disposition of the intestinal *villi* would be completely altered by drying, it is indispensable that the preparation should be mounted in fluid, in a cell deep enough to prevent any pressure on its surface. Either Goadby's solution or weak spirit answers the purpose very well.

FIG. 328.



Villi of Small Intestine of Monkey.

438. In no part of the circulating apparatus, however, does the disposition of the capillaries present more points of interest, than it does in the Respiratory organs.

Fig. 330



In *Fishes*, the respiratory surface is formed by an outward extension into fringes of *gills*, each of which consists of an arch with straight laminae hanging down from it; and every one of these laminae (Fig. 330) is furnished with a double row of leaflets, which is most minutely supplied with bloodvessels, their network (as seen at A) being so close, that its meshes (indicated by the dots in the figure) cover less space than the vessels themselves. The gills of Fish are not ciliated on their surface like those of Mollusks and of the larva of the Water-newt; the necessity for such a mode of renewing the fluid in contact with them, being superseded by the muscular apparatus with which the gill-chamber is furnished. But in *Reptiles*, the respiratory surface is formed by the walls of an internal cavity, that

Fig. 330. A. Capillaries of the Gill of the Fish. B. The internal lamellae.—A. Enlarged view of the capillaries, B. Enlarged view of the network of the lamellae.

in *Reptiles*, these organs, however, are constructed on a plan different from that which they present in higher Vertebrata, the great extension of surface which is effected in the latter by the subdivision of the cavity, not being here necessary. Thus (for example) the cavity of each lung is undivided:

Fig. 331




Fig. 331. Capillaries of the Lung of Frog.

its walls, which are thin and membranous at the lower part, there present a simple smooth expanse; and it is only at the upper part, where the extensions of the tracheal cartilage form a network over the interior, that its surface is depressed into sacculi, whose lining is crowded with bloodvessels (Fig. 331). In this manner, a set of air-cells is formed in the thickness of the upper wall of the lung, which communicates with the general cavity, and very much increases the surface

area over which the blood comes into relation with the air; but each sacculus has a capillary network of its own, which lies on

one side against its wall, so as only to be exposed to the air on its free surface. In the elongated lung of the Snake, the same general arrangement prevails; but the cartilaginous reticulation of its upper part projects much further into the cavity, and encloses in its meshes (which are usually square, or nearly so) several layers of air-cells, which communicate, one through another, with the general cavity. The structure of the lungs of Birds presents us with an arrangement of a very different kind, the purpose of which is to expose a very large amount of capillary surface to the influence of the air. The entire mass of each may be considered as subdivided into an immense number of "lobules" or "lunglets" (Fig. 332), each of which has its own bronchial tube (or

FIG. 332.




Interior structure of Lung of *Fowl*, as displayed by a section, A, passing in the direction of a bronchial tube, and by another section, B, cutting it across.

subdivision of the windpipe), and its own system of bloodvessels, which have very little communication with those of other lobules. Each lobule has a central cavity, which closely resembles that of a Frog's lung in miniature; having its walls strengthened by a network of cartilage derived from the bronchial tube, in the interstices of which are openings leading to sacculi in their substance. But each of these cavities is surrounded by a solid plexus of bloodvessels, which does not seem to be covered by any limiting membrane, but which admits air from the central cavity freely between its meshes; and thus its capillaries are in immediate relation with air on all sides, a provision that is obviously very favorable to the complete and rapid aeration of the blood they contain. In the lung of *Man* and *Mammals*, again, the plan of structure differs from the foregoing, though the general effect of it is the same. For the whole interior is divided up into minute air-cells, which freely communicate with each other, and with the ultimate ramifications of the air-tubes into which the trachea (windpipe) subdivides; and the network of bloodvessels (Fig. 333) is so disposed in the partitions between these cavities, that the blood is exposed to the air on both sides. It has been calculated that the number of these air-cells grouped around the

termination of each air-tube in man, is not less than 18,000; and that the total number in the entire lungs is *six hundred millions*.

439. The following list of the parts of the bodies of Verte-

FIG. 333.



Arrangement of the Capillaries of the air-cells of the Human Lung.

brata, of which Injected preparations are most interesting as Microscopic objects, may be of service to those who may be inclined to apply themselves to their production. *Alimentary Canal*; Stomach, showing the orifices of the gastric follicles, and the rudimentary villi near the pylorus; Small Intestine, showing the villi and the orifices of the follicles of Lieberkühn, and at its lower part the Peyerian glands; Large Intestine, showing the

various glandular follicles:—*Respiratory Organs*; Lungs of Mammals, Birds, and Reptiles; Gills and Swimming-bladder of Fish:—*Glandular Organs*; Liver, Gall-bladder, Kidney, Parotid:—*Generative Organs*; Oviduct of Bird and Frog; Mammalian Placenta; Uterine and Fœtal Cotyledons of Ruminants:—*Organs of Sense*; Iris, Choroid, and Ciliary processes of Eye, Pupillary Membrane of fœtus; Papillæ of Tongue; Mucous Membrane of Nose; Papillæ of Skin of finger:—*Tegumentary Organs*; Skin of different parts, hairy and smooth, with vertical sections showing the vessels of the Hair-follicles, Sebaceous glands, and Papillæ; Matrix of nails, hoofs, &c.:—*Tissues*; Fibrous, Muscular, Adipose, Sheath of Tendon.

440. *Development*.—The study of the Embryological development of Vertebrated animals has been pursued of late years with great zeal and success by the assistance of the Microscope; but as this is a department of inquiry which needs for its successful pursuit a thoroughly scientific culture, and is only likely to be taken up by a professed Physiologist, no good purpose seems likely to be served by here giving such an imperfect outline of the process, as could alone be introduced into a work like the present; and the reader who may desire information upon it, will find no difficulty in obtaining this through systematic treatises on Physiology.¹

¹ The Author takes the liberty of referring to his "Principles of Comparative Physiology," 4th Ed. chap. xi. as containing a general view of the whole subject, with references to the principal sources of more detailed information.

CHAPTER XIX.

APPLICATIONS OF THE MICROSCOPE TO GEOLOGICAL INVESTIGATION.

441. THE utility of the Microscope is by no means limited to the determination of the structure and actions of the Organized beings at present living on the surface of the earth; for a vast amount of information is afforded by its means to the geological inquirer, not only with regard to the minute characters of the many Vegetable and Animal remains that are entombed in the successive strata of which its crust is composed, but also with regard to the essential nature and composition of many of those strata themselves. We cannot have a better example of its value in both these respects, than that which is afforded by the results of microscopic examination of *lignite* or fossilized wood, and of ordinary *coal*, which there is every reason to regard as a product of the decay of wood.

442. Specimens of *Fossilized Wood*, in a state of more or less complete preservation, are found in numerous strata of very different ages,—more frequently, of course, in those whose materials were directly furnished by the dry land, and were deposited in its immediate proximity, than in those which were formed by the deposition of sediments at the bottom of a deep ocean. Generally speaking, it is only when the wood is found to have been penetrated by *silex*, that its organic structure is well preserved; but instances occur every now and then, in which penetration by *carbonate of lime* has proved equally favorable. In either case, transparent sections are needed for the full display of the organization; but such sections, though made with great facility when lime was the fossilizing material, require much labor and skill when *silex* has to be dealt with. Occasionally, however, it has happened that the infiltration has filled the cavities of the cells and vessels, without consolidating their walls; and as the latter have undergone decay without being replaced by any cementing material, the lignite, thus composed of the internal “casts” of the woody tissues, is very friable, its fibres separating from each other like those of asbestos; and their laminae split asunder with a knife, or isolated fibres separated by rubbing down between the fingers, exhibit the characters of the woody structure extremely well, when mounted in Canada

balsam. Generally speaking, the lignites of the Tertiary strata present a tolerably close resemblance to the wood of the existing period; thus the ordinary structure of *Dicotyledonous* and *Monocotyledonous* stems, may be discovered in such lignites in the utmost perfection; and the peculiar modification presented by *Coniferous* wood, is also most distinctly exhibited (Fig. 171). As we descend, however, through the strata of the Secondary period, we more and more rarely meet with the ordinary dicotyledonous structure; and the lignites of the earliest deposits of these series are, almost universally, either gymnosperms or palms.¹ Descending into the Palæozoic series, we are presented in the vast *Coal* formations of our own and other countries, with an extraordinary proof of the prevalence of a most luxuriant vegetation in a comparatively early period of the world's history; and the microscope lends the geologist essential assistance, not only in determining the nature of much of that vegetation, but also in demonstrating, what has been suspected on other grounds, that Coal itself is nothing else than a mass of decomposed vegetable matter, chiefly derived from the decay of *Coniferous* wood. The determination of the characters of the *Ferns*, *Sigillaria*, *Lepidodendra*, *Calamites*, and other kinds of vegetation whose forms are preserved in the shales and sandstones that are interposed between the strata of coal, must be chiefly based on their external characters; since it is very seldom that any of the specimens present any such traces of minute internal structure, as can be subjected to microscopic elucidation. But notwithstanding the general absence of any definite *form* in the masses of decomposed wood of which Coal itself consists (these having apparently been reduced to a pulpy state by decay, before the process of consolidation by pressure, aided perhaps by heat, commenced), the traces of *structure* revealed by the Microscope, are sufficient not only to determine its vegetable origin, but, in some cases, to justify the Botanist in assigning the characters of the vegetation from which it must have been derived. Different specimens of Coal exhibit these structural characters in very different degrees of distinctness; but they uniformly indicate, with a clearness proportionate to their distinctness, that such vegetation must have been *Coniferous* in its nature, and that it probably approximated most nearly to that group of existing Coniferae, to which the *Araucaria* belong. These inferences are based upon the fact, that the woody structure consists of woody fibres without interposed vessels; upon the presence of glandular dots on the woody fibres; and upon the peculiar arrangement of these dots in two or more rows, alternating one with another (§§ 231, 238).

443. In examining the structure of Coal, various methods may be followed. Of those kinds which have sufficient tenacity, thin sections may be made; but the opacity of the substance requires

¹ Under this head are included the *Cycadææ*, along with the ordinary *Conifera* or pine and fir tribe.


that such sections should be ground extremely thin, before they become transparent; and its friability renders this process one of great difficulty. Any section must either cross the woody tissue transversely, so that the appearance it presents will resemble that of Fig. 170; or it must traverse it vertically, in which case the fibrous structure will be brought into view, either as in Fig. 171, or as in Fig. 172; or it must pass in an intermediate direction. The following method, which would seem not only to be more simple, but also to give more satisfactory results, is recommended by the authors of the "Micrographic Dictionary" (p. 150):— "The coal is macerated for about a week in a solution of carbonate of potash; at the end of that time, it is possible to cut tolerably thin slices with a razor. These slices are then placed in a watch-glass with strong nitric acid, covered, and gently heated; they soon turn brownish, then yellow, when the process must be arrested by dropping the whole into a saucer of cold water, or else the coal would be dissolved. The slices thus treated appear of a darkish amber color, very transparent, and exhibit the structure, when existing, most clearly. The specimens are best preserved in glycerine, in cells; we find that spirit renders them opaque, and even Canada balsam has the same effect." When the coal is so friable, that no sections can be made of it by either of these methods, it may be ground to fine powder, and the particles may then, after being mounted in Canada balsam, be subjected to microscopic examination; the results which this method affords, are by no means satisfactory in themselves; but they will often enable the organic structure to be sufficiently determined, by the comparison of the appearances presented by such fragments, with those which are more distinctly exhibited elsewhere. Valuable information may often be obtained, too, by treating the ash of an ordinary coal-fire in the same manner, or (still better) by burning to a white ash a specimen of coal that has been previously boiled in nitric acid, and then carefully mounting the ash in Canada balsam; for mineral casts of vegetable cells and fibres may often be distinctly recognized in such ash; and such casts are not unfrequently best afforded by samples of coal, in which the method of section is least successful in bringing to light the traces of organic structure, as is the case, for example, with the "anthracite" of Wales.¹

444. Passing on now to the Animal kingdom, we shall first cite some parallel cases in which the essential nature of deposits that form a very important part of the Earth's crust, has been determined by the assistance of the microscope; and shall then select a few examples of the most important contributions which it has afforded, to our acquaintance with types of Animal life long since extinct. It is an admitted rule in Geological science, that the past history of the Earth is to be interpreted, so far as may be found possible, by the study of the changes which are

¹ See Prof. Quekett's Memoir on the Minute Structure of the Torbane-hill Mineral, in "Transact. of the Microsc. Societ." Ser. 2, vol. ii, p. 41, *et seq.*

still going on. Thus, when we meet with an extensive stratum of fossilized *Diatomaceæ* (§ 191) in what is now dry land, we can entertain no doubt that this siliceous deposit originally accumulated either at the bottom of a fresh-water lake, or beneath the waters of the ocean; just as such deposits are formed at the present time, by the production and death of successive generations of these bodies, whose indestructible casings accumulate in the lapse of ages, so as to form layers whose thickness is only limited by the time during which this process has been in action (§ 190). In like manner, when we meet with a limestone-rock

entirely composed of the calcareous shells of *Foraminifera*, some of them entire, others broken up into minute particles, we interpret the phenomenon by the fact, that the dredgings obtained from certain parts of the ocean-bottom consists almost entirely of remains of existing *Foraminifera*, in which entire shells, the animals of which may be yet alive, are mingled with the debris of others that have been reduced by the action of the waves to a fragmentary state.¹ Now in the fine white mud which is brought up from almost every part of the sea-bottom of the Levant, where it forms a stratum that is continually undergoing a slow but steady increase in thickness, the microscopic researches of Prof. Williamson² have shown that not only are there multitudes of minute remains of living or-



Microscopic Organisms in Levant Mud:—A, D, siliceous spicules of *Tethya*; B, H, spicules of *Geodia*; C, Sponge-spicule (unknown); E, calcareous spicule of *Grantia*; F, G, M, O, portions of calcareous skeleton of Echinodermata; H, I, calcareous spicule of *Gorgonia*; K, L, N, siliceous spicules of *Halichondria*; P, portion of prismatic layer of shell of *Pinna*.


¹ Such a deposit, consisting chiefly of *Orbitolites* (§ 287) is at present in the act of formation on certain parts of the shores of Australia, as the Author is informed by Mr. J. Beete Jukes; thus affording the exact parallel to the stratum of *Orbitolites* (belonging, as the Author's investigations have led him to believe, to the very same species) that forms part of the "Calcaire Grossier" of the Paris basin.

² "Memoirs of the Manchester Literary and Philosophical Society," vol. viii.

ganisms, both animal and vegetable, but that it is entirely or almost wholly composed of such remains. Among these were about 26 species of Diatomaceæ (siliceous), 8 species of Foraminifera (calcareous), and a miscellaneous group of objects (Fig. 334), consisting of calcareous and siliceous spicules of Sponges and Gorgoniæ, and of fragments of the calcareous skeletons of Echinoderms and Mollusks.

445. Now almost exactly the same collection of forms, with the exception of the siliceous Diatomaceæ, is found in many parts of the "Calcaire Grossier" of the Paris basin, as well as in other extensive deposits of the same early tertiary period. And there is little doubt that a large proportion of the great Cretaceous (chalk) formation has a like composition; for many parts

FIG 335.



Microscopic Organisms in Chalk from Gravesend; *a, b, c, d*, *Textularia globulosa*; *e, e, e*, *Rotalia aspera*; *f*, *Textularia aculeata*; *g*, *Planularia hexas*; *h*, *Navicula*.

of it consist in great part of the minuter kinds of Foraminifera, whose shells are imbedded in a mass of apparently amorphous particles, many of which, nevertheless, present indications of being the worn fragments of similar shells, or of larger calcareous organisms. In the Chalk of some localities, Foraminifera constitute the principal part of the minute organisms which can be recognized with the microscope (Figs. 335, 336); in other instances, the disintegrated prisms of *Pinna* (§ 336) or other large shells of the like structure (as *Inoceramus*) constitute the great bulk; whilst in other cases, again, the chief part is made up of the shells of *Cytherina*, a marine form of Entomostracous Crustacean (§ 367). Different specimens of Chalk vary greatly in the

proportion which the distinctly organic remains bear to the amorphous particles, and which the different kinds of the former bear to each other; and this is quite what might be anticipated, when we bear in mind the predominance of one or another tribe of animals or plants in the several parts of a large area. True Chalk seems to differ from the Levant Mud, in the small proportion which the siliceous remains of Diatomaceæ bear, in the former, to that which is mingled in the latter with the calcareous

FIG. 336.



Microscopic Organisms in *Chalk* from Meudon; partly seen as opaque, and partly as transparent objects.

shells of Foraminifera, &c.; and it seems doubtful to what extent they were present in the seas of that epoch. Such remains are found in abundance, however, forming marly strata which alternate with those of a chalky nature, in the South of Europe and the North of Africa (Fig. 101); and it is surmised by Prof. Ehrenberg, that the layers of *flint* which the British Chalk contains, have been derived by some metamorphic process from similar layers of siliceous Diatomaceæ which have disappeared. It is now certain, however, that the deposits referred to by Prof. Ehrenberg are of an age later than that of the great Chalk formation; so that little support is furnished by their phenomena to his hypothesis. But whatever may have been the origin of the siliceous material, it may be stated as a fact beyond all question, that nodular flints and other analogous concretions (such as agates) may generally be considered as fossilized Sponges or Alcyonian Zoophytes; since not only are their external forms and their superficial markings often highly cha-

racteristic of those organisms, but, when sections of them are made sufficiently thin to be transparent, a spongy texture may be most distinctly recognized in their interior.¹ It is curious that many such sections contain well-preserved specimens of *Xanthidia*, which are Desmidiaceæ whose divided body is covered with long spinous projections, often cleft, and sometimes furnished with hooks at their extremities; and we occasionally also find upon their surface, or even imbedded in their substance, Foraminiferous shells (especially *Rotaliæ*), in which not only the substance of the shell has undergone silicification, but also that of the soft animal body, the shrunken form of which may be recognized in the dark carbonaceous hue imparted to the central portion of the silex which fills each chamber.

446. In examining Chalk or other similar mixed aggregations, whose component particles are easily separable from each other, it is desirable to separate, with as little trouble as possible, the larger and more definitely organized bodies, from the minute amorphous particles; and the mode of doing this will depend upon whether we are operating upon the large or upon the small scale. If the former, a quantity of soft chalk should be rubbed to powder with water, by means of a soft brush; and this water should then be proceeded with, according to the method of levigation already directed for separating the Diatomaceæ (§ 192). It will usually be found that the first deposits contain the larger Foraminifera, fragments of shell, &c., and that the smaller Foraminifera and Sponge spicules fall next; the fine amorphous particles remaining diffused through the water after it has been standing for some time, so that they may be poured away. The organisms thus separated should be dried, and mounted in Canada balsam. If the smaller scale of preparation be preferred, as much chalk scraped fine as will lie on the point of a knife, is to be laid on a drop of water on the glass slide, and allowed to remain there for a few seconds; the water, with any particles still floating on it, should then be removed; and the sediment left on the glass should be dried and mounted in balsam. For examining the structure of flints, such chips as may be obtained with a hammer will commonly serve very well: a clear translucent flint being first selected, and the chips that are obtained being soaked for a short time in turpentine (which increases their transparency), those which show organic structure, whether Sponge tissue or *Xanthidia*, are to be selected and mounted in Canada balsam. The most perfect specimens of sponge-structure, however, are only to be obtained by slicing and polishing,—a process which is best performed by the lapidary.

447. There are various other deposits, of less extent and importance than the great Chalk formation, which are, like it, composed in great part of microscopic organisms, chiefly minute

¹ See Mr. Bowerbank's Memoirs in the "Transact. of the Geolog. Societ." 1840, and in the "Ann. of Nat. Hist." 1st Ser. vols. vii, x.


Foraminifera; and the presence of animals of this group may be recognized, by the assistance of this instrument, in sections of calcareous rocks of various dates, whose chief materials seem to have been derived from Corals, Eocrinite stems, or Molluscous shells. Thus in the "Crag" formation (tertiary) of the eastern coast of England, the greater portion of which is perceived by the unassisted eye to be composed of fragments of Shells, Corals (or rather Polyzoaria, § 325), and Echinodermata, the microscope enables us to discover Foraminifera, minute fragments of shells and corals, and spicules of Sponges; the aggregate being such as is at present in process of formation on many parts of our shores, and having been, therefore, in all probability, a "littoral" formation, whilst the Chalk (with other formations chiefly consisting of Foraminifera) was deposited at the bottom of deeper waters. Many parts of the Oolitic (secondary) formation have an almost identical character, save that the forms of organic life give evidence of a different age; and in those portions which exhibit the "roe-stone" arrangement from which the rock derives its name (such as is beautifully displayed in many specimens of Bath-stone and Portland-stone), it is found by microscopic examination of transparent sections, that each rounded concretion is composed of a series of concentric spheres enclosing a central nucleus, which nucleus is often a Foraminiferous shell. In the Carboniferous (palæozoic) limestone, again, well-preserved specimens of Foraminifera present themselves; and there are certain bands of limestone of this epoch in Russia, varying in thickness from fifteen inches to five feet, and frequently repeated through a vertical depth of two hundred feet, which are almost entirely composed of Foraminiferous shells belonging to a genus now extinct, the *Fusulina*.

448. It is not only, however, in the condition of organisms of microscopic size, that the Foraminifera have contributed in an important degree to the formation of the solid crust of the earth; for the Nummulitic limestone,¹—which forms a band, often 1800 miles in breadth, and frequently of enormous thickness, that may be traced from the Atlantic shores of Europe and Africa, through Western Asia to Northern India and China, and over vast areas of North America likewise,—is composed of an aggregation of larger bodies belonging to the same type; the "matrix," or rock-substance, in which these are imbedded, being itself usually made up (as microscopic examination of their sections demonstrates) of the comminuted particles of similar organisms, and of smaller Foraminifera; although it is sometimes composed (as in the British beds of London Clay which include Nummulites) of accumulations of clayey or other inorganic particles. The structure of the *Nummulite* itself, as elucidated by microscopic examination, presents some extremely remarkable modifications of the ordinary Foraminiferous type. It is composed

¹ The Pyramids of Egypt are made of this material.

of a series of chambers, symmetrically disposed in a spiral round a centre, so that a section through its median plane would present very much the appearance of Fig. 209; but each whorl invests all the preceding whorls, so as to form a new layer over the entire surface of the disk; and this layer is usually separated from that which it covers, by an intervening space, which is divided into smaller spaces of more or less regular form, by prolongations from the partitions that divide the chambers of the central plane. These prolongations are very differently arranged in different species; thus in some, as *Nummulites distans*, they keep their own separate course, tending towards the centre; whilst in others, as *N. levigata*, they inosculate with each other, so as to divide the space that intervenes between one layer and another into an irregular network. Hence in a vertical section, such as that of which a part is shown in Fig. 337, we

FIG. 337.



Vertical Section of portion of *Nummulites levigata* :—*a*, margin of external whorl; *b*, one of the outer row of chambers; *c, c*, whorl invested by *a*; *d*, one of the chambers of the fourth whorl from the margin; *e, e'*, marginal portions of the enclosed whorls; *f*, investing portion of outer whorl; *g, g*, spaces left between the investing portions of successive whorls; *h, h*, sections of the partitions dividing these.

see not only the succession of chambers along the central plane, each of them having its own roof and floor, and its own lateral partitions dividing it from other chambers of the same whorl, but we also see the superposition of layers over the inner whorls; so that any chamber *d* in a whorl that is surrounded by three others, is shut in above and below, not only by its proper shelly covering, but by three additional layers formed by the prolongation of the shelly investments of the external whorls; and in like manner, the innermost of the chambers here represented (that nearest *e*) is enclosed by nine layers above and below, in addition to that by which it is itself covered, these nine layers being extensions of the covering of the nine whorls that surround it. Notwithstanding that the inner chambers are thus so deeply buried in the mass of investing whorls, yet there is evidence that the segments of sarcode which they contained, were not cut off from communication with the exterior; but that they may have retained their vitality to the last. The shell itself is almost everywhere minutely porous, being penetrated by parallel tubuli,

which pass from one surface to the other like those of dentine. These tubes are shown, as divided lengthways by a vertical section, in Fig. 338 (*a, a*); whilst the appearance they present when

FIG. 338.



Portion of a thin section of *Nummulites lavigata*, taken in the direction of the preceding, highly magnified to show the minute structure of the shell:—*a, a*, portions of the ordinary shell-substance traversed by parallel tubuli; *b, b*, portions forming the marginal wall, traversed by diverging and larger tubuli; *c*, one of the chambers laid open; *d, d, d*, pillars of solid substance not perforated by tubuli.


cut across in a horizontal section is shown in Fig. 339, the transparent shell-substance, *a, a, a*, being closely dotted with minute punctations which mark their orifices. In that portion of the shell, however, which forms the margin of each whorl (Fig. 338, *b, b*), the tubes are larger, and diverge from each other at greater intervals; whilst at certain other points, *d, d, d*, the shell-substance is not perforated by tubes, but is peculiarly dense in its texture, forming solid pillars which seem to strengthen the other parts. In *Nummulites* whose surfaces have been much exposed to attrition, it commonly happens that the pillars of the superficial layer, being harder than the ordinary shell-substance, and being consequently less worn down, are left as prominences; the presence of which has often been accounted (but erroneously) as a specific character. The successive chambers of the same whorl communicate with each other by a passage left between the inner edge of the partition that separates them, and the margin of the preceding whorl that forms their inner boundary; this passage is sometimes a single large broad aperture, but is more commonly formed by the more or less complete coalescence of several separate perforations, as is seen in Fig. 337, *b*. Such marked differences in this respect are observable in the several parts of one and the same specimen, that it is obvious that very little account should be taken of differences in the form of aperture, as affording specific or generic distinctions among Foraminifera of this type. But besides the foregoing means of communication, by which the segments of sarcode included in the inner chambers were enabled to continue receiving supplies of nutriment, we meet in *Nummulites* with a remarkable development of that system of "interseptal" canals, one of the most characteristic examples of which among recent Foraminifera is

presented in *Faujasina*, as already described (§ 291). These canals are frequently found to be filled up in Nummulites by the fossilizing material; but a careful examination will generally disclose traces of them in the middle of the partitions that divide the chambers (Fig. 339, *b, b*), while from these may be seen to proceed the lateral branches (*c, c*) which, after burrowing (so to speak) in the walls of the chambers, enter them by large orifices (*d*). As the general distribution of this system of canals in the Nummulite is the same as that shown in *Faujasina* (Fig. 209), and as the canals, although smaller, are far more numerous, it is obvious that through its means the segments of sarcode occupying the chambers of the most internal walls

could send their pseudopodial extensions at once to the exterior. Of all Foraminifera, the Nummulite is undoubtedly one of the most highly developed types; and its extraordinary multiplication at the earliest part of the Tertiary period, is a very curious feature in the Earth's history. It is commonly considered that this type is now extinct; but the Author, in common with Prof. Williamson, is disposed to question whether there is any essential difference between Nummulites and the existing genus *Nonionina*, which is very abundant in certain localities; since in many species of Nummulites, as in *Nonionina*, the investing layers of the successive whorls are in immediate contact with those that have preceded them, instead of being separated, as in Fig. 337, by spaces prolonged from the cavities of the chambers.


449. The same Nummulitic limestone also contains, in certain localities (as the southwest of France, north-eastern India, &c.) a vast abundance of discoidal bodies termed *Orbitoides*, which are so similar to Nummulites as to have been taken for them, but which, while

FIG. 339.



Portion of Horizontal Section of *Nummulite*, showing the structure of the walls and of the septa of the chambers:—*a, a*, portion of the wall covering three chambers, the punctations of which are the orifices of tubuli; *b, b*, septa between these chambers, containing canals which send out lateral branches, *c, c*, entering the chambers by larger orifices, one of which is seen at *d*.

FIG. 340.



Section of *Orbitoides Prattii*, parallel to the surface; traversing at *a, a*, the superficial layer, and at *b, b*, the median layer.

still Foraminiferous, are formed upon a plan of structure altogether different. On account of the minuteness of their parts, and the completeness of their fossilization, their structure can only be elucidated by sections thin enough to be examined by the microscope with transmitted light; and it is consequently to the assistance afforded by this instrument, that we are indebted for our knowledge of the curious type of organization which it presents. When one of these disks (which vary in size, in different species, from that of a four-penny piece to that of half a crown) is rubbed down so as to display its internal organization, two different kinds of structure are usually seen in it; one being composed of chambers of very definite form, quadrangular in some species, circular in others, arranged with a general but not constant regularity in concentric circles (Figs. 340, 341, *b, b*); the other, less transparent, being formed of minuter cells which have no such constancy of form, but which might almost be taken for the pieces of a dissected map (Figs. 340, 341, *a, a*). In the upper

FIG. 341.

Portions of the same section, more highly magnified:—*a*, superficial layer; *b*, median layer.

and lower walls of these last, minute punctations may be observed, which seem to be the orifices of connecting tubes whereby they are perforated. The relations of these two kinds of structure to each other, are made evident by the examination of a vertical section (Fig. 342); which shows that the portion *a*, Figs.

FIG. 342.

Vertical Section of *Orbitoides Prattii*, showing the large central cell at *a*, and the median layer surrounding it, covered above and below by the superficial layers.

340, 341, forms the central plane, its concentric circles of cells being arranged round a large central cell *a*, as in Orbitolite (Fig. 206); whilst the cells of the portion *b* are irregularly superposed one upon the other, so as to form several layers, which are most numerous towards the centre of the disk, and thin away gradually towards its margin. By the perforations in these layers, the pseudopodia proceeding from the central plane of chambers may have found their way direct to the surface, or at any rate would have been brought into connection with the segments lying nearer to it. No organisms precisely resembling the Orbitoides,

are known to exist at present; but there are some which differ from it so little, that a knowledge of their structure helps materially to elucidate points, which would otherwise be rendered obscure in it through the changes induced by fossilization.¹

450. The foregoing details, taken in addition to the facts of like nature that have been mentioned in previous parts of this work (as, for example, in § 294), will serve as examples of the essential importance of microscopic investigation, in determining, on the one hand, the real character of various stratified deposits, and, on the other, in elucidating the nature of the organic remains which these may include. The former of these lines of inquiry has not yet attracted the attention which it deserves; since, as is very natural, the greater number of Microscopists are more attracted by those definite forms which they can distinctly recognize, than by the amorphous sediments which present no definite structural characters. Yet it is a question of extreme interest to the Geologist, to determine how far these had their origin in the disintegration of organic structures; and much light may often be thrown upon this question by careful microscopic analysis. Thus the author having been requested by Mr. Chas. Darwin, about twelve years since, to examine into the composition of the extensive calcareous deposit which covers the surface of the Pampas region of South America, and to compare it with that of the calcareous tufa still in process of formation along the coast of Chili, was able to state that their constituents were in all probability essentially the same, notwithstanding the difference in their mode of aggregation. For the Chilian tufa is obviously composed in great part of fragments of shells, distinguishable by the naked eye; the dense matrix in which these are imbedded is chiefly made up of minuter fragments, only distinguishable as such by the microscope; while through the midst of these is diffused an aggregation of amorphous particles, that present every appearance of having originated in the yet finer reduction of the same shells, either by attrition or by decomposition. In the Pampas deposit, on the other hand, the principal part was found to be composed of amorphous particles, so similar in aspect to those of the Chilian rock that their identity could scarcely be doubted; and scattered at intervals through these were particles of shell, distinctly recognizable by the microscope, though invisible to the naked eye. Thus, although the evidence afforded by the larger fragments of shell was altogether wanting in the Pampas deposit, it could not be doubted that the materials of both were the same, those of the Pampean formation having been subject to greater comminution than those of the Chilian; and this view served to

¹ See the Author's Memoir on the Microscopic Structure of Nummulite, Orbitolite, and Orbitoides, in the "Quart. Journ. of the Geolog. Society," for Feb., 1850; and the admirable "Description des Animaux Fossiles du Groupe Nummulitique de l'Inde," by MM. D'Archiac and Jules Haime.

confirm, whilst it was itself confirmed by, the idea thought most probable on other grounds by Mr. Darwin, that the Pampean formation was slowly accumulated at the mouth of the former estuary of the Plata, and in the sea adjoining it.¹ A similar line of inquiry has been of late systematically pursued by Mr. R. C. Sorby; who has applied himself to the microscopic study of the composition of fresh-water marls and limestones, by ascertaining the characters and appearances of the minute particles into which shells resolve themselves by decay, and by estimating the relative proportions of the organic and inorganic ingredients of a rock, by delineating on paper (by means of the camera lucida) the outlines of the particles visible in thin sections, then cutting them out, and weighing the figures of each kind.²

451. It is obvious that, under ordinary circumstances, only the hard parts of the bodies of animals that have been entombed in the depths of the earth, are likely to be preserved; but from these a vast amount of information may be drawn; and the inspection of a microscopic fragment will often reveal, with the utmost certainty, the entire nature of the organism of which it formed part. In the examination of the minuter fossil Corals, and of those Polyzoaries (§ 325) which are commonly ranked with them, the assistance of the microscope is indispensable. Minute fragments of the "test" or "spines" of Echinodermata, and of all such Molluscous shells as present distinct appearances of structure (this being especially the case with the Brachiopoda, and with the families of Lamellibranchiate bivalves most nearly allied to them), may be unerringly identified by its means, when the external form of these fragments would give no assistance whatever. In the study of the remarkable ancient group of Trilobites, not only does a microscopic examination of the

Fig. 343.



Eye of Trilobite.

casts which have been preserved of the surface of their eyes (Fig. 343), serve to show the entire conformity in the structure of these organs to the "composite" type which is so remarkable a characteristic of the higher Articulata (§ 383), but it also brings to light certain peculiarities

which help to determine the division of the great Crustacean series with which this group has most alliance.³ It is in the case of the Teeth, the Bones, and the Dermal skeleton of Vertebrated animals, however, that the value of Microscopic inquiry becomes most apparent; since the structure of these presents so many characteristics that are subject to well-marked variations in their several classes, orders, and families, that a knowledge of these

¹ See Mr. C. Darwin's "Geological Observations on South America," p. 32.


² See "Quart. Journ. of Geolog. Science," 1853, p. 344.

³ See Prof. Burmeister "On the Organization of the Trilobites," published by the Ray Society, p. 19.

characters frequently enables the Microscopist to determine the nature of even the most fragmentary specimens, with a positiveness which must appear altogether misplaced to such as have not studied the evidence.

452. It was in regard to *Teeth*, that the possibility of such determinations was first made clear by the laborious researches of Prof. Owen;¹ and the following may be given as examples of their value:—A rock formation extends over many parts of Russia, whose mineral characters might justify its being likened either to the *Old* or to the *New Red Sandstone* of this country, and whose position relatively to other strata is such, that there is great difficulty in obtaining evidence from the usual sources, as to its place in the series. Hence the only hope of settling this question (which was one of great practical importance, since, if the formation were *new red*, Coal might be expected to underlie it, whilst if *old red*, no reasonable hope of coal could be entertained) lay in the determination of the Organic remains which this stratum might yield; but unfortunately these were few and fragmentary, consisting chiefly of teeth, which are seldom perfectly preserved. From the gigantic size of these teeth, together with their form, it was at first inferred that they belonged to Saurian Reptiles, in which case the sandstone must have been considered as *New Red*; but microscopic examination of their intimate structure unmistakably proved them to belong to a genus of Fishes (*Dendrodus*) which is exclusively Palæozoic, and thus decided that the formation must be *Old Red*. So again, the microscopic examination of certain fragments of teeth found in a Sandstone of Warwickshire, disclosed a most remarkable type of tooth-structure (shown in Fig. 344), which was also ascertained to exist in certain teeth that had been discovered in the “*keuper-sandstein*” of Wirtemberg; and the identity or close resemblance of the animals to which these teeth belonged having been thus established, it became almost certain that the Warwickshire and Wirtemberg sandstones were equivalent formations, a point of much geological importance. The next question arising out of this discovery, was the nature of the animal (provisionally termed *Labyrinthodon*, a name expressive of the most peculiar feature in its

FIG. 344.

Section of Tooth of *Labyrinthodon*.

¹ See his magnificent “*Oolontography*.”

dental structure) to which these teeth belonged. They had been referred, from external characters merely, to the order of Saurian Reptiles; but these characters were by no means conclusive; and as the nearest approaches to their peculiar internal structure are presented by Fish-Lizards and Lizard-like Fish, it might be reasonably expected that the Labyrinthodon would combine with its reptilian characters an affinity to fish. This has been clearly proved to be the case, by the subsequent discovery of parts of its skeleton in which such characters are very obvious; and by a very beautiful chain of reasoning, Prof. Owen succeeded in establishing a strong probability, that the Labyrinthodon was a gigantic Frog-like animal five or six feet long, with some peculiar affinities to Fishes, and a certain mixture also of Crocodilian characters; and that it made the well-known foot-prints which have been brought to light, after an entombment whose duration can scarcely be conceived (much less estimated), in the Stourton quarries of Cheshire.

453. The more recent researches of Prof. Quekett on the minute structure of *Bone*,¹ promise to be scarcely less fruitful in valuable results. From the average size and form of the "lacunæ," their disposition in regard to each other and to the Haversian canals, and the number and course of the canaliculi, he feels assured that the nature of even a minute fragment of bone may be determined with a considerable approach to certainty; and the following examples, among many which might be cited, appear to justify such assurance. Dr. Falconer, the distinguished investigator of the fossil remains of the Himalayan region, and the discoverer of the gigantic fossil Tortoise of the Sivalik Hills, having met with certain small bones about which he was doubtful, placed them in the hands of Prof. Quekett for minute examination; and was informed, on microscopic evidence, that they might certainly be pronounced Reptilian, and probably belonged to an animal of the tortoise tribe; and this determination was fully borne out by other evidence, which led Dr. Falconer to conclude that they were toe bones of his great tortoise. Some fragments of bone were found, some years since, in a chalk-pit; which were considered by Prof. Owen to have formed part of the wing-bones of a long-winged sea-bird allied to the Albatross. This determination, founded solely on considerations derived from the very imperfectly preserved external forms of these fragments, was called in question by some other palæontologists; who thought it more probable that these bones belonged to a large species of the extinct genus *Pterodactylus*, a flying-lizard, whose wing was extended upon a single immensely prolonged digit. No species of *Pterodactyle*, however, at all comparable to this in dimensions, was at that time known; and the characters

¹ See his Memoir on the "Comparative Structure of Bone," in the "Transac. of the Microsc. Societ." Ser. 1, vol. ii; and the "Catalogue of the Histological Museum of the Roy. Coll. of Surgeons," vol. ii.

furnished by the configuration of the bones, not being in any degree decisive, the question would have remained unsettled, had not an appeal been made to the Microscopic test. This appeal was so decisive, by showing that the minute structure of the bone in question corresponded exactly with that of Pterodactyle bone, and differed essentially from that of every known Bird, that no one who placed the least reliance upon that evidence could entertain the slightest doubt on the matter. By Prof. Owen, however, the validity of that evidence was questioned, and the bone was still maintained to be that of a bird; until the question was finally set at rest, and the value of the microscopic test triumphantly confirmed, by the discovery of undoubted Pterodactyle bones of corresponding and even of greater dimensions, in the same and other chalk quarries.¹

¹ See Prof. Owen's Monograph on the British Fossil Reptiles of the Chalk Formation, p. 80, *et seq.*

CHAPTER XX.

INORGANIC OR MINERAL KINGDOM.—POLARIZATION.

454. ALTHOUGH by far the most numerous and most important applications of the Microscope, are those by which the structure and actions of Organized beings are made known to us, yet there are many Mineral substances which constitute both interesting and beautiful objects; being remarkable either for the elegance of their forms, or for the beauty of their colors, or for both combined. The natural forms of inorganic substances, when in any way symmetrical, are so in virtue of that peculiar arrangement of their particles which is termed *crystallization*; and each substance which crystallizes at all, does so after a certain type or plan,—the identity or difference of these types furnishing characters of primary value to the Mineralogist. It does not follow, however, that the form of the crystal shall be constantly the same for each substance; on the contrary, the same plan of crystallization may exhibit itself under a great variety of forms; and the study of these, in such minute crystals as are appropriate subjects for observation by the microscope, is not only a very interesting application of its powers, but is capable of affording some valuable hints to the designer. This is particularly the case with crystals of *Snow*, which belong to the "hexagonal system," the basis of every figure being a hexagon of six rays; for these rays "become incrustated with an endless variety of secondary formations of the same kind, some consisting of thin laminae alone, others of solid but translucent prisms heaped one upon another, and others gorgeously combining laminae and prisms in the richest profusion;"¹ the angles by which these figures are bounded, being invariably 60° or 120° . Beautiful arborescent forms are not unfrequently produced by the peculiar mode of aggregation of individual crystals; of this we have often an example on a large scale on a frosted window; but microscopic crystallizations sometimes present the same curious phenomenon (Fig. 345). In the following list are enumerated some of the most interesting natural specimens, which the Mineral kingdom affords as microscopic objects; these should be viewed by reflected light, under a very low power:—

¹ See Mr. Glaisher's Memoir on "Snow-Crystals in 1855," with a number of beautiful figures, in "Quart. Journ. of Micros. Sci." vol. iii, p. 179.

Antimony, sulphuret.
Asbestos.
Aventurine.
ditto, artificial.
Copper, native.
arsenate.
malachite-ore.
peacock-ore.
pyrites (sulphuret).
ruby-ore.

Iron, ilvaite or Elba-ore.
pyrites (sulphuret).
Lapis lazuli.
Lead, oxide (minium).
sulphuret (galena).
Silver, crystallized.
Tin, crystallized.
oxide.
sulphuret.
Zinc, crystallized.

455. The actual process of the *Formation of Crystals* may be watched under the microscope with the greatest facility; all that

is necessary being to lay on a slip of glass, previously warmed, a saturated solution of the salt, and to incline the stage in a slight degree, so that the drop shall be thicker at its lower than at its upper edge. The crystallization will speedily begin at the upper edge, where the proportion of liquid to solid is more speedily reduced by evaporation, and will gradually extend downwards. If it should go on too slowly, or should cease altogether, whilst yet a large proportion of the liquid remains, the slide may be again warmed, and the part already solidified may be redissolved; after

which the process will recommence with increased rapidity. This interesting spectacle may be watched under any microscope; and the works of Adams and others among the older observers, testify to the great interest which it had for them. It becomes far more striking, however, when the crystals, as they come into being, are made to stand out bright upon a dark ground, by the use of the spotted lens, the paraboloid, or any other form of black-ground illumination; still more beautiful is the spectacle when the Polarizing apparatus is employed, so as to invest the crystals with the most gorgeous variety of hues. The following list specifies the salts and other mineral substances, whose crystalline forms are most interesting. When these are viewed with polarized light, some of them exhibit a beautiful variety of colors of their own, whilst others require the interposition of the selenite plate for the development of color.

Acetate of Copper.
of Manganese.
of Soda.
of Zinc.
Agate (transparent sections).
Alum.

Arragonite (transparent sections).
Arsenate of Potass.
Bicarbonate of Potass.
Bichromate of Potass.
Bichloride of Mercury.
Boracic acid.

FIG. 345.



Crystallized Silver.

Borate of Ammonia.	Nitrate of Uranium.
of Soda (borax).	Oxalic acid.
Carbonate of Lime (from urine of horse).	Oxalate of Ammonia.
of Potass.	of Chromium.
of Soda.	of Lime.
Chlorate of Potass.	of Potass.
Chloride of Barium.	of Soda.
of Cobalt.	Oxalurate of Ammonia.
of Sodium.	Phosphate of Ammonia.
Cholesterine.	Ammoniac-Magnesian (triple, of urine).
Chromate of Potass.	of Soda.
Citric Acid.	Prussiate of Potass (red).
Cyanide of Mercury.	(yellow).
Granite (transparent sections).	Salicine.
Hypermanganate of Potass.	Sulphate of Ammonia.
Iodide of Potassium.	of Cadmium.
of Quinine.	of Copper.
Mannite.	of Copper, ammoniated.
Murexide.	of Iron.
Muriate of Ammonia.	of Magnesia.
Nitrate of Ammonia.	of Potassa.
of Barytes.	of Soda.
of Bismuth.	of Zinc.
of Copper.	Tartaric Acid.
of Potass.	Uric Acid.
of Soda.	Urate of Ammonia.
of Strontian.	of Soda.

456. It not unfrequently happens that a remarkably beautiful specimen of crystallization develops itself, which the observer desires to keep for display. In order to do this successfully, it is necessary to exclude the air; and Mr. Warington recommends castor-oil as the best preservative. A small quantity of this should be poured on the crystallized surface, a gentle warmth applied, and a thin glass cover then laid upon the drop, and gradually pressed down; and after the superfluous oil has been removed from the margin, a coat of marine glue or other varnish is to be applied.

457. Although most of the objects furnished by Vegetable and Animal structures, which are advantageously shown by Polarized light, have been already noticed in their appropriate places, it will be useful here to recapitulate the principal, with some additions.

<i>Vegetable.</i>	
Cuticles, Hairs, and Scales, from Leaves (§§ 220, 246).	Polypidoms of Hydrozoa (§ 305).
Fibres of Cotton and Flax.	Polyzoaries (§ 330).
Raphides (§ 230).	Spicules of Gorgonide (§ 309).
Spiral cells and vessels (§§ 228, 232).	Tongues (Palates) of Gasteropods (§ 347).
Starch grains (§ 229).	Scales of Fishes (§§ 408, 409).
Wood, longitudinal sections of, mounted in balsam (§ 239).	Sections of Hairs (§ 411).
<i>Animal.</i>	of Quills (§ 412).
Fibres and Spicules of Sponges (§ 296).	of Horns (§ 413).
	of Shells (§ 336).
	of Skin (§ 418).
	of Teeth (§§ 406, 407).
	of Tendon, longitudinal.

APPENDIX.

THE MICROSCOPE AS A MEANS OF DIAGNOSIS.

APPENDIX BY THE EDITOR.

THE MICROSCOPE AS A MEANS OF DIAGNOSIS.

Value of the Microscope in the Diagnosis of Disease.—During the past few years the Microscope, in the hands of the physician, has become an indispensable auxiliary in the detection and diagnosis of disease. The anatomist in his researches into the structure and functions of various organs, and the physiologist in his attempts to unveil the mysterious phenomena of life, alike find in this instrument a valuable coadjutor. Indeed, the invention of the microscope has added to the already extensive list of the sciences another—Histology—full of importance and interest, constituting as it necessarily does, the basis of pathology. The results flowing from its application to medical inquiries are so important, that it has, at length, been assigned a place in the same category as the stethoscope, pleximeter, speculum, and other well-tried instruments employed in physical exploration. By its aid such an extensive acquaintance with the intimate structure of the tissues of the animal economy, both in health and disease, has been obtained, that the practitioner can now pursue his difficult profession, with far more accurate and rational views of the nature and pathology of the various affections which he is called upon to treat, than were enjoyed by his predecessors. Individual cases are constantly occurring where long-known and well-attested methods of investigation have signally failed to elicit the information necessary to a rational treatment. Upon many such cases the microscope casts a flood of light. We have but to glance over the rapidly increasing literature of our profession, to discover many proofs of the obligations of practical and scientific medicine to the invention and judicious application of the microscope. By its aid the impositions so frequently practised upon the physician have been often detected. One of the most common of these attempts at deception consists in mixing various substances with urine, the patient pretending that he voided them by the urethra. Some of these, such as starch, flour, and sand, are readily detected by subjecting the fluid to

microscopic examination; and even where milk has been added, it can be distinguished by the presence of oil-globules, from the so-called chylous urine, in which the fatty matter is found in a molecular state. In the same way have been ascertained the nature and origin of many strange and unusual substances discharged from the bowels. Drs. Bennett, Todd, Quekett, and others have placed upon record, numerous instances of the value of the microscope, in detecting impositions and establishing a certain diagnosis in obscure cases of disease.

"Some years ago, I was summoned to see a Dispensary patient laboring under bronchitis, who was spitting florid blood. On examining the sputum with a microscope, I found that the colored blood-corpuscles were those of a bird. On my telling her that she had mixed a bird's blood with the expectoration, her astonishment was unbounded, and she confessed that she had done so for the purpose of imposition."¹

The malignant or non-malignant character of certain suspicious tumors has, on many occasions, been positively settled by recourse to the microscope, as the following example will show.

"An eminent surgeon, in London, was treating a case of what he considered to be pharyngeal abscess. Before opening it, however, he scraped off a little of the matter on its surface with his nail, and took it to Mr. Quekett, who told me that on examining it with a microscope, he found it to contain numerous cancer-cells. The tumor was allowed to progress uninterruptedly; and on the death of the individual, some months afterwards, the bones at the base of the cranium were found to be enlarged, from a cancerous growth."²

That medico-legal science has been greatly enriched and rendered far more certain in its results by the aid of the microscope, few persons will deny. The ends of justice have sometimes depended solely upon its power of detecting spermatozoa in cases of rape, of distinguishing between the stains of blood and those of colored fluids, or of pointing out the difference between human hair and that of animals.

The microscope has afforded valuable assistance to the pathologist, in disclosing the obscure processes by which changes or alterations in nutrition have gradually produced, in some of the most complex tissues of the body, the peculiar morbid condition known as fatty degeneration. In the hands of skilful observers, this instrument has taught us that apoplexy is not always dependent upon a plethoric or hyperæmic state of the cerebral vessels, but is, in many instances, the result of altered nutrition affecting the structure of these vessels, impairing their strength and elasticity, and otherwise altering their properties and functions. A brief microscopic examination of the urine is not unfrequently sufficient, as the laborious researches of Dr. John-

¹ Bennett's Introduction to Clinical Medicine.

² Bennett, *op. cit.*

son have shown, to reveal, during life, the existence of this peculiar pathological condition in the kidney.

But while, on the one hand, we urge upon the student the importance of the study of microscopy, and direct his attention, by way of proof, to the brilliant labors of the German School of Histologists; on the other, we must caution him against its exclusive cultivation, to the neglect of the other established and reliable modes of investigation of which the intelligent physician avails himself, in the daily routine of business. "You must not suppose," writes Dr. Bennett,¹ "that an additional method of gaining information implies abandonment of those, the utility of which has stood the test of experience. Men must learn the every-day use of their senses; must know how to feel, hear, and see, in the same manner as they did before instruments were invented. We don't see the stars less clearly with our naked sight, because the telescope is necessary for an astronomer. Neither should a physician observe the symptoms of a disease less accurately because he examines the chest with a stethoscope, or a surgeon be less dexterous with the knife, because it is only by means of the microscope he can determine with exactitude the nature of a tumor." "We should learn to distinguish between the mechanical means necessary for arriving at truths, and those powers of observation and mental processes which enable us to recognize, compare, and arrange the truths themselves. In short, rather endeavor to observe carefully and reason correctly on the facts presented to you, than waste your time in altering the fashion and improving the physical properties of the means by which facts are ascertained. At the same time, these are absolutely necessary; and perhaps no kind of knowledge has been so much advanced in modern times by the introduction of instruments and physical means of investigation, as that of medicine. These enable the practitioner to extend the limits to which otherwise his senses would be limited. I do not say employ one to the exclusion of the other, but be equally dexterous in the use of all. Do not endeavor to gain a miserable reputation as a microscopist, or as a stethoscopist; but by the appropriate application of every instrument and means of research, seek to arrive at the most exact diagnosis and knowledge of disease, so as to earn for yourselves the title of enlightened medical practitioners."

As with all other mechanical aids to the senses, the microscope, to be successfully applied in medicine, requires a degree of skill in its manipulation not to be acquired at once, but by repeated and persevering practice. Let the student, therefore, not be discouraged by the many failures and uncertainties which, to a greater or less extent, must necessarily accompany his early efforts. Let him remember that exact and accurate habits of observation are acquired slowly and almost insensibly, and that, in attempting to obtain proficiency in any practical art or science,

¹ Introduction to Clinical Medicine.

a methodical and systematic procedure is always requisite. So many unknown objects, so many strange and unusual forms, so many structural peculiarities are revealed to the eyes of the tyro in microscopy, that he is at once plunged into profound confusion, from which he can extricate himself only by adopting the most laborious and rigid system of observation. He should examine with the utmost care the physical appearance and character of the ultimate structures; he should note the exact shape of the object, whether round or oval, globular or flat, &c.; the peculiarities of its edge or border, whether fine and brilliantly illuminated, or dark and abrupt, whether smooth or rough, regular or irregular, serrated or beaded, &c. Peculiarities of color produced by strong and faint, and by reflected and transmitted light, should next claim his attention. The size of the object should, in all cases, be obtained by actual measurement, and all variations in diameter noted. The transparency must also be observed,—whether the body be opaque or diaphanous. If opaque, the degree of opacity must be stated, its causes, and the effects upon the transmission of the luminous rays. The superficial and deep-seated layers, and in the case of cellular and tubular bodies, the contents should also be investigated; and lastly, the effects of various reagents upon these physical properties must be ascertained with the same care and patience.

It will thus be seen that the successful application of the microscope to the diagnosis of disease, requires a very considerable acquaintance with the healthy appearance and structure of both the animal and vegetable tissues. Armed with this preliminary knowledge, the student will be surprised at the facility with which he will be enabled to distinguish from each other the various animal solids and fluids, different morbid products, the matters constituting food, &c., whether these be unchanged, or in a state of disintegration from the processes of mastication, digestion, ulceration, &c.

In view, however, of the great difficulty experienced in demonstrating accurately the histological character of the healthy tissues, and the still greater difficulty of making out the characteristics of morbid growths, the student should exercise great caution and deliberation in pronouncing an opinion upon the nature of any morbid tissues examined by him.

The student should early acquire the habit of recording all his observations in a note-book kept expressly for the purpose. He should exercise himself also in making drawings, as exact as possible, of all the objects he examines. Such a practice, though laborious, will leave upon his mind a more vivid and lasting impression of the various objects of his research, and gradually render him a very close and reliable observer.

EXAMINATION OF THE NERVOUS SYSTEM.

Brain.—As the nerve-fibres rapidly undergo change, the brain should be examined very soon after death, by depositing minute portions upon a perfectly clean slip of glass, and moistening them with serum or a weak saccharine solution. If it is desired to examine the distribution and arrangement of nerve-fibres, the brain should be placed in a solution of chromic acid; by the hardening thus produced it can be easily cut into thin slices by means of a Valentin's knife. The addition of water to a portion of white cerebral matter changes the natural appearance of these fibres, by separating the oily and albuminous contents of the tubular sheath. The oily matter collects into globules, giving a beaded appearance to the fibres. (Fig. 346, *g, g'*.) The nerve-tubes may be rendered very distinct by the addition of a dilute solution of caustic soda.

Small portions of the meninges of the brain may be examined in the same manner.


To examine the cerebral vessels, a thin section must be well washed, and subjected to gentle pressure. The vessels are thus deprived of their investing neurine, and may now be rendered more distinct by the dilute caustic soda.

The corpora amylacea, or gritty particles found in the pineal gland and other parts of the brain (Fig. 347), must be separated from the nervous tissue for examination by repeated washing in water.

Spinal Cord.—To examine the spinal cord with advantage, it should first be hardened in a solution of chromic acid, or in spirits of wine. The structure of thin sections is thus rendered quite conspicuous. The following method was employed by Mr. J. S. Clarke.¹


“A perfectly fresh cord was hardened in spirits of wine, so that extremely thin sections, in various directions, could be made by means of a very sharp knife. A section so made was placed on a glass slide, and treated with a mixture composed of one part of acetic acid and three of spirits of wine, which not only makes the nerves and fibrous portion more distinct and conspicuous,

FIG. 346.



but renders also the gray substance much more transparent. The section was then covered with thin glass, and viewed first

FIG. 347.




The lower figure represents a choroid plexus with several small tumors at * * *, supposed at first to have been tubercular; they proved to consist of aggregations of concentric corpuscles, cholesterol, and pure oil, united by areolar tissue; the concentric corpuscles which are shown above the plexus are magnified 100 diameters.

by reflected light with low magnifying powers, and then by transmitted light with higher ones.

“According to the second method, the section is first macerated for an hour or two in the mixture of acetic acid and spirit. It is then removed into pure spirit, and allowed to remain there for about the same space of time. From the spirit it is transferred to oil of turpentine, which expels the spirit in the form of opaque globules, and shortly (sometimes immediately) renders the section perfectly transparent. The preparation is then put up in Canada balsam, and covered with thin glass. By this means the nerve-fibrils and vessels become so beautifully distinct, that they may be clearly seen with the highest powers of the microscope. If the section be removed from the turpentine when it is only semi-transparent, we sometimes obtain a good view of the arrangement of the bloodvessels. This mode of preparation succeeds best in cold weather; for in summer, the cord, fresh when immersed in the spirit, remains more or less spongy, instead of becoming firm and dense in the course of five or six days. The spirit should be diluted with an equal quantity of water during the first day, after which it should

FIG. 348.



be used pure. Certain modifications of this mode of preparation

may be sometimes employed with advantage by a practised hand."

Nerves.—The structure and arrangement of nerve-fibres are best studied in the mesentery of small animals, as the newt; though with a little care in manipulation they can be very well displayed in any part of the nervous system. Their ultimate distribution, however, presents greater difficulties. Phosphoric acid, and solutions of caustic soda, and iodine of different strengths, are of great use in rendering these fibres more distinct. According to Dr. Waller, the tongue of the frog is best adapted for examining the arrangement of nerve-fibres in papillæ. When the nerve-fibres are not quite fresh, or have been soaked in water, and where they have been stretched or subjected to pressure for some time, their structure will be found to have undergone certain peculiar changes, as complete conversion into fibrous tissue, fatty degeneration, &c. (Fig. 348.)

EXAMINATION OF THE MUSCULAR SYSTEM.

Muscular Fibre.—*Sarcolemma.*—Muscular fibre is of two kinds, —the striated, voluntary, or muscular fibre of animal life; and the unstriated, involuntary, or muscular fibre of organic life.

The voluntary muscles of man and the lower animals furnish specimens of the striated fibre. They may be prepared for examination by cutting out a small slice from a muscle, separating the fibres with fine needles, and placing them upon a glass slide, and adding a drop or two of water. Muscles which have been boiled or hardened in chromic acid, corrosive sublimate, or spirits of wine, yield excellent sections for examination. The general anatomy of voluntary muscular fibre is well displayed in the thin slips of muscle lying just beneath the skin of small animals, as the frog; while the general arrangement and form of the fibres is well shown, according to Beale, in a transverse section of the pectoral muscle of a teal (*Querquedula crecca*), which has been put upon the stretch, and allowed to become perfectly dry.

The ultimate fibrillæ may be studied with advantage upon the muscular tissue of the eel and pig. The fibrillæ are separated and rendered distinct by maceration in chromic acid. From the back of the throat, after a meal of meat, in the discharges of cholera patients and in vomited matters, admirable specimens, showing the transverse striæ, may often be obtained. In examining the arrangement of the nuclei, solutions of caustic soda and acetic acid will be found very useful.

In the tongue of the frog, as shown by Kölliker, and in the upper lip of the rat, according to Huxley, peculiar fibres, known as branched muscular fibres, may be found. To obtain specimens of these fibres, the tongue of a frog is boiled for a short time in water. A piece of the mucous membrane is then dissected away,

and a very small portion of the submucous tissue cut from the edge of the tongue with a pair of sharp scissors. This is torn into fragments with very fine needles, and then placed in the field of a quarter-inch object-glass. If the tongue is boiled very long, the fibres become too brittle for separation and examination.

The crustacea, mollusca, and insecta present peculiarities in the structure of their voluntary muscular fibre which separate them in a marked manner from the higher divisions of the vertebrata.

FIG. 349.



The involuntary, smooth, or non-striated muscular fibres, though appearing like flattened bands (Fig. 350), in reality, according to Kölliker, consist of elongated cells. They are found in various situations, as in the alimentary canal, the large and small arteries, veins and lymphatics, the trabecular tissue of the spleen, the uterus, bladder, and urethra, &c.

FIG. 350.



“The contractile fibre-cells have been arranged in three classes:—

1. Short rounded or flattened cells, somewhat resembling epithelium.
2. Flattened bands, with fringed edges.
3. Long rounded or fusiform fibres, slightly wavy, and terminating at each end in a point.

FIG. 351.



“The first two varieties are obtained from the bloodvessels. The last form is met with in the intestinal canal, uterus, &c. These cells may be readily isolated by macerating small pieces of the muscular coat of the alimentary canal, &c., in dilute nitric acid, containing about 20 per cent. of strong acid. By a little

tearing, with the aid of fine needles, separate cells may be readily obtained."¹ (Fig. 350.)

The sarcolemma is best seen in the long muscular fibres of the fin of the skate, by tearing them apart with delicate needles and spreading them out upon a piece of glass. In the heart the sarcolemma is so thin that it can scarcely be detected. Some observers doubt its existence (Fig. 351).


EXAMINATION OF THE RESPIRATORY ORGANS.

Healthy Lung.—The mucous membrane of the trachea and bronchial tubes, and the parenchymatous structure of the lung may be readily examined by cutting thin sections with a very sharp knife, moistening them with water and spreading them out upon slips of glass in the ordinary manner. To examine the ciliated epithelium, and the characteristic movements of the cilia, the air-passages should be scraped and the matters thus obtained softened with serum instead of water, and deposited upon glass. The ciliary motion is well displayed in the branchiæ of the mollusca, as the oyster. The yellow elastic tissue of the lung is rendered quite distinct by the addition of acetic acid (Fig. 352).

To examine the vascular tissue, the lung should be injected with a somewhat thick solution of transparent gelatine. This oozing through the walls of the vessels, fills and expands the air-cells so that their forms and arrangement can be easily detected, while the vessels are seen in their natural position, and apparently deprived of epithelium (Fig. 353).

An excellent idea of the characteristic appearance of the trachææ of insects, may be obtained by separating with fine needles the viscera of a fly, placing them on a glass slide, and adding a few drops of water.

FIG. 352.



a, epithelium; b, elastic trabeculae; c, membranous wall, with fine elastic fibre.

¹ Beale. The Microscope, and its application to clinical medicine.

Morbid Lung.—Diseased lung is examined in the manner described above. Emphysematous, tuberculous and pneu-

FIG. 353.



Arrangement of the Capillaries of the air-cells of the Human Lung.

mous, tuberculous and pneu-
monic lungs present points of the highest interest to the student of pathological histology.

“A small fragment of tolerably firm miliary tubercle, squeezed between glasses with a drop of water, and examined under a magnifying power of 250 diameters linear, presents a number of irregular shaped bodies, approaching a round, oval, or triangular form, varying in their longest diameters from the 1-4000th to the

1-2000th of an inch. These are the so-called *tubercle-corpuscles*. They are composed of a distinct wall, containing generally three or more granules without any distinct nucleus, and are mixed with numerous granules and molecules, varying in size from a point scarcely measurable to the 1-6000th of an inch in diameter (Fig. 354, *a*). If we add to these bodies a drop of weak acetic acid, all the corpuscles become more transparent, but are otherwise unchanged, and many of the granules disappear, as in Fig. 354, *b*.”

FIG. 354.



Yellow tubercle treated in the same manner presents similar corpuscles imbedded in a molecular and granular mass (Fig. 355). Sometimes these corpuscles are observed to be larger and rounder, resembling in this respect those of scrofulous pus (Figs. 356, 357). Occasionally tuberculous matter will be seen to consist almost entirely of granules, and at other times of very minute molecules.

Yellow tubercle treated in the same manner presents similar corpuscles imbedded in a molecular and granular mass (Fig. 355). Sometimes these corpuscles are observed to be larger and rounder, resembling in this respect those of scrofulous pus (Figs. 356, 357). Occasionally tuberculous matter will be seen to consist almost entirely of granules, and at other times of very minute molecules.

FIG. 355.




FIG. 356.



FIG. 357.



A thin section of the gray, semi-transparent granulation is very different in appearance from ordinary tubercle. The constituent elements, though more transparent, are less distinct (Fig. 358). In the cretaceous and calcareous forms of tubercle

the corpuscles and granules are mixed with gritty saline particles of an irregular form and size. Crystals of cholesterine are sometimes found in the cretaceous and cheesy varieties of tubercle (Fig. 359). If no crystals can be detected, a small quantity

FIG. 358.





FIG. 359.



FIG. 360.



Section of gray granulations after addition of acetic acid, showing air-vesicles filled with tubercles.

of alcohol may be added to a portion of the tuberculous mass, and then evaporated. As the evaporation proceeds, the crystals will be formed.

Thin sections of calcareous lung present a granular appearance, in consequence of the close aggregation of the minute earthy particles. Fragments of the calcareous mass may be broken off, and examined with the low powers of the microscope, as in the case of opaque objects generally. A drop of acetic acid added to these fragments causes them to dissolve with effervescence, showing the presence of carbonate. If this solution be treated with excess of ammonia, phosphate of lime

FIG. 361.




FIG. 362.



will be precipitated; solution of the oxalate of ammonia will also detect the presence of lime.

Irregular, black masses of pigmentary matter, consisting of exceedingly minute molecules, are also frequently found, mixed with tubercle (Fig. 360), giving the tissues a black or bluish

tinge. As the tubercle becomes older, the pigmentary matter generally increases in quantity. It also varies in chemical composition according to its situation. That obtained from the lungs and bronchial glands is pure charcoal and chemically indestructible; that found in the peritoneum is destroyed by the action of alcohol and the mineral acids.

Gulliver, Vogel, and Schroeder Van der Kolk affirm, that nucleated cells may be observed in miliary tubercle; but both Lebert and Bennett deny it. (Figs. 361, 362.)

It is oftentimes very difficult to distinguish tubercle from fibrinous exudations and from cancerous growth. "If we are asked," writes Prof. Bennett, "to determine what is positively tubercle, as distinguished from all other morbid products, we must answer, that deposition which is composed of the peculiar corpuscles and granules described and figured in the preceding pages. From pus-corpuscles they are readily distinguished by the action of acetic acid, which in them causes no granular nucleus to appear. From plastic corpuscles they may be separated by their irregular form, smaller size, and the absence of primitive filaments. With the granular corpuscle they can scarcely ever be confounded, on account of its large size, brownish or blackish color, and nucleated or granular structure. The cells of cancer are large, transparent, and distinctly nucleated, and, consequently, easily distinguished from the small, non-nucleated corpuscles of tubercle." "The only other structure likely to be confounded with tubercle is the reticulum of cancer, which not only presents a yellowish appearance closely resembling it, but is composed of nuclei and molecular matter resulting from the disintegration of cancer-cells. But, as this reticulum is always associated with cancerous formation, it may at once be distinguished by the cell-elements which accompany it. It should further be noticed that every form of exudation, at a certain period, presents a molecular and granular structure throughout, and that then it becomes impossible to determine its nature, unless it be associated with the more characteristic



Fig. 363. A, Tubercle corpuscle from lung; B, Pus corpuscles. Fig. 364. Plastic, or pyoid corpuscles. Fig. 365. Granular corpuscles from cerebral softening. Fig. 366. Corpuscles in reticulum of cancer.

elements distinctive of the simple, tubercular, or cancerous exudations." The accompanying figures will illustrate these distinctions. (Figs. 363, 364, 365, 366.)

Where a cavity is found in the lung, separate examinations should be made of its contents, the surface of the walls, and the subjacent tissue.

In emphysema of the lungs, the investing membrane will be often found full of very small holes; the vessels elongated, and the interspaces much enlarged; and the yellow elastic fibres stretched to such an extent as to be deprived of their elasticity.

In the first stage of pneumonia, the vesicular walls contain here and there collections of minute granules. The epithelial cells are separated from the basement membrane; and while the nuclei are unaltered, the cell-contents have become a little more granular. The minute capillaries are very much congested. As the engorgement proceeds, granules collect in greater quantities in the air-cells; while in the effused serum may be detected distinct blood-corpuscles, small nucleated cells, and exudation-corpuscles. These changes are shown in Fig. 367.




FIG. 367.

In the second or stage of red hepatization, the cavity of the air-vesicle is filled with cells about 0.018 of a millimetre in size, of very varied shapes, and differing in respect to

FIG. 368.





FIG. 369.



their contents. These are mixed with concrete albumen, oil-granules, and free fat. (Figs. 368, 369.)

In the third stage—that of gray hepatization—large well-marked cells may be seen, containing granules and oil. Most

of these cells are without nuclei. Occasionally they are found associated with large masses of pus-cells. The epithelium is wanting, and free fat-molecules and globules are abundant. (Fig. 370.)¹

FIG. 370.



EXAMINATION OF THE GLANDULAR SYSTEM.

Liver.—The relation to each other of the constituent elements of

FIG. 371.



Transverse section of a lobule of the human liver, highly magnified, and presenting to view the reticulated structure of the biliary tubes. In the centre of the figure is seen the hepatic vein cut across and several small branches terminating in it. Where the injecting matter did not run freely, it is seen standing in dots along the course of the vessels. At the periphery are seen branches of the hepatic artery, vena portarum, and hepatic duct.—After Leidy.

the liver, may be easily demonstrated upon thin sections cut out of the fresh liver of a pig, with a Valentin's knife. The arrangement of the minute vessels are best seen in the injected liver of a frog. The larger vessels can be studied in thin sections, from which the cells have been washed away by a stream of water, and dilute caustic soda afterwards applied. Excellent illustrative specimens of the vessels may be obtained, by injecting the liver of some of the lower animals, as the frog, with two different colors; throwing one into the portal vein, and the other into the hepatic artery or vein, so that the two shall meet in the capillaries. (Fig. 371.)

The hepatic cells lie in the

¹ See a valuable paper by Dr. Da Costa, of Philadelphia, on the Pathological Anatomy of Acute Pneumonia, in Amer. Journ. of Med. Sciences, Oct. 1855.

matrix or network formed by the union of the vascular and the condensed cellular tissues. They are of different sizes, though generally very small, and contain a few oil-globules. They may be obtained by scraping a freshly-cut surface; placed upon a slip of glass and moistened with a few drops of water, they are ready for examination. These cells undergo various changes in disease. Sometimes they are withered and shrunk; sometimes filled with pale granules, as in diabetic cases, and at others, gorged with fat to such an extent as to obliterate the cell-walls, and give to the liver the appearance of ordinary adipose tissue. (Fig. 372.)

FIG. 372.



Kidney.—The general arrangement of the straight and convoluted tubes, and the varying appearance of their epithelium, may be well shown upon thin sections cut out of the cortical and medullary portions of the kidney, by means of a Valentin's knife. By scraping the freshly-cut surface, epithelial cells may be obtained, mixed, however, with Malpighian tufts and fragments of tubes. (Fig. 373.) In the convoluted tubes the epithelium is thick and glandular; in the straight it presents a scaly appearance. Care must be taken not to confound the tubes, as they bend in and out through the matrix, with circumscribed cysts. The kidney of the mouse and of many of the other rodentia, are better adapted for demonstrating the matrix than the human kidney. Indeed, its existence in the latter has been doubted by some, not-

FIG. 373.



A, Portion of uriniferous tube lined by epithelium. B, Epithelial cells, highly magnified. C, Portion of tube from medullary substance, deprived of epithelium.

withstanding that Goodsir, Kölliker, and Johnson have described and figured it. "With care," says Mr. Beale, "I believe it may always be demonstrated in the healthy human kidney; and in some specimens of fatty kidney, as well as in the small contracted kidney of chronic nephritis, it may very readily be seen by washing a thin section with a stream of water, in order to remove the epithelium and remaining portion of the tubes. The matrix appears to consist of very fine fibres, amongst which no indications of the yellow element can be detected. By the addition of acetic acid it becomes more transparent, and a few granules are developed, but no other change is produced."

In the uriniferous tubes of the frog and newt, ciliary motion is beautifully shown. The arrangement of the vessels of the Malpighian tufts can be studied with facility and success in the large tufts of the kidney of the horse.

Sections of kidney are best kept in large thin glass-cells containing a solution of creasote or weak spirit and water.

Morbid Kidney.—In disease the kidney oftentimes becomes quite opaque. This condition may result from a variety of causes; as, hypertrophy of the matrix, deposition of fatty matter, and unusual accumulation of epithelium or oil-globules in the tubes. To examine such a kidney, therefore, the sections should be exceedingly thin. The matrix and the vessels, which are often much thickened in chronic nephritis, should be prepared for examination by the addition of acetic acid, or dilute caustic soda, and subsequent washing with clear water to remove the epithelium and granular matter from the tubes.

Salivary Glands.—To examine the structure of the salivary glands, thin sections of the fresh gland should be treated first with acetic acid, and then with caustic soda in excess. The epithelium of the ultimate lobules is rendered distinct by soaking the section in acetic acid. In the ducts, large cells filled with oil-globules may sometimes be detected.

Thymus and Thyroid Glands.—Sections of the recent glands should, before being submitted to examination, be washed with water to clean away the soft and pulpy portions, which are apt to obscure the structure of the tissues. The relation of the lobules and other constituents is best shown in sections hardened with chromic acid; though it must be borne in mind that this process alters the natural appearance of the cellular tissue. The membranous thyroid gland of small animals is well adapted for microscopic examination.

Adipose Tissue.—To examine adipose tissue, small masses of fat-cells may be taken from the subcutaneous areolar tissue, and exposed to reflected light under a low power; or thin sections, moistened with water, may be placed between two pieces of glass, and examined by means of transmitted light. The arrangement of vessels can only be demonstrated upon an injected preparation. Small acicular crystals, of margaric acid and margarine,

disposed in a stellate (Fig. 374, *a*) form, are sometimes observed, especially in specimens obtained from emaciated subjects. In disease, the fat-cells are sometimes found degenerated, and containing a serous fluid, in which the nucleus is quite distinctly seen, amidst numerous granules. Sometimes the cell is shrivelled and irregular in form; frequently it assumes an angular shape.

Fatty degeneration consists in the conversion of healthy structures into true adipose tissue. The muscular system seems to be most liable to this change.

Prior to the appearance of fatty degeneration in voluntary muscle, the transverse striæ disappear. According to Mr. Quekett, the first trace of this disease is marked by a disturbance of the particles of myoline, which appear as so many very minute granules scattered irregularly within the sarcolemma, leading one to suppose that the delicate cell around each particle had given way, thereby allowing the myoline to escape, and destroying all regularity both of the transverse and longitudinal markings. As the disease progresses, the myoline is replaced by minute highly-refracting globules of oil, until at last the whole sheath is full of them.¹

The muscular fibres of the heart, and especially those of the *musculi pectinati*, afford frequent instances of the change in question. Muscles which have been long disused, as in cases of paralysis, club foot, &c., exhibit this species of degeneration in a striking degree. "Fatty degeneration appears also to occur in osseous tissues, and indeed the disease termed *Mollities ossium* is of this nature. All bones so affected have thin walls, are always more or less soft, and contain an abundance of oil. I have examined the bones in several cases, and found that the disease first commences in the bone-cells, the cell itself becoming larger and larger, its canaliculi disappearing, and several of these cells uniting to form a cavity, in which oil-globules soon make their appearance, all the parts of the bone in the neighborhood of the cells becoming at the same time thin and transparent from the removal of the granules of earthy matter." (*Quekett*.) In the *Lancet*, for 1850, the student will find an interesting paper, by Mr. Canton, on the *Arcus Senilis*, produced by fatty degeneration of the cornea. In fatty degeneration of the kidney, the epithelial cells become filled with numerous oil-globules, to such an extent sometimes as to burst and be discharged in fragments, leaving the surface of the tubules in some places almost bare.

FIG. 374.




¹ Lectures on Histology.

EXAMINATION OF THE VASCULAR AND ABSORBENT SYSTEMS.

Vessels.—The examination of the minute vessels requires but little previous preparation. A piece of the pia mater, or the mesentery of a young child, or a small artery from which the cerebral neurine has been gently washed, may be placed for this purpose under the microscope. The epithelial cells of the lining membrane, and the contractile fibre-cells may be rendered distinct by the addition of acetic acid. To demonstrate the fibre-cells of the contractile coat, Dr. Beale recommends that an artery of moderate size, and not quite fresh, be slit up, and its lining membrane removed by careful scraping; the subjacent elastic tissue is then to be removed and torn to pieces with fine needles, and finally placed upon a glass slide and moistened with a few drops of water. The spindle-shaped or muscular fibre-cells are readily obtained from the renal veins.

FIG. 375.



It is highly important that the student should make himself well acquainted with the healthy appearance of the minute arteries of the brain, since they suffer remarkable changes in disease. In white softening of the brain, they undergo a sort of fatty degeneration, numerous minute oil-globules being aggregated together at short intervals along their walls. These oily masses are readily detected by their high refracting power. (Fig. 375.)

The vessels of the kidney are also, from the changes they suffer in disease, worthy of especial investigation. To prepare them for examination it is only necessary to wash out from a thin section the epithelium of the renal tubes, and add a few drops of acetic acid to render them more distinct. The distinct arrangement of the nuclei of the circular and longitudinal fibres, and the greater thickness of their walls, will serve to distinguish the arteries from the veins. The coats of the latter are very thin.

Thickening of the arterial coats of the corpora Malpighiana is well seen, according to Dr. Johnson, in the small, contracted drunkard's kidney.¹ The normal thickness of the Malpighian artery is about one-fifth or one-sixth the diameter of the vessel; in this disease it is increased to one-third.

Some observers have thought that they detected epithelial cells upon the external surface of the Malpighian vessels; but the researches of Mr. Bowman negative this opinion.

¹ Diseases of the Kidneys.

Plates and rings of bones, and atheromatous deposits, containing oil-globules, granules, and crystals of cholesterine, are all

FIG. 376.




FIG. 377.



sometimes found in the larger vessels. (Fig. 376, 377.)

Lymphatics.—It is very difficult to examine the minute structure of the lymphatic glands. The fluid which exudes from a freshly cut surface may be examined by mixing it with water and placing it between two pieces of glass. It is very important to distinguish between lymphatic cells on the one hand, and pus-globules and white corpuscles of the blood on the other; not only as regards their appearance, but also in relation to the different effects produced by reagents.

EXAMINATION OF THE SKIN, MUCOUS AND SEROUS MEMBRANES, ETC.

Skin.—A good view of all the structures composing the skin may be obtained by making a vertical section. In this manner the relation of the various layers, the arrangement of the hair-bulbs and sebaceous follicles, and the position and course of the sweat-ducts, may all be demonstrated.

The following method of procedure is recorded by Dr. Beale, as quoted from Giraldis by Kölliker:

“The skin should be perfectly fresh, and a piece about two inches square, or rather less, is to be stretched with the outer surface downwards upon a thick deal board by means of numerous pins. If the sudoriferous glands are to be included in the preparation care must be taken to leave sufficient of the cellular tissue adhering to the skin. The piece of skin is allowed to dry by exposure to the air. Several small pieces taken from various parts of the body may be pinned out on the same board, care being taken to attach a label to each. Specimens may be taken from the scalp, eyelids, chin, mamma, axilla, arm or leg, palm of the hand, tips of the fingers, scrotum, and sole of the foot. With these, the varying thickness of the epidermis and other peculiarities in the different regions may be demonstrated.

“The portion of skin being quite dry, it is to be removed from the board, and, after cutting off the edge, several thin sections may be made by the aid of a very sharp knife through the whole thickness. In order to obtain a good specimen of the spiral portion of the sweat-ducts, the skin of the heel should be selected, and the section should be made parallel with the furrows, and in a slightly slanting direction, instead of at a right angle with the surface.

“The sections may next be placed in a watch-glass, with a few drops of clean water, and in the course of a short time it will be found that they have again attained the original thickness of the skin, in consequence of the absorption of water. They may now be submitted to examination, and after selecting a satisfactory specimen, it may be mounted in weak spirit and water, Goadby's solution, or other preservative fluid; or, the specimen may be washed in water, placed upon a slide, and allowed to dry slowly by spontaneous evaporation (when it will be found to have adhered tightly to the glass), and mounted in Canada balsam, with the usual precautions.”

Any opacity of the preparation may be removed by a weak solution of potash or caustic soda. Soaking in ether will dissipate the fat. The sweat-glands are made more distinct by soaking the tissue in a mixture of one part of nitric acid and two of water.

Large flakes of cuticle may be obtained for examination by exposing a small piece of skin to a moist atmosphere for several days. The superficial cells of the cuticle are brought into view by scraping the surface of the skin with a knife. These cells are flattened and adherent, and present a scaly appearance. The deeper-seated epidermic cells are more or less round, and appear to rest upon a layer of very minute granules mixed with coloring matter. The deep cells are soluble in acetic acid; the superficial are not. On the under surface of the cuticle are found a number of depressions, which receive the tactile papillæ of the cutis vera.

The papillæ may be studied either upon a vertical section made in the manner above described, or upon a section of the cutis itself, the cuticle having been first removed. From their large size, the papillæ of the skin of the dog's foot are well adapted for examination. The papillary vessels are best seen in an injected specimen, while the nerves and “axis-corpuscles” are sometimes brought into view by the addition of acetic acid or a weak solution of caustic soda.

The pigmentary cells are best seen in the skin of the negro, in that of some of the lower animals, and also in freckled surfaces.

Cutaneous Eruptions, Growths, Ulcers, &c.—Corns, callosities, and condylomatous warts consist of condensed epidermic scales. In *Veruca achrocordon*, the scales are collected around a central canal supplied with bloodvessels. Small cutaneous tumors are sometimes formed by the thickening of the subjacent areolar

tissue. This appears to be the case in elephantiasis, where the hypertrophy is increased by the effusion of plastic lymph into the areolæ and its subsequent organization.

The squamous eruptions of the skin,—ichthyosis, pityriasis, psoriasis,—all consist of collections of epidermic scales. In pityriasis they are placed loosely together; in psoriasis they are more aggregated; and in ichthyosis very much condensed.

A number of epidermic scales, arranged into the form of a capsule, constitute a favous crust. This capsule is lined by a mass of very fine granules, from which sprout cryptogamic plants in the greatest abundance.

Upon healthy granulating surfaces may be seen pus-corpuscles, fibre-cells in various stages of development, and newly-formed fibres. (Fig. 378.) In scrofulous and unhealthy sores, the broken-down pus bears some resemblance to tubercle-corpuscles. The epithelial ulcer or cancer, as it is commonly, but erroneously called, generally commences on the lip as a small induration or wart, which soon softens in the centre, while the hardened edges extend over the cheek and chin. This softened matter consists of epithelial and fibre or fibro-plastic cells. (Fig. 381.) Sometimes the cells are large and flat, and contain numerous fat-molecules and granules. (Fig. 380.) According to Bennett, the so-called

FIG. 378.



Fibre-cells passing into fibres.

FIG. 379.



Fibrous tissue formed from fibre-cells.

FIG. 380.




FIG. 381.




Fig. 380. Altered epithelial cells, from ulcer of lip. Fig. 381. Epithelial and fibre-cells, from ulcer of lip.

chimney-sweeps' cancer of the scrotum is essentially a similar formation, consisting externally of flattened epithelial scales passing into fibres; and, deeper-seated, either groups of younger

cells, or concentric layers of aggregated scales. The distinctions between these growths and true cancer will be pointed out hereafter.

Mucous Membrane.—The submucous areolar tissue may be examined upon a small piece cut from the under surface of the mucous membrane, and torn up with needles. Between the proper muscular coat of the small intestine and the basement-membrane, and in close apposition with the latter, Brucke has demonstrated a thin layer of pale muscular fibres, which is known as the muscular layer of the mucous coat. The contractile fibre-cells of this coat are disposed in circular and longitudinal directions. The villi are best seen in a vertical section of the membrane. By washing off the epithelium, and adding a solution of acetic and nitric acid, composed of about one part of acid to four of water, the muscular fibres of the villi will be brought into view. The villi situated around intestinal ulcers are often found to be very much elongated. In the examination of such ulcers it is important in all cases to ascertain if the muscular coat has suffered from the ulcerative action or not. The presence or absence of non-striated fibres in the base of the ulcer will determine this point.

Serous and Synovial Membranes.—Serous membranes consist mainly of condensed areolar tissue, containing an abundance of yellow elastic fibres. At the surface this areolar tissue is very dense; the deeper layers are less dense, and often contain fat-cells. Portions of recent membrane are generally necessary to demonstrate the delicate surface-cells. The fibres of the sub-basement tissue, and often the vessels and nerves, are well seen in the peritoneum of the mouse and other small animals. The vessels of the synovial membranes should be injected before examination. In an injected specimen, the distribution of the vessels in the fringe-like processes which dip down into the joint, is displayed to great advantage. In some cases of disease, as in ascites, and pleurisy of long standing, great alterations take place in the structure of the serous membrane, such as the deposition of a thick cellular layer over the whole surface. Cells of a similar character are also found in the fluid contained in the cavity.

Epithelium.—Epithelium may be obtained for examination by scraping a mucous or serous surface with a sharp knife. It should then be placed upon a slip of glass, and moistened with water. Very delicate cells should be treated with serum, syrup, or a mixture of glycerine and water, in preference to pure water, as the rapidity of endosmose is checked, and the liability to rupture diminished. Acetic and nitric acids, tincture of iodine, and solutions of potash and soda of different strengths, are the most useful reagents in examining epithelium. Various kinds of epithelium are described by histologists.

Scaly epithelium may be procured from the vagina, the mouth, &c.; a modified form exists in the epidermis, in nails, and in hair. The vaginal epithelium consists of large, flat, ragged, and

very irregular cells, folded over each other, and perhaps creased in different directions. The epithelial cells of the mouth have very distinct nuclei, which are made to disappear under the action of acetic acid.

Tessellated or pavement epithelium is beautifully shown in the epidermis of the frog. It may be examined upon the choroid coat of the eye, the lining membrane of the heart, arteries, veins, and pelvis of the kidney, and upon serous surfaces generally.

Glandular or spheroidal epithelium consists of round cells, which occasionally become polyhedral from mutual pressure. The nucleus is usually well marked, and sometimes seems to be surrounded with numerous minute granules and oil-globules. This variety of epithelium is well shown in the sweat-glands, in the convoluted tubes of the kidney, in the follicles of the stomach, pancreas, liver, &c.

Columnar, prismatic, or cylindrical epithelium, may be obtained from the gall-bladder, the ureters, the urethra, the intestinal villi, and the follicles of Lieberkühn.

Ciliated epithelium is found in the human body in the following situations:—On the surface of the ventricles of the brain, and on the choroid plexuses; on the mucous membrane of the nose and its sinuses; on the upper and posterior part of the soft palate, and in the Eustachian tube; in the cavity of the tympanum; on the membrane lining the frontal and sphenoidal sinuses; on the inner surface of the lachrymal sac and lachrymal canal; on the mucous membrane of the larynx, trachea, and bronchial tubes; upon the os uteri; within the cavity of the uterus; throughout the whole length of the Fallopian tubes, and upon their fimbriated extremities (*Beale*).

For examination, this variety of epithelium may be obtained from the back part of a frog's mouth, or from the branchiæ of an oyster or mussel.

EXAMINATION OF THE EYE.

The cornea is examined by dividing the ball transversely with a sharp knife, washing the anterior half, and removing the ciliary processes. It is then pinned out flat and allowed to dry; thin sections are next made with very fine scissors; these sections are moistened with water, and finally treated with acetic acid, in order to render the structures distinct.

The elements composing the retina are most satisfactorily examined in microscopic sections made at right angles to the surface of the membrane, after maceration in dilute solution of chromic acid. Viewed in this manner, according to Prof. Good-sir, it exhibits from the peripheral to the central margin of a successful section a series of strata, which may be distinguished as the bacillary, white cellular, gray cellular, filamentary, and liminary layers.

There are several methods of preparing the crystalline lens for examination. Minute portions of the recent lens may be moistened with water and placed under the microscope. The lens may be hardened in chromic acid, or soaked in oil for some time, and thick sections then made. To examine the fibres of the lens, the latter should be boiled, and the fibres torn off and separated with needles. In cases of cataract, the soft, pulpy, external portion of the lens will be seen to contain numerous oil-globules, consisting, according to Beale, chiefly of cholesterine dissolved in an oily fat. Larger globules of a different character are also observed.

EXAMINATION OF THE HARD TISSUES.

Bone.—For the methods of cutting, grinding, and polishing thin sections of the hard tissues, the student is referred to the more elaborate works on the application of the microscope to clinical medicine, and to the chapter on this subject, in the preceding pages, by the author.

A thin section of bone, viewed by transmitted light and with a low power, presents numerous round or oval apertures. These are the orifices of the Haversian canals. In the flat bones these canals are radiating and parallel to the surfaces; in the long bones, they are parallel to the axis. They communicate with each other by transverse and oblique branches, and vary in size from about 1-1000 to 1-200".

Under a higher power (200) the lacunæ, bone-corpuscles, or bone-cells, and the canaliculi or calcigerous canals, become visible, appearing like a number of dark spots with delicate lines radiating from their sides. Their dark appearance is due to the contained air, which is readily dissipated by immersion in oil of turpentine. They are oblong and flattened, and vary exceedingly in size.

The laminated structure of the cartilage or osseous basis of bone is best seen by previously digesting sections in hydrochloric acid diluted with water (one part to twenty), in order to remove the inorganic matter. The laminæ have a fibrous appearance, and are arranged either parallel to the surfaces of the bone, or in concentric layers around the Haversian canals.

The nuclei of cartilage-cells may be rendered very distinct by boiling the cartilage for two or three minutes in water or a solution of caustic soda.

The different steps in the development of bone may be very well studied upon the long bones of young animals, and also upon the ossifying bones of the cranium.

Thin sections of bone may be preserved in the dry state; thick specimens should be mounted in thickened Canada balsam. Sections of cartilage are best kept in weak spirits of wine, or in weak solutions of creasote.

Teeth.—Owing to the great hardness and brittleness of the

enamel, and its tendency to chip off, considerable difficulty is experienced in obtaining sections of teeth for examination. Sections having been obtained, they may be moistened with water, turpentine, and Canada balsam, and examined by transmitted light with low powers. A tooth that has been macerated in strong acid for several days can be very readily cut in any direction.

The dentinal tubules are microscopic canals, pursuing a waving and anastomosing course through the whole thickness of the dentine, from the wall of the pulp-cavity to the cement and enamel. According to Kölliker, each canal has a special wall, rather less in thickness than its diameter, which can only be observed in transverse sections, as a narrow yellowish ring surrounding the cavity; in longitudinal sections, on the other hand, it is almost entirely invisible. These tubules may be isolated from each other by long maceration in acid, and subsequent soaking in dilute caustic soda or potash.

The enamel consists of hexagonal or pentagonal prisms, one extremity of which is attached to Nasmyth's membrane, the other to the dentine. These prisms are very closely united, having no intervening substance between them.

The development of the teeth may be studied in the lower animals, or in human embryos at different ages.

Nails.—By maceration in water the nail may be separated from the skin and thin sections made by means of a sharp knife.

A nail consists of two layers, the upper or horny one forming the true nail, and marked with ridges on its lower surface, while the under soft one is continuous with the rete mucosum of the skin, from which it differs by the cells being elongated and arranged perpendicularly. The horny layer is composed of flattened epidermic cells aggregated into laminae. The addition of caustic potash or soda in solution causes the nuclei to be developed.

Hair.—A white hair is best adapted for demonstrating the intimate structure of the shaft and other parts. For this purpose, solutions of soda and potash and strong sulphuric and acetic acids are found to be the most useful reagents.

The cortex, which constitutes a large part of the shaft, presents upon its surface a great number of longitudinal striæ or interrupted dark lines and dots. Upon treating it with sulphuric acid at a gentle heat, as recommended by Kölliker, it is first changed into plates and fibres, varying in length and breadth, and afterwards converted into elliptical or spindle-shaped cells, which become flattened and angular from pressure. These cells contain elongated, dark-looking nuclei and pigment-granules, to which the color of the hair is due. The addition of caustic potash or soda will isolate these pigment-granules, and sometimes cause them to exhibit molecular motion. The presence of small spaces containing air, and the unequal refraction of light by dif-

ferent parts of the cells, give a striated or dotted appearance to the shaft.

The medulla consists of numerous angular or rounded cells, containing granules or globules of fat, and arranged in one or more linear series.

The cells of the cuticular coat are somewhat flattened, and quadrangular in form, the margins are black, and they are without a nucleus.

The fibrous layer of the hair-follicle consists of an outer membrane composed of areolar tissue, having longitudinal fibres, with long, spindle-shaped nuclei, and an inner delicate membrane, consisting of transverse fibres, with long and narrow nuclei. The pulp of the hair consists of fibrous areolar tissue, containing nuclei and granules of fat, but no cells.

EXAMINATION OF MORBID GROWTHS.

Morbid growths are of common occurrence, and are found growing in different parts of the body; externally upon the surface, and internally in the solid parenchyma of the viscera.

Sometimes these growths, as in the case of fatty and certain fibrous tumors, bony exostoses, &c., consist of a simple hypertrophy or rapid development of the tissues of the part in which they are located; very frequently, again, they are highly complex in structure, and differ more or less from the surrounding tissues. It is exceedingly difficult to classify them, even for the purposes of study, since the microscopic characters of one run into and blend almost imperceptibly with those of another. Thus the so-called benign tumors present various shades of transition into the malignant forms. It will be seen, therefore, that the varieties of these growths are numerous.

Sometimes the cutaneous epithelium undergoes an unusual development, constituting warts. The subcutaneous areolar tissue of the foot, leg, scrotum, and other parts, may be hypertrophied to such an extent as to occasion the most serious results.

In making an examination of morbid growths, "the secretion, if such exists on the surface of the tumor, should be first separately examined: secondly, the microscopical characters of the juice which exudes from the freshly-cut surface should be ascertained; and, lastly, a thin section ought to be made, in order to determine the relation of the constituents of the tumor to each other, and especially the proportions in which the different elements are present. Its connection with surrounding structures may be seen by examining a thin section, which should include a portion of the adjacent texture; and these observations should be made first with low powers, and afterwards with a power of about two hundred diameters." (*Beale.*)

The arrangement and direction of fibres; the form, size, and contents of cells; the presence or absence of nuclei, granules,

&c., and the effects of different reagents, should be carefully observed, and noted in a book kept for that purpose.

Cancer and Cancroid Growths.—“Cancer, or carcinoma, is a vascular, morbid production, characterized by a form of organic cell, which is peculiar, and never enters as a constituent in any normal tissue. It is usually deposited in the form of tumors, but occasionally as an infiltration, in any of the organs of the body, and the circumstances which give rise to its development are yet unknown to us.

There are several varieties of cancer, and the physical elements which ordinarily enter into their composition are as follow :

1. The characteristic cancer-cells, which are spherical, ovoid, irregularly polyhedral, and frequently exhibit caudate prolongations. They average about the .02 mil. in diameter, and possess finely granular contents, with a round or oval nucleolated nucleus, as large or larger than a pus-corpuscle. Sometimes cancer-cells are double the ordinary size, or more, and not unfrequently contain several nuclei, or even other cells, constituting parent or endogenous cells.

2. Nuclei, which are spheroid or oval, and resemble those within cancer-cells.

3. Granules, and amorphous liquid or semi-solid matter.

4. Fusiform or fibro-plastic cells. These are liable to be confounded with the characteristic cancer-cell, but usually may be distinguished by the smaller nucleus, and the disposition to elongate at opposite extremities, and pass from this condition into the form of bands or fibres.

5. Fibrous tissue; most usually of the white variety, but not unfrequently mingled with elastic fibres.

6. Black pigment, in granules, or contained within cells.

7. Fat, in granules, globules, and in the form of adipose cells.

8. Vessels.

The varieties of cancer are encephaloid, or medullary carcinoma, scirrhus, colloid, melanosis, and fungus hæmatodes.

Encephaloid is that form in which the cancer-cell predominates over every other constituent. Occasionally, the cancer-cells exist in it to the exclusion of all other matters, except liquid, granules, and vessels.

In scirrhus, fibrous matter predominates, and encloses the cancer-cells within the areolæ.

Colloid is composed of a fibrous stroma, with loculi, filled with a gelatinoid matter and cancer-cells.

Melanotic cancer consists of any of the preceding forms, combined with black pigment.

Fungus hæmatodes is a term applied to an unusually vascular form of cancer, or to any of the other varieties when they are ulcerated and liable to bleed.¹

¹ Prof. J. Leidy, in Gluge's Atlas of Histological Pathology, Amer. Ed. p. 69.

The cancer-cells may be demonstrated in the thick, opaque juice which exudes from the freshly-cut surface of a cancerous tumor, while the relation of the constituent elements of the mass are best shown upon a thin section, made with a Valentin's knife. The fluid portions containing the cells will be found, in the areolæ or interspaces formed by the crossing of the fibres. Both the fibres and cells vary much in appearance in different specimens. Sometimes the cells are round, sometimes elongated into fibres, and occasionally they present very irregular forms. The nuclei also differ considerably in numbers and size.

Dr. Walshe divides cancerous tumors into three varieties, according as the viscous juice, fibrous, or cellular elements predominate.

Very great difficulty is often experienced in arriving at an accurate opinion as to the cancerous or non-cancerous nature of a tumor; because there is no single element which can be regarded as pathognomonic of true cancer. "Neither the character of the cells," says Dr. Beale, "nor the nature of the matrix, nor the arrangement of the elementary constituents, can separately determine the point, and it is only by carefully noting the collective appearances observed upon a microscopic examination that we shall be enabled to decide."

Enchondroma, epithelioma, certain fibrous tumors containing spindle-shaped cells, &c., resemble true cancer so strongly that they have been called canceroid growths. For the peculiarities of each of these varieties of tumor, the student is referred to the works of Lebert, Bennett, Walshe, and other writers upon this subject. In the following table, taken from Dr. Lionel Beale's work on the medical applications of the microscope, will be found enumerated the most important characters which distinguish the true cancer-cells from those of canceroid tumors.

Cancerous.

Cells not connected with the matrix in a regular manner, or forming laminae.

Cells differing much from each other in size and form.

Cells readily separable from each other.

Cells not connected together at their margins, their edges seldom forming straight lines.

Cells containing several smaller cells in their interior often met with.

Nuclei varying much in size and number in different cells.

Juice scraped from the cut surface containing many cells floating freely in the fluid, and not connected with each other.

Canceroid.

Cells connected with the matrix, often forming distinct laminae.

Cells resembling each other in size and general outline.

Cells often cohering by their edges, which generally form straight lines, three or four cells being frequently found united together.

Cells usually containing one nucleus.

Nuclei not varying much in size in different cells.


Juice scraped from the cut surface, containing small collections of cells, which are often connected with each other.

Dr. Donaldson¹ asserts positively that true cancer can be dis-

¹ American Journal of Medical Sciences, January, 1853.

tinguished from all other tissue, normal or pathological, by certain clear and well-defined elements. If we "compare the physical characters of cancer with those of the simple tissues, such as the muscular, areolar, cartilaginous, osseous, &c., or with those of the compound, as the glandular, the synovial, the mucous, &c., the difference will be very apparent. Its greater or less firmness, its homogeneous fibrous aspect, with its lactescent infiltrated juice, are very characteristic. The presence of this peculiar fluid is, of itself, a point of differential diagnosis of great value; the microscope always detecting in it, when found, the presence of cancer-cells, &c. No matter what organ is the seat of the disease, this fluid can generally be scraped from the cut surface, or squeezed out by gentle pressure. It is particularly abundant in encephaloid, and frequently oozes out in drops, having a white cloudy appearance, of the consistence of cream, and very much of its color, being slightly tinged with yellow. It may sometimes, on superficial inspection, be confounded with light-colored pus, which has, however, with the yellow, a slightly greenish tinge. If, from the conditions of its formation, there can be any doubt, an appeal to the microscope will at once settle it by giving us the characteristic pus-globule" (Figs. 382, 383).

FIG. 382.



Pus-Corpuscles.

"The element of cancer consists of three parts, *cell*, *nucleus*, and *nucleolus*, all of which are peculiar to it.

"In all the varieties of cancerous tissue, nuclei are to be found either enveloped by a cell, or floating free, generally more or less of both; in some specimens there exist a large number of free nuclei with only an occasional cell. The form and appearance of these nuclei are the most constant and unvarying of all cancer elements. They are (Fig. 384, *a*) ovoid, or more or less round; the latter are found more particularly when the eye or the lymphatic glands are the organs diseased. Sometimes (as in *b*) we find little pieces of the wall of the nuclei apparently nicked out, but evidently it is purely accidental, and the proper shape can easily be recognized. They have, ordinarily, in width, a diameter of from 1-100th of a millimetre, or (a millimetre being equal to .039th of an inch) of .0039th of an inch to 1-66th of a millimetre; in one instance we met with one as wide as 1-38th of a millimetre; in length they measure from 1-133d to 1-100th of a millimetre. Their contour is dark and well defined, with the interior containing very minute dark granulations; indeed, when the specimen is perfectly fresh, they have a homogeneous aspect, the granulations being so small as to give the appearance of a mere shading (Fig. 384, *c*); if the specimen is kept a day or two we find


the interior filling up with large granulations (as in *d*). Within these nuclei, when they have not been obscured by granular or fatty degeneration, there is found, habitually, a small body, or *nucleolus*, averaging in diameter about 1-500th of a millimetre.

FIG. 383.



Pus-corpuscles after acetic acid.

FIG. 384.



Free cancer nuclei.

These nucleoli have somewhat of a yellowish tinge, with a brilliant centre and dark borders, refracting light like the fat-vesicles. We would call attention, particularly, to the peculiar brilliancy of the centres of these nucleoli, which, we think, is characteristic; it can almost invariably be noticed, if the focus is varied. Their large size, in proportion to the nuclei, should also be noticed, together with the great variableness of their position, sometimes being near the centre, and again in close contact with the walls (see Fig. 384, *e*). Ordinarily, in other elements, they are found almost constantly in the centre. Very frequently two or three nucleoli are found within the same nucleus. M. Robin¹ mentions the action of acetic acid upon cancer-nuclei and their nucleoli as differing from that on other elements, particularly epithelial; it renders the nucleus gradually paler, together with the cell, destroying neither, but the nucleolus is perfectly untouched by it; whereas, in epithelial cells, where generally in those of the skin the nucleoli are wanting, the action of acetic acid destroys the cell, leaving the nucleus unaltered."


The polygonal shape may be regarded as the typical form of the cancer-cell. "In hard firm tumors, particularly those of the breast and ovaries, the cells found are exceedingly irregular, sometimes nearly triangular. (Fig. 385, *f*.) The ovoid or spherical are more frequently met with in soft or medullary cancer (Fig. 386, *g*), where there is but little pressure, although its juice appears often to be but one mass of cells. It is rare, however, that perfectly round cells are met with, but very generally the angles are well rounded in those which appear to be derived directly from

¹ MS. notes of his Cours de Histologie, 1850.

the *polygonal* form, the diameter of which is very variable, ordinarily from 1-75th to 1-25th of a millimetre. One peculiarity of this, as of the other forms of cancer-cell, is the presence of the granulations of different sizes in their interior; whereas, in epi-

FIG. 385.

FIG. 386.




thelial cells, the interior is generally, when fresh, of course, homogeneous. In cancer we find the three varieties of granulations given by M. Robin.¹ First, the very fine black dots, found in all organic elements, and named by the French, very appropriately, *poussière organique*. Secondly, the gray granulations, a form somewhat larger; and, lastly, the fat granulations, distinguished by the refraction of the light. This first variety of cells contains nuclei, having in their interior invariably one or more nucleoli, both of which retain the characteristic points described above. The large size of the nucleus, in proportion to the diameter of its cell, will at once strike the eye of the careful observer. The variable position, also, of the nucleus within the enclosure, appears to us to be peculiar to cancer; in cells of other structures, the rule is to find the nucleus very nearly in the centre, except with fibro-plastic cells, where the nuclei appear to have a peculiar affinity for the walls. All the varieties of cancer-cells contain very frequently two or more nuclei; whereas, the epithelial, more particularly those found on the surface of the body (where there is most danger of confusion and doubt), but rarely have more than one. Moreover, the cell of epithelium is much larger than that of cancer, yet the cancer-nucleus is twice as large

¹ Tableaux d'Anatomie, &c. par Ch. Robin, Paris, 1851.

as that of epithelium, as is also the nucleolus, compared with that found in epithelium."

The caudated cells are of common occurrence in cancerous

FIG 387.



tumors; in cancer of the bladder they are invariably present (Fig. 387.) The fusiform cells are most frequently met with in cancer of the bones. (Fig. 388.) The concentric cancer-cell is

FIG. 388.



best seen, according to Robin, in cancer of the uterus, breast, and ovaries. (Fig. 389.) Examples of the compound or mother-cell

of cancer and the agglomerated nuclei are shown in Figs. 390, 391 (after Donaldson).

Fig. 389.




Fig. 390.





Fig. 391.



EXAMINATION OF ANIMAL FLUIDS.

Blood.—To examine blood microscopically it is only necessary to press a small drop between two pieces of glass, until it is flattened out into a thin layer. A number of yellow, round, bi-concave disks will then be seen, varying in diameter from the 1-5000th to the 1-3000th of an inch, the average size being about 1-4000th of an inch. These disks have a bright margin, and a dark centre; or a bright centre and a dark margin, according to

Fig. 392.




Fig. 393.



the focus in which they are placed. (Fig. 392.) Exposure to the atmosphere causes the edges of the corpuscles to lose their smooth outline, and become irregularly notched or serrated, and sometimes beaded. (Fig. 393 B.)

The corpuscles vary in size in different animals. Thus in birds, fishes, and reptiles they are elliptical and flattened, and exhibit a distinct nucleus, which is generally oval. In the camel tribe they are elliptical and bi-convex.

The red globules are diminished to half their size by prolonged maceration in serum. Water renders them spherical and deprives them of their color. In strong syrup they shrink very much, from the rapid exosmosis which takes place.

Acetic acid renders the membrane of the corpuscle so transparent, as to be almost invisible, while nitric acid causes the outline to become darker and thicker. Acid urine and the acid of the gastric juice produces a similar effect, as is seen in cases of hemorrhagic effusion into the stomach and bladder.

The colorless or lymph-corpuscles of the blood are spherical, highly refractive, and vary in diameter from the 1-2500th to the 1-2000th of an inch. They have a granular appearance, which is lost by immersion in water, and are specifically lighter than the colored corpuscles. Within the cell-wall are numerous granules and molecules of different sizes, and one or more nuclei. (Fig. 394.) These nuclei are rendered distinct by acetic acid, while

FIG. 394.



Colorless corpuscles.

FIG. 395.



Blood in Leucocythemia.

the granules are dissolved. It has been estimated that the colorless corpuscles constitute about 1-50th part of the corpuscular element of healthy blood. In enlargement of the spleen and lymphatic glands they increase in numbers, producing the condition called by Bennett, Leucocythemia or "white-cell blood." (Fig. 395.) In some dropsical and cancerous affections, a slight increase of these white globules has been observed. In certain extreme cases they equal the red corpuscles in number. Their nuclei have occasionally been seen quite naked by Virchow and Bennett. Dr. Beale speaks of finding in the blood of some cholera cases, very large white cells, containing oil-globules collected together in one part, while the remainder of the cell was quite transparent.

Where the blood is thickened from an excess of fibrine the colored corpuscles become caudate or flask-like in shape, and aggregate themselves in irregular masses, instead of in the form of rouleaux. (Fig. 396.)

According to Funke and Kölliker yellowish crystals are sometimes seen in the blood-corpuscles of the spleen of the dog, perch, and other animals. Oil-granules have also been observed in the blood, giving it a lactescent or creamy appearance. The

microscopic changes which the blood undergoes in plethora, fever, and various other diseases, have yet to be accurately determined.

Milk.—A microscopic examination of milk reveals a number of spherical bodies, having dark, smooth, and well-defined margins, and a transparent and highly refractive centre, and varying from a mere point up to the 1-4000th or 1-3000th of an inch in diameter. These bodies consist of oil-globules invested with a covering of albumen, which prevents them from running together and forming larger globules when pressed between two pieces of glass. If this albuminous membrane be dissolved with a little acetic acid

or carbonate of soda, the oil is separated, and may be readily collected. Under this treatment the smaller globules may be made, by the slightest pressure, to run together and form larger ones. These globules collecting on the surface of milk, in virtue of their lighter specific gravity, constitute cream. An excess of ether effects the solution of the milk-globules, while water causes them to swell out.

The colostrum or first milk of the human female is yellow in color, and contains many large cells filled with oil, and mingled with a number of compound granular bodies, which disappear about the fifth or sixth day after delivery. (Fig. 397.)

In fresh and healthy milk, the globules are more or less uniform in size, and move freely in the surrounding fluid, showing no tendency to aggregate in masses. (Fig. 398.)

The microscope readily detects adulterations of milk.

The addition of water causes the globules to be separated farther from each other. The presence of flour is determined by large starch-corpuses, which strike a blue color with iodine. Gritty particles, soluble in the mineral acids, indicate the addition

FIG. 396.



FIG. 397.




FIG. 398.



of chalky matters. In milk to which sheep's brains have been added, fine nerve-tubes will be seen in the field of the microscope mingled with oil-globules. A tendency on the part of the globules to collect in masses is an indication of acidity. Pus and blood-corpuscles are easily distinguished by their peculiar characteristics.


The richness of milk is dependent upon the number of globules. In determining the quality of the milk of different animals the greatest care in manipulation is necessary.

Saliva.—A drop of saliva placed upon a glass slide and covered with a thin piece of glass, presents for examination epithelial scales from the buccal mucous membrane, salivary corpuscles, and numerous molecules and granules.

The epithelial scales are flat plates, varying in length from the 1-800th to the 1-500th of an inch. They are generally oblong in shape, sometimes square, and occasionally very irregular. The edges are curled up and often adherent to those of other scales. With a magnifying power of 250 diameters linear, a round or oval nucleus may be seen in the substance of these scales, surrounded with a great number of molecules and granules. The addition of acetic acid renders the nucleus more distinct, at the same time increasing the transparency of the scale. Water produces little or no effect.

“The salivary corpuscles are colorless, spherical bodies, with smooth margins, varying in size from the 1-3000th to the 1-1800th of an inch in diameter. They contain a round nucleus, varying in size, but generally occupying a third of the cell; and between this nucleus and the cell-wall are numerous molecules and granules, which communicate to the entire corpuscle a finely molecular aspect. The addition of water causes these bodies to swell out and enlarge from endosmosis; while acetic acid somewhat dissolves the cell-wall, and it becomes more transparent; while the nucleus appears more distinct as a single, double, or tripartite body. Both water and acetic acid also, produce coagulation of the albuminous matter contained in the fluid of the saliva, which assumes the form of molecular fibres, in which the corpuscles and epithelial scales become entangled, and present to the naked eye a white film.”¹ (Fig. 399.)

FIG. 399.



Salivary corpuscles, epithelial scales and granules.

The salivary corpuscles are accompanied with a quantity of molecular and granular matter, which undergoes an increase in ulceration of the mucous

¹ Dr. Bennett's Introduction to Clinical Medicine.

membrane of the mouth. The debris of various articles taken as food are also often found in the saliva, rendering its study more difficult to the unpractised eye.

Sputum.—To examine sputum it should first be placed in water, in order that any dense cretaceous or tubercular portions may be deposited at the bottom of the vessel, and thus separated from the lighter portions, which, on account of the confined air, will generally float upon the surface. The different constituent elements may then be isolated by breaking up small detached masses with a glass rod, and spreading them out upon a glass slide. Parts presenting any peculiar appearances, as dark spots, fibrous tissue, &c., should be removed by means of broad-pointed forceps and scissors, and examined separately and with much care.

The various matters which enter into the composition of sputum, and which complicate its study, are thus enumerated by Prof. Bennett.

“1st. All the tissues which enter into the composition of the lung, such as filamentous tissue, young and old epithelial cells, blood-corpuscles, &c. 2d. Mucus from the œsophagus, fauces, or mouth. 3d. Morbid growths, such as pus, pyoid and granular cells, tubercle-corpuscles, granules and amorphous molecular matter, pigmentary deposits of various forms, and parasitic vegetations, which are occasionally found in the lining membrane of tubercular cavities. 4th. All the elements that enter into the composition of the food, whether animal or vegetable, which hang about the mouth or teeth, and which are often mingled with the sputum, such as pieces of bone or cartilage, muscular fasciculi, portions of esculent vegetables, as turnips, carrots, cabbages, &c. ; or of grain, as barley, tapioca, sago, &c. ; or of bread and cakes ; or of fruit, as grapes, apples, oranges, &c.” It has lately been asserted by Shroeder Van der Kolk, that fragments of pulmonary tissue may be detected in the sputum before the existence of ulceration can be revealed by physical exploration. This, however, is doubted by Prof. Bennett.

A very common appearance in sputum under the microscope is represented by small masses of molecular and granular matter.

In the sputum of phthisis small lumps of softened tubercle may often be found, mingled with purulent mucus, at the bottom of the vessel. They have a yellowish, cheesy appearance, and consist of small and somewhat transparent cells, round, oval, or triangular, in shape, and varying from the 1-120th to the 1-75th of a millimetre in their longest diameter. They are known as tubercle-corpuscles, contain a number of granules, and are surrounded by free granular matter and oil-globules. (Fig. 400.)

In the sputum which accompanies the cretaceous or calcareous transformation of tubercle in the lungs, small, hard or gritty, and white masses will be found, consisting of amorphous aggregations of phosphate and carbonate of lime mixed with some

animal matter. (Fig. 401.) Pus and blood-corpuscles are often observed in the sputum, and occasionally crystals of cholesterine.

FIG. 400.



FIG. 401.



The sputum of acute pneumonia often contains minute fibrinous casts of the bronchial tubes, together with large cells filled with oil-globules, and numerous finely granular cells, similar to pus-globules. (Fig. 402.)

FIG. 402.



The thickened sputum expectorated in the morning on first rising, consists of epithelial cells pressed more or less closely together, and varying somewhat both in shape and size. The dark color is due to the numerous carbonaceous granules contained in the cells.

Mucus.—The gelatinous material known as mucus, presents different appearances, according to the peculiar cell-structures and pigmentary matters which it contains. It possesses no essential morphological element, the so-called mucus-corpuscles being in all probability merely pus-cells, or modifications of epithelium. According to Prof. Bennett, irritation of a mucous surface causes the exudation which is poured out to be transformed into pus-corpuscles by mixing with the gelatinous secretion. The thick white gelatinous mucus secreted by the lining-membrane of the os uteri contains numerous epithelial cells, while gonorrhœal or catarrhal mucus abounds in pus-cells. The viscid albuminous substance in which these cells are contained manifests a marked tendency to coagulate in the form of minute fibres, and constitutes the most characteristic element of mucus. An increase of the cell-elements, and a diminution of viscosity, are indications of disease; while an increase of the albuminous matter, and a decrease in the number of cells, are marks of a more healthy mucus.

Pus.—Normal pus, placed between two glasses and examined with a magnifying power of two hundred diameters, exhibits numerous granular corpuscles floating in a clear fluid, called *liquor puris*. These corpuscles are larger than blood-globules, have a smooth margin and a finely tuberculated surface, and vary in diameter from the 1-1300th to the 1-1000th of an inch. Many of them contain a round or oval nucleus, which is rendered more distinct on the addition of water, and is liberated in the

form of granules, having a central dark spot, by the addition of strong acetic acid, which dissolves the cell-wall. (Fig. 403.) In

FIG. 403.




FIG. 404.



scrofulous pus and in various unhealthy discharges, these corpuscles lose their globular form, and are found surrounded with numerous molecules and granules. (Fig. 404.) In gangrenous and ichorous pus they are mixed with broken-down blood-globules, remains of tissue, &c.

Dr. Beale makes the following judicious remarks with regard to these pus-corpuscles. "The cells above referred to have been considered as characteristic of pus, and much trouble was taken, in the earlier days of microscopical research, to assign definite characters to them, by which they might be distinguished from the so-called mucus-corpuscle, and other cells which they much resemble. Such a distinction, however, cannot be made, for, in the first place, cells may be obtained which present various stages, apparently intermediate between an ordinary epithelial cell, and a pus-globule; secondly, cells agreeing in their microscopical characters with the pus-globule, are not unfrequently formed on the surface of a mucous membrane, without its function being seriously impaired, and certainly without the occurrence of those preliminary changes which usually precede the formation of pus; and, thirdly, cells are found in the lymph, in the blood, in the lymphatic glands, in the serous fluid in the interior of cysts, and in many other situations, which in their size, form, and general appearance so much resemble the globules found in true pus, that it is quite impossible to assign characters by which they may be distinguished. The figures of these cells, as they appear before and after treatment by acetic acid, often could not be distinguished from the figures of pus-cells, treated in a similar manner, given by the same authors.

"Cases occur in which it appears almost useless to attempt to decide as to the presence or absence of pus, if only a few globules are to be found (nor do I think that if such were possible, it would be of any advantage), because no characters by which the globules can be distinguished individually have been laid down.

"At the same time it must not be supposed that the diagnosis of pus is a matter of secondary importance; and all that is intended in introducing these observations is to impress upon the student the importance of not stating that pus has been found in

any particular locality, or in any particular fluid, merely because a few cells having all the characters of a pus-globule have been observed. To say that 'pus had been found in the blood,' or that 'the casts of the uriniferous tubes contained pus,' would lead to a very different inference from that derived from the statement that 'cells having all the characters of pus-globules had been found in the blood,' or that the 'casts of the tubes contained cells resembling those of pus.' The former will be true in extremely few cases; the latter in a vast number that fall under the observation of every practitioner. If, however, we find a considerable number of globules under the field of the microscope, of nearly uniform size, agreeing in general characters with the pus-globule, and upon the addition of acetic acid exhibiting the characteristic reaction, we shall seldom be wrong in inferring that they are really pus-cells."

Fæces.—In the fæces may be found, says Prof. Bennett, "1st. All the parts which compose the structure of the walls of the alimentary canal; 2d. All kinds of morbid products; and 3d. All the elements which enter into the composition of food."

In severe attacks of dysentery, epithelial scales, membranous flakes, shreds of fibrous matter, pus-globules, and blood-corpuses are all observed in the discharges, mingled sometimes with crystals of the triple phosphate, and occasionally with numerous torule and sarcine. In ulceration of the intestines, pus-globules may readily be detected upon the surface of the fecal masses. Perfectly formed pus and blood-globules originate low down in the rectum, near the anus; broken-down globules originate higher up in the bowel. The white flocculi composing the stools of cholera patients consist of epithelial cells imbedded in mucus. Sometimes the sheaths of the villi are also found, together with free nuclei.

In the stools of typhus and other fevers of a low type great quantities of crystallized phosphates and carbonates are found.

Dr. Farre, Prof. Bennett, and others, have observed collections of confervoid growths in the matters discharged from the bowels.

Urine.—The student should early accustom himself to the examination of the various constituents of healthy urine. He cannot be too familiar with the different appearances presented by different specimens of this important secretion. Sometimes the urine will appear utterly structureless or homogeneous, offering absolutely nothing for examination. At other times, even the well-educated eye, will with difficulty identify the various and dissimilar objects crowded together in the field of the microscope, and which have accidentally found their way into the urine, or been introduced by the patient for purposes of deception. Starch-granules, woody fibres, hair, fragments of wool, cotton, feathers, &c., have all in their turn, and very frequently, been mistaken for some of the ordinary matters deposited by the urine. We take from Dr. Beale's work on the

Microscope, the following table of the most common extraneous matters constantly met with in urinary deposits.

Fragments of human hair.	Fragments of tea-leaves, or separated spiral vessels and cellular tissue.
Cats' hair.	Fibres of coniferous or other wood swept off the floor.
Hair from blankets.	Particles of sand.
Portions of feathers.	Oily matter in distinct globules, arising from the use of an oiled catheter, or from the accidental presence of milk or butter.
Fibres of worsted of various colors.	
Fibres of cotton of various colors.	
Fibres of flax.	
Potato starch.	
Rice starch.	
Wheat starch, bread crumbs.	

Great care, therefore, should be exercised in the collection and preparation of urine for microscopic examination. It is a very good practical rule to examine a portion of the urine an hour or two after it is voided; and another portion after standing a day, or in some instances two days. Occasionally it will be necessary to institute the examination immediately upon the passage of the secretion, as in those instances where there is a strong and rapid tendency to decomposition upon exposure to the atmosphere, or where this change has already taken place in the bladder. Urine containing lithic acid or oxalate of lime should be allowed to stand for some time in order that it may be deposited.

Perfectly healthy urine, after standing for about twelve hours, exhibits a slight cloudy deposition, consisting of epithelial scales, some crystals of the triple phosphate, and granular fragments of the urate of ammonia.

Urinary deposits are best obtained for examination by pouring the fluid into a tall wine-glass, or wide-mouthed bottle, capable of containing several ounces, and allowing it to stand for a few hours. The clear supernatant liquid should then be poured off, and the thick turbid under-stratum of urine, containing the deposit, emptied into a small test-tube. The sediment will thus be obtained in a small bulk, and can readily be examined by being deposited upon a glass slide after the liquid portions have been removed by means of a pipette.

Large crystals of uric acid require an inch object-glass to be seen distinctly; when very small a fourth of an inch glass is necessary, as is the case also with the octohedral crystals of the oxalate of lime, epithelium, casts of the uriniferous tubes, &c. The eighth of an inch object-glass is best adapted for the examination of spermatozoa.


In examining the deposits small portions should be brought into the field of the microscope at a time, and in this careful manner the different constituents, and their relation to each other, may first be studied to considerable advantage with the lower magnifying powers. Afterwards the nature and structure of each object may be more minutely investigated by subjecting it to the higher glasses. Sometimes it will be found advisable

to examine the deposit in various fluids, such as water, mucilage, spirit, Canada balsam, turpentine, &c. Occasionally it is necessary to resort to certain chemical reagents, before the deposit can be examined satisfactorily. Thus, in some amorphous sediments containing lithic acid alone, or combined with alkaline bases, the familiar rhomboidal crystals cannot be detected until the mass be first dissolved with potash, and then treated with excess of acetic acid.

ORGANIC CONSTITUENTS OF URINARY DEPOSITS.

In healthy urine the mucus settles towards the bottom, as a flocculent, transparent, and somewhat bulky cloud. In this transparent substance the microscope detects

FIG. 405.



Mucous corpuscles and epithelium.

merely a few granular cells, larger than blood-corpuscles, of very delicate structure and surrounded by a few minute granules. In disease the mucous deposition often loses its transparency, becoming viscid and thick from the addition of various kinds of epithelium. The peculiar appearance of the epithelial cells will indicate the part of the genito-urinary mucous membrane from which the mucus was secreted. (Fig. 405.) In some diseases of the bladder, as cystitis, a thick, glairy, and gelatinous deposit will often appear, simulating inspissated mucus. This is pus chemically changed by contact with the carbonate of ammonia generated in the decomposition of urea by the alkaline mucus. Very minute octohedral crystals of oxalate of lime are sometimes found like dark square-shaped specks imbedded in the mucus. A power of two hundred is generally sufficient to distinguish them. Fragments of cotton-fibre, hair, &c., are sometimes found incrustated with these crystals.

FIG. 406.



A. Portion of a secreting canal from cortical portion. B. Epithelium gland cells, magnified 700 times. C. Portion of a canal from medullary substance of kidney.

The epithelium found in the urine differs in different specimens, according to the part of the urinary apparatus from which it is derived. In the convoluted portion of the tubuli uriniferi it is glandular, and forms a thick layer upon the basement membrane. (Fig. 406.)

forms a thick layer upon the basement membrane. (Fig. 406.)

In the straight portion of the tubes it is flatter, and more like the scaly variety. In the pelvis of the kidney, it is tessellated or pavement-like, consisting of thin, flat scales united at their edges. In the ureter it is columnar or cylindrical in shape, having a large and distinct nucleus. In the fundus of the bladder columnar epithelium is found mixed with large oval cells; flattened cells having a distinct nucleus and nucleolus abound in the trigone.

FIG. 407.




Mucus, pus, blood, and epithelium from leucorrhœa.

In the mucous follicles columnar epithelium is found; on the surface between them, the scaly variety. The columnar variety prevails in the posterior part of the urethra; anteriorly it becomes scaly. The vaginal epithelium found in the urine of females, is also of the scaly variety. (Fig. 407.)

Occasionally casts, consisting of moulds of the uriniferous tubes, are observed in the urine. They furnish valuable aids in arriving at a correct diagnosis as to the pathological changes which may be going on in the kidney. They consist mainly of oily granules, or epithelial cells abundantly supplied with these granules. (Fig. 408 A.) Prof. Bennett divides them into two distinct varieties, namely,—“ 1st. Fibrinous or exudation casts, which are most commonly found in the urine at critical periods of acute inflammations, especially in scarlatina, small-pox, pneumonia, &c. 2d. Casts, with oily granules, indicative of chronic disease, and especially of Bright’s disease. (Fig. 409’.) At the

FIG. 408.



same time, it should be understood that they may be more or less associated together, and that the rule is not invariable." Dr. Beale divides them into three classes according to their diameter; namely, 1st. Casts of medium diameter, about the

FIG. 409.



1-700th of an inch. These contain granular matter with epithelial debris, oil-globules, and occasionally blood and pus-corpuscles. In the urine of a cholera patient, Dr. Beale once detected dumb-bell and octohedral crystals of oxalate of lime in one of these casts. 2d. Casts of considerable diameter, about the

1-500th of an inch. These are transparent and have a smooth, glistening, or waxy appearance. Sometimes they are granular. 3d. Casts of small diameter about the

FIG. 410.



Tube containing an homogeneous cast.

1-1000th of an inch. According to Dr. Johnson these originate in cases in which there is no tendency on the part of the epithelium to desquamate, as in non-desquamative nephritis.

Spermatozoa are sometimes found in the urine when examined soon after it has been passed. They can be seen with a power of two hundred diameters, though in demonstrating them it is better to employ a power of four hundred diameters. (Fig.

FIG. 411.



411.) The presence of these bodies in the urine must not be regarded as a sign of spermatorrhœa, unless accompanied with the symp-

toms of that disease. In urine, which has been allowed to stand for some time, vibriones and certain forms of vegetable fungi or torulæ are gradually developed. With a power of two hundred diameters, vibriones, looking like minute lines, may be seen writhing about in the mucus. They are always to be found in decomposing urine, and are sometimes generated in the bladder.

The species of fungi vary in different specimens of urine, and appear after different intervals of time. Dr. Hassall, however, considers the different species to be merely the successive stages of development of the same fungus; the stage of development being dependent upon the degree of acidity of the urine, and the length of time in which it has been exposed to the air. In acid urine, containing nitrogenous matter, and exposed to the atmosphere, a peculiar fungus is generated known as the *penicilium glaucum*,—the same fungus which is developed in lactic acid fermentation.

Dr. Hassall has shown that in urine containing even very minute traces of sugar, a peculiar fungus is developed, which may be regarded as the characteristic test of the presence of sugar, since it is found in no other condition of urine. In diabetic urine torulæ are often very rapidly developed. (Fig. 412.)



FIG. 412.

The fat-cells found in the urine are usually epithelial cells loaded with oil. According to Dr. Beale, fatty matter may occur in the urine in three conditions. 1st. "As distinct and separate globules, resembling those which are produced by intimately mixing oil with water by the aid of mucilage, &c. (Fig. 413.) When fatty matter occurs in this state only in urine, its presence is 'usually accidental. 2d. In the form of globules enclosed within a cell-wall, or in casts. 3d. In the so-called 'chylous urine,' the fatty matter is suspended in an exceedingly minute state of division."




FIG. 413.




FIG. 414.

Blood-globules generally form a brownish-red granular sediment at the bottom of the vessel. If the urine has been standing long, the globules will appear very much changed in shape. Acid urine containing blood has a smoky hue, the globules

appearing of a dark brown color. Where the liquid has a neutral, or slightly alkaline reaction, the globules are red.

Pus-globules are also occasionally found in the urine, more or less changed in shape, according to the length of time they have remained in the liquid. After long soaking they completely disintegrate. Deposits of pus are often accompanied with crystals of the triple phosphate. This is especially the case when the pus is derived from the bladder.

Large and small organic globules, exudation cells, spherical cells containing nuclei and granular matter, &c., are also found in the urine. For descriptions of these bodies, the student is referred to the work of Dr. Golding Bird, on Urinary Deposits.

INORGANIC CONSTITUENTS OF URINARY DEPOSITS.

Lithic or Uric acid is one of the most common urinary deposits. In color it varies from a light fawn to a deep orange-red, the usual hue of the crystals being yellow. Sometimes they are almost colorless. They assume a great variety of forms, the most common and most characteristic of which is the rhomboidal, Fig. 413,—the form usually generated when the lithates are decomposed by means of an acid. Fig. 414 represents the peculiar forms of uric acid found in the urine of patients laboring under acute and scarlatinous dropsy. Sometimes the crystals are square and lozenge-shaped.

Other varieties are seen to consist of adhering masses and

FIG. 415.



FIG. 416.



flat scales, with transverse and longitudinal markings. Some

times they are six-sided, resembling crystals of cystine, but distinguishable from the latter, by two of their sides being longer than the others. Occasionally they appear as truncated or rounded columns.


Cystine crystallizes in flat, hexagonal plates, with irregularly hexagonal markings on their surface. Sometimes radiating lines pass from an opaque centre to the margins. (Fig. 416.) A deposit of cystine may be readily distinguished from that of the pale urates, which it resembles in appearance, by not dissolving when heated.

Oxalate of Lime occurs in octohedral crystals, having one axis shorter than the other two. It is important to remember that these crystals differ in appearance, according to the position in which they are viewed, as shown in Figs. 417, 418. When large they may be seen with the



FIG. 417.

FIG. 418.



Various forms of oxalate of lime.

unassisted eye as minute glistening points imbedded in the sediment. Where crystals of the oxalate of lime are associated with and obscured by the pale lithates, the addition of a drop or two of solution of potash, by dissolving the latter, will render the former more apparent.

Occasionally oxalate of lime assumes the form of dumb-bells. Dr. Golding Bird considers these to consist of oxalurate of lime, on account of their polarizing influence upon light. (Fig. 419.) The dumb-bell crystals are probably compound, appearing to be formed of collections of minute acicular crystals. They are generally accompanied by the octohedral form, and, according to Dr. Beale, their appearance is sometimes preceded and succeeded by the presence of the circular, oval, and less regular forms of crystals. They are formed in the kidney,




FIG. 419.

Dumb-bell crystals.

having been met with in the uriniferous tubes after death, and in fibrinous casts of those tubes. Phosphate of lime and lithic acid also assume the dumb-bell form; but the solubility in potash, and the different refracting power of these latter crystals, distinguish them from those of the oxalate.

FIG. 420.



Lithates constitute the so-called "lateritious deposits" in urine. According to Heintz these buff-colored or deep red sediments consist mainly of lithate of soda mixed with small portions of the lithates of ammonia and lime, and a trace of the lithate of magnesia. Under the microscope they appear as minute granules in different states of aggregation. (Fig. 420.) The lithates of ammonia and soda sometimes occur in spherical masses, adhering to thin films of the phosphates. The ammoniacal lithate occasionally assumes a stellate form; Prof. Bennett has seen it arranged in such a manner as strongly to resemble in appearance an organic membrane.

The triple phosphate, or ammonio-phosphate, of magnesia, occurs in the form of triangular prisms, occasionally truncated,

FIG. 421.



FIG. 422.



and sometimes with terminal facets. (Fig. 421.) If the urine be very ammoniacal, they present a star-like or foliaceous appearance (Fig. 422). Carbonate of lime is sometimes associated with the earthy phosphates in human urine, but rarely in a crystalline form. It generally appears in small round masses, or as an amorphous powder.

Urinary deposits may be preserved either in the dry way, in Canada balsam, turpentine, oil, and similar fluids, or in aqueous solutions. Only large crystals of the oxalate of lime, lithic acid, and some of the phosphates and lithates, can be preserved in the dry way. Dr. Beale gives the following directions for the preservation of urinary deposits: "After the crystals have been allowed to collect at the bottom of a conical glass vessel, the clear supernatant fluid is to be poured off, and the crystals are to be washed with a little dilute alcohol, or with a very weak solution of acetic acid. When the process of washing has been repeated two or three times, a small quantity of the deposit is to be transferred by means of a pipette to a glass slide, and the greater part of the fluid soaked up with a small piece of blotting paper. The crystals are next to be spread a little over the glass with the aid of a fine needle, in order to separate the individual crystals from each other; and the slide is to be placed in a warm place, or in the sun, until quite dry; but care must be taken that the drying is not carried on too rapidly, and that too great a degree of heat is not employed. A narrow rim of paper, or cardboard, is next to be gummed on the slide so as to include the crystals in a sort of shallow cell; and lastly, the glass cover is to be put on and kept in its place either by anointing the edges with a little gum-water, or by pasting it down with narrow strips of paper, which may be variously arranged and ornamented according to taste.

"If the crystals of lithic acid are to be mounted in Canada balsam, they should be carefully dried first, as above directed, and afterwards over sulphuric acid, and then moistened with a small drop of spirits of turpentine. The slide is now to be slightly warmed, in order to volatilize the greater part of the turpentine, and a drop of Canada balsam is to be dropped upon the preparation from the end of a wire, which may be readily effected by holding the wire with the balsam over the lamp or hot brass plate for a minute or two in order to soften it. The slide is next to be held over a lamp, in order to keep the balsam fluid until any air-bubbles which may be present have collected into one spot on the surface of the liquid balsam, an operation which is expedited by gently moving the slide from side to side. The air-bubbles may now be removed by touching them with a fine-pointed wire. Lastly, the glass cover is to be taken up with a pair of forceps, slightly warmed over a lamp, and one edge is allowed to touch the balsam. The surface is permitted to fall gradually upon the balsam, so that it is wetted by it regularly, and only by very slow degrees, for otherwise air-bubbles would yet be included in the preparation. The glass slide with the preparation may now be set aside to cool."

There are many substances, however, which cannot be preserved to advantage in Canada balsam, or by the dry method. Such are epithelium, casts, torulæ, confervæ, fat-cells, pus, mucus, &c. Such substances should be placed in shallow glass-cells,

and covered with aqueous solutions, varying in character and strength to suit the specimen. The best preservative fluids are weak spirit, glycerine diluted with water, solutions of gelatine, creasote, naphtha, &c. The gelatine solution answers very well for the preservation of dumb-bell crystals of oxalate of lime, while the creasote and naphtha solutions are better adapted for the preservation of epithelium, tubular casts, &c. Crystals of the triple phosphate are best kept in aqueous solutions of ammonia; for cystine dilute acetic acid answers very well.

Vomited matters consist of articles of food variously altered by the digestive processes, epithelium and mucus from the mouth, fauces, pharynx, œsophagus, and stomach, gastric juice, bile, and the various matters generated in disease. As different portions of vomit contain different ingredients, small portions taken from points considerably separated, should successively be subjected to examination. The various transitions which alimentary substances undergo in the stomach, must often necessarily render the determination of the exact composition of the vomited matters a point of extreme difficulty.

Starch-granules are often met with in abundance, but sometimes so changed as to require the addition of tincture of iodine to detect them. Fig. 423 represents the appearance of starch

FIG. 423.



corpuscles after partial digestion in the stomach. The epithelium is also frequently found to be more or less altered from endosmosis, and partial digestion. Vibriones and various species of torulæ are also observed in vomit. The *sarcina ventriculi*, a peculiar fungus discovered by Mr. Goodsir, in matters ejected from the stomach, has also been observed in the feces, in the urine, and in an abscess of the lung. The fluid of waterbrash consists mainly of epithelial

scales and small oil-globules. In the rice-water vomit of cholera patients numerous flocculi of epithelial cells are found. The coffee-ground vomit appears to consist mainly of the coloring matter of the blood reduced to a finely granular state and mingled with disintegrated blood-corpuscles. The sediment deposited by the black-vomit of yellow fever upon standing, in all probability is chiefly composed of blood-globules in various stages of disintegration. The epithelial cells of this fluid "vary in respect to their abundance, size, and shape, and while stated by some to have presented themselves in all the specimens examined, they have, in some instances, been found wanting. Of the six specimens reported upon by Dr. Leidy, two were deficient in this particular. The size and shape of these cells, as observed by Dr. Riddell, have already been referred to. In the hands of Dr. Michel, the *scaly*, *columnar*, and *spheroidal*, have, at different times, been plainly made out with their nuclei

and nucleoli, but in very different proportions—the scaly or lamellar cells being always most numerous.”¹

Uterine and Vaginal Discharges present quite different characters in different specimens. The examination should be instituted as soon after they are collected as possible, and without the addition of water, as this may affect the natural appearance of the constituent elements. Epithelial cells and blood-globules in varying quantities compose the menstrual discharge. In leucorrhœa many of the epithelial cells are filled with oil-granules, and mingled to a greater or less extent with pus-corpuscles. Blood-globules are also observed very much altered in shape. (See Fig. 407.) In cancer of the uterus the microscopic examination of the discharges, becomes highly important in arriving at an accurate diagnosis. Cancer-cells, in such cases, may often be detected in the discharges.

FIG. 424.




When they are broken down or considerably altered in form, not a little difficulty will be experienced in assigning to them their true value. The student should be careful, also, not to confound the columnar epithelium of the ureter, with the spindle-shaped cancer-cells. Fig. 424 represents the microscopic appearances of some cancerous juice squeezed from the uterus; that to the left is the natural appearance, the other after the addition of acetic acid.

SEROUS AND DROPSICAL FLUIDS.

The sedimentary matters should be collected from serous fluids, and examined in the same way as urinary deposits.

An examination of the freshly effused fluid of ascites reveals only a few cells floating in a clear liquid. In chronic ascites, however, numerous granular and spherical cells, mostly non-nucleated and varying in size, are observed. The sediment consists of delicate fibres, interlaced, and having cells in the meshes or interstices, together with plates of cholesterine. Occasionally blood and pus-corpuscles may also be detected. The hydrocelic fluid consists of some delicate cells, and oil-globules, and occasionally some spermatozoa and plates of cholesterine.

FIG. 425.



¹ Dr. R. La Roche on the Nature and Composition of Black Vomit. Amer. Jour. of Med. Sciences, April, 1854.

Cells, oil-globules, free granular matter, and occasionally blood-corpuses and crystals of cholesterine are the principal constituents of the deposit obtained from ovarian fluid. Sometimes masses of gelatinous or colloid matter are mixed with these elements. Minute fibres are sometimes observed crossing each other in various directions, and contain in their meshes thus formed, a transparent jelly filled with round or oval corpuses. (See Fig. 424.) The cells are either small, transparent, granular, and non-nucleated, or large, opaque, and filled with oil-globules. Fig. 376 represents fatty granules, mixed with plates of cholesterine from an ovarian tumor, after Prof. Bennett.

INJECTIONS.

In studying the vascularity of tissues, injected specimens are of great utility. The student should give some attention, therefore, to the practice of injecting the different organized structures which he may desire to examine. Dr. Beale, in his admirable little work on the application of the microscope to clinical medicine—a work of which we have availed ourselves freely in the construction of this chapter—gives the following practical directions as to the time, mode, &c., of making injections.

“Generally, it may be remarked that we should not attempt to inject while the *rigor mortis* lasts. Many days may in some cases with advantage be allowed to elapse, particularly if the weather is cold, while in warm weather we are compelled to inject soon after death. As a general rule, the more delicate the tissue, and the thinner the vessels, the sooner should the injection be performed. Many of the lower animals, annelids, mollusca, &c., and fishes, should be injected soon after death. In making minute injections of the brain, only a short time should be allowed to elapse after the death of the animal, before the injection is commenced. Injections of the alimentary canal of the higher animals should be performed early—not more than a day or two after.

“Minute injections of the papillæ of skin, particularly of the fingers and toes, cannot be successfully made until the cuticle has become somewhat softened by allowing the preparation to remain in a damp cloth, or to soak in water, for some days. In these situations the vessels are strong, and in their ordinary state, the injection will not traverse them, in consequence of the cuticle preventing their gradual distension by the injecting fluid. A similar plan must be followed in making injections of the tongue, and other parts where the epithelial covering is unusually dense, and firmly adherent to the vascular surface beneath.

“If the subject be a small animal, it is better to take out part of the sternum, and fix the pipe in the aorta. If only part of an animal is to be injected, the largest artery supplying the part should be selected, and all the other open vessels may be tied or stopped with the small forceps.

“A small portion of intestine can be injected by cutting out the corresponding portion of mesentery attached to it; and after searching for a large vessel, all the others may be tied, together with the open ends of the alimentary tube.” “A pipe of somewhat smaller diameter than the vessel should be selected, and an opening may then be made in the vessel of sufficient size to admit the pipe, which can now be inserted. The needle, charged with thread or silk, is then carefully passed round the vessel, the thread seized with forceps, and the needle withdrawn over the thread. This operation is sufficiently simple where the vessel is large and strong; but where thin and easily torn, it requires great care. The thread is now tied tightly round the vessel close to the extremity of the pipe, and then attached to the two projecting wires, to prevent the possibility of slipping.

“In injecting from veins a similar method is pursued, taking care to choose a vein in which the valves are not numerous, or in which they are altogether absent. The portal vein can be reached by opening the abdominal cavity, care being taken not to tear any of the branches below the point where the pipe is inserted.

“Greater care is required to fix the pipe in the vessels of fish, in consequence of their being so readily torn. Excellent injections of fish may frequently be made as follows: The tail is cut off with a sharp knife at a short distance posterior to the anus, and if the cut surface be examined the ventral artery may be easily found situated immediately beneath the bodies of the vertebrae. A pipe is carefully introduced and pushed down some distance, so as to prevent the injection from coming out, or the end of the vessel may sometimes be separated from the surrounding parts and tied in the usual way. By this simple proceeding capital injections can often be made very easily.

“Minute injections of the branchiae of some of the mollusca may often be made by very carefully placing the pipe in the largest vessel that can be found, and slowly injecting. The extreme delicacy of the vessels prevents any attempt being made to tie them to the pipe, and, of course, much injection will be lost. From the large size of the vessels, however, much will run into the capillaries. In this way I have easily succeeded in injecting the branchiae of the *Pinna ingens*, and fresh-water mussel (*Anodon*), both of which form beautiful microscopical objects.

“In order to inject the smaller gasteropods (slugs, snails, &c.), we must pursue a different method. In the muscular foot of these are situated many large lacunae, or cavities, which communicate with the vascular system, or, in fact, form the vessels which are distributed to this organ. If the injection can be forced into any of these lacunae, it may be made to traverse the whole vascular system. To introduce the pipe a small hole is made obliquely in the foot, taking care not to force the instrument too far. A small pipe is next inserted, and when the pre-

paration is warm enough the injection of size and vermilion is very slowly and carefully forced in, and the progress which is made can be seen by observing the vessels distributed to the respiratory organs. When a sufficient quantity of injection has been introduced, the pipe may be withdrawn, and the hole plugged with a piece of wood cut to the proper size, to prevent the injection again escaping before the size has had time to set.

“The vascular system of insects may sometimes be partially injected by forcing the injection into the abdominal cavity, from which it finds entrance into the dorsal vessel, and from thence is distributed to various parts of the body. The injection in the cavity of the abdomen is then allowed to escape.

“It is very important that the size and vermilion, or other injection which is to be thrown into the vessels, should be thoroughly mixed and well strained before being used. The coloring matter, properly powdered, should be placed in a small earthenware mortar, and the melted size or other fluid carefully added by degrees, the whole being constantly stirred until well mixed. When the proper color has been obtained, the whole must be strained through muslin, or through a fine perforated strainer, into another vessel, which should be kept warm. The injection should be well stirred with a wooden stick previous to filling the syringe.

“We can judge of the intensity of the color by removing a drop of the solution with a stirring-rod, and allowing it to fall on a white plate so as to form a thin stratum, which should have a pretty deep color. It is always better to have too large a quantity of the coloring matter rather than too little.

“Of vermilion, about two ounces will be sufficient for a pint of size; but it is better in all cases to regulate the quantity by examining the intensity of the color in the manner just mentioned.

“When the preparation is warmed through, the injection properly strained, and the pipe fixed in the vessel, we may proceed carefully to inject, taking care that the injection is kept at a proper temperature, by allowing it to remain in the warm water-bath during the operation.

“The air should be first withdrawn from the upper part of the vessel by means of the syringe, after which the stop-cock is turned off and left attached to the pipe. The syringe is then disconnected, and after being washed out once or twice with warm water, is nearly filled with injection, which must be well stirred up immediately before it is taken. The syringe should not be quite filled, in order that the air in the pipe may be made to rise into the syringe through the injection, by the ascent of the piston, before any of the latter is forced into the vessel. The end of the syringe is then to be pressed firmly into the upper part of the stop-cock with a slightly screwing movement.

“The piston is now very gently forced down by the thumb,

until the syringe has been nearly emptied, when the stop-cock must be turned off, and the syringe refilled with warm injection as before.

“Care must always be taken to keep the syringe in the inclined position, so that any air which may be in it, may remain in the upper part; and for the same reason, all the injection should not be forced out, for fear of the enclosed air entering the vessels, in which case all chance of obtaining a successful injection would be destroyed.

“After a certain quantity of fluid has been injected, it will be necessary to use a greater amount of force, which, however, must be increased very gradually, and should only be sufficient to depress the piston very slowly. If too great force be employed, extravasation will be produced before the capillaries are half filled. Gentle and very gradually-increased pressure, kept up for a considerable time, will cause the minute vessels to become slowly distended without giving way to any great extent. At the same time it must be borne in mind that extravasation frequently occurs at various points in a successful injection; but the longer this event can be kept off, the more likely are we to succeed.

“When the injection begins to flow from the large veins mixed with the blood contained in these vessels, and the surface of the injected preparation looks of a red color, and has a somewhat velvety appearance, we may infer that the injection has been completed. This occurs at different periods. Sometimes the first or second syringeful causes a general redness of surface, while in other instances a considerable time will elapse before more than a slight blush appears. As a general rule, it is better to proceed slowly and cautiously, and to use as little force as possible, which should not be more than sufficient to produce an observable depression on the piston. Many minute injections will require an hour or more to complete.

“When the injecting is completed, and all the openings by which any of the injection could escape during cooling are closed, the preparation should be placed in cold water, and allowed to remain until the size has set, which will require twelve hours or more in hot weather.”

In the *Medical Examiner* of Philadelphia, for December, 1849, Dr. P. B. Goddard details the following method of making minute ethereal injections, originally employed by him.

“For the purpose of making such an injection, the anatomist must provide himself with a small and good syringe; some vermilion, *very finely* ground in oil; a glass-stoppered bottle, and some sulphuric ether. The prepared vermilion paint must be put into the ground-stoppered bottle, and about twenty or thirty times its bulk of sulphuric acid added; the stopper must then be put in its place, and the whole well shaken. This forms the material of the injection. Let the anatomist now procure the organ to be

injected (say a sheep's kidney, which is very difficult to inject in any other way, and forms an excellent criterion of success), and fix his pipe in the artery, leaving the vein open. Having given his material a good shake, let him pour it into a cup, and fill the syringe. Now inject with a *slow, gradual, and moderate* pressure. At first, the matter will return by the vein, colored, but in a few moments this will cease, and nothing will appear except the clear ether, which will distil freely from the patulous vein. This must be watched, and when it ceases, the injection is complete. The kidney is now to be placed in warm water of 120° Fahrenheit, for a quarter of an hour, to drive off the ether, when it may be sliced and dried, or preserved in alcohol, Goadby's solution, or any other antiseptic fluid. For glands, as the kidney, liver, &c., it is better to dry and mount the sections in Canada balsam, but for membranous preparations, stomach, intestine, &c., the plan of mounting in a cell filled with antiseptic solution is preferable."

MICROSCOPES OF AMERICAN MANUFACTURE.

Microscopes of great excellence are manufactured in this country. These, from their comparative cheapness, and the facility with which they can be procured, offer inducements to students and others to procure them at home, and thus save time to themselves, at the same time that they stimulate the manufacturers to make increased efforts to attain even greater excellence. A brief description of some of the best will enable the reader to form some comparative estimate of their value.


Mr. CHARLES A. SPENCER, of Canastota, New York, has manufactured a microscope of great excellence, the objectives of which will bear comparison with the best of foreign construction. His common angle of aperture for $\frac{1}{4}$ inch objectives is 135°; for $\frac{1}{8}$ inch, 170°, and for $\frac{1}{12}$ and $\frac{1}{16}$ inch, 176°. This is believed to be the largest angle ever given to an object-glass, and for sharpness of definition and power of penetration, they are unexcelled by any of foreign make.

"To Mr. Spencer is due the credit of having first resolved, with lenses of his own construction, the fine markings on the *Navicula Spencerii* and *Grammatophora Subtilissima*: these minute shells have since been adopted by microscopists as test-objects for the highest powers. The *Navicula Spencerii*, will exhibit one set of lines with Mr. Spencer's $\frac{1}{4}$ th-inch object-glass: both sets with the $\frac{1}{8}$ th-inch. The *Grammatophora Subtilissima* is a good test for a $\frac{1}{12}$ th or $\frac{1}{16}$ th.

Of several microscopes made by Mr. Spencer, two or three only will be here noticed. His first-class or best instrument is mounted on trunnions, and embraces all the acknowledged improvements, in form and stage, whereby the greatest steadiness and freedom from tremor are secured. The price of this instru-

ment, with all the accessories and full sets of object-glasses, will approach \$350. (Fig. 426.)

FIG. 426.



Spencer's Trunnion Microscope.

The second-class instruments, complete as to object-glasses and accessories, but mounted less expensively, cost from \$200 to \$250.

A very efficient microscope, is one known as the "Pritchard form:" this instrument has been somewhat modified by Mr. Spencer, and where a less expensive instrument than either of the others is desired, this one will be found a good working instrument, and available for all purposes of anatomical study. The cost of this form, with object-glasses as high as the $\frac{1}{4}$ th, with the usual accessories, is from \$125 to \$150.

Mr. Spencer also makes some simpler forms of instruments,

and yet very efficient working ones, with objectives as high as $\frac{1}{4}$ th, the price of which does not exceed \$75.¹

Mr. JAS. W. QUEEN, of Philadelphia, has prepared one of the most convenient and portable microscopes for the use of students that has been made. It is all contained in a mahogany box, six inches square and eleven inches high, with a small drawer within for objects, forceps, &c. The stand is of cast iron, with two uprights supporting the stage and body of the instrument; between the uprights is an axis upon which the whole upper part of the instrument turns, enabling it to take a horizontal or vertical position, or any intermediate angle most convenient for observation. The movable part, consisting of the stage, the body, and bar containing the adjustment, is fixed to the axis, on which it readily turns. There is an inner tube for increasing power by extending the distance between the eye and object-glass. Within the quadrangular bar, to the top of which is attached the body of the instrument, is the fine adjustment for foci; the milled head on the top of the bar operates upon the spring within, and carries the body and object-glass from, or to, the object being observed, with

the greatest facility. The stage upon which objects are placed to be viewed is made double, the upper portion being movable by a small lever which produces rectilinear or eccentric motion, thus giving great facilities to the microscopist for viewing any portion of the object he may desire; the slide containing the object is kept in place by two spring clips attached to the stage; the under portion of the stage carries a revolving plate to which the polarizing prism is attached when in use; a diaphragm with various apertures is beneath the stage; the mirror is made to slide up and down on the tube

FIG. 427.



J. W. Queen's Microscope.

that supports it, and can be turned to any angle for light that may be desired; a condensing lens for opaque objects, mounted with a jointed arm, is attached to the collar through which the body passes. Two eye-pieces and two object-glasses are furnished with the instrument, also the polarizing prisms. The lowest power is about 50 diameters, and that can be increased to 400 or 500. A drawing-prism can also be furnished when desired. (Fig. 427.)

Messrs. J. & W. GRUNOW, of New Haven, Connecticut, have in-

¹ See Hassall's Microscopic Anatomy, edited by H. Vanarsdale, M.D.

vented an admirable Student's Microscope, which commends itself to all who desire efficiency, cheapness, and portability. The accompanying cut explains its appearance and mode of action. It is mounted on a tripod base, with uprights of japanned cast iron. It has a quick and a slow movement, with draw-tube and a stage 3 by 4 inches, movable by a lever. This movement the Messrs. G. have nearly perfected, and at the suggestion of Dr. Goddard, of this city, an eminent microscopist, they have so arranged it that the stage follows the hand instead of taking the opposite direction, as is usually the case in other instruments having the lever stage. (Fig. 428.)

FIG. 428.



J. & W. Grunow's Student's Microscope.

Messrs. Grunow have also improved their stand in the mode of adaptation of accessory apparatus, especially of that which is attached below the stage; this is held in place by a bayonet

catch, and when once centred it may be detached and reattached with the greatest facility, without requiring a readjustment.

Fig. 429 represents a smaller and less expensive form of student's microscope, made by the Messrs. Grunow, and possessing the same objectives as that represented in Fig. 428, but with a non-

FIG 429.



movable stage. Messrs. Grunow also manufacture a first class microscope, with brass mountings and all the accessories. (Fig. 430.) Their instruments have given great satisfaction wherever they have been used.

THE INVERTED MICROSCOPE OF DR. J. LAWRENCE SMITH.

This instrument was invented and brought to the notice of the *Société de Biologie* of Paris, in 1850, by Dr. Smith, for whom

the first one was constructed by Nacet, of Paris. Its design is to enable the chemist to carry on observations under the microscope without the risk of injuring his objectives, or at least having his view obscured, by the fumes arising from liquids experimented upon. This has always been a serious difficulty in

FIG. 430.



micro-chemical research, and, together with the difficulty of manipulating in the limited space between the object-glass and the stage, has hitherto prevented this branch of scientific inquiry from being fully illustrated. "The only way," says Dr. Smith, "by which these difficulties can be surmounted, is by putting the object-glass beneath the stage and the object above it, with an optical arrangement of such a nature as to permit observation." It was with this view that M. Chevalier made a chemical

support, to go with his general instrument, and so constructed it, that it could be inverted so as to have the stage above, and the objective below the object; but every one who has used it knows how awkward it is for manipulation, although exceedingly ingenious. Dr. Smith, impressed with these difficulties, was led to the construction of the instrument, which, in *this* country, bears his name.

It was important for the arrangement in question, so to have the relative position of the stage and eye-piece, that the eye, while on a level with the latter, could readily see the former, and guide the required manipulation.

The most important part of the instrument is a four-sided prism, Fig. 430, with the angles ab , bc , cd , and da , respectively 55° , $107\frac{1}{2}^\circ$, $52\frac{1}{2}^\circ$, 145° , the angles being of such dimensions that a ray of light passing into the prism in the direction Ea , and perpendicular to the upper surface of the prism, after undergoing total reflection from the inner surfaces (on both of which the light strikes at an angle much less than forty-five degrees), will pass out perpendicular to the surface connected with the body of the instrument.

It will be readily seen how a ray of light, entering the object-glass, descends into the prism, and passes out of it upwards, through the eye-glass, the tube of which is inclined to the perpendicular 35° . The other parts of the instrument will be understood by looking at the figure. (Fig. 431.) The illumination of the object is effected by a prism, instead of a mirror.

In examining an object with this microscope, the object is arranged in the ordinary way; when liquid, it is placed in a watch-glass, or such glass cells as are convenient to use. In employing reagents, they can be added and watched immediately, for it is readily seen how the eye guides the manipulation on the stage and looks into the instrument almost at one and the same time.

In observing with high powers, as the object-glass is beneath the glass supporting the object, and as the glass is usually of a certain thickness, the method of observation must be changed. For all powers resorted to in chemical examination this difficulty never occurs, and in using high powers it is easily obviated. When the object is already mounted and dry, the thin glass can be readily turned downwards; but where it is moist, as for in-

FIG. 431.



stance, in examining fresh *Desmidiæ* and *Diatomaceæ*, the following plan is resorted to, namely: to use a cell, made of a thin piece of brass or glass, perforated with a hole, about a half an inch in diameter; it is best to give the hole a considerable bevel in one direction, as it facilitates the cleaning of it. Over the small end

FIG. 432.



of the hole a piece of thin glass is stuck, with balsam or other cement. When used, the object to be examined is placed within, and a cover of thin glass placed above. When brass is used to make the cell, it may be as thin as the twentieth of an inch. Dr. Smith remarks that for all observation with high powers, the Inverted Microscope is decidedly superior to the ordinary forms of mounting, for in the latter case, when an object-glass of a $\frac{1}{4}$ th or $\frac{1}{8}$ th inch focus is used, the focus is too short to admit of the use of cells; whereas, in the inverted form, as the object is looked at from beneath, the cell may be as thick as one pleases. Another thing connected with this class of observations, is that the *Diatomaceæ* and *Desmidiæ* can be observed to much greater advantage from beneath than from above, for reasons that will be obvious to persons accustomed to observe these classes of objects.

Another advantage possessed by this instrument calculated to extend its use for general purposes, is its great capacity for every variety of illumination, without sacrificing the ease and freedom

from fatigue belonging to the use of this form of microscope; for when placed on a table, rather higher than the one commonly used, and a foot or two from the edge, the observer can recline on his arms, and observe for hours without the slightest sensation of fatigue.¹ Messrs. J. & W. Grunow, New Haven, Conn., are the manufacturers of this instrument.


DR. J. LAWRENCE SMITH'S GONIOMETER AND MICROMETER.

Professor Smith has also invented a Goniometer for measuring the angles of crystals under the microscope. It is also combined with a Micrometer. The following is a description of the instrument with the method of using. (Figs. 433-4.)

E is the upper end of the draw-tube of the microscope, with the ring *k* soldered to it. Over this ring *k*, screws another ring *F*, which serves as a support and as a centre to the graduated circle *D*, which freely, but without shaking, revolves upon the same. Into the bore of the ring *F*, fits by its lower conical end *h*, the tube *G*, which is held in it by a screw-ring *y*, that prevents its being taken out. Into the tube *G*, which also has a free revolving movement, fits the positive eye-piece *a*, *d* being the field-lens, *s* the eye-lens.

The slide *b b*, on opposite sides of *G*, admits of the micrometer with its mounting *B*, *B* being introduced into *G*, and the graduation being brought into the field of the eye-piece.

FIG. 433.




C is an index, attached to *G* by the screw *c*; it may be taken off, when the apparatus is not used as a goniometer.

¹ See American Journal of Science and Arts, for September, 1852, in which the entire paper of Prof. Smith will be found.

USE OF THE GONIOMETER.

Bring the object into focus near the centre of the field of the micrometer, apply your finger to the knob *K*, and revolve the micrometer, till the lines of its graduation are parallel to one

FIG. 434.



side of the angle to be measured. Revolve then separately the graduated circle, till zero is brought to agree with the point of the index *C*. Then revolve again the micrometer by the knob *K*, until the graduation lines are parallel to the other side of the angle to be measured, when the index *C* will show the value of this angle. (The references are the same in both cuts.)

The graduated lines of the micrometer are generally $\frac{1}{250}$ of the American inch apart. But their relative value as micrometer, with the different object-glasses and eye-pieces, must be ascertained by a glass stage-micrometer, and recorded in a table.



INDEX.

- Aberration, chromatic, 73, 74.**
 spherical, 71, 72.
Acalephs, see Medusa.
Acarida, 581, 582.
Achyla prolifera, 315, 316.
Acineta-form of Vorticella, 419, 420.
Achnanthes, 297.
Achromatic Condenser, 131-133, 136; use
 of, 172, 173.
Achromatic correction, 41; principle of, 73,
 74; application of, 74-76.
Actiniform Zoophytes, 474-476.
Actinocyclus, 289, 290.
Actinophrys, 407-409.
Adipose tissue, 607, 608, 664.
Adjustment of Focus, 86, 87; uses of, 162-
 165.
 of Object-glass, for thickness
 of cover, 76, 77; mode of
 making, 165-168.
Agates, sponge-remains in, 633.
Air-bubbles, microscopic appearances of,
 184.
Alcyonian Zoophytes, 472-474.
Alcyonidium, 496, 497.
ALGÆ, higher, microscopic structure of,
 328-333 (see Protophyta).
Amaroucium, structure of, 501, 502.
Amici's Prism for oblique illumination, 137.
Amoeba, 405-407.
Anacharis alsinastrum, rotation in, 364,
 365.
Anagallis, petal of, spiral vessels in, 396.
Angular aperture of object-glasses, 71-73,
 76; real value of, 188-192.
Animalcule-Cage, 148, 168.
ANIMALCULES, early researches on, 38;
 Ehrenberg's observations on, 47-49, 411-
 424; Dujardin's researches on, 49; see
 Infusoria, Rhizopoda, and Rotifera.
Animals, Distinction of, from Plants, 247,
 248, 409, 410.
ANNELIDA, 535; marine, circulation in, 535,
 536; metamorphoses of, 536, 537; lumi-
 nosity of, 537; fresh-water, 537, 538.
Annular Condenser, Shadbolt's, 139, 140.
ANNULOSA, 529; see Entozoa, Turbellaria,
 and Annelida.
Anomia, structure of shell of, 512.
Antennæ of Insects, 564, 565.
Anthers, structure of, 396, 397.
Antheridia of Cryptogamia, discovery of,
 45, 46; see Antherozoids.
Antherozoids, of Vaucheria, 314; of Sphæ-
 roplea, 318; of Characæ, 325, 326; of
 Fuci, 330; of Floridæ, 331; of Mar-
 chantia, 347; of Mosses, 349; of Ferns,
 355, 356.
Aphides, non-sexual reproduction of, 580,
 581.
Anthozoa, 457, 472-476.
Apple, cuticle of, 389, 390.
Apus, 544.
Aquatic Box, 148, 169.
ARACHNIDA, microscopic forms of, 581, 582;
 eyes of, 582; respiratory organs of, 582,
 583; feet of, 583; spinning apparatus of,
 583.
Arachnoidiscus, 290, 291.
Archegonia, of Marchantia, 344-347; of
 Mosses, 349; of Ferns, 356-358.
Arcus Senilis, 665.
Areolar tissue, 603.
Argulus, 547, 548.
Artemia, 545.
Ascaris, 531, 532.
Asci of Lichens, 334; of Fungi, 335.
Ascidians, compound, 50, 500; social, 502-
 505.
Asphalte, use of, 207.
Aspidium, fructification of, 353-355.
Aspidisca-form of Trichoda, 421.
Asteriada, skeleton of, 482, 483; metamor-
 phoses of, 486-488.
Asteroida, 472-474.
Auditory vesicles of Mollusks, 527, 528;
 development of, 525.
Avicularia of Polyzoa, 498, 499.
Azure blue butterfly, scales of, 195, 558.
Bacillaria paradoxa, 296; movements of,
 287, 288.
Bailey, Prof., his diatomaceous tests, 199.
Balani, metamorphoses of, 548-550.
Bark, structure of, 384, 385.
Barnacles, metamorphoses of, 548-550.
Bat, hair of, 195, 595, 596.
Battle-door scale of Polyommatus, 195, 558.
Batrachospermæ, 321, 322.
Bee, eyes of, 561; proboscis of, 567;
 wings of, 574; sting of, 578.
Berg-mehl, 303, 304.
Beroe, 470, 471.
Biddulphia, 299; markings on, 281; self-
 division of, 284.
Binocular Microscopes, 114-117.
Bird, Dr. Golding, on preparation of Zoo-
 phytes, 474.

- BIRDS, bone of, 587; feathers of, 597, 598; blood of, 599, 600; lungs of, 624-626.
 Bird's-head processes of Polyzoa, 498, 499.
 Black-Japan varnish, 219.
 Black-Ground Illuminators, 138-140, 174, 175.
Blenny, viviparous, scales of, 591.
 Blood-Disks of Vertebrata, 183, 598-601; mode of examining and preserving, 601, 602; circulation of, mode of examining, 616-619, 691.
 Bloodvessels, injected preparations of, 619-623; disposition of, in different parts, 623-626; morbid, 666.
 Bone, structure of, 183, 584-587; mode of making sections of, 207-211, 587.
 Bones, Fossil, examination of, 207-643.
Botrytis bassiana, 336.
Botryllans, 502.
Bowerbankia, 50, 496, 500.
Brachionus, 427, 429, 434.
Brachiopoda, structure of shell of, 513-515.
Branchiopoda, 542-545.
Branchipus, 545.
 Brooke, Mr., his Object-glass holder, 127.
 Brunswick-black, use of, 218.
Buccinum, tongue of, 521; egg-capsules of, 522; development of, 523.
 Bugs and their allies, 648; wings of, 676.
Bugula avicularia, 498, 499.
 Bull's-Eye Condenser, 143; use of, 178.
 Busk, Mr. G., on Volvox, 260, 265.
 Butterflies, see Lepidoptera.
- Cabinets for microscopic objects, 243.
Cactus, raphides of, 371.
Calycanthus, stem of, 385.
 Camera Lucida, 125, 126; use of, in Micrometry, 126.
Campanularida, 463-465.
Campylodiscus, 292.
 Canada balsam, use of, as cement, 208, 209, 219, 220; mounting of objects in, 224-232.
 Canal-system of Foraminifera, 446, 447.
 Canaliculi of bone, 585-587.
 Cancer cells, 675.
 Capillaries, circulation in, 616-619; injection of, 619-622; distribution of, 623-626.
Carp, scales of, 592.
 Cartilage, structure of, 608, 609.
 Cells for mounting objects, of cement, 235; of thin glass, 236; of plate-glass, 237; shallow, 237; deep and built up, 238-240; mounting objects in, 240-242.
 Cells, Animal, nature of, 404, 405.
 Cells, Vegetable, 42; nature of, 249; rotation in, 363-367; thickening deposits in, 367, 368; spiral deposits in, 369; starch-grains in, 370, 371; raphides in, 371.
 Cellular Tissue, ordinary form of, 360-362; stellate, 362; cubical, 363; dimensions of, 363; component cells of, 363.
 Cement-Cells, mode of making, 235, 236.
 Cementum of Teeth, 589-591.
 CEPHALOPODS, shell of, 516, 517; chromatophores of, 528.
Ceramiacea, 331.
Chatophoraceae, 321, 322.
 Chalk, Foraminifera, &c., of, 631-633.
Characeae, 322, 323; rotation of fluid in, 47, 324; multiplication of, by zoospores, 324; sexual apparatus of, 325, 326.
 Cherry-stone, cells of, 368.
 Chemical Reagents, use of, in Microscopic research, 211-213.
Chilodon, teeth of, 415; self-division of, 418.
Chirodota, calcareous skeleton of, 486.
 Chloride of Calcium, as mounting medium, 233.
 Choroid, pigment of, 605.
 Chromatic Aberration, 71, 72; means of reducing and correcting, 73, 74.
 Chromatophores of Cephalopods, 527.
Cidaris, spines of, 481.
 Ciliary action, nature of, 424, 425; on gills of Mollusks, 527.
 Ciliated epithelium, 606, 607.
 Circulation of Blood in Vertebrata, 616-619; in Insects, 569, 570; alternating, in Tunicata, 504, 505.
Cirrhipeda, metamorphoses of, 548-550.
Clavellinida, 503.
 Clarke, Mr. J. L., his mode of preparing sections of Spinal Cord, 616.
 Cleanliness, importance of, to Microscope, 158-160; in mounting objects, 242.
Closterium, movement of fluid in, 267, 268; duplicative subdivision of, 263, 269; multiplication of, by gonidia, 275; conjugation of, 274.
Clypeaster, spines of, 481.
 Coal, nature of, 627-629.
 Coarse-adjustment, 86-88; uses of, 162, 163.
Cockchaffer, cellular integument of, 556; eyes of, 562; antenna of, 565; spiracle of larva of, 572.
 Cockle of wheat, 531, 532.
 Coddington lens, 80.
 Cohn, Dr., his account of various states of Protococcus, 253-259; his researches on Infusoria, 425, note; on reproduction of Rotifera, 430, 431.
Coleoptera, mouth of, 565-567.
 Collection of objects, general directions for, 243-245.
Collomia, spiral fibres of, 368-370.
 Colostrum, 683.
 Columella of Mosses, 351.
Comatula, metamorphosis of, 490.
 Compound Microscope, principle of, 81-85; various forms of, 96-117.
 Compressorium, 150-152; use of, 168.
 Concave lenses, refraction by, 69.
 Conceptacles of Marchantia, 345, 346.
 Condenser, Achromatic, 131-133, 136; use of, 172, 173.
 Annular, 139.
 for Opaque Objects, ordinary, 144; bull's-eye, 144; mode of using, 177-179.
Confervaceae, 317; self-division of, 318; zoospores of, 318; sexual reproduction of, 318, 319.
Coniferae, peculiar woody fibre of, 372; absence of ducts in, 375, 382; fossil, 382, 628, 629.
 Conjugation of Palmoglaea, 252; of Desmidiaceae, 274-276; of Diatomaceae, 284-286; of Conjugatae, 319, 320; (supposed) of Actinophrys, 408.

- Contractile vesicle of Volvox**, 260; of *Amœba*, 405; of *Actinophrys*, 407; of *Infusoria*, 417, 418.
Convex lenses, refraction by, 65-69; formation of images by, 69.
Coquilla nut, cells of, 368.
Coral, red, 473; stony, 473, 474.
Corallines (zoophytic), 38, 49; true, 331, 332.
Cordylophora, 461, 462.
Cork, 380.
Corpuscles of Blood, 598-601.
Corti, Abbé, on rotation in *Chara*, 47.
Corynida, 461, 462.
Coscinodiscus, 289, 290.
Cosmarium, self-division of, 269; conjugation of, 274.
Crab, shell-structure of, 550, 551; metamorphoses of, 551, 552.
Cricket, gastric teeth of, 569; sounds produced by, 575.
Crinoidea, skeleton of, 482.
Cruata petrosa of Teeth, 589-591.
CRUSTACEA, 539; lower forms of, 539-541; entomostracous, 541-546; suctorial, 547, 548; cirriped, 548-550; decapod, shell of, 550, 551; metamorphoses of, 551, 552.
ΚΡΥΠΤΟΓΑΜΙΑ, microscopic study of, 43-46; movements of, 44; sexuality of, 45, 46, 359. (See *Protophyta*, *Algæ*, *Mosses*, &c.)
Crystals of snow, 644.
Crystallization, Microscopic, 645.
Ctenoid scales of Fish, 594.
Curculionida, 556-559.
Cutaneous eruptions, 668.
Cuticle of Equisetaceæ, 358; of Flowering Plants, 388-391.
Cuttle-fish, shell of, 516, 517.
Cycloid scales of Fish, 592.
Cyclops, 542, 543; fertility of, 546.
Cydippe, 470, 471.
Cypris, 541, 542.
Cypræa, structure of shell of, 515.
Cystic Entozoa, 530.
Cysticercus, 530.
Cythere, 542.

Dalrymple, Mr., on *Notommata*, 429.
Dalyell, Sir J. G., on development of *Medusæ*, 468-470.
Daphnia, 544; ephippial eggs of, 546, 547.
Darwin, Mr., on Pampean formation, 639, 640.
Deane's Gelatine, 232-234.
Decapod Crustacea, shell of, 550, 551; metamorphoses of, 551, 552.
Defining power of object-glasses, 187, 188.
Demodex folliculorum, 582.
Dentine of Teeth, 588, 589.
Dendrodus, teeth of, 641.
Dermestes, hair of, 195, 559.
Desmidiaceæ, general structure of, 265-267; movement of fluid in, 267, 268; duplicative subdivision of, 268-270; formation of gonidia by, 270-273; organization and multiplication of varieties in, 273, 274; conjugation of, 274-276; collection of, 276, 277.
Deutzia, stellate hairs of, 390, 391.
Development, Von Baer's law of, 52.
Animal, application of Microscope to, 50-57.
Vegetable, application of Microscope to, 42-47.
Diamond-beetle, 556-560; foot of, 576.
Diaphragm-Plate, 130, 131.
Diatoma, 298.
Diatomaceæ, vegetable nature of, 277; cohesion of frustules of, 278, 279; siliceous envelope of, 277-280; markings of, 280-283; self-division of, 283, 284; gonidia of, 284; conjugation of, 284-286; limits of species of, 286, 287, 301; movements of, 287, 288; classification of, 288-301; general habits of, 301-303; fossilized deposits of, 303, 304, 629-633; collection of, 304-306; mounting of, 306, 307; their value as tests, 195-199.
Dicotyledonous Stems, structure of, 378-386.
Dictyochalix, 453.
Didymoprium, self-division of, 269; conjugation of, 274, 275.
Diffraction of Light, errors arising from, 181, 182.
Dipping-Tubes, 152.
Dissecting Microscope, Quekett's, 94-96. Smith and Beck's, 103-105.
Dissection, microscopic, 201-203.
Distoma, 532, 533.
Dog, foot of, epidermis of, 606.
Doris, spicular skeleton of, 516.
Dotted Ducts, 374.
Doublet, microscopic, 79.
Draw-Tube, 118, 119.
Dropsical fluids, examination of, 699.
Dry-mounting of objects, 221-225.
Ducts of Plants, 374, 375.
Dujardin, M., on *Rotifera*, 432-435.
Duplicative subdivision of Palmoglaæ, 252; of *Protococcus*, 255; of *Desmidiaceæ*, 268-270; of *Diatomaceæ*, 283, 284; of *Confervaceæ*, 318, 319; of *Protozoa*, 407-409; of *Infusoria*, 418.
Dusidea, skeleton of, 455.
Dytiscus, foot of, 577, 578.

Eagle Ray, teeth of, 589.
Echinida, shell of, 477, 478; ambulacral disks of, 479; spines of, 479, 480; pedicellariæ of, 481, 482; teeth of, 482; metamorphosis of, 488-491.
ECHINODERMATA, skeleton of, 477-486; metamorphoses of, 486-491.
Educational Microscopes, 93-97, 104, note.
Educational Value of Microscope, 57-64.
Eel, scales of, 591, 592.
Eels, of Paste and Vinegar, 531.
Eggs of Insects, 579, 580.
Egg-shell, fibrous structure of, 602.
Ehrenberg, Prof., his researches on *Animalcules*, 47-49, 411-424, 428-435; on *Polycystina*, 449-452.
Elaters of Marchantia, 349.
Elementary Tissues: see *Tissues*.
Elytra, of Beetles, 575.
Embryo, vegetable, development of, 400.
Enamel of Teeth, 589-591.
Encrinites, 482.
Encysting process of Infusoria, 419-422.
Endochrome of vegetable cell, 249.

- Enterobryus*, 338-340.
Entomostracous Crustacea, 541; their classification, 541-545; reproduction of, 545-547.
Entozoa, 529; simplest form of, 530, 531; cystic, 530; nematoid, 531, 532; trematode, 532, 533.
Ephemera, larva of, 555-569, 573.
 Epidermis of Animals, structure of, 604, 605.
 of Plants, nature of, 391.
 Epithelium, 606, 607; ciliated, 607, 608.
Equisetaceæ, 358; cuticle of, 358; spores of, 359.
 Erector, 119, 120.
 Ether injection, 703.
Euryale, skeleton of, 482.
 Eye, examination of, 671.
 Eyes, care of, 158.
 Eyes, of Mollusks, 526, 527; of Insects, 561-564.
 Eye-piece, 82; Huyghenian, 83, 84; Ramsden's, 84, 85; Achromatic, 85; Meniscus, 85; Micrometric, 122-124, 713.
 Fallacies of Microscopists, 39-41, 180-186.
 Falconer, Dr., on bones of fossil Tortoise, 642.
 Fat-cells, 607, 608; capillaries of, 623, 624.
 Fatty degeneration, 665.
Faujasiina, 439; canal-system of, 446, 447.
 Feathers, structure of, 597, 598.
 Feet of Insects, 576, 578.
 FERNS, 352, 253; fructification of, 353, 354; spores of, 354, 355; prothallium of, 355; antheridia of, 355, 356; archegonia of, 356, 357; generation and development of, 357, 358.
 Fertilization of ovule, in flowering plants, 399-401.
 Fibre-cells of anthers, 396, 397.
 Fibres, Muscular, 611-614.
 Nervous, 614-616.
 Fibrillæ of Muscle, structure of, 612, 613.
 Fibrous tissues, 602-604.
 Field's Compound Microscope, 97-99.
 Simple Microscope, 93, 94.
 Filiferous capsules of Zoophytes, 474-476.
 Finder, 129, 130.
 Fine Adjustment, 86, 87; uses of, 163, 164.
 FISHES, bone of, 587; teeth of, 588; scales of, 591-593; blood of, 598-600; circulation in, 619; gills of, 624.
 Fishing-Tubos, 132.
 Flatness of field of object-glasses, 189.
 Flint, organic structure in, 633; examination of, 634.
Floridaea, 331.
Floucularians, 432, 433.
 Flowers, small, as microscopic objects, 394, 395.
 Fluid, mounting objects in, 234-242.
Fluke, 532, 533.
Flustra, 49, 492-497.
 Fly, number of objects furnished by, 553; tongue of, 566, 567; spiracle of, 572; wing of, 574; foot of, 576, 577.
 Focal Adjustment, 86, 87; precautions in making, 162-165; errors arising from imperfection of, 183, 184.
Foraminifera, 48, 436; their relation to Rhizopoda, 437; their general structure, 437-440; peculiar forms of, 440-445; relation of, to Sponges, 446; canal-system of, 446, 447; collection and mounting of, 446-449; fossil deposits of, 629-634.
 Forceps, 153; stage, 146.
 Fossil Bone, 642.
 Diatomaceæ, 303, 304, 629-633.
 Fossil Foraminifera, 629-639.
 Polycystina, 451, 452.
 Sponges, 632.
 Teeth, 641.
 Wood, 388, 627-629.
 Fowl, lung of, 625.
 Frog, blood of, 599, 600; circulation in web of, 616-618; in tongue of, 618, 619; lung of, 624.
Fucaceæ, 328, 329; sexual apparatus of, 328-331; development of, 331.
Fungi, simplest forms of, 334, 335; in bodies of living animals, 335-340; in substance, or on surface of plants, 340-343; higher forms of, 343, 344.
Fusulina, 634.
 Furze, Mr., his arrangement for illumination, 143.
God-flies, ovipositor of, 579.
Gaillonella, 299.
 Gairdner's Simple Microscope, 92, 93.
Gall-flies, ovipositor of, 579.
 Ganoid scales of Fish, 593.
 GASTEROPODA, structure of shell of, 515, 516; tongues of, 517-521; development of, 521-525; organs of sense of, 526-528.
 Gastric teeth of Mollusks, 521; of Insects, 569.
 Gelatine, Deane's, 232-234.
 Gelatinous nerve-fibres, 614, 615.
 Geology, applications of Microscope to, 627.
Geranium-petal, peculiar cells of, 395.
 Gills, of Mollusks, ciliary motion on, 526; of Fishes, distribution of vessels in, 624; of Water-newt, circulation in, 629; ciliary movement in, 624.
 Gillett's Condenser, 132.
 Glands, structure of, 609-611.
 Glass Slides, 214.
 Thin, 214-217.
 Glue, liquid, use of, 218.
 marine, use of, 219-221.
 Glycerine, use of, in mounting objects, 232.
Gnats, larvæ of, 569, 570.
 Goadby's Solution, 233.
 Gold-size, use of, 218.
 Goniometer, 124.
Gomphonema, 294, 295.
 Gonidia, multiplication by, in Desmidiaceæ, 270-272; in Diatomaceæ, 284; in Hydrodictyon, 300; in Chara, 325; in Floridaea, 331; in Lichens, 333.
Gorgonia, spicules of, 474.
 Gosse, Mr., on Melicerta, 433; on thread-cells of Zoophytes, 474-476.
Grammatophora, 298; its use as test, 199.
Grantia, structure of, 309-317.
Grasses, silicified cuticle of, 390, 391.
Gregarina, 530, 531.
 Grinding and Polishing of Sections, 207-211.

- Gromia*, 437, 438.
Growing-Slide, 147.
Grubb's Prism for oblique illumination, 137.
Guano, Diatomaceæ of, 304-306.
- Hairs**, of Mammals, 183; structure of, 593-596; mode of mounting, 597; of Insects, 559-561.
Hairs, of vegetable cuticles, 390; rotation of fluid in, 365, 366.
Halichondria, spicules of, 454.
Harvest-bug, 582.
Haversian Canals of bone, 585.
Heliopelta, 290, 291.
Haustellate Mouth, 567, 568.
Henfrey, Mr., on development of pollen-grains, 396.
Hepatica, 344; see *Marchantia*.
Herapathite, 140.
Hepworth, Mr., on feet of Insects, 577.
Highley's Hospital Microscope, 99, 100.
Hippocrepian Polyzoa, 497-500.
Histology, 54.
Hamatococcus, its relations to *Protococcus*, 254.
sanguineus, 308.
Hogg, Mr., on development of *Lymnæus*, 523.
Holland, Mr., his triplet, 80.
Holothurida, skeletons of, 485, 486.
Hoofs, structure of, 598.
Hooker, Dr. Jas., on Antarctic Diatomaceæ, 301, 302.
Horns, structure of, 598.
Huxley, Mr., on Rotifera, 430, 431, 435; on Noctiluca, 472; on blood of Annelida, 536.
Huyghenian eye-piece, 82-84.
Hydatina, 434; reproduction of, 431.
Hydra, discovery of, 37; structure of, 457-461.
Hydrodictyon, 316, 317.
Hydrozoa, 461-470.
Hymenoptera, proboscis of, 566, 567; wings of, 574; stings and ovipositors of, 578-579.
- Jackson, Mr.**, his eye-piece micrometer, 122, 123.
Jelly-fish, development of, 467-470.
- Ice-plant**, cuticle of, 390.
Ichneumonida, ovipositor of, 578.
Illumination of opaque objects, 176-180; of transparent objects, 172-175.
Illuminators, Black-ground, 138-140, 174, 175.
 Oblique, 135-138, 173, 174.
 White Cloud, 134, 173.
- Images**, formation of, by convex lenses, 69.
Indicator, 124.
Indusium of Ferns, 352-354.
Inflection of Light, errors arising from, 181, 182.
Infundibulate Polyzoa, 497-500.
Infusorial Earths, 303-306, 629.
INFUSORIA, 48, 411, 412; structure of, 413-417; movements of, 415, 416; reproduction of, 418-423; peculiar forms of, 423, 424.
Injections of bloodvessels, mode of making, 619-622; mode of mounting, 622, 623, 700.
Inorganic substances as objects, 644.
INSECTS, great number of objects furnished by, 553, 554; microscopic forms of, 554, 555; antennæ of, 564, 565; circulation of blood in, 569, 570; eggs of, 579-581; eyes of, 561-564; feet of, 576-578; gastric teeth of, 569; hairs of, 559-561; integument of, 555, 556; mouth of, 565-568; ovipositors of, 578, 579; scales of, 556-561; spiracles of, 572-574; stings of, 578, 579; trachææ of, 570-574; wings of, 574-576.
Iris, structure of leaf of, 392-394.
Isthmia, 298; markings on, 280, 281; self-division of, 283.
Ith-Acarus, 587.
Integument of Insects, structure of, 555, 556.
- Kidneys**, structure of, 610; morbid, 664.
- Labelling** of objects, 242, 243.
Labyrinthodon, tooth of, 641.
Lacunæ of bone, 585-588.
Laguncula, 492-495, 500.
Lamps, microscope, 154-157.
Lealand, Mr., his preparations of muscular fibre, 195, 614.
Leaves, structure of, 392-394; mode of examining, 394.
Leech, 538; teeth of, 538.
Legg, Mr., on collection of Foraminifera, 447-449.
Lepidoptera, scales of, 556-561; proboscis of, 567, 568; wings of, 574, 575; eggs of, 564.
Lepidosteus, bony scales of, 586, 587, 593.
Lepisma, scales of, 559.
Lepralia, 492-497.
Lever of Contact, 216.
Levant Mud, microscopic organisms of, 630-633.
Lever-Stage, 128, 707.
Lichens, 333, 334.
Licnophora, 295, 296.
Lieberkühn, speculum of, 146; mode of using, 179, 180.
Light, suitable for Microscope, 154-157; position of, 157; arrangement of, for transparent objects, 169-176; for opaque objects, 177-180.
Liquid Glue, use of, 218.
Lister, Mr., his improvements in Achromatic lenses, 75, 76.
Liver, structure of, 609; morbid, 663.
Lungs of Reptiles, 624; of Birds, 624, 625; of Mammals, 626; morbid, 658.
Lymnæus, development of, 521, 522.
- Magnetic Stage**, 130.
Magnifying power, mode of determining, 199, 200; augmentation of, 161; of different objectives, 192-196.
Mahogany, section of, 383.
MAMMALS, bone of, 587; teeth of, 589-591; hairs, &c., of, 593-598; blood of, 598-602; lungs of, 625.

- Man*, hair of, 595, 596.
Mandibulate mouth of Insects, 565-567.
Marchantia, general structure of, 344, 345; stomata of, 345, 346; conceptacles of, 346; sexual apparatus of, 347, 348.
Margaritacea, shells of, 506-508, 509.
Marine Glue, uses of, 219-221.
Matogloia, 299-301.
Medullary Rays, 381-384.
Medusa, development of, from Zoophytes, 52, 53, 467-470.
Megatherium, teeth of, 590.
Melicertians, 432-434.
Meloseira, 299; self-division of, 284; conjugation of, 286.
Menelaus, scale of, 557.
Meniscus eye-piece, 85.
Meridion, 296, 297.
Metamorphoses of Animals, 50; of Cirripeds, 50, 548-550; of Crustacea, 52, 551, 552; of Echinoderms, 52, 486-491; of Infusoria, 419-422; of Mollusks, 51, 504, 505; of Annelids, 52, 536, 537.
Micrometer, Cobweb, 120-122; eye-piece, 122-124.
Micrometry, by Micrometer, 120-124; by Camera Lucida, 126.
Microscope, early history of, 35-41; later history of, 41-57; educational uses of, 57-64. optical principles of construction of, 77-85; mechanical principles of construction of, 86-90.
Compound, 81-85, 96, 97.
Field's, 97-99.
Highley's, 99, 100.
Nacht's, 100-102.
Ditto, binocular, 114-117.
Powell and Lealand's, 110-112.
Ross's, 108-110.
Smith and Beck's large, 112-114.
Ditto dissecting, 103-105.
Ditto student's, 102, 103.
Ditto educational, 104, *note*.
Warington's universal, 105-108.
Simple, 77-81, 90.
Field's, 93, 94.
Gairdner's, 92, 93.
Grunow's, 706.
Queen's, 706.
Quekett's, 94-96.
Ross's, 90-92.
Spencer's, 704.
support required for, 154; care of, 158-160; general arrangement of, 160; for transparent objects, 168-172; for opaque objects, 175-177.
Microscopic Dissections, 201-203.
Microscopists, fallacies of, 39-41.
Microtome, 203, 204.
Mildev, fungous vegetation of, 340, 341.
Milk, 683.
Millon's test for albuminous substances, 213.
Mineral Objects, 645.
Minnow, circulation in, 619.
Molecular movement, 185, 186.
MOLLUSCA, shells of, 506-517; tongues of, 517-521; development of, 521-525; ciliary motion on gills of, 526; organs of sense of, 527, 528.
Monocotyledonous Stems, structure of, 376-378.
Morbid growths, examination of, 674.
Morpho Menelaus, scale of, 194, 557.
Mosses, structure of, 348, 349; sexual apparatus of, 347-350; urns of, 350, 351; peristome of, 351; development of spores of, 352.
Moths; see Lepidoptera.
Mould, fungous vegetation of, 340, 341.
Mounting of objects; see Objects.
Mounting-Plate, 219.
Mouse, Ear of, cartilage of, 608.
Mouth of Insects, 565-568.
Mucous Membranes, structure of, 604; capillaries of, 623; ulcers of, 670.
Müller, O. F., his researches on Animalcules, 38.
Müller, Prof., his researches on Echinoderm-larvæ, 486-491.
Muscardine, or silk-worm disease, 44, 45, 336.
Muscular fibre, structure of, 611-613; value of, as test-object, 195; mode of examining and preparing, 613, 614, 655; capillaries of, 623, 624.
Musk-deer, hair of, 595.
Mussel, ciliary action on gills of, 526.
Mya, structure of hinge-tooth of, 509.
Mycelium of Fungi, 340-343.
Myliobates, teeth of, 589.
Nacht's Microscope, 100-102; his Binocular Microscope, 114-117; his prism for oblique illumination, 136, 137.
Nacre, structure of, 509-511.
Nais, 537, 538.
Nassula, teeth of, 415.
Navicula, 293, 294; movements of, 287.
Needles for dissection, mode of mounting, 203.
Nematoid Entozoa, 531, 532.
Nepenthes, spiral vessels of, 374.
Nervous Tissue, structure of, 614, 615, 653; mode of examining, 615, 616, 653.
Newt, circulation in larva of, 619.
Nicol-Prism, 140.
Nobert's Test, 196, 197.
Noctiluca, 471, 472.
Nostochacea, 312, 313.
Notommata, 429-435.
Nucleus of Vegetable cells, 251, 365; of Infusoria, 418.
Nudibranchs, development of, 521, 522.
Nummulite, structure of, 446, 634-637.
Nuphar lutea, parenchyma of, 362.
Object-Finder, 129, 130.
Object-glasses, achromatic, principle of, 73, 74; construction of, 74-76; adjustment of, for covering of object, 76, 77, 165-168; defining power of, 187, 188; penetrating power of, 188; resolving power of, 188, 189; flatness of field of, 189; comparative value of, 188-192; different powers of, 193-198; tests for, 193-199;

- determination of magnifying power of, 199, 200.
Object-glass holder, Mr. Brooke's, 127.
Object-Marker, 127, 128.
Objects, mode of mounting, dry, 221-224; in Canada balsam, 224-232; in preservative fluids, 232-242; see **Opaque and Transparent Objects**.
Oblique Illuminators, 135-138, 173, 174.
Ocelli of compound eyes of Insects, 561.
Edogonium, zoospore of, 318; sexual reproduction of, 319.
Oil-globules, microscopic appearances of, 184.
Oleander, cuticle of, 389.
Oncidium, spiral cells of, 368.
Onion, raphides of, 371.
Oolite, structure of, 634.
Opaque Objects, arrangement of Microscope for, 175-177; various modes of illuminating, 177-180; modes of mounting, 121-124.
Opercula of Mosses, 350.
Ophiocoma, 482.
Ophiurida, skeleton of, 482, 483.
Ophrydina, 423.
Orbitoides, structure of, 637-639.
Orbitolite, structure and development of, 440-445.
Ornithorhynchus, hair of, 595.
Oscillatoria, 311-313.
Ovipositors, 578, 579.
Ovules of Phanerogamia, 399; fertilization of, 48, 399, 400; mode of studying, 400, 401.
Owon, Prof., on Fossil Teeth, 641, 642; on Fossil Bone, 642, 643.
Ozytricha-form of Trichoda, 421, 422.
- Palm**, stem of, 377, 378.
Palmellacea, 307-309.
Palmoglaea macrococca, life-history of, 251-253.
Pampean deposit, organic composition of, 639.
Papillæ of skin, structure of, 605-615; of tongue, 615.
Parabolic Illuminator, 139, 140.
Paramecium, 413, 414; contractile vesicle of, 417; multiplication of, 418.
Parasitic fungi, 335-443.
Pearls, structure of, 509-511.
Pecari, hair of, 595.
Pecten, eyes of, 527; tentacles of, 528.
Pediastrum, multiplication and development of, 270-273.
Pedicellariæ of Echinoderms, 481, 482.
Penetrating power of object-glasses, 188.
Pentacrinus, skeleton of, 482; larval, 491.
Peristome of Mosses, 350, 351.
Perophora, 502-505.
Petals of Flowers, structure of, 395, 396.
Pettenkofer's test, 212.
PHANEROGAMIA, elementary tissues of, 360-376 (see **Tissues of Plants**); stems and roots of, 376-388; cuticle and leaves of, 388-394; flowers of, 394-401; seeds of, 401-403.
Phytelephas, cells of, 368.
Phyllopora, 544.
- Pigment-cells**, 605.
Pith, structure of, 378, 379.
Pinna, structure of shell of, 506-508.
Pinnularia, 294.
Placoid scales of Fish, 593.
Planaria, 533-535.
Plants, distinction of from Animals, 247, 248, 409, 410.
Plantago, rotation in hairs of, 366.
Pleurosigma, 294; nature of markings on, 281, 282; value of, as tests, 195-198.
Pluteus-larva of Echinus, 488-490.
Pneumonia, pathological condition in, 661.
Podura, scale of, 195, 556, 557.
Polarization, objects suitable for, 645.
Polarizing Apparatus, 140-143.
Pollen-grains, development of, 397; structure and markings of, 194, 397-399.
Polycystina, nature of, 449, 450; distribution of, 450-452.
Polygastrica; see **Infusoria**.
Polyommatus argus, scale of, 557, 558.
Polypes; see **Hydra** and **Zoophytes**.
Polypodium, fructification of, 353.
POLYZOA, 49, 492; general structure of, 492-496; molluscan nature of, 496; classification of, 497-500.
Potato-disease, 342.
Powell and Lealand's Microscope, 110-112; their **Achromatic Condenser**, 133.
Preservative Liquids, 232-234.
Primordial utricle, 248.
Prismatic shell-substance, 506-508.
Prisms, Dujardin's, 133; 134, 173; Nachet's, 136; Amici's, 137; Grubb's, 137; polarizing, 140, 141.
Proboas of Bee, 566, 567; of Butterfly, 567, 568; of Fly, 566, 567.
Proteus, 405.
Protococcus pluvialis, life-history of, 253-257; various forms of, 257, 258; conditions influencing changes of, 258, 259.
Protoplasm of vegetable cell, 248.
ПРОТОФЫТА, general characters of, 247-249.
PROTOZOA, 404-409; their relations to **Protophyta**, 247, 248, 409, 410.
Pseudopodia, of Rhizopods, 407-411.
Pterodactyle, bone of, 642.
Puccinia, 341.
Purpura, development of, 521-525.
Pycnogonida, 539-541.
- Quekett**, Mr., his **Dissecting Microscope**, 94-96; his **Indicator**, 124; on **Structure of Bone**, 642, 643.
Querquedula crecca, 655.
- Rainey**, Mr., his **moderator**, 156.
Ramsden's eye-piece, 84, 85.
Raphides, vegetable, 371, 372.
Reade, Rev. J. B., on **black-ground illumination**, 138.
Reagents, Chemical, use of, in **Microscopic research**, 211-213.
Red Corpuscles of Blood, 599, 600.
Red Snow, 307.
Refraction, laws of, 65, 66; by **convex lenses**, 65-69; by **concave and meniscus lenses**, 69.

- Reindeer*, hair of, 595.
REPTILES, bone of, 587; teeth of, 591; scales of, 593; blood of, 598-600; lungs of, 624.
 Resolving power of object-glasses, 187-189.
 Rete Mucosum, 604.
 Reticulated Ducts, 374.
Rhinoceros, horn of, 598.
RHIZOPODA, 47, 410, 411.
Rhubarb, raphides of, 371.
Rhynchonellida, structure of shell of, 513-515.
 Rice-Paper, 362.
Rochea falcata, 390.
 Rocks, structure of, 639.
 Roots, structure of, 386; mode of making sections of, 386-388.
Rosalina, 439.
 Ross, Mr., on adjustment of object-glass, 76, 77.
 his Compound Microscope, 108-110; his Achromatic Condenser, 131-133; his Simple Microscope, 90-92; his lever of contact, 216; his side reflector, 145; his eye-piece Micrometer, 122.
 Rotation of fluid in Chara, 324; in cells of Phanerogamia, 363-367.
Rotifer, anatomy of, 428-430; reproduction of, 431; desiccation of, 432; occurrence of, in leaf-cells of Sphagnum, 426.
ROTIFERA, 57, 412; general structure of, 426-430; reproduction of, 430-432; desiccation of, 432; classification of, 432-434.
Rush, stellate parenchyma of, 362.
 Rust, of Corn, 341.
- Sable*, hair of, 595.
 Saliva, 684.
 Sarcode, 405-409.
 Sarcolemma, 655.
Saw-flies, ovipositor of, 578.
 Scalariform ducts of Ferns, 352, 353, 374.
 Scales, of cuticle of Plants, 390.
 of Fish, 591-593.
 of Insects, 556-561; their use as test-objects, 194.
 of Reptiles and Mammals, 594.
 Schacht on fertilization of ovule, 399-401.
 Schleiden, Prof., his researches, 42; on fertilization of ovule, 399-401.
 Schultz, Prof., his experiments on development of Infusoria, 422, 423.
 Schultz's solution, 212.
 Schwann, doctrines of, 56.
 Scissors for microscopic dissection, 201-203; for cutting thin sections, 204.
 Sclerogen, deposit of, on walls of cells, 367.
Sea-Anemone, 474-476.
 Sealing-wax varnish, 219.
 Section-Instrument, 205-207.
 Sections, thin, mode of making, of soft substances, 204, 205; of substances of medium hardness, 205-207; of hard substances, 207-211; of Wood, 386-388; of Echinus-spines, 483, 484; of Bones and Teeth, 587; of Hairs, 597.
 Seeds, microscopic characters of, 401-403.
 Segmentation of Yolk-mass, 521.
 Selenite-Plate, 142.
- Self-division of cells; see Duplicative Sub-division.
 Serous membranes, structure of, 604.
Sertularida, 463-465.
 Shadbolt, Mr., on Arachnoidiscus, 290.
 his Annular condenser, 139; his Turn-table, 236.
Shark, teeth of, 588, 589; scales, &c., of, 593.
 Shell, of Crustacea, 550, 551; of Echinida, 477, 478; of Mollusca, 506-517.
 Side Reflector, Ross's, 145.
 Siliceous Cuticles, 358, 391.
 Silk-worm disease, 336.
 Simple Microscope, principle of, 77-81; various forms of, 90-96.
Siphonacea, 313-317.
 Skin, structure of, 604, 605; papillæ of, 605-615.
 Slider-forceps, 226.
 Slides, glass, 214.
 wooden, 221, 222.
Slug, rudimentary shell of, 516.
 Smith's, Prof. J. L., inverted microscope, 710; goniometer and micrometer, 712.
 Smith, Prof. W., on Diatomacea, 301, 302.
 Smith and Beck's Dissecting Microscope, 103-105; their Large Compound Microscope, 112-114; their Achromatic Condenser, 133; their Student's Microscope, 102, 103; their Educational Microscopes, 89.
Snake, lung of, 624.
 Snow-crystals, 644.
Sole, skin and scales of, 591, 592.
 Sollitt's Achromatic Condenser, 136.
 Sorby, Mr., his microscopic examination of rocks, 640.
Spatangus, spines of, 481.
 Speculum-disk for drawing, 125.
 Spencer, Mr., his object-glass, 51, *note*; his microscopes, 704.
 Spermata of Lichens, 334; of Fungi, 344.
 Spermogonia of Lichens, 333; of Fungi, 344.
Sphacclaria, 328.
Sphagnum, peculiar cells of, 348, 349; occurrence of Rotifer in leaf-cells of, 426.
Sphæria, development of, within animals, 338.
Sphaeroplea, sexual reproduction of, 318.
Sphaerosira volvox, 264.
 Spherical Aberration, 70; means of reducing and correcting, 71, 72.
 Spicules of Sponges, 453-455; of Aleyonian Zoophytes, 472-474; of Tunicated Mollusks, 500; of Doris, 516.
Spiders, eyes of, 582; respiratory organs of, 583; feet of, 583; spinning apparatus of, 583.
 Spinal Cord, mode of preparing sections of, 616.
 Spines of Echinus, &c., 479, 480; mode of making sections of, 482-484.
 Spinning-apparatus of Spiders, 583.
 Spiracles of Insects, 571-573.
 Spiral Vessels, 373, 374.
SPONGES, their structure, 452, 453; ciliary action in, 453; skeleton of, 453-455; reproduction of, 455; examination of, 455; fossil, 633.

- Sporangia**, of Desmidiaceæ, 274, 275; of Diatomaceæ, 284-286; of Fuci, 330.
- Spores of Hepaticæ**, 347; of Mosses, 351, 352; of Ferns, 355; of Equisetaceæ, 358.
- Spotted Lens**, 138.
- Spring-press**, 227.
- Spirogyra**, 241.
- Sputum**, 685.
- Squirrel**, hair of, 594, 595.
- Stage**, lever, 128, 129; magnetic, 130. movements of, 109, 110.
- Stage-Forceps**, 146.
- Stage-Plate**, glass, 147.
- Stanhope lens**, 80, 81.
- Star-Anise**, cells of seed-coat of, 367, 368.
- Starch Granules** in cells, 370, 371; appearance of, by polarized light, 370, 371.
- Staurastrum**, prominences of, 265, 266; self-division of, 269; varieties of, 273.
- Stein, Dr.**, his researches on Infusoria, 419, 420.
- Stellate cells** of Rush, 362; of water-lily, 362.
- Stemmata of Insects**, 563.
- Stems**, Endogenous, structure of, 376-378. Exogenous, structure and development of, 378-386. Mode of making sections of, 386-388.
- Stentor**, 423.
- Stephanoceros, Eichornii**, 433.
- Stickleback**, circulation in, 619.
- Stigmata of Insects**, 571-573.
- Stings**, structure of, 578, 579.
- Stomata of Marchantia**, 345, 346; of Flowering Plants, 391, 392.
- Suctorial Crustacea**, 547, 548.
- Suminski, Count**, on development of Ferns, 46.
- Surirella**, 293; conjugation in, 284.
- Swarming**, of Desmidiaceæ, 270.
- Synapta**, calcareous skeleton of, 485, 486.
- Tadpole**, pigment-cells of, 604, 605; mode of viewing circulation in, 619.
- Tænia**, 529, 530.
- Tardigrada**, 435; desiccation of, 432.
- Teeth**, of Echinida, 482; of Mollusks, 517-521; of Leech, 538; of Vertebrata, structure of, 588-591; fossil, 641, 642; mode of making sections of, 207-211, 587.
- Terebella**, circulation and respiration in, 535, 536.
- Terebratula**, structure of shell of, 513-515.
- Test-Bottles**, 212.
- Test-Liquids**, 212, 213.
- Test-Objects**, 192-199.
- Tetraspores of Floridæ**, 331, 332.
- Thalassicolla**, 445.
- Thaumantia**, 467.
- Thecæ of Fungi**, 334, 335; of Ferns, 352, 353.
- Thin Glass**, 214-216.
- Thompson, Mr. J. V.**, on Polyzoa, 492; on metamorphosis of Cirrhipeds, 548-550; on metamorphoses of Crustacea, 551, 552.
- Thread-cells of Zoophytes**, 474-476.
- Thread**, glutinous, of spider's web, 582, 583.
- Thrush**, fungous vegetation of, 340.
- Thwaites's fluid** for Algæ, 232.
- Thymus and thyroid gland**, 664.
- Ticks**, 581.
- Tinea favosa**, fungous vegetation of, 340.
- Tissues, Elementary, of Animals**, microscopic study of, 53-55, 584; independent life of, 56, 57; see Blood, Bone, Capillaries, Cartilage, Epidermis, Epithelium, Fat, Feathers, Fibrous Tissues, Glands, Hair, Horn, Mucous Membranes, Musculo, Nervous Tissues, Pigment-cells, Scales, Serous Membranes, Teeth.
- Tissues, Elementary, of Plants**, 360; cellular, 360-363; varieties of, 367-371; woody, 372; fibro-vascular, 372; vascular, 373, 374; vasiform, 374, 375; dissection of, 375, 376.
- Tongues of Gasteropods**, 517-520; of Insects, 565-567.
- Torula cerevisiæ**, 334, 335.
- Tracheæ of Insects**, 570-572; mode of preparing, 573, 574.
- Tradescantia**, rotation in hairs of, 365, 366.
- Transparent Objects**, arrangement of Microscope for, 168-172; various modes of illuminating, 172-175.
- Trematode Entozoa**, 532, 533.
- Trembley**, his researches on the Hydra, 37.
- Triceratium**, 292; markings on, 281.
- Trychoda lynceus**, bristles of, 415; metamorphoses of, 420-422.
- Trilobite**, eye of, 640.
- Triplet, Microscopic**, 80.
- Trout**, circulation in young, 619.
- Tubercle**, microscopic appearance of, 658.
- Tubularidæ**, 462, 463, 465.
- Tubuleæ**, nervous, 614, 615.
- TUNICATA**, Compound, general organization of, 500-502; different types of, 502-504; alternating circulation in, 504, 505; development of, 505.
- Turbellaria**, 533-535.
- Turn-table**, Shadbol's, 236.
- Ulvaceæ**, 309-311.
- Unionidæ**, shells of, 508-512.
- Uredo**, 342.
- Urinary deposits**, 688.
- Urns of Mosses**, 350, 351.
- Vacuoles**, microscopic appearances of, 185.
- Valentin's knife**, 205.
- Vallisneria spiralis**, rotation in, 363, 364.
- Varnishes useful to Microscopists**, 216-219.
- Vasiform Tissue**, 374, 375.
- Vaucher**, his researches on Confervæ, 38.
- Vaucheria**, 313; zoospores of, 314, 315; sexual reproduction of, 315, 316.
- Vegetable Organization**, general nature of, 248; see Plants.
- Vegetable Cell**, nature of, 248, 249.
- Vegetable Ivory**, 368.
- VERTEBRATA**, elementary structure of, 585-616 (see Tissues); blood of, 598-601; circulation in, 616-619; development of, 626.
- Vesicles of Nervous tissue**, 614, 615.
- Vibracula of Polyzoa**, 498, 499.
- Villi of intestine**, injections of, 622.
- Vine disease**, 342.
- Volvox**, structure of, 259, 260; development of, 261-265.

- Vomited matters, 698.
 Vomit, black, 698.
 Von Baer's law of development, 52.
Vorticella, 416-423; encysting process in, 419, 420.
- Warington's Universal Microscope, 105-108.
 Water-vascular system, of Rotifera, 429, 430; of Entozoa, 530.
 Wenham, Mr., his Parabolic Speculum, 139.
 his observations on rotation, 364-367.
- Whalebone, structure of, 598.
Wheel-Animalcules; see Rotifera.
 White-Cloud Illuminator, 134, 173.
 White Corpuscles of blood, 598-601.
 White Fibrous tissue, 603.
 Williamson, Prof., on Volvox, 261-265.
 Wings of Insects, 574-576.
 Winter-eggs of Rotifera, 431; of Hydra, 61; of Entomostraca, 546.
- Wood, of Exogenous stems, 378-384.
 Woody Fibre, 372; glandular, of Coniferae, 373.
- Xanthidia* of Flints, 633.
- Yeast, a vegetable substance, 45; production of, 45, 334, 335.
 Yellow fibrous tissue, 603, 604.
 Yellow Water-lily, parenchyma of, 362.
Yucca, stomata of, 391, 392.
- Zoophyte-Trough, 149, 150.
 ZOOPHYTES, 457; Hydraform, 461-465; preparation of, for microscope, 465-467; development of Medusæ from, 52, 53, 467-470; Alcyonian, 472-474; Actiniform, 474-476.
 Zoospores, formation of, by Ulvaceæ, 310, 311; by Vaucheria, 314, 315; by Achlya, 315, 316; by Confervaceæ, 318, 319; by Chætophoraceæ, 321; by Fucaceæ, 330, 331.

THE END.

CATALOGUE

OF

BLANCHARD & LEA'S

MEDICAL AND SCIENTIFIC PUBLICATIONS.

PHILADELPHIA, JULY 1, 1854.

TO THE MEDICAL PROFESSION.

In submitting the following catalogue of our publications in medical and other sciences, we beg to remark, that no exertions are spared to render the issues of our press worthy a continuance of the confidence which they have thus far enjoyed, both as regards the high character of the works themselves, and in respect to every point of typographical accuracy and mechanical execution. Gentlemen desirous of adding to their libraries from our list, can in almost all cases procure the works they wish from the nearest bookseller, who can readily order any which he may not have on hand. From the great variation in the expenses of transportation through territories so extensive as those of the United States, prices cannot be the same in all sections of the country, and therefore we are unable to affix retail prices to our publications. Information on this point may be had of booksellers generally, or from ourselves, and we shall be always happy to answer inquiries.

Catalogues of our miscellaneous and educational publications will be forwarded by mail on application.

BLANCHARD & LEA.

PHILADELPHIA, July 1, 1854.

INDEX TO CATALOGUE.

	Page		Page
Abel and Bloxam's Handbook of Chemistry	52	Carson's Synopsis of Materia Medica	51
American Journal of the Medical Sciences	3	Chapman on Eruptive Fevers	18
Arnott's Elements of Physics	60	Chelius's System of Surgery, by South	32
Ashwell on Diseases of Females	42	Christison and Griffith's Dispensatory	50
Barlow's Practice of Medicine	19	Churchill's System of Midwifery	39
Bartlett on Fevers of the United States	22	Churchill on Diseases of Females	43
Beale on the Laws of Health	55	Churchill on Diseases of Children	43
Benedict's Compendium of Chapman's Lectures	18	Churchill on Puerperal Fever	43
Bennett (Henry) on Diseases of the Uterus	42	Clymer on Fevers	18
Bennett (J. Hughes) on Pulmonary Tuberculosis	4	Colombat de l'Isère on Females, by Meigs	41
Billing's Principles of Medicine	18	Coudie on Diseases of Children	44
Bird on Urinary Deposits	21	Cooper's (Bransby B.) Lectures on Surgery	37
Bird's Natural Philosophy	60	Cooper (Sir Astley P.) on Dislocations and Fractures	33
Blakiston on the Chest	18	Cooper (Sir Astley P.) on Hernia	33
Bonaparte's American Ornithology	57	Cooper (Sir Astley P.) on the Testis and Thymus Gland	33
Bowman's (John E.) Practical Chemistry	54	Cooper (Sir Astley P.) on the Breast	33
Bowman's (John E.) Medical Chemistry	54	Copland on Palsy and Apoplexy	18
Bowman's (Wm.) and Todd, Physiological Anatomy	17	Cyclopedia of Practical Medicine	19
Brewster's Elements of Optics	53	Dana on Corals and Zoophytes	61
Broderip's Zoological Recreations	57	Dana's Atlas to do.	61
Brodie's Select Surgical Works	37	Dana's Classification of Zoophytes	61
Brodie's Clinical Lectures on Surgery	37	Day on Diseases of Old Age	20
Buckler on Bronchitis	19	De Jongh on Cod-Liver Oil	49
Budd on Diseases of the Liver	22	De la Beche's Geological Observer	59
Burrows on Cerebral Circulation	18	Dewees's System of Midwifery	44
Carpenter's Principles of Human Physiology	12	Dewees on Diseases of Females	44
Carpenter's Elements of Physiology	13	Dewees on Diseases of Children	44
Carpenter's Comparative Physiology	14	Druitt's Modern Surgery	30
Carpenter on the Microscope and its Revelations	13	Dufton on Diseases of the Ear	37
Carpenter on Use and Abuse of Alcohol	45	Dunglison's Medical Dictionary	6
		Dunglison's Human Physiology	15
		Dunglison on New Remedies	48

	Page		Page
Dunglison's Therapeutics and Materia Medica	48	Meigs on Diseases of the Uterus	40
Dunglison's Practice of Medicine	21	Meigs on Diseases of Children	40
Dunglison's Human Health	55	Meigs on Puerperal Fever	40
Dunglison's Cyclopaedia of Practical Medicine	19	Mohr, Redwood, and Procter's Pharmacy	51
Ellis's Medical Formulary, by Thomas	49	Muller's Physics and Meteorology	61
Erichsen's System of Surgery	26	Neill and Smith's Compendium of Medical Science	7
Evans's Millwright's Guide	62	Neligan on Diseases of the Skin	23
Fergusson's Operative Surgery	31	Owen on the Skeleton and Teeth	11
Fownes's Elementary Chemistry	53	Paget's (Kirkes and) Manual of Physiology	16
Frick on Renal Affections	18	Pancoast's Operative Surgery	32
Gardner's Medical Chemistry	52	Parker's Modern Treatment of Syphilis	24
Gluge's Pathological Histology, by Leidy	21	Peaslee's Human Histology	8
Graham's Elements of Chemistry	54	Pereira's Materia Medica and Therapeutics	46
Gregory's Letters on Animal Magnetism	55	Philips on Scrofula	18
Griffith's (R. E.) Universal Formulary, by Thomas	45	Pirrie's System of Surgery	29
Griffith's (R. E.) Medical Botany	50	Popular Physiology of Animal and Vegetable Life	16
Griffith's (Thomas) Chemistry of the Four Seasons	57	Quain and Sharpey's Anatomy, by Leidy	9
Griffith's (John W.) Manual on the Blood, &c.	18	Ramsbotham on Parturition	38
Gross on Urinary Organs	35	Rees on Blood and Urine	18
Gross on Foreign Bodies in the Air-Passages	35	Ricord and Hunter on Venereal, by Bumstead	24
Gross's Principles and Practice of Surgery	35	Ricord's Illustrations of Syphilis, by Betton	24
Hale's Ethnography and Philology	57	Ricord's Letters on Syphilis, by Lattimore	24
Harrison on the Nervous System	12	Ricord on Venereal, by Doane	24
Hasse's Pathological Anatomy	18	Rigby's Principles and Practice of Midwifery	39
Herschel's Outlines of Astronomy	59	Rokitanski's Pathological Anatomy	18
Hoblyn's Medical Dictionary	5	Royle's Materia Medica and Therapeutics	49
Hope on Diseases of the Heart	18	Sargent's Minor Surgery	34
Horner's Anatomy and Histology	11	Schædler & Medlock's Book of Nature	56
Horner and Smith's Anatomical Atlas	10	Sharpey and Quain's Anatomy, by Leidy	9
Hughes on Auscultation and Percussion	20	Sibson's Medical Anatomy	11
Humboldt's Aspects of Nature	57	Sievekking and Jones, Pathological Anatomy	18
Hunter and Ricord on Venereal, by Bumstead	24	Simon's General Pathology	18
Hunter's Complete works	24	Skey's Operative Surgery	33
Johnson's Gardener's Dictionary	64	Smith (H. H.) and Horner's Anatomical Atlas	10
Johnston's Physical Atlas	60	Smith (F. G.) and Neill's Compendium of Medicine	7
Jones's (T. Wharton) Ophthalmic Medicine and Surgery	37	Smith (Tyler) on Parturition	42
Jones (C. Handfield) and Sievekking's Pathological Anatomy	18	Solly on Anatomy and Diseases of the Brain	17
Kirkes and Paget's Physiology	16	Somerville's Physical Geography	60
Knapp's Chemical Technology	63	Stanley on the Bones	37
Lallemand on Spermatorrhœa	18	Stille's General and Special Therapeutics	49
Lardner's Natural Philosophy and Astronomy	58	Taylor's Medical Jurisprudence	55
La Roche on Pneumonia, &c.	22	Taylor on Poisons	55
Lawrence on the Eye, by Hays	36	Thomson on the Sick Room	20
Lawrence on Ruptures	36	Todd and Bowman's Physiological Anatomy	17
Lee's Clinical Midwifery	42	Tomes's Dental Practice	37
Leidy and Gluge's Pathological Histology	21	Walshe on the Lungs and Heart	20
Leidy's Sharpey and Quain's Anatomy	9	Watson's Practice of Physic	20
Liston and Mutter's Operative Surgery	37	Weisbach's Mechanics and Engineering	62
Longet's Physiology, by Smith	15	West on Diseases of Children	44
Maclise's Surgical Anatomy	25	West on Ulceration of Os Uteri	5
Malgaigne's Operative Surgery, by Brittan	32	What to Observe in Medical Cases	19
Markwick's Examination of Urine	18	Whitehead on Abortion and Sterility	42
Matteucci on Living Beings	15	Wilde on Diseases of the Ear	34
Mayne's Dispensatory and Formulary	51	Williams's Principles of Medicine	18
Medical News and Library	3	Williams on Respiratory Organs	18
Miller's Practice of Surgery	27	Wilson's Human Anatomy	8
Miller's Principles of Surgery	28	Wilson's Dissector	9
Meigs's Obstetrics, the Science and the Art	40	Wilson on Diseases of the Skin	23
Meigs's Letters on Diseases of Women	41	Wilson on Healthy Skin	23
		Wilson on Syphilis	23
		Youatt and Skinner on the Horse	64
		Youatt on the Dog	64

THE AMERICAN MEDICAL JOURNAL.

TWO MEDICAL PERIODICALS, FREE OF POSTAGE,
FOR FIVE DOLLARS PER ANNUM.

THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES, subject to postage,
when not paid for in advance - - - - - \$5 00

THE MEDICAL NEWS AND LIBRARY, invariably in advance - - - - - 1 00
OR, BOTH PERIODICALS furnished, FREE OF POSTAGE, for Five Dollars remitted in
advance.

THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES,

EDITED BY ISAAC HAYS, M. D.,

is published Quarterly, on the first of January, April, July, and October. Each number contains at least two hundred and eighty large octavo pages, appropriately illustrated, wherever necessary, by engravings on copper, stone, or wood. It has now been issued regularly for a period of THIRTY-FIVE years, during a quarter of a century of which it has been under the control of the present editor. Throughout this long space of time, it has maintained its position in the highest rank of medical periodicals both at home and abroad, and has received the cordial support of the entire profession in this country. Its list of Collaborators will be found to contain a large number of the most distinguished names of the profession in every section of the United States, rendering the department devoted to

ORIGINAL COMMUNICATIONS

full of varied and important matter, of great interest to all practitioners.

As the aim of the Journal, however, is to combine the advantages presented by all the different varieties of periodicals, in its

REVIEW DEPARTMENT

will be found extended and impartial reviews of all important new works, presenting subjects of novelty and interest, together with very numerous

BIBLIOGRAPHICAL NOTICES,

including nearly all the medical publications of the day, both in this country and Great Britain, with a choice selection of the more important continental works. This is followed by the

QUARTERLY SUMMARY,

being a very full and complete abstract, methodically arranged, of the

IMPROVEMENTS AND DISCOVERIES IN THE MEDICAL SCIENCES.

This department of the Journal, so important to the practising physician, is the object of especial care on the part of the editor. It is classified and arranged under different heads, thus facilitating the researches of the reader in pursuit of particular subjects, and will be found to present a very full and accurate digest of all observations, discoveries, and inventions recorded in every branch of medical science. The very extensive arrangements of the publishers are such as to afford the editor complete materials for this purpose, as he not only regularly receives

ALL THE AMERICAN MEDICAL AND SCIENTIFIC PERIODICALS,

but also twenty or thirty of the more important Journals issued in Great Britain and on the Continent, thus enabling him to present in a convenient compass a thorough and complete abstract of everything interesting or important to the physician occurring in any part of the civilized world.

An evidence of the success which has attended these efforts may be found in the constant and steady increase in the subscription list, which renders it advisable for gentlemen desiring the Journal, to make known their wishes at an early day, in order to secure a year's set with certainty, the publishers having frequently been unable to supply copies when ordered late in the year. To their old subscribers, many of whom have been on their list for twenty or thirty years, the publishers feel that no promises are necessary; but those who may desire for the first time to subscribe, can rest assured that no exertion will be spared to maintain the Journal in the high position which it has occupied for so long a period.

By reference to the terms, it will be seen that in addition to this large amount of valuable and practical information on every branch of medical science, the subscriber, by paying in advance, becomes entitled, without further charge, to

THE MEDICAL NEWS AND LIBRARY,

a monthly periodical of thirty-two large octavo pages. Its "NEWS DEPARTMENT" presents the current information of the day, while the "LIBRARY DEPARTMENT" is devoted to present-

ing standard works on various branches of medicine. Within a few years, subscribers have thus received, without expense, the following works which have passed through its columns:—
 WATSON'S LECTURES ON THE PRACTICE OF PHYSIC;
 BRODIE'S CLINICAL LECTURES ON SURGERY;
 TODD AND BOWMAN'S PHYSIOLOGICAL ANATOMY AND PHYSIOLOGY OF MAN,
 724 pages, with numerous wood-cuts, being all that has yet appeared in England;
 WEST'S LECTURES ON THE DISEASES OF INFANCY AND CHILDHOOD;
 MALGAIGNE'S OPERATIVE SURGERY, with wood-cuts, and
 SIMON'S LECTURES ON GENERAL PATHOLOGY.

While the year 1854, will contain

BENNETT ON PULMONARY TUBERCULOSIS,
 WITH HANDSOME ILLUSTRATIONS ON WOOD,

AND

WEST ON ULCERATION OF THE OS UTERI,

of which new and valuable works a more detailed account is subjoined.

THE

PATHOLOGY AND TREATMENT OF PULMONARY TUBERCULOSIS,

And on the Local Medication of Pharyngeal and Laryngeal Diseases,
 frequently mistaken for or associated with Phthisis.

BY J. HUGHES BENNETT, M. D.,

Professor of Clinical Medicine in the University of Edinburgh, &c.

With Handsome Illustrations on Wood.



Section of Lung, showing cicatrization of tubercular excavation.

To say that we have been pleased with the book would be too faint praise; we have been more than pleased, it has deeply interested us, and has elucidated some points hitherto obscure, and given us a pathology of Tuberculosis, on the whole, more satisfactory than any we have met with; and, although, as the author

remarks, it is probable that some of the details may require modification, it will be found to possess a general harmony with all known facts, while it suggests to the medical practitioner resources the advantages of which continued experience is daily rendering more evident.—*Medical Examiner*, April, 1854.

We commend the book to those who can procure it as the latest, the most concise, able, and positive information to be obtained upon the subject of pulmonary tuberculosis.—*Buffalo Medical Journal*, April, 1854.

How it may be most effectually carried into practice, our readers will learn from Dr. Bennett's pages, especially from the histories of the valuable and in-

teresting cases which he records. Indeed, if the author had only reported these cases he would have benefited his profession, and deserved our thanks. As it is, however, his whole volume is so replete with valuable matter, that we feel bound to recommend our readers, one and all, to peruse it.—*London Lancet*.

The elegant little treatise before us shows how faithfully and intelligently these investigations have been pursued, and how successfully the author's studies have resulted in clearing up some of the most doubtful points and conflicting doctrines hitherto entertained in reference to the history and treatment of pulmonary tuberculosis.—*N. Y. Journal of Medical and Collateral Science*, March, 1854.

AN ENQUIRY INTO THE PATHOLOGICAL IMPORTANCE OF ULCERATION OF THE OS UTERI.

Being the Croonian Lectures for the year 1854.

BY CHARLES WEST, M. D.,

Author of "Lectures on the Diseases of Children," &c. &c.

The great experience and reputation of the author, and the importance of the subject, cannot fail to attract much attention to this work. Although just issued, it has already met with the general commendation of the English medical press.

Advance-paying subscribers to the Journal for 1854, will therefore receive, without charge, these two valuable works, and those who continue their subscriptions may rely that no relaxation will occur in the efforts to provide them with a succession of works of equally high character.

It will thus be seen that for the small sum of FIVE DOLLARS, paid in advance, the subscriber will obtain a Quarterly and a Monthly periodical,

EMBRACING ABOUT FIFTEEN HUNDRED LARGE OCTAVO PAGES.

mailed to any part of the United States, free of postage.

These very favorable terms are now presented by the publishers with the view of removing all difficulties and objections to a full and extended circulation of the Medical Journal to the office of every member of the profession throughout the United States. The rapid extension of mail facilities, will now place the numbers before subscribers with a certainty and dispatch not heretofore attainable; while by the system now proposed, every subscriber throughout the Union is placed upon an equal footing, at the very reasonable price of Five Dollars for two periodicals, without further expense.

Those subscribers who do not pay in advance will bear in mind that their subscription of Five Dollars will entitle them to the Journal only, without the News, and that they will be at the expense of their own postage on the receipt of each number. The advantage of a remittance when ordering the Journal will be thus apparent.

As the Medical News and Library is in no case sent without advance payment, its subscribers will always receive it free of postage.

It should also be borne in mind that the publishers will now take the risk of remittances by mail, only requiring, in cases of loss, a certificate from the subscriber's Postmaster, that the money was duly mailed and forwarded.

☞ Funds at par at the subscriber's place of residence received in payment of subscriptions.

Address, BLANCHARD & LEA, PHILADELPHIA.

Subscribers for 1853, who do not possess the commencement of Todd and Bowman's Physiology, can obtain it, in a handsome octavo volume, of 552 pages, with over 150 illustrations, by mail, free of postage, on a remittance of \$2 50 to the publishers.

HOBLYN'S MEDICAL DICTIONARY—(New Edition, Nearly Ready.)

A DICTIONARY OF THE TERMS USED IN MEDICINE, AND THE COLLATERAL SCIENCES.

BY RICHARD D. HOBLYN, A. M.

SECOND AND IMPROVED AMERICAN EDITION, REVISED, WITH ADDITIONS,

BY ISAAC HAYS, M. D.

In one large royal 12mo. volume, of over four hundred double-columned pages.

In passing this work a second time through the press, the editor has subjected it to a very thorough revision, making such additions as the progress of science has rendered desirable, and supplying any omissions that may have previously existed. As a concise and convenient Dictionary of Medical Terms, at an exceedingly low price, it will therefore be found of great value to the student and practitioner.

THE AMERICAN MEDICAL DICTIONARY—(New Edition, Just Issued.)

MEDICAL LEXICON;

A DICTIONARY OF MEDICAL SCIENCE,

CONTAINING

A Concise Explanation of the Various Subjects and Terms of

PHYSIOLOGY, PATHOLOGY, HYGIENE, THERAPEUTICS, PHARMACOLOGY,
OBSTETRICS, MEDICAL JURISPRUDENCE, &c.

WITH THE FRENCH AND OTHER SYNONYMES;

NOTICES OF CLIMATE AND OF CELEBRATED MINERAL WATERS;

FORMULÆ FOR VARIOUS OFFICINAL, EMPIRICAL, AND DIETETIC PREPARATIONS, ETC.

BY ROBLEY DUNGLISON, M. D.,

Professor of Institutes of Medicine in Jefferson Medical College, &c. &c.

Eleventh Edition, Revised.

In one large and handsome octavo volume, of over nine hundred double-columned pages, strongly bound in leather, with raised bands.

Every successive edition of this work bears the marks of the industry of the author, and of his determination to keep it fully on a level with the most advanced state of medical science. Thus nearly FIFTEEN THOUSAND WORDS have been added to it within the last few years. As a complete Medical Dictionary, therefore, embracing over FIFTY THOUSAND DEFINITIONS, in all the branches of the science, it is presented as meriting a continuance of the great favor and popularity which have carried it, within no very long space of time, to an eleventh edition.

Every precaution has been taken in the preparation of the present volume, to render its mechanical execution and typographical accuracy worthy of its extended reputation and universal use. The very extensive additions have been accommodated, without materially increasing the bulk of the volume by the employment of a small but exceedingly clear type, cast for this purpose. The press has been watched with great care, and every effort used to insure the verbal accuracy so necessary to a work of this nature. The whole is printed on fine white paper; and, while thus exhibiting in every respect so great an improvement over former issues, it is presented at the original exceedingly low price.

We welcome it cordially; it is an admirable work, and indispensable to all literary medical men. The labor which has been bestowed upon it is something prodigious. The work, however, has now been done, and we are happy in the thought that no human being will have again to undertake the same gigantic task. Revised and corrected from time to time, Dr. Dunglison's "Medical Lexicon" will last for centuries.—*British and Foreign Med. Chirurg. Review.*

A miracle of labor and industry in one who has written able and voluminous works on nearly every branch of medical science. There could be no more useful book to the student or practitioner, in the present advancing age, than one in which would be found, in addition to the ordinary meaning and derivation of medical terms—so many of which are of modern introduction—concise descriptions of their explanation and employment; and all this and much more is contained in the volume before us. It is therefore almost as indispensable to the other learned professions as to our own. In fact, to all who may have occasion to ascertain the meaning of any word belonging to the many branches of medicine. From a careful examination of the present edition, we can vouch for its accuracy, and for its being brought quite up to the date of publication; the author states in his preface that he has added to it about four thousand terms, which are not to be found in the preceding one.—*Dublin Quarterly Journal of Medical Sciences.*

One of the most complete and copious known to the cultivators of medical science.—*Boston Med. Journal.*

The most comprehensive and best English dictionary of medical terms extant.—*Buffalo Medical Journal.*

Whilst it is a wonderful monument of its author's erudition and industry, it is also a work of great practical utility, as we can testify from our own experience; for we keep it constantly within our reach, and make very frequent reference to it, nearly always finding in it the information we seek.—*British and Foreign Med. Chirurg. Review.*

It has the rare merit that it certainly has no rival in the English language for accuracy and extent of references. The terms generally include short physiological and pathological descriptions, so that, as the author justly observes, the reader does not possess in this work a mere dictionary, but a book, which, while it instructs him in medical etymology, furnishes him with a large amount of useful information. No medical library will be complete without it.—*London Med. Gazette.*

It is certainly more complete and comprehensive than any with which we are acquainted in the English language. Few, in fact, could be found better qualified than Dr. Dunglison for the production of such a work. Learned, industrious, persevering, and accurate, he brings to the task all the peculiar talents necessary for its successful performance: while, at the same time, his familiarity with the writings of the ancient and modern "masters of our art," renders him skilful to note the exact usage of the several terms of science, and the various modifications which medical terminology has undergone with the change of theories or the progress of improvement.—*American Journal of the Medical Sciences.*

The Student's Manual.

AN ANALYTICAL COMPENDIUM
OF THE VARIOUS BRANCHES OF MEDICAL SCIENCE.
For the Use and Examination of Students.

BY

JOHN NEILL, M. D.,
Surgeon to the Pennsylvania
Hospital, Professor of
Surgery in
Penna. Medical College, &c.,

AND

F. G. SMITH, M. D.,
Professor of Institutes of Medi-
cine in the Pennsylvania
Medical College.

SECOND EDITION, REVISED AND
IMPROVED.

In one very
large royal 12mo. volume
of 1000
pages, beautifully printed, with
240 engravings
on wood.



Application of the Vectis.

This work comprehends the whole range of medical science, presenting the leading and important facts of each department in a clear, methodical, and concise manner. Embracing as it does, the latest discoveries and developments in each subject, it is admirably suited for the student, as an introduction to the course of reading, or as a remembrancer at the close of his studies, and also to the practitioner who may desire to keep himself on a level with the progress of science, while his leisure may not admit of the perusal of extended treatises on special subjects. It is divided as follows:—

ANATOMY, 196 pages, 159 illustrations.
PHYSIOLOGY, 144 pages, 43 illustrations.
SURGERY, 134 pages, 50 illustrations.
OBSTETRICS, 124 pages, 41 illustrations.

MATERIA MEDICA AND THERAPEUTICS,
130 pages, 29 illustrations.
CHEMISTRY, 106 pages, 19 illustrations.
PRACTICE OF MEDICINE, 172 pages.

The speedy sale of a large impression of this work has afforded to the authors gratifying evidence of the correctness of the views which actuated them in its preparation. In meeting the demand for a second edition, they have therefore been desirous to render it more worthy of the favor with which it has been received. To accomplish this, they have spared neither time nor labor in embodying in it such discoveries and improvements as have been made since its first appearance, and such alterations as have been suggested by its practical use in the class and examination-room. Considerable modifications have thus been introduced throughout all the departments treated of in the volume, but more especially in the portion devoted to the "Practice of Medicine," which has been entirely rearranged and rewritten. The authors therefore again submit their work to the profession, with the hope that their efforts may tend, however humbly, to advance the great cause of medical education.

Notwithstanding the enlarged size and improved execution of this work, the price has not been increased, and it is confidently presented as one of the cheapest volumes now before the profession.

In the rapid course of lectures, where work for the students is heavy, and review necessary for an examination, a compend is not only valuable, but it is almost a *sine qua non*. The one before us is, in most of the divisions, the most unexceptionable of all books of the kind that we know of. The newest and soundest doctrines and the latest improvements and discoveries are explicitly, though concisely, laid before the student. Of course, it is useless for us to recommend it to all last course students, but there is a class to whom we very sincerely commend this cheap book as worth its weight in silver—that class is the graduates in medicine of more than ten years' standing, who have not studied medicine since. They will perhaps find out from it that the science is not exactly now what it was when they left it off.—*The Stethoscope*.

Having made free use of this volume in our examinations of pupils, we can speak from experience in recommending it as an admirable compend for students, and as especially useful to preceptors who examine their pupils. It will save the teacher much labor by enabling him readily to recall all of the points upon which his pupils should be examined. A work of this sort should be in the hands of every one who takes pupils into his office with a view of examining them; and this is unquestionably the best of its class. Let every practitioner who has pupils provide himself with it, and he will find the labor of refreshing his knowledge so much facilitated that he will be able to do justice to his pupils at very little cost of time or trouble to himself.—*Transylvania Medical Journal*.

A SYSTEM OF HUMAN ANATOMY,

GENERAL AND SPECIAL.

BY ERASMUS WILSON, M. D.,
Lecturer on Anatomy, London.Fourth American from a late
London Edition.

EDITED BY P. B. GODDARD, M. D.

With
two hundred and fifty-one
handsome
illustrations on wood.
In one
beautifully printed octavo volume
of nearly
six hundred pages.

In many, if not all the Colleges of the Union, it has become a standard text-book. This, of itself, is sufficiently expressive of its value. A work very desirable to the student; one, the possession of which will greatly facilitate his progress in the study of Practical Anatomy.—*New York Journal of Medicine.*

Its author ranks with the highest on Anatomy.—*Southern Medical and Surgical Journal.*

It offers to the student all the assistance that can be expected from such a work.—*Med. Examiner.*

The most complete and convenient manual for the student we possess.—*American Journal of Medical Science.*

This is probably the prettiest medical book ever published, and we believe that its intrinsic merits are in keeping with its exterior advantages; it may be recommended to the student as no less distinguished by its accuracy and clearness of description than its typographical elegance. The woodcuts are exquisite.—*British and Foreign Medical Review.*

As a satisfactory proof that the praise we bestowed on the first edition of this work was not unmerited, we may observe it has been equally well thought of in foreign countries, having been reprinted in the United States and in Germany. In every respect, this work, as an anatomical guide for the student and practitioner, merits our warmest and most decided praise.—*London Med. Gazette.*



The Great Sympathetic Nerve.

A NEW TEXT-BOOK ON HISTOLOGY—(In Press.)

HUMAN HISTOLOGY,

IN ITS APPLICATIONS TO PHYSIOLOGY AND GENERAL PATHOLOGY.

Designed as a Text-book for Medical Students.

BY E. R. PEASELEE, M. D.,

Professor of Anatomy and Physiology in Dartmouth Medical College.

In one handsome volume, royal 12mo., with numerous illustrations.

The subject of this work is one, the growing importance of which, as the basis of Anatomy and Physiology, demands for it a separate volume. The book will therefore supply an acknowledged deficiency in medical text-books, while the name of the author, and his experience as a teacher for the last thirteen years, is a guarantee that it will be thoroughly adapted to the use of the student


HUMAN ANATOMY.

BY JONES QUAIN, M. D.

EDITED BY

RICHARD QUAIN, F. R. S., AND WILLIAM SHARPEY, M. D., F. R. S.,
Professors of Anatomy and Physiology in University College, London.

From the fifth London Edition.



Heart and Great Vessels.

It is indeed a work calculated to make an era in anatomical study, by placing before the student every department of his science, with a view to the relative importance of each; and so skilfully have the different parts been interwoven, that no one who makes this work the basis of his studies, will hereafter have any excuse for neglecting or undervaluing any important particulars connected with the structure of the human frame; and whether the bias of his mind lead him in a more especial manner to surgery, physic, or physiology, he will find here a work at once so comprehensive and practical as to defend him from exclusiveness on the one hand, and pedantry on the other.—*Monthly Journal and Retrospect of the Med. Sciences.*

We have no hesitation in recommending this treatise on anatomy as the most complete on that subject in the English language; and the only one, perhaps, in any language, which brings the state of knowledge forward to the most recent discoveries.—*The Edinburgh Med. and Surg. Journal.*

Admirably calculated to fulfil the object for which it is intended.—*Provincial Medical Journal.*

The most complete Treatise on Anatomy in the English language.—*Edinburgh Medical Journal.*

There is no work in the English language to be preferred to Dr. Quain's Elements of Anatomy.—*Lond. Journal of Medicine.*

EDITED BY JOSEPH LEIDY, M. D.,

Professor of Anatomy in the University of Pennsylvania, &c.

In two large and handsome octavo volumes, of thirteen hundred pages, beautifully illustrated with over five hundred engravings on wood.

THE DISSECTOR;


OR, PRACTICAL AND SURGICAL ANATOMY.

BY ERASMUS WILSON, M. D.,

Author of "A System of Human Anatomy."

EDITED BY P. B. GODDARD, M. D.

A new and improved edition, with one hundred and fifty handsome illustrations. In one large and handsome royal 12mo. volume.



Muscles of the Eye ball.

In passing this work again through the press, the editor has made such additions and improvements as the advance of anatomical knowledge has rendered necessary to maintain the work in the high reputation which it has acquired in the schools of the United States, as a complete and faithful guide to the student of practical anatomy.

A number of new illustrations have been added, especially in the portion relating to the complicated anatomy of Hernia. In mechanical execution, the work will be found superior to former editions.

ANATOMICAL ATLAS,

ILLUSTRATIVE OF THE STRUCTURE OF THE HUMAN BODY.

BY HENRY H. SMITH, M. D.

UNDER THE SUPERVISION OF
WILLIAM E. HORNER, M. D.

Professor of Anatomy in the University of Pennsylvania, &c.

In one imperial octavo volume, extra cloth, with about six hundred and fifty beautifully executed illustrations on wood.



Epidermis and Cutis Vera magnified.

The great advantages possessed by wood-engraving for scientific illustration, in the clearness and distinctness of its minute details, are fully shown in this work, containing as it does, in the compass of a single convenient volume, the general and special anatomy of all the component parts of the body. Commencing with the Bones and Ligaments, it follows with the Muscular and Dermoid Systems, the Organs of Digestion and Generation, Respiration and Circulation, and concludes with the Nervous System and the Senses. Not only is every separate organ and portion of the human frame thus brought distinctly and separately before the eye of the student, but he is also presented with the results of the recent microscopical investigations into the minute anatomy of the various tissues; while the plan adopted, giving the plate and references on the same page, enables him to obtain a more definite impression of the objects, by avoiding the annoyance and interruption of turning backwards and forwards. As a specimen of art, nothing superior to it has yet been produced, while the exceedingly low rate at which it is offered, places it within the reach of every member of the profession. To country practitioners and students it will be found especially useful, as supplying in a great measure the place of skeletons and subjects.

These numbers complete the series of this beautiful work, which fully merits the praise bestowed upon the earlier numbers. We regard all the engravings as possessing an accuracy only equalled by their beauty, and cordially recommend the work to all engaged in the study of anatomy.—*N. Y. Journal of Medicine and Surgery.*

A more elegant work than the one before us could not easily be placed by a physician upon the table of his student.—*Western Journal of Medicine and Surgery.*

This is an exquisite volume, and a beautiful specimen of art. We have numerous Anatomical Atlases, but we will venture to say that none equal it in cheap-



Muscles of the Back.

ness, and none surpass it in faithfulness and spirit. We strongly recommend to our friends, both urban and suburban, the purchase of this excellent work, for which both editor and publisher deserve the thanks of the profession.—*Medical Examiner.*

The plan of this Atlas is admirable, and its execution superior to anything of the kind before published in this country. It is a real labor-saving affair, and we regard its publication as the greatest boon that could be conferred on the student of anatomy. It will be equally valuable to the practitioner, by affording him an easy means of recalling the details learned in the dissecting-room, and which are soon forgotten.—*American Medical Journal.*



Arteries of the Chest and Neck.

In
two large
and handsome
octavo
volumes of over
one thousand
pages,
with more
than
three hundred
beautiful
illustrations.

SPECIAL ANATOMY AND HISTOLOGY.

BY WILLIAM E. HORNER, M. D.,
Professor of Anatomy, University of Pennsylvania, &c. &c.
Eighth and Improved Edition.

This work has enjoyed a thorough and laborious revision on the part of the author, with the view of bringing it fully up to the existing state of knowledge on the subject of general and special anatomy. To adapt it more perfectly to the wants of the student, he has introduced a large number of additional wood-engravings, illustrative of the objects described, while the publishers have endeavored to render the mechanical execution of the work worthy of the extended reputation which it has acquired. The demand which has carried it to an EIGHTH EDITION is a sufficient evidence of the value of the work, and of its adaptation to the wants of the student and professional reader.

(PREPARING.)

MEDICAL ANATOMY,

Illustrating the Form, Structure, and Position of the Internal Organs in Health and Disease. By FRANCIS SIBSON, M. D., Physician to St. Mary's Hospital. In large imperial quarto, with splendid colored plates, to match "Maclise's Surgical Anatomy."

A NEW WORK—(Now Ready.)

PRINCIPAL FORMS OF THE SKELETON, AND STRUCTURE OF THE TEETH.

BY PROFESSOR R. OWEN.

In one neat royal 12mo. volume, with numerous illustrations.



Skeleton of the Crocodile.

This little work, embodying the latest and most correct views on comparative osteology, and presenting a complete outline of the subject, cannot fail to prove of great interest to all students of comparative anatomy, to whom the name of its distinguished author is sufficient guarantee for all the facts and opinions advanced.

New and much Enlarged Edition.

PRINCIPLES OF HUMAN PHYSIOLOGY,

WITH THEIR CHIEF APPLICATIONS TO

PSYCHOLOGY, PATHOLOGY, THERAPEUTICS, HYGIENE, AND FORENSIC MEDICINE.

BY WILLIAM B. CARPENTER, M. D., F. R. S.,

Examiner in Physiology and Comparative Anatomy in the University of London.

FIFTH AMERICAN FROM THE FOURTH AND ENLARGED ENGLISH EDITION.

EDITED, WITH ADDITIONS, BY FRANCIS GURNEY SMITH, M. D.,

Professor of the Institutes of Medicine in the Pennsylvania Medical College, &c.

With 314 beautiful illustrations.

In one very large and handsome octavo volume, of eleven hundred pages, strongly bound in leather, with raised bands.



Transverse Section of Medulla Oblongata, magnified.

This edition has been printed from sheets prepared for the purpose by the author, who has introduced nearly one hundred illustrations not in the London edition; while it has also enjoyed the advantage of a careful superintendence on the part of the editor, who has added notices of such more recent investigations as had escaped the author's attention. Neither care nor expense has been spared in the mechanical execution of the work, to render it superior to former editions, and it is confidently presented as in every way one of the handsomest volumes as yet placed before the medical profession in this country.

The most complete work on the science in our language.—*American Medical Journal*.

A complete cyclopaedia of this branch of science.—*N. Y. Med. Times*.

We have thus adverted to some of the leading "additions and alterations" which have been introduced by the author into this edition of his physiology. These will be found, however, very far to exceed the ordinary limits of a new edition, "the old materials having been incorporated with the new, rather than the new with the old." It now certainly presents the most complete treatise on the subject within the reach of the American reader.—*Medical Examiner*.

The most complete work now extant in our language.—*N. O. Med. Register*.

The most complete exposition of physiology which any language can at present give.—*British and For. Medico-Chirurgical Review*.

The greatest, the most reliable, and the best book on the subject which we know of in the English language.—*Stethoscope*.

The best text-book in the language on this extensive subject.—*London Med. Times*.

The standard of authority on physiological subjects. * * * In the present edition, to particularize the alterations and additions which have been made, would require a review of the whole work, since scarcely a subject has not been revised and altered, added to, or entirely remodelled.—*Charleston Med. Journal*.

AN ESSAY TOWARDS A CORRECT THEORY OF THE NERVOUS SYSTEM. By JOHN HARRISON, M. D. In one octavo volume, 292 pages.

Second
American
from a new
and
Revised
London
edition.



Vessels of Dental Papilla.

With
one hundred
and ninety
illustrations.

In one
very
handsome
octavo
volume of
over 500
pages.

ELEMENTS OF PHYSIOLOGY,

INCLUDING PHYSIOLOGICAL ANATOMY.

BY WILLIAM B. CARPENTER, M. D., F. R. S.,

Author of "Principles of Human Physiology."

To say that it is the best manual of Physiology now before the public, would not do sufficient justice to the author.—*Buffalo Medical Journal.*

In his former works, it would seem that he had exhausted the subject of Physiology. In the present, he gives the essence, as it were, of the whole.—*N. Y. Journal of Medicine.*

Those who have occasion for an elementary treatise on Physiology, cannot do better than to possess themselves of the manual of Dr. Carpenter.—*Med Examiner.*

The best and most complete exposé of modern Physiology, in one volume, extant in the English language.—*St. Louis Medical Journal.*

With such an aid in his hand, there is no excuse for the ignorance often displayed respecting the subjects of which it treats. From its unpretending dimensions, it may not be so esteemed by those anxious to make a parade of their erudition; but whoever masters its contents will have reason to be proud of his physiological acquirements. The illustrations are well selected and finely executed.—*Dublin Med. Press.*

A NEW WORK BY DR. CARPENTER.

Preparing.

THE MICROSCOPE

AND ITS

REVELATIONS.

In one very handsome volume, with numerous beautiful illustrations.

The importance of the Microscope as a means of investigation and diagnosis, and its constantly increasing practical applications, render exceedingly desirable some accurate manual to guide the student in its use. Few could be found more competent than Dr. Carpenter to supply this desideratum. The present work may therefore be confidently expected to at once assume the position of the standard text-book on the subject.



Thoroughly Revised Edition—(Now Ready.)

In one
large and
handsome
octavo
volume, of
seven
hundred and
fifty pages.



With
over three
hundred
exquisite
illustrations.

Paludina Vivipara laid open.

PRINCIPLES OF COMPARATIVE PHYSIOLOGY.

INTENDED AS AN INTRODUCTION TO THE STUDY OF HUMAN PHYSIOLOGY, AND
AS A GUIDE TO THE PHILOSOPHICAL PURSUIT OF NATURAL HISTORY.

BY WILLIAM B. CARPENTER, M. D., F. R. S.,

Author of "Principles of Human Physiology," &c.

New and Much Improved Edition.

The delay which has occurred in the appearance of this edition, has arisen from the very thorough revision and remodelling which it has undergone at the hands of the author, who has spared no pains to render it a complete and trustworthy exposition of the present advanced condition of Comparative Physiology. A very large number of new illustrations have been introduced, and the publishers confidently present it as worthy of its exalted reputation in every point of mechanical and artistic finish.

*Section of Sphynx Ligustri.*

A FEW NOTICES OF THE FORMER EDITION ARE APPENDED.

Without pretending to it, it is an Encyclopedia of the subject, accurate and complete in all respects—a truthful reflection of the advanced state at which the science has now arrived.—*Dublin Quarterly Journal of Medical Science.*

A truly magnificent work—in itself a perfect physiological study.—*Ranking's Abstract.*

This work stands without its fellow. It is one few men in Europe could have undertaken; it is one no man, we believe, could have brought to so successful an issue as Dr. Carpenter. It required for its production a physiologist at once deeply read in the labors of others, capable of taking a general, critical, and

unprejudiced view of those labors, and of combining the varied, heterogeneous materials at his disposal, so as to form an harmonious whole. We feel that this abstract can give the reader a very imperfect idea of the fulness of this work, and no idea of its unity, of the admirable manner in which material has been brought, from the most various sources, to conduce to its completeness, of the lucidity of the reasoning it contains, or of the clearness of language in which the whole is clothed. Not the profession only, but the scientific world at large, must feel deeply indebted to Dr. Carpenter for this great work. It must, indeed, add largely even to his high reputation.—*Medical Times.*



Interior of the Larynx.



Erectile Tissue of the Corpus Cavernosum.

Seventh Edition.

Thoroughly Revised and extensively
Modified and Enlarged, with nearly Five Hundred
Illustrations.

HUMAN PHYSIOLOGY.

BY ROBLEY DUNGLISON, M. D.,

Professor of the Institutes of Medicine in Jefferson Medical College, Philadelphia.

In two large and handsomely printed octavo volumes, containing nearly 1450 pages.

The merits of this work as a text-book for the student, and a work of reference for the professional man, are too well known to require exposition. The fact of its having passed to a SEVENTH EDITION is sufficient evidence of its appreciation by those best able to form an opinion; and this appreciation the author has endeavored to merit by unsparing labor in its improvement as it has successively passed through his hands. On no previous edition, however, has this revision been so thorough as on the present. Many portions have been almost rewritten, and the numerous discoveries and investigations made since the previous impression, have been carefully noted and inserted. It can therefore be recommended as completely on a level with the existing state of science. The very numerous illustrations with which it abounds have been thoroughly revised, a large number have been added, and many new and superior engravings have been substituted for less satisfactory ones. The mechanical execution has also been materially improved, and the work is confidently presented as eminently worthy a continuance of the very remarkable favor with which it has been received.

It has long since taken rank as one of the medical classics of our language. To say that it is by far the best text-book of physiology ever published in this country, is but echoing the general testimony of the profession.—*N. Y. Journal of Medicine.*

There is no single book we would recommend to the student or physician, with greater confidence than the present, because in it, will be found a mirror of almost every standard physiological work of the day. We most cordially recommend the work to every member of the profession, and no student should be without

it. It is the completest work on Physiology in the English language, and is highly creditable to the author and publishers.—*Canadian Medical Journal.*

The most complete and satisfactory system of Physiology in the English language.—*Amer. Med. Journal.*

The best work of the kind in the English language.—*Silliman's Journal.*

The most full and complete system of Physiology in our language.—*Western Lancet.*

A NEW WORK ON PHYSIOLOGY—(Preparing.)

A TREATISE ON PHYSIOLOGY. By F. A. LONGET. Translated from the French by FRANCIS G. SMITH, M. D., Professor of Institutes of Medicine in the Pennsylvania Medical College.

LECTURES ON THE PHYSICAL PHENOMENA OF HUMAN BEINGS. By CARLO MATTEUCCI. Edited by JONATHAN PEREIRA, M. D. In one neat volume, royal 12mo.

Second
American
from
the Second and
Improved
London Edition.



Capillary Bloodvessels and Lymphatics.

With
one hundred and
sixty-five
illustrations.
—
In one
large and handsome
royal 12mo.
volume
of
five hundred and
fifty pages.

A MANUAL OF PHYSIOLOGY.

BY WILLIAM SENHOUSE KIRKES, M. D.,
Demonstrator of Morbid Anatomy at St. Bartholomew's Hospital, &c.

AND

JAMES PAGET, F. R. S.,
Author of "Lectures on Surgical Pathology," &c.

In the present edition, the Manual of Physiology has been brought up to the actual condition of the science, and fully sustains the reputation which it has already so deservedly attained. We consider the work of MM. Kirkes and Paget to constitute one of the very best handbooks of Physiology we possess—presenting just such an outline of the science, comprising an account of its leading facts and generally admitted principles, as the student requires during his attendance upon a course of lectures, or for reference whilst preparing for examination. The text is fully and ably illustrated by a series of very superior wood-engravings, by which a comprehension of some

of the more intricate of the subjects treated of is greatly facilitated.—*American Journal of Med. Science.*

We need only say, that, without entering into discussions of unsettled questions, it contains all the recent improvements in this department of medical science. For the student beginning this study, and the practitioner who has but leisure to refresh his memory, this book is invaluable, as it contains all that it is important to know, without special details, which are read with interest only by those who would make a specialty, or desire to possess a critical knowledge of the subject.—*Charleston Medical Journal.*

POPULAR PHYSIOLOGY—(Now Ready.)

THE

PHYSIOLOGY OF ANIMAL AND VEGETABLE LIFE.


In one
handsome
royal 12mo.
volume of about
two hundred
pages.



With
one hundred
illustrations on
wood.

Head and Eyes of Bee

Though designed for general readers, this little work will be found of high scientific interest, as it presents the latest views and investigations in the various branches of physiological inquiry. Written in a popular and intelligible manner, it possesses great recommendations for the student commencing this subject, as well as for District Libraries and Academies, for which purposes it is more expressly designed.



Section of Lung, magnified.

THE PHYSIOLOGICAL ANATOMY AND PHYSIOLOGY OF MAN.

BY ROBERT B. TODD, M.D., F. R. S., AND WILLIAM BOWMAN, F. R. S.

With several Hundred Beautiful Illustrations on Wood.

PARTS I, II, III, lately issued in one octavo volume of 552 pages, with 156 illustrations. PART IV, completing the work, was commenced in the Medical News for 1853. The conclusion of it may be shortly expected in London, when it will immediately be issued here, in a form enabling those who possess the first portion to complete their sets. Subscribers for 1853, who have not got the first portion, can obtain it free by mail on remitting \$2 50 to the publishers.

It cannot fail to recommend itself to students by the clearness of description and the accuracy of the illustrations met with in every chapter; and we do not hesitate to express our opinion that it supplies a void which has hitherto existed between works especially devoted to Anatomy and Physiology.—*London Medical Gazette.*

We must urgently recommend those among our readers who are not already in possession of this Physiological Anatomy, no longer to delay adding it to their library. * * * This treatise is quite indispensable to all who aspire to keep up their knowledge with the high standard of the present day.—*Medico-Chirurgical Review.*

THE HUMAN BRAIN; ITS STRUCTURE, PHYSIOLOGY, AND DISEASES.

WITH A DESCRIPTION OF THE TYPICAL FORMS OF THE BRAIN IN THE ANIMAL KINGDOM.

BY SAMUEL SOLLY, F. R. S.

FROM THE SECOND LONDON EDITION.

In one handsome octavo volume of 500 pages, with 118 illustrations.



Longitudinal Section of the Pons Varolii and Crus Cerebri.

NEW AND IMPROVED EDITION—(Now Ready.)

PRINCIPLES OF MEDICINE;
 COMPRISING
GENERAL PATHOLOGY AND THERAPEUTICS,
 AND A BRIEF GENERAL VIEW OF
ETIOLOGY, NOSOLOGY, DIAGNOSIS, PROGNOSIS, AND HYGIENICS,

BY C. J. B. WILLIAMS, M. D., F. R. S.,

Professor of Clinical Medicine in University College, London.

EDITED, WITH ADDITIONS, BY MEREDITH CLYMER, M. D.

Fourth American Edition, Revised.

In one octavo volume, of about five hundred pages.

This edition has been thoroughly revised and brought up by the editor.

FEVERS, THEIR DIAGNOSIS, PATHOLOGY, AND TREATMENT. Prepared and edited, with large additions, from the Essays on Fever in Tweedie's Library of Practical Medicine. By Meredith Clymer, M. D. In one octavo volume, of six hundred pages.

A PRACTICAL TREATISE ON DISEASES OF THE RESPIRATORY ORGANS: including Diseases of the Larynx, Trachea, Lungs, and Pleuræ. By C. J. B. Williams, M. D. Edited, with numerous Additions and Notes, by Meredith Clymer, M. D. With woodcuts. In one octavo volume, of five hundred pages.

NEARLY READY.

A MANUAL OF PATHOLOGICAL ANATOMY.



Aneurism of the Left Ventricle.

WITH NUMEROUS ENGRAVINGS ON WOOD.

BY C. HANDFIELD JONES, M. D.

AND

EDWARD H. SIEVEKING, M. D.,

Assistant Physician to St. Mary's Hospital.

In one

handsome octavo volume,

beautifully illustrated.

GENERAL PATHOLOGY;

As Conducive to the Establishment of Rational Principles for the Prevention and Cure of Disease. A Course of Lectures delivered at St. Thomas's Hospital. By JOHN SIMON, F. R. S., &c. In one neat octavo volume.

THE CAUSES, SYMPTOMS, AND TREATMENT OF SPERMATORRHOEA. By M. LALLEMAND. Translated and Edited by HENRY J. McDOUGAL. Second American edition. In one octavo volume, of 326 pages. (Just Issued.)

ROKITANSKI'S PATHOLOGICAL ANATOMY.—Translated from the last German edition. (Preparing.)

BURROWS ON CEREBRAL CIRCULATION.—On Disorders of the Cerebral Circulation, and on the Connection between Affections of the Brain and of the Heart. In one 8vo. vol., colored plates, pp. 216.

BLAKISTON ON THE CHEST.—Practical Observations on Diseases of the Chest, and on the Principles of Auscultation. In one volume, 8vo., pp. 384.

BENEDICT'S CHAPMAN.—Compendium of Chapman's Lectures on the Practice of Medicine. One neat volume, 8vo., pp. 258.

HOPE ON THE HEART.—A Treatise on the Diseases of the Heart and Great Vessels. Edited by Pennock. In one volume, 8vo., with plates, pp. 572.

FRICK ON THE URINE.—Renal Affections, their Diagnosis and Pathology. In one handsome volume, royal 12mo., with illustrations.

CHAPMAN'S LECTURES on Fevers, Dropsy, Gout, Rheumatism, &c. In one 8vo. vol., pp. 450.

HASSE'S PATHOLOGICAL ANATOMY.—An Anatomical Description of the Diseases of Respiration and Circulation. In one volume, 8vo., pp. 379.

COPLAND ON PALSY.—Of the Causes, Nature, and Treatment of Palsy and Apoplexy. In one volume, royal 12mo.

PHILIPS ON SCROFULA.—Scrofula: its Nature, its Prevalence, its Causes, and the Principles of its Treatment. In one vol. 8vo., with a plate, pp. 350.

MANUALS ON THE BLOOD AND URINE.—By John William Griffith, M. D., G. Owen Reese, M. D., and Alfred Markwick. In one large 12mo. volume, extra cloth, of 460 pages, with numerous illustrations.

BILLING'S PRINCIPLES OF MEDICINE.—The Principles of Medicine. 2d American from the 5th London Edition. In one volume, 8vo., extra cloth, 260 pp.

THE
CYCLOPÆDIA OF PRACTICAL MEDICINE;

COMPRISING

Treatises on the Nature and Treatment of Diseases, Materia Medica and Therapeutics,
Diseases of Women and Children, Medical Jurisprudence, &c. &c.

EDITED BY

JOHN FORBES, M. D., F. R. S., ALEXANDER TWEEDIE, M. D., F. R. S., AND
JOHN CONNOLLY, M. D.

REVISED, WITH ADDITIONS,

BY ROBLEY DUNGLISON, M. D.

COMPLETE, IN FOUR LARGE SUPER-ROYAL OCTAVO VOLUMES,

Containing Thirty-two Hundred and Fifty-four unusually large Pages in Double Columns,
printed on good paper, with a new and clear type.

THE WHOLE WELL AND STRONGLY BOUND WITH RAISED BANDS AND DOUBLE TITLES.

This work contains no less than FOUR HUNDRED AND EIGHTEEN DISTINCT TREATISES,

By Sixty-Eight Distinguished Physicians.

The most complete work on Practical Medicine extant; or, at least, in our language.—*Buffalo Medical and Surgical Journal*.

For reference, it is above all price to every practitioner.—*Western Lancet*.

One of the most valuable medical publications of the day—as a work of reference it is invaluable.—*Western Journal of Medicine and Surgery*.

It has been to us, both as learner and teacher, a work for ready and frequent reference, one in which modern English medicine is exhibited in the most advantageous light.—*Medical Examiner*.

We rejoice that this work is to be placed within the reach of the profession in this country, it being un-

questionably one of very great value to the practitioner. This estimate of it has not been formed from a hasty examination, but after an intimate acquaintance derived from frequent consultation of it during the past nine or ten years. The editors are practitioners of established reputation, and the list of contributors embraces many of the most eminent professors and teachers of London, Edinburgh, Dublin, and Glasgow. It is, indeed, the great merit of this work that the principal articles have been furnished by practitioners who have not only devoted especial attention to the diseases about which they have written, but have also enjoyed opportunities for an extensive practical acquaintance with them, and whose reputation carries the assurance of their competency justly to appreciate the opinions of others, while it stamps their own doctrines with high and just authority.—*American Med. Journal*.

~~~~~  
NOW READY.

ON THE ETIOLOGY, PATHOLOGY, AND TREATMENT OF FIBRO-BRONCHITIS AND RHEUMATIC PNEUMONIA, By T. H. BUCKLER, M. D., formerly Physician to the Baltimore Almshouse Infirmary. In one neat volume, extra cloth.

~~~~~  
A NEW MANUAL OF PRACTICE—(Preparing.)

A MANUAL OF THE PRINCIPLES AND PRACTICE OF MEDICINE. By GEORGE H. BARLOW, M. D., Physician to Guy's Hospital, London. In one volume.

~~~~~  
WHAT TO OBSERVE

AT THE BEDSIDE AND AFTER DEATH,  
IN MEDICAL CASES.

PUBLISHED UNDER THE AUTHORITY OF THE LONDON SOCIETY FOR MEDICAL OBSERVATION.

In one very handsome royal 12mo volume, extra cloth.

We are convinced that the possession of the work will be almost necessary to every member of the profession—that it will be found indispensable to the practical physician, the pathologist, the medical jurist, and above all to the medical student.—*London Medical Times*.

We hail the appearance of this book as the grand desideratum.—*Charleston Medical Journal*.

This little work, if carefully read by even old practitioners, cannot fail to be productive of much good; as a guide to the younger members of the profession in directing their attention specially to the best mode of investigating cases so as to arrive at correct diagnosis, it will prove exceedingly valuable. The great difficulty with beginners, who have not been under the immediate training of an experienced physician,

is continually found to be in the appreciation of the true condition of the organs and tissues. Let such provide themselves with this work and study it thoroughly, and they will find much of the difficulty removed.—*Southern Medical and Surgical Journal*.

This is truly a very capital book. The whole medical world will reap advantages from its publication. The medical journals will soon show its influence on the character of the "Reports of Cases" which they publish. Drs. Ballard and Walke have given to the world, through a small but useful medical organization, a cheap but invaluable book. We do advise every reader of this notice to buy it and use it. Unless he is so vain as to imagine himself superior to the ordinary human capacity, he will in six months see its inestimable advantages.—*Stethoscope*.

## The Student's Text-Book.

LECTURES ON THE  
PRINCIPLES AND PRACTICE OF PHYSIC.

BY THOMAS WATSON, M. D.,

Late Physician to the Middlesex Hospital, &amp;c.

Third American, from a late London Edition.

REVISED, WITH ADDITIONS, BY D. FRANCIS CONDIE, M. D.,

Author of "A Practical Treatise on the Diseases of Children," &amp;c.

*In one very large octavo volume of nearly eleven hundred closely printed pages, strongly bound in leather, with raised bands.*

One of the most practically useful books that ever was presented to the student—indeed, a more admirable summary of general and special pathology, and of the application of therapeutics to diseases, we are free to say, has not appeared for very many years. The lecturer proceeds through the whole classification of human ills, *a capite ad calcem*, showing at every step an extensive knowledge of his subject, with the ability of communicating his precise ideas, in a style remarkable for its clearness and simplicity.—*N. Y. Journal of Medicine.*

The style is correct and pleasing, and the matter worth the attention of all practitioners, young and old.—*Western Lancet.*

We are free to state that a careful examination of this volume has satisfied us that it merits all the commendation bestowed on it in this country and at home. It is a work adapted to the wants of young practitioners, combining, as it does, sound principles and substantial practice. It is not too much to say, that it is a representative of the actual state of medicine as taught and practised by the most eminent physicians of the present day, and as such we would advise every one about embarking in the practice of physic to provide himself with a copy of it.—*Western Journal of Medicine and Surgery.*

To say that it is the very best work on the subject now extant, is but to echo the sentiment of the medical press throughout the country.—*N. O. Med. Journal.*

Of the text-books recently republished, Watson is very justly the principal favorite.—*Holmes's Report to Nat. Med. Association.*

By universal consent, the work ranks among the very best text-books in our language.—*Illinois and Indiana Med. Journal.*

Regarded on all hands as one of the very best, if not the very best, systematic treatise on practical medicine extant.—*St. Louis Med. Journal.*

Confessedly one of the very best works on the principles and practice of physic in the English or any other language.—*Med. Examiner.*

As a text-book it has no equal; as a compendium of pathology and practice no superior.—*N. Y. Annalist.*

We know of no work better calculated for being placed in the hands of the student, and for a text-book; on every important point the author seems to have posted up his knowledge to the day.—*American Medical Journal.*

DISEASES OF THE  
HEART, LUNGS, AND APPENDAGES;

THEIR SYMPTOMS AND TREATMENT. By W. H. WALSHE, M. D., Professor of the Principles and Practice of Medicine in University Medical College, London. In one large and handsome royal 12mo. volume, of 512 pages.

We consider this as the ablest work in the English language, on the subject of which it treats; the author being the first stethoscopist of the day.—*Charleston Medical Journal.*

The examination we have given the above work, convinces us that it is a complete system or treatise upon the great specialty of Physical Diagnosis. To give the reader a more perfect idea of what it con-

tains, we should be glad to copy the whole table of contents and make some extracts from its pages, but our limits forbid. We have no hesitation in recommending the work as one of the most complete on this subject in the English language; and yet it is not so voluminous as to be objectionable on this account, to any practitioner, however pressing his engagements.—*Ohio Medical and Surgical Journal.*

## NEW AND IMPROVED EDITION—(Just Ready.)

## A CLINICAL INTRODUCTION TO THE PRACTICE OF AUSCULTATION,

And other Modes of Physical Diagnosis, in Diseases of the Lungs and Heart. By H. M. HUGHES, M. D., Assistant Physician to Guy's Hospital, &c. Second American, from the Second and Improved London edition. In one neat volume, small royal 12mo., extra cloth.

## FROM THE PREFACE.

It has been carefully revised throughout. Some small portions have been erased; much has been, I trust, amended; and a great deal of new matter has been added; so that, though fundamentally it is the same book, it is in many respects a new work.

DAY ON OLD AGE.—A Practical Treatise on the Domestic Management and more important Diseases of Advanced Life. With an Appendix on a new and successful mode of treating Lumbago and other forms of Chronic Rheumatism. One vol. 8vo., pp. 226.

THOMSON ON THE SICK ROOM.—Domestic Management of the Sick-room, necessary in all of Medical Treatment for the cure of Diseases. Edited by R. E. Griffith, M. D. In one large royal 12mo. volume, with wood-cuts, pp. 360.

THE  
PRACTICE OF MEDICINE.

A TREATISE ON  
SPECIAL PATHOLOGY AND THERAPEUTICS.

BY ROBLEY DUNGLISON, M. D.,

Professor of the Institutes of Medicine in Jefferson Medical College, Philadelphia.

THIRD AND REVISED EDITION.

In two large octavo volumes, of fifteen hundred pages.

Upon every topic embraced in the work, the latest information will be found carefully posted up.—*Med. Examiner.*

The student of medicine will find in these two elegant volumes, a mine of facts, a gathering of precepts and advice from the world of experience, that will nerve him with courage, and faithfully direct him in

his efforts to relieve the physical sufferings of the race.—*Boston Medical and Surgical Journal.*

It is certainly the most complete treatise of which we have any knowledge.—*Western Journal of Medicine and Surgery.*

One of the most elaborate treatises of the kind we have.—*Southern Med. and Surg. Journal.*

New and Improved Edition—(Now Ready.)

URINARY DEPOSITS;  
THEIR DIAGNOSIS, PATHOLOGY, AND THERAPEUTICAL INDICATIONS.

BY GOLDING BIRD, M. D.,

Assistant Physician at Guy's Hospital, London, &c. &c.

Third American, from the Fourth Revised and Enlarged London Edition

In one handsome royal 12mo. volume, with numerous illustrations.

The opportunity afforded, of another revision of his work, has not been lost upon the author. The most recent information will be found embodied in its appropriate place, and an additional chapter has been introduced. It will, therefore, fully maintain the position it has acquired as an admirable manual for both the student and practitioner, on a subject of daily increasing importance.



Urate of Soda.



Hippuric Acid.

It can scarcely be necessary for us to say anything of the merits of this well-known Treatise, which so admirably brings into practical application the results of those microscopical and chemical researches regarding the physiology and pathology of the urinary secretion, which have contributed so much to the increase of our diagnostic powers, and to the extension

and satisfactory employment of our therapeutic resources. In the preparation of this new edition of his work, it is obvious that Dr. Golding Bird has spared no pains to render it a faithful representation of the present state of scientific knowledge on the subject it embraces.—*British and Foreign Med. Chir. Review.*

ATLAS OF PATHOLOGICAL HISTOLOGY.

BY GOTTLIEB GLUGE, M. D.,

Professor of Physiology and Pathological Anatomy in the University of Brussels, &c.

TRANSLATED, WITH NOTES AND ADDITIONS,

BY JOSEPH LEIDY, M. D.,

Professor of Anatomy in the University of Pennsylvania, &c.

In one handsome volume, large imperial quarto, with 320 Figures, plain and colored, on twelve copperplate engravings.

This being, as far as we know, the only work in which pathological histology is separately treated of in a comprehensive manner, it will, we think, for this reason, be of infinite service to those who desire to investigate the subject systematically, and who have felt the difficulty of arranging in their mind the un-

connected observations of a great number of authors. The development of the morbid tissues, and the formation of abnormal products, may now be followed and studied with the same ease and satisfaction as the best arranged system of Physiology.—*American Medical Journal.*



THE HISTORY, DIAGNOSIS, AND TREATMENT  
OF THE  
**FEVERS OF THE UNITED STATES.**

BY ELISHA BARTLETT, M. D.,  
Professor of Materia Medica, &c., in the College of Physicians and Surgeons, New York.  
Third Edition, Revised and Improved.

*In one octavo volume, of six hundred pages, beautifully printed and strongly bound.*

The masterly and elegant treatise by Dr. Bartlett is invaluable to the American student and practitioner.—*Dr. Holmes's Report to the Nat. Med. Association.*

We regard it, from the examination we have made of it, the best work on fevers extant in our language, and as such cordially recommend it to the medical public.—*St. Louis Medical and Surgical Journal.*

Take it altogether, it is the most complete history of our fevers which has yet been published, and every practitioner should avail himself of its contents.—*The Western Lancet.*

Of the value and importance of such a work, it is needless here to speak; the profession of the United States owe much to the author for the very able volume which he has presented to them, and for the careful and judicious manner in which he has executed his task. No one volume with which we are acquainted contains so complete a history of our fevers as this. To Dr. Bartlett we owe our best thanks for the very able volume he has given us, as embodying certainly the most complete, methodical, and satisfactory account of our fevers anywhere to be met with.—*The Charleston Med. Journal and Review.*

NOW READY.

**PNEUMONIA;**

Its supposed Connection, Pathological and Etiological, with Autumnal Fevers;

INCLUDING

AN INQUIRY INTO THE EXISTENCE AND MORBID AGENCY OF MALARIA.

BY R. LA ROCHE, M. D., &c.

*In one very handsome octavo volume of 500 pages, extra cloth.*

A more simple, clear, and forcible exposition of the groundless nature and dangerous tendency of certain pathological and etiological heresies, has seldom been presented to our notice.—*N. Y. Journal of Medicine and Collateral Science*, March, 1854.

This work should be carefully studied by Southern physicians, embodying as it does the reflections of an original thinker and close observer on a subject peculiarly their own.—*Virginia Med. and Surg. Journal.*

The author had prepared us to expect a treatise from him, by his brief papers on kindred topics in the

periodical press; and yet, in the work before us, he has exhibited an amount of industry and learning, research and ability, beyond what we are accustomed to discover in modern medical writers; while his own extensive opportunities for observation and experience have been improved by the most laudable diligence, and display a familiarity with the whole subject in every aspect, which commands both our respect and confidence. As a corrective of prevalent and mischievous error, sought to be propagated by novices and innovators, we could wish that Dr. La Roche's book could be widely read.—*N. Y. Medical Gazette.*

JUST ISSUED.

**ON DISEASES OF THE LIVER.**

BY GEORGE BUDD, M. D., F. R. S.,

Professor of Medicine in King's College, London.

SECOND AMERICAN, FROM THE SECOND AND ENLARGED LONDON EDITION.

In one  
handsome volume  
of 468 pages.  
With four  
exquisite colored  
plates  
and numerous  
wood-cuts.



*Echinococci from Liver.*

The full digest we have given of the new matter introduced into the present volume, is evidence of the value we place on it. The fact that the profession has required a second edition of a monograph such as that before us, bears honorable testimony to its usefulness. For many years, Dr. Budd's work must be the authority of the great mass of British practitioners on the hepatic diseases; and it is satisfactory that the subject has been taken up by so able and experienced a physician.—*British and Foreign Med.-Chirurgy Review.*

New and Improved Edition—(Lately Issued.)

## ON DISEASES OF THE SKIN.

BY ERASMUS WILSON, M. D.,

Author of "A System of Human Anatomy."

THIRD AMERICAN, FROM THE THIRD AND IMPROVED LONDON EDITION.

*In one neat octavo volume, of about five hundred pages, extra cloth.*

*Also, to be had done up with Fifteen beautiful Steel Plates, of which Eight are exquisitely colored; representing the Normal and Pathological Anatomy of the Skin, together with accurately colored delineations of more than sixty varieties of disease, most of them the size of nature.*

THE PLATES ARE ALSO FOR SALE SEPARATE, DONE UP IN BOARDS.

The "Diseases of the Skin," by Mr. Erasmus Wilson, may now be regarded as the standard work in that department of medical literature. The plates by which this edition is accompanied leave nothing to be desired, so far as excellence of delineation and perfect accuracy of illustration are concerned.—*Medico-Chirurg. Review.*

Of these plates it is impossible to speak too highly. The representations of the various forms of cutaneous disease are singularly accurate, and the coloring exceeds almost anything we have met with in point of delicacy and finish.—*British and Foreign Med. Review.*

BY THE SAME AUTHOR.

## ON CONSTITUTIONAL AND HEREDITARY SYPHILIS, AND ON SYPHILITIC ERUPTIONS.

*In one beautifully printed octavo volume, with four exquisite colored plates, presenting more than thirty varieties of syphilitic eruptions.*

Dr. Wilson's views on the general subject of Syphilis appear to us in the main sound and judicious, and we commend the book as an excellent monograph on the subject. Dr. Wilson has presented us a very faithful and lucid description of Syphilis, and has cleared up

many obscure points in connection with its transmissibility, pathology, and sequelæ. His facts and references will, we are satisfied, be received as decisive, in regard to many questions vexatæ. They appear to us entitled to notice at some length.—*Med. Examiner.*


BY THE SAME AUTHOR. (Just Ready.)

## HEALTHY SKIN.

A Treatise on the Management of the Skin and Hair in Relation to Health.

Second American, from the Fourth and Revised London Edition.

*In one royal 12mo. volume, with illustrations.*



Magnified View of various kinds of Hair.

The student will be delighted to find his labors so much facilitated; and a few hours of agreeable society with a most pleasantly-written book, will do more to

make him acquainted with a class of obscure diseases than all that has been previously written on the subject.—*Lancet.*

## A PRACTICAL TREATISE ON DISEASES OF THE SKIN.

BY J. MOORE NELIGAN, M. D., M. R. I. A., &c. &c.

*In one neat royal 12mo. volume of 334 pages.*

We know of no other treatise on this interesting and important class of diseases that so happily meets the urgent wants of the great mass of physicians.—*N. Y. Journal of Medicine.*

The greatest value of Dr. Neligan's treatise consists in the plain and thoroughly practical exposition he has given of this class of maladies.—*British and Foreign Med.-Chirurg. Review.*

RICORD'S LATEST WORK ON SYPHILIS—(Just Issued.)

## A TREATISE ON THE VENEREAL DISEASE.

BY JOHN HUNTER, F. R. S., &c.

WITH COPIOUS ADDITIONS, BY PH. RICORD, M. D.,

EDITED, WITH NOTES, BY FREEMAN J. BUMSTEAD, M. D.

*In one large and handsome octavo volume, of over five hundred pages, with plates.*

"M. Ricord's annotations to *Hunter's Treatise on the Venereal Disease* were first published at Paris, in 1840, in connection with Dr. G. Richelot's translation of the work, including the contributions of Sir Everard Home and Mr. Babington. In a second edition, which has recently appeared, M. Ricord has thoroughly revised his part of the work, bringing it up to the knowledge of the present day, and so materially increasing it that it now constitutes full one-third of the volume. This publication has been received with great favor by the French, both because it has placed within their reach an important work of Hunter, and also because it is the only recent practical work which M. Ricord has published, no edition of his *Traité des Maladies Vénéériennes* having appeared for the last fifteen years."—*Translator's Preface.*

Every one will recognize the attractiveness and value which this work derives from thus presenting the opinions of these two masters side by side. But, it must be admitted, what has made the fortune of the book, is the fact that it contains the "most complete embodiment of the veritable doctrines of the Hôpital du Midi," which has ever been made public. The doctrinal ideas of M. Ricord, ideas which, if not universally adopted, are incontestably dominant, have heretofore only been interpreted by more or less skillful

secretaries, sometimes accredited and sometimes not. In the notes to Hunter, the master substitutes himself for his interpreters, and gives his original thoughts to the world, in a summary form it is true, but in a lucid and perfectly intelligible manner. In conclusion, we can say that this is incontestably the best treatise on syphilis with which we are acquainted, and, as we do not often employ the phrase, we may be excused for expressing the hope that it may find a place in the library of every physician.—*Va Med. and Surg. Journ.*

Also—HUNTER'S COMPLETE WORKS. In four octavo volumes, with numerous Illustrations.

## LETTERS ON SYPHILIS,

Addressed to the Chief Editor of the *Union Médicale*. By PH. RICORD, M. D. With an Introduction by Amédée Latour. Translated by W. P. Lattimore, M. D. In one neat octavo volume.

To those who have listened to the able and interesting lectures of our author at the Hôpital du Midi, this volume will need no commendation; while to those who have not had the pleasure to which we allude, the book will commend itself by the truths it contains, told as they are in the same inimitable style in which M. Ricord delivers his clinical lectures.—*Translator's Preface.*

BY THE SAME AUTHOR.

A PRACTICAL TREATISE ON VENEREAL DISEASES. With a Therapeutical Summary and Special Formulary. Translated by SIDNEY DOANE, M. D. Fourth edition. One volume, octavo, 340 pages.

## ILLUSTRATIONS OF SYPHILITIC DISEASE.

By PH. RICORD, M. D. Translated from the French, by THOMAS F. BETTON, M. D. With the addition of a History of Syphilis, and a complete Bibliography and Formulary of Remedies, collated and arranged, by PAUL B. GODDARD, M. D. With fifty large quarto plates, comprising one hundred and seventeen beautifully colored illustrations. In one large and handsome quarto volume, extra cloth.

*Now Ready.*

## THE MODERN TREATMENT OF SYPHILITIC DISEASES, BOTH PRIMARY AND SECONDARY.

COMPRISING THE

Treatment of Constitutional and confirmed Syphilis by a safe and successful Method.

WITH NUMEROUS CASES, FORMULÆ, AND CLINICAL OBSERVATIONS.

BY LANGSTON PARKER,

Surgeon to the Queen's Hospital, Birmingham.

From the Third and entirely rewritten London Edition,

*In one neat octavo volume, of over 300 pages.*

The third edition of Mr. Parker's work constitutes, we must say, an excellent practical treatise, the subject is remarkably well arranged; indeed, the author has contrived to embody in it everything of importance respecting syphilis and its treatment.—*Dublin Medical Press*, March, 1854.

# SURGICAL ANATOMY.

BY JOSEPH MACLISE, SURGEON.

FORMING ONE VOLUME, VERY LARGE IMPERIAL QUARTO,

With Sixty-eight large and splendid Plates, drawn in the best style, and beautifully colored.

Containing one hundred and ninety Figures, many of them the size of life.

TOGETHER WITH COPIOUS AND EXPLANATORY LETTER-PRESS.

Strongly and handsomely bound in extra cloth, being one of the cheapest and best executed Surgical works as yet issued in this country.

Copies can be sent by mail, in five parts, done up in stout covers.

This great work being now concluded, the publishers confidently present it to the attention of the profession as worthy in every respect of their approbation and patronage. No complete work of the kind has yet been published in the English language, and it therefore will supply a want long felt in this country of an accurate and comprehensive Atlas of Surgical Anatomy to which the student and practitioner can at all times refer, to ascertain the exact relative position of the various portions of the human frame towards each other and to the surface, as well as their abnormal deviations. The importance of such a work to the student in the absence of anatomical material, and to the practitioner when about attempting an operation, is evident; while the price of the book, notwithstanding the large size, beauty, and finish of the very numerous illustrations, is so low as to place it within the reach of every member of the profession. The publishers therefore confidently anticipate a very extended circulation for this magnificent work.

One of the greatest artistic triumphs of the age in Surgical Anatomy.—*British American Medical Journal*.

Too much cannot be said in its praise; indeed, we have not language to do it justice.—*Ohio Medical and Surgical Journal*.

The most admirable surgical atlas we have seen. To the practitioner deprived of demonstrative dissections upon the human subject, it is an invaluable companion.—*N. J. Medical Reporter*.

The most accurately engraved and beautifully colored plates we have ever seen in an American book—one of the best and cheapest surgical works ever published.—*Buffalo Medical Journal*.

It is very rare that so elegantly printed, so well illustrated, and so useful a work, is offered at so moderate a price.—*Charleston Medical Journal*.

Its plates can boast a superiority which places them almost beyond the reach of competition.—*Medical Examiner*.

Every practitioner, we think, should have a work of this kind within reach.—*Southern Medical and Surgical Journal*.

No such lithographic illustrations of surgical regions have hitherto, we think, been given.—*Boston Medical and Surgical Journal*.

As a surgical anatomist, Mr. MacLise has probably no superior.—*British and Foreign Medico-Chirurgical Review*.

Of great value to the student engaged in dissecting, and to the surgeon at a distance from the means of keeping up his anatomical knowledge.—*Medical Times*.

The mechanical execution cannot be excelled.—*Transylvania Medical Journal*.

A work which has no parallel in point of accuracy and cheapness in the English language.—*N. Y. Journal of Medicine*.

No practitioner whose means will admit should fail to possess it.—*Rankin's Abstract*.

Country practitioners will find these plates of immense value.—*N. Y. Medical Gazette*.

To all engaged in the study or practice of their profession, such a work is almost indispensable.—*Dublin Quarterly Medical Journal*.

We are extremely gratified to announce to the profession the completion of this truly magnificent work, which, as a whole, certainly stands unrivalled, both for accuracy of drawing, beauty of coloring, and all the requisite explanations of the subject in hand. To the publishers, the profession in America is deeply indebted for placing such a valuable, such a useful work, at its disposal, and at such a moderate price. It is one of the most finished and complete pictures of Surgical Anatomy ever offered to the profession of America. With these plates before them, the student and practitioner can never be at a loss, under the most desperate circumstances. We do not intend these for commonplace compliments. We are sincere; because we know the work will be found invaluable to the young, no less than the old, surgeon.—*The New Orleans Medical and Surgical Journal*.

This is by far the ablest work on Surgical Anatomy that has come under our observation. We know of no other work that would justify a student, in any degree, for neglect of actual dissection. A careful study of these plates, and of the commentaries on them, would almost make an anatomist of a diligent student. And to one who has studied anatomy by dissection, this work is invaluable as a perpetual remembrancer, in matters of knowledge that may slip from the memory. The practitioner can scarcely consider himself equipped for the duties of his profession without such a work as this; and this has no rival, in his library. In those sudden emergencies that so often arise, and which require the instantaneous command of minute anatomical knowledge, a work of this kind keeps the details of the dissecting-room perpetually fresh in the memory. We repeat that no medical library, however large, can be complete without MacLise's Surgical Anatomy. The American edition is well entitled to the confidence of the profession, and should command, among them, an extensive sale. The investment of the amount of the cost of this work will prove a very profitable one, and if practitioners were to qualify themselves thoroughly with such important knowledge, as is contained in works of this kind, there would be fewer of them sighing for employment.—*The Western Journal of Medicine and Surgery*.

The very low price at which this work is furnished, and the beauty of its execution, require an extended sale to compensate the publishers for the heavy expenses incurred.

A NEW SURGICAL TEXT-BOOK—(New Ready.)

In one  
very large  
and handsome  
octavo volume,  
of  
over nine hundred  
pages.



Illustrated  
with  
three hundred  
and eleven  
beautiful  
engravings on  
wood.

Plastic Operation of the Face.

**THE SCIENCE AND ART OF SURGERY;**  
BEING  
**A TREATISE ON SURGICAL INJURIES, DISEASES, AND OPERATIONS**  
BY JOHN ERICHSEN,  
Professor of Surgery in University College.  
EDITED BY JOHN H. BRINTON, M. D.

From Prof. S. D. Gross, Louisville, May 17, 1854.

"Decidedly the ablest treatise on the subject since the days of Benjamin Bell."

The aim of Mr. Erichsen appears to be to improve upon the plan of Samuel Cooper; and by connecting in one volume the science and art of Surgery, to supply the student with a text-book and the practitioner with a work of reference, in which scientific principles and practical details are alike included. It must raise

the character of the author, and reflect great credit upon the College to which he is Professor, and we can cordially recommend it as a work of reference, both to students and practitioners.—*Med. Times and Gazette.*

The volume before us gives a very admirable practical view of the science and art of surgery of the present day, and we have no doubt that it will be highly valued as a surgical guide as well by the surgeon as by the student of surgery.—*Edinburgh Med. and Surg. Journal.*



Amputation of Hip.

Just Issued.

# THE PRACTICE OF SURGERY.

BY JAMES MILLER, F. R. S. E.,

Professor of Surgery in the University of Edinburgh: author of "Principles of Surgery," &c. &c.

Third American, from the Second Edinburgh Edition.

EDITED, WITH ADDITIONS, BY F. W. SARGENT, M. D.,

One of the Surgeons to Wills Hospital, &c.

WITH THREE HUNDRED AND NINETEEN BEAUTIFUL ILLUSTRATIONS ON WOOD.

In one very large and handsome octavo volume, of 720 pages.

This new edition will be found greatly improved and enlarged, as well by the addition of much new matter as by the introduction of a large and complete series of handsome illustrations. An equal improvement exists in the mechanical execution of the work, rendering it in every respect a companion volume to the "Principles."

No encomium of ours could add to the popularity of Miller's Surgery. Its reputation in this country is unsurpassed by that of any other work, and, when taken in connection with the author's *Principles of Surgery*, constitutes a whole, without reference to which no conscientious surgeon would be willing to practise his art. The additions, by Dr. Sargent, have materially enhanced the value of the work.—*Southern Medical and Surgical Journal*.

It is seldom that two volumes have ever made so profound an impression in so short a time as the "Principles" and the "Practice" of Surgery, by Mr. Miller—or so richly merited the reputation they have acquired. The author is an eminently sensible, practical, and well-informed man, who knows exactly what he is talking about and exactly how to talk it.—*Kentucky Medical Recorder*.

The two volumes together form a complete exposé of the present state of Surgery, and they ought to be on the shelves of every surgeon.—*N. J. Medical Reporter*.


By the almost unanimous voice of the profession, his works, both on the principles and practice of surgery have been assigned the highest rank. If we were limited to but one work on Surgery, that one should be Miller's, as we regard it superior to all others.—*St. Louis Med. and Surg. Journal*.

The author, distinguished alike as a practitioner and writer, has in this and his "Principles," presented to the profession one of the most complete and reliable systems of Surgery extant. His style of writing is original, impressive, and engaging, energetic, concise, and lucid. Few have the faculty of condensing so much in small space, and at the same time so persist-



Resection of the Elbow

ently holding the attention; indeed, he appears to make the very process of condensation a means of producing attractions. Whether as a text-book for students or a book of reference for practitioners, it cannot be too strongly recommended.—*Southern Journal of Medical and Physical Science*.



Ligatures of the Anterior Tibial



Structure of Colloid Cancer.

## PRINCIPLES OF SURGERY.

BY JAMES MILLER, F. R. S. E.,

Professor of Surgery in the University of Edinburgh.

THIRD AMERICAN, FROM THE SECOND AND REVISED EDINBURGH EDITION.

REVISED, WITH ADDITIONS, BY F. W. SARGENT, M. D.,

Demonstrator of Anatomy in the University of Pennsylvania; author of "Minor Surgery," &c.



Cysto-Sarcoma.

In one large  
and very  
beautiful volume  
of  
seven hundred and  
fifty pages,  
with  
two hundred and forty  
exquisite  
illustrations on wood.



Neuromata, after Amputation.

The publishers have endeavored to render the present edition of this work, in every point of mechanical execution, worthy of its very high reputation, and they confidently present it to the profession as one of the handsomest volumes as yet issued in this country.

This edition is far superior, both in the abundance and quality of its material, to any of the preceding. We hope it will be extensively read, and the sound principles which are herein taught treasured up for future application. The work takes rank with Watson's Practice of Physic; it certainly does not fall behind that great work in soundness of principle or depth of reasoning or research. No physician who values his reputation, or seeks the interest of his clients, can acquit himself before his God and the world, without making himself familiar with the sound and philosophical views developed in the foregoing book.—*New Orleans Medical and Surgical Journal.*

Without doubt, the ablest exposition of the principles of that branch of the healing art in any language.

This opinion, deliberately formed after a careful study of the first edition, we have had no cause to change on examining the second. This edition has undergone thorough revision by the author; many expressions have been modified, and a mass of new matter introduced. The book is got up in the finest style, and is an evidence of the progress of typography in our country.—*Charleston Medical Journal and Review.*

We recommend it to both student and practitioner, feeling assured that, as it now comes to us, it presents the most satisfactory exposition of the modern doctrines of the principles of surgery to be found in any volume in any language.—*N. Y. Journal of Medicine.*

Lately Issued.

**THE PRINCIPLES AND PRACTICE OF SURGERY.**

BY WILLIAM PIRRIE, F. R. S. E.,

Regius Professor of Surgery in the University of Aberdeen.

EDITED, WITH ADDITIONS, BY JOHN NEILL, M. D.,

Surgeon to the Pennsylvania Hospital, Professor of Surgery in Penna. Medical College, &c. &c.

In one large and handsome octavo volume, of nearly 800 pages, with 316 beautiful illustrations on wood.



*Excursion of Spine.*


However well it may be adapted for a text-book (and in this respect it may compete with the best of them), of this much our reading has convinced us, that, as a systematic treatise, it is carefully and ably written, and can hardly fail to command a prominent position in the library of practitioners; though not complete in the fullest sense of the word, it nevertheless furnishes the student and practitioner with as chaste and concise a work as exists in our language. The additions to the volume by Dr. Neill, are judicious; and while they render it more complete, greatly enhance its practical value, as a work for practitioners and students.—*N. Y. Journal of Medicine.*

We know of no other surgical work of a reasonable size, wherein there is so much theory and practice, or where subjects are more soundly or clearly taught.—*The Stethoscope.*

There is scarcely a disease of the bone or soft parts, fracture, or dislocation, that is not illustrated by accurate wood-engravings.—Then, again, every instrument employed by the surgeon is thus represented. These engravings are not only correct, but really beautiful, showing the astonishing degree of perfection to which the art of wood-engraving has arrived. Prof. Pirrie, in the work before us, has elaborately discussed the principles of surgery, and a safe and effectual practice predicated upon them. Perhaps no work upon this subject heretofore issued is so full upon the science of the art of surgery.—*Nashville Journal of Medicine and Surgery.*

We can pronounce it to be one of the best treatises on surgery in the English language. We very strongly recommend this excellent work, both to the practitioner and student.—*Cunala Medical Journal.*

Our impression is that, as a manual for students, Pirrie's is the best work extant.—*Western Medical and Surgical Journal.*



*Amputation of the Thigh.*





*Staphylopharynx.*

In one  
very handsome  
octavo volume, of nearly  
six hundred pages,  
illustrated  
with  
one hundred and ninety-three  
engravings  
on  
wood.



*Stricture of Esophagus.*

THE  
PRINCIPLES AND PRACTICE OF MODERN SURGERY.

BY ROBERT DRUITT,

Fellow of the Royal College of Surgeons.

A new American, from a late and improved London Edition.

EDITED BY F. W. SARGENT, M. D.,

Author of "Minor Surgery," &c.



*Flap Operation of the Thigh.*


Dr. Drutt's researches into the literature of his subject have been not only extensive, but well directed; the most discordant authors are fairly and impartially quoted, and, while due credit is given to each, their respective merits are weighed with an unprejudiced hand. The grain of wheat is preserved, and the chaff is unmercifully stripped off. The arrangement is simple and philosophical, and the style, though clear and interesting, is so precise, that the book contains more information condensed into a few words than any other surgical work with which we are acquainted.—*London Medical Times and Gazette*, Feb. 18, 1854.

An unsurpassable compendium, not only of Surgical, but of Medical Practice.—*London Med. Gazette*.

Probably no work of the kind has ever been more cordially received and extensively circulated than this. The fact that it comprehends in a comparatively small compass, all the essential elements of theoretical and practical surgery—that it is found to contain reliable and authentic information on the nature and treatment of nearly all surgical affections, is a sufficient reason for the liberal patronage it has obtained. It abounds everywhere with spirited and life-like illustrations, which to the young surgeon, especially, are of no minor consideration. Every medical man frequently needs just such a work as this, for immediate reference in moments of sudden emergency, when he has not time to consult more elaborate treatises.—*Ohio Med. and Surg. Journal*.

New and much Improved Edition—(Lately Issued.)

In  
one very  
beautiful  
octavo volume  
of over  
six hundred  
pages,



With  
nearly  
four hundred  
handsome  
illustrations

*Salter's Swing-Splint.*

## A SYSTEM OF PRACTICAL SURGERY.

BY WILLIAM FERGUSSON, F. R. S.,

Professor of Surgery in King's College, London, &c.

FOURTH AMERICAN FROM THE THIRD AND ENLARGED LONDON EDITION.

The most important subjects in connection with practical surgery which have been more recently brought under the notice of, and discussed by, the surgeons of Great Britain, are fully and dispassionately considered by Mr. Fergusson, and that which was before wanting has now been supplied; so that we can now look upon it as a work on practical surgery, instead of one on operative surgery alone, which many have hitherto considered it to be. And we think the author has shown a wise discretion, in making the additions on surgical disease which are to be found in the present volume, and has very much enhanced its value; for, besides two elaborate chapters on the diseases of bones and joints, which were wanting before, he has headed each chief section of the work by a

general description of the surgical disease and injury of that region of the body which is treated of in each, prior to entering into the consideration of the more special morbid conditions and their treatment. There is also, as in former editions, a sketch of the anatomy of particular regions. We have now pointed out some of the principal additions in this work. There was some ground formerly for the complaint before alluded to, that it dwelt too exclusively on operative surgery; but this defect is now removed, and the book is more than ever adapted for the purposes of the practitioner, whether he confines himself more strictly to the operative department, or follows surgery on a more comprehensive scale.—*Medical Times and Gazette.*

No work was ever written which more nearly comprehended the necessities of the student and practitioner, and was more carefully arranged to that single purpose than this.—*N. Y. Medical and Surgical Journal.*

The addition of many new pages makes this work more than ever indispensable to the student and practitioner.—*Ranking's Abstract.*

For the general practitioner, who does not make a speciality of surgery, it is certainly invaluable. The style is concise, pointed, and clear. The descriptions of the various operations are concentrated and accurate, so that in cases of emergency, the principles of the most difficult operations may be obtained by a reference of a few moments to its pages.—*Western Lancet.*

Among the numerous works upon surgery published of late years, we know of none we value more highly than the one before us. It is perhaps the very best we have for a text-book and for ordinary reference, being concise and eminently practical.—*Southern Medical and Surgical Journal.*



*Extraction of Foreign Bodies from the Oesophagus.*

# A SYSTEM OF SURGERY.

BY J. M. CHELIUS, M. D.,

Professor of Surgery in the University of Heidelberg, &c.

*Translated from the German, and accompanied with additional Notes and References,*

BY JOHN F. SOUTH.

In three very large octavo volumes, of nearly twenty-two hundred pages, strongly bound in leather, with raised bands.

We do not hesitate to pronounce it the best and most comprehensive system of modern surgery with which we are acquainted.—*Medico-Chirurgical Review.*

The fullest and ablest digest extant of all that relates to the present advanced state of surgical pathology.—*American Medical Journal.*

The most learned and complete systematic treatise now extant.—*Edinburgh Medical Journal.*

A complete encyclopædia of surgical science—a very complete surgical library—by far the most complete and scientific system of surgery in the English language.—*N. Y. Journal of Medicine.*

## OPERATIVE SURGERY;

OR,

A DESCRIPTION AND DEMONSTRATION OF THE VARIOUS PROCESSES OF THE ART. INCLUDING ALL THE NEW OPERATIONS,

And exhibiting the state of Surgical Science in its present advanced condition.

BY JOSEPH PANCOAST, M. D.,

Professor of Anatomy in the Jefferson Medical College of Philadelphia.

*In one large and handsome royal quarto volume.*

With eighty large and splendid Plates, comprising four hundred and eighty-six Illustrations.

Blanchard & Lea having become the publishers of this important book, have much pleasure in offering it to the profession.

This excellent work is constructed on the model of the French Surgical Works by Velpeau and Malgaigne; and, so far as the English language is con-

cerned, we are proud as an American to say that, of ITS KIND, IT HAS NO SUPERIOR.—*New York Journal of Medicine.*

## OPERATIVE SURGERY;

BASED ON NORMAL AND PATHOLOGICAL ANATOMY.

BY J. F. MALGAIGNE.

TRANSLATED FROM THE FRENCH, BY FREDERIC BRITTAN, M. D.

In one  
neat  
octavo  
volume,  
of  
over  
five hundred  
and fifty  
pages.




With  
handsome  
illustrations  
from  
designs  
by  
Dr.  
Westmacott.

*Surgical Anatomy of the Perineum.*

We have long been accustomed to refer to it as one of the most valuable text-books in our library.—*Buffalo Med. and Surg. Journal.*

Certainly one of the best books published on operative surgery.—*Edinburgh Medical Journal.*

To express in a few words our opinion of Malgaigne's work, we unhesitatingly pronounce it the very best guide in surgical operations that has come before the profession in any language.—*Charleston Medical and Surgical Journal.*



*Reduction of Dislocation.*

## OPERATIVE SURGERY.

BY FREDERICK C. SKEY, F. R. S., &c.

With about 100 Illustrations on Wood.

*In one handsome octavo volume, of over six hundred and fifty pages.*

Its literary execution is superior to most surgical treatises. It abounds in excellent moral hints, and is replete with original surgical expedients and suggestions.—*Buffalo Med. and Surg. Journal.*

With high talents, extensive practice, and a long experience, Mr. Skey is perhaps competent to the task of writing a complete work on operative surgery.—*Charleston Med. Journal.*

We cannot withhold from this work our high commendation. Students and practitioners will find it an invaluable teacher and guide upon every topic connected with this department.—*New York Medical Gazette.*

A work of the very highest importance—a work by itself.—*London Med. Gazette.*

## A TREATISE ON Dislocations and Fractures of the Joints.

BY

SIR ASTLEY P. COOPER, BART., F. R. S., &c.

EDITED BY

BRANSBY B. COOPER, F. R. S., &c.

With additional Observations.

BY PROFESSOR J. C. WARREN.

A new American edition.

In one handsome octavo volume, illustrated

with numerous wood-cuts.



*Dislocation of the Ilium.*

### WORKS BY THE SAME AUTHOR.

COOPER (SIR ASTLEY) ON THE ANATOMY AND TREATMENT OF ABDOMINAL HERNIA. One large vol., imperial 8vo., with over 130 lithographic figures.

COOPER ON THE STRUCTURE AND DISEASES OF THE TESTIS, AND ON THE THYMUS

GLAND. One vol., imperial 8vo., with 177 figures on 29 plates.

COOPER ON THE ANATOMY AND DISEASES OF THE BREAST, with Twenty-five Miscellaneous and Surgical Papers. One large vol., imperial 8vo., with 262 figures on 36 plates.

These three volumes complete the surgical writings of Sir Astley Cooper. They are very handsomely printed, with a large number of lithographic plates, executed in the best style, and are presented at exceedingly low prices.

Now Ready.

*Catheterism of the Eustachian Tube.*

**PRACTICAL OBSERVATIONS ON AURAL SURGERY,  
AND THE NATURE AND TREATMENT OF  
DISEASES OF THE EAR.**

With Illustrations.

BY WILLIAM R. WILDE,

Surgeon to St. Mark's Ophthalmic Hospital, Dublin.

*In one very handsome octavo volume of nearly five hundred pages.*

So little is generally known in this country concerning the causes, symptoms, and treatment of aural affections, that a practical and scientific work on that subject, from a practitioner of Mr. Wilde's great experience, cannot fail to be productive of much benefit, by attracting attention to this obscure class of diseases, which too frequently escape attention until past relief. The immense number of cases which have come under Mr. Wilde's observation for many years, have afforded him opportunities rarely enjoyed for investigating this branch of medical science, and his work may therefore be regarded as of the highest authority.

This work certainly contains more information on the subject to which it is devoted than any other with which we are acquainted. We feel grateful to the author for his manful effort to rescue this department of surgery from the hands of the empirics who nearly monopolize it. We think he has successfully shown that aural diseases are not beyond the resources of

art; that they are governed by the same laws, and amenable to the same general methods of treatment as other morbid processes. The work is not written to supply the cravings of popular patronage, but it is wholly addressed to the profession, and bears on every page the impress of the reflections of a sagacious and practical surgeon.—*Va. Surg. and Med. Journal.*

**ON BANDAGING, AND OTHER POINTS OF MINOR SURGERY.**

BY F. W. SARGENT, M. D.,

Surgeon to Wills Hospital, Philadelphia; Demonstrator of Anatomy in the University of Pennsylvania, &amp;c.

*In one very neat royal 12mo. volume, of about four hundred pages.*

With one hundred and twenty-eight illustrations.

The very best manual of Minor Surgery we have seen; an American volume, with nearly four hundred pages of good practical lessons, illustrated by about one hundred and thirty wood-cuts.—*Buffalo Med. Jour.*

We have carefully examined this work, and find it well executed and admirably adapted to the use of the student.—*The Charleston Medical Journal.*

*Gibson's Splint for Fracture of Femur.*


**A PRACTICAL TREATISE ON THE DISEASES AND INJURIES  
OF THE  
URINARY BLADDER, THE PROSTATE GLAND, AND THE URETHRA.**

BY S. D. GROSS, M. D.,

Professor of Surgery in the University of Louisville, &c.

WITH ONE HUNDRED AND SIX ILLUSTRATIONS.

In one large and very handsome octavo volume, of over seven hundred pages.



*Division of the Prostate in the Lateral Operation of Lithotomy.*

A volume replete with truths and principles of the utmost value in the investigation of those diseases.—*American Medical Journal.*

Dr. Gross has brought all his learning, experience, tact, and judgment to the task, and has produced a work worthy of his high reputation. We feel perfectly safe in recommending it to our readers as a monograph unequalled in interest and practical value by any other on the subject in our language.—*The Western Journal of Medicine and Surgery.*


It has remained for an American writer to wipe away this reproach; and so completely has the task been fulfilled, that we venture to predict for Dr. Gross's treatise a permanent place in the literature of surgery, worthy to rank with the best works of the present age. Not merely is the matter good, but the getting up of the volume is most creditable to transatlantic enterprise: the paper and print would do credit to a first-rate London establishment; and the numerous

wood-cuts which illustrate it, demonstrate that America is making rapid advances in this department of art. We have, indeed, unfeigned pleasure in congratulating all concerned in this publication, on the result of their labors; and experience a feeling something like what animates a long-expectant husbandman, who, oftentimes disappointed by the produce of a favorite field, is at last agreeably surprised by a stately crop which may bear comparison with any of its former rivals. The grounds of our high appreciation of the work will be obvious as we proceed.—*British and Foreign Medico-Chirurgical Review.*

Whoever will peruse the vast amount of valuable practical information it contains, and which we have been unable even to notice, will, we think, agree with us, that there is no work in the English language which can make any just pretensions to be its equal.—*N. Y. Journal of Medicine.*

BY THE SAME AUTHOR—(Nearly Ready.)

**A PRACTICAL TREATISE ON  
FOREIGN BODIES IN THE AIR-PASSAGES.**



With illustrations, in one handsome octavo volume, of about 450 pages.

BY THE SAME AUTHOR—(Preparing.)

**A SYSTEM OF SURGERY;  
DIAGNOSTIC, PATHOLOGICAL, THERAPEUTIC, AND OPERATIVE.**

NOW READY.

*Purulent Ophthalmia.**Extraction of Cataract.*

With about  
two hundred and fifty  
handsome illustrations on wood.

## A TREATISE ON THE DISEASES OF THE EYE.

BY W. LAWRENCE, F. R. S.,  
Surgeon Extraordinary to the Queen.

NEW EDITION, WITH NUMEROUS ADDITIONS.

BY ISAAC HAYS, M. D.,  
Surgeon to Wills Hospital, &c.

In one beautiful octavo volume of 950 pages, strongly bound in leather, with raised bands.

This work is so universally recognized as the standard authority on the subject, that the publishers in presenting this new edition have only to remark that in its preparation the editor has carefully revised every portion, introducing additions and illustrations wherever the advance of science has rendered them necessary or desirable. In this manner it will be found to contain over one hundred pages more than the last edition, while the list of wood-engravings has been increased by sixty-seven figures, besides numerous improved illustrations substituted for such as were deemed imperfect or unsatisfactory. The various important contributions to ophthalmological science, recently made by Dalrymple, Jacob, Walton, Wilde, Cooper, &c., both in the form of separate treatises and contributions to periodicals, have been carefully examined by the editor, and, combined with the results of his own experience, have been freely introduced throughout the volume, rendering it a complete and thorough exponent of the most advanced state of the subject. Among the most important additions may be mentioned a full account of the recent microscopical investigations into the structure and pathology of the eye; the description of several affections not treated of in the original; an account of the catoptric investigation of the eye, and of its employment as a means of diagnosis; a description of recently invented instruments for illuminating the retina, and of some new methods for examining the interior structures of the eye. Very great improvements will likewise be found in the typographical and mechanical execution of the work.

In a future number we shall notice more at length this admirable treatise—the safest guide and most comprehensive work of reference, which is within the reach of all classes of the profession.—*Stethoscope*, March, 1854.

This standard text-book on the department of which it treats, has not been superseded, by any or all of the numerous publications on the subject heretofore issued. Nor with the multiplied improvements of Dr. Hays, the American editor, is it at all likely that this great work will cease to merit the confidence and preference of students or practitioners. Its ample extent, nearly one thousand large octavo pages, has enabled

both author and editor to do justice to all the details of this subject, and condense in this single volume the present state of our knowledge of the whole science in this department, whereby its practical value cannot be excelled. We heartily commend it, especially as a book of reference, indispensable in every medical library. The additions of the American editor very greatly enhance the value of the work, exhibiting the learning and experience of Dr. Hays, in the light in which he ought to be held, as a standard authority on all subjects appertaining to this specialty, to which he has rendered so many valuable contributions.—*N. Y. Medical Gazette*.

BY THE SAME AUTHOR.

A TREATISE ON RUPTURES. From the fifth London Edition. In one octavo volume, sheep, 480 pages.

THE PRINCIPLES AND PRACTICE OF  
OPHTHALMIC MEDICINE AND SURGERY.


BY T. WHARTON JONES, F. R. S., &c.

EDITED BY ISAAC HAYS, M. D.



*Sclerotic Staphyloma*

In one very neat  
royal 12mo. volume,  
of over  
five hundred pages,  
with  
four plates,  
plain or colored,  
and  
ninety-eight  
wood-cuts.



*Iridauzesis.*

The work amply sustains, in every point the already high reputation of the author as an ophthalmic surgeon as well as a physiologist and pathologist. The book is evidently the result of much labor and research, and has been written with the greatest care and attention. We entertain little doubt that this book will become what its author hoped it might be-

some, a manual for daily reference and consultation by the student and the general practitioner. The work is marked by that correctness, clearness, and precision of style which distinguish all the productions of the learned author.—*British and Foreign Medical Review.*

PREPARING.

A MANUAL OF DENTAL PRACTICE.

BY JOHN TOMES, F. R. S.

Illustrated with numerous Engravings on Wood.

LECTURES ON THE

PRINCIPLES AND PRACTICE OF SURGERY.

BY BRANSBY B. COOPER, F. R. S.

Senior Surgeon to Guy's Hospital, &c.

*In one large octavo volume of seven hundred and fifty pages.*

For twenty-five years Mr. Bransby Cooper has been surgeon to Guy's Hospital; and the volume before us may be said to consist of an account of the results of his surgical experience during that long period. We cordially recommend Mr. Bransby Cooper's Lectures

as a most valuable addition to our surgical literature, and one which cannot fail to be of service both to students and to those who are actively engaged in the practice of their profession.—*The Lancet.*

CLINICAL LECTURES ON SURGERY. By Sir BENJAMIN C. BRODIE, M. D., &c. One vol. 8vo., cloth. 350 pp.

BY THE SAME AUTHOR.

SELECT SURGICAL WORKS. One vol. 8vo. sheep, containing Clinical Lectures on Surgery, Diseases of the Joints, and Diseases of the Urinary Organs.

A TREATISE ON DISEASES OF THE BONES. By EDWARD STANLEY. In one octavo volume, extra cloth.

THE NATURE AND TREATMENT OF DEAFNESS AND DISEASES OF THE EAR, and the Treatment of the Deaf and Dumb. By William Duffon, M. R. C. S. In one small 12mo. volume, cloth.

and on Diseases and Accidents requiring Operations. By ROBERT LISTON. Edited, with numerous Additions and Alterations, by T. D. MUTTER, M. D. In one large and handsome octavo volume of 566 pages, with 216 wood-cuts.

LECTURES ON THE OPERATIONS OF SURGERY,



## AN ILLUSTRATED LIBRARY OF OBSTETRICAL SCIENCE.

Illustrated  
with  
more than  
fifty  
large and  
splendid plates  
and  
wood-cuts  
through  
the text.  
In  
one large  
imperial octavo  
volume,  
strongly bound.



Foot Presentation.

THE PRINCIPLES AND PRACTICE OF  
OBSTETRIC MEDICINE AND SURGERY.

IN REFERENCE TO THE PROCESS OF PARTURITION.

BY FRANCIS H. RAMSBOTHAM, M. D.,  
Physician to the Royal Maternity Charity, &c. &c.

A NEW AND IMPROVED AMERICAN, FROM THE LAST AND REVISED LONDON EDITION.

*From Prof. Hodge of the University of Pennsylvania.*

This work needs no commendation from me, receiving as it does, the unanimous recommendation of the British periodical press, as the standard work on Midwifery; "chaste in language, classical in composition, happy in point of arrangement, and abounding in most interesting illustrations."

To the American public, therefore, it is most valuable, from its intrinsic undoubted excellence, and as being the best authorized exponent of British Midwifery. Its circulation will, I trust, be extensive throughout our country.

All the organs concerned in the process of parturition, and every step of this process, in all its different forms, are illustrated with admirable plates. We recommend the student who desires to master this difficult subject with the least possible trouble, to possess himself at once of a copy of this work.—*American Journal of the Medical Sciences.*

This new edition of Dr. Rambotham's work forms one of the most complete and thoroughly useful treatises on midwifery with which we are acquainted. One of the best guides that the student or young practitioner can follow.—*British and Foreign Medical Review.*

It is the book of midwifery for students; clear, but not too minute in its details and sound in its practical instructions. It is so completely illustrated by plates (admirably chosen and executed), that the student must be stupid indeed who does not understand the details of this branch of the science, so far at least as description can make them intelligible.—*Dublin Journal of Medical Science.*

This work is by far the best that has appeared in this country for those who seek practical information upon midwifery, conveyed in a clear and concise style.—*London Medical Journal.*

The Student's Text-Book of Midwifery.

ON THE  
THEORY AND PRACTICE OF MIDWIFERY.

BY FLEETWOOD CHURCHILL, M. D., M. R. I. A., &c. &c.

WITH NOTES AND ADDITIONS,

BY D. FRANCIS CONDIE, M. D.,  
Author of "A Practical Treatise on Diseases of Children," &c.

A new  
American,  
from the  
last improved  
Dublin  
edition.



In one  
very handsome  
octavo .  
volume, of  
over  
five hundred pages,  
with  
one hundred and  
thirty-nine  
beautiful  
illustrations  
on wood.

*Nerves of the Gravid Uterus.*

To bestow praise on a book that has received such marked approbation would be superfluous. We need only say, therefore, that if the first edition was thought worthy of a favorable reception by the medical public, we can confidently affirm that this will be found much more so. The lecturer, the practitioner, and the student, may all have recourse to its pages, and derive from their perusal much interest and instruction in everything relating to theoretical and practical midwifery.—*Dublin Quarterly Journal of Medical Science.*

Few treatises will be found better adapted as a text-book for the student, or as a manual for the frequent consultation of the young practitioner.—*American Medical Journal.*

No work holds a higher position, or is more deserving of being placed in the hands of the tyro, the advanced student, or the practitioner.—*Medical Examiner.*

Previous editions, under the editorial supervision of Prof. R. M. Huston, have been received with marked favor, and they deserved it; but this, reprinted from a very late Dublin edition, carefully revised and brought up by the author to the present time, does present an unusually accurate and able exposition of every important particular embraced in the department of midwifery. \* \* The clearness, directness, and precision of its teachings, together with the great amount of statistical research which its text exhibits, have served to place it already in the foremost rank of works in this department of remedial science.—*N. O. Medical and Surgical Journal.*

This is certainly the most perfect system extant. It is the best adapted for the purposes of a text-book, and that which he whose necessities confine him to one book, should select in preference to all others.—*Southern Medical and Surgical Journal.*

A SYSTEM OF MIDWIFERY.

BY EDWARD RIGBY, M. D.,

Physician to the General Lying-in Hospital.

SECOND AMERICAN EDITION.

In one octavo volume, with Notes and Additional Illustrations.

# OBSTETRICS; THE SCIENCE AND THE ART.

BY CHARLES D. MEIGS, M. D.,


Professor of Midwifery, and Diseases of Women and Children, in the Jefferson Medical College of Philadelphia.  
SECOND EDITION REVISED, WITH ONE HUNDRED AND THIRTY-ONE ILLUSTRATIONS.

*In one large octavo volume, of 750 pages.*



*Use of the Fillet.*

The rapid demand for a second edition of this work is a sufficient evidence that it has supplied a desideratum of the profession, notwithstanding the numerous treatises on the same subject which have appeared within the last few years. Adopting a system of his own, the author has combined the leading principles of his interesting and difficult subject, with a thorough exposition of its rules of practice, presenting the results of long and extensive experience and of familiar acquaintance with all the modern writers on this department of medicine. As an American Treatise on Midwifery, which has at once assumed the position of a clas-



*Turning.*

sic, it possesses peculiar claims to the attention and study of the practitioner and student, while the numerous alterations and revisions which it has undergone in the present edition are shown by the great enlargement of the work, which is not only increased as to the size of the page, but also in the number. Among other additions may be mentioned.

A NEW AND IMPORTANT CHAPTER ON "CHILDBED FEVER."

By the same Author—(Now Ready.)

## A TREATISE ON ACUTE AND CHRONIC DISEASES OF THE NECK OF THE UTERUS.

With numerous Plates, drawn and colored from Nature, in the highest style of art.

*In one very handsome octavo volume, extra cloth.*

The object of the author in this work has been to present in a small compass the practical results of his long experience in this important and distressing class of diseases. The great changes introduced into practice, and the accessions to our knowledge on the subject, within the last few years, resulting from the use of the metroscope, brings within the ordinary practice of every physician numerous cases which were formerly regarded as incurable, and renders of great value a work like the present, combining practical directions for diagnosis and treatment with an ample series of illustrations, copied accurately from colored drawings made by the author, after nature. No such accurate delineations of the pathology of the neck of the uterus have heretofore been given, requiring, as they do the rare combination of physician and artist, and their paramount importance to the physician in whose practice such cases are frequent, is too evident to be dwelt upon, while in artistic execution they are far in advance of anything of the kind as yet produced in this country.

BY THE SAME AUTHOR.

OBSERVATIONS ON CERTAIN OF THE DISEASES OF YOUNG CHILDREN. In one handsome octavo volume, of 214 pages.

BY THE SAME AUTHOR. (*Preparing.*)

ON THE NATURE, SIGNS, AND TREATMENT OF PUERPERAL FEVER. In one handsome octavo volume.

*New and Improved Edition—(Now Ready.)*

## WOMAN; HER DISEASES AND REMEDIES.

A SERIES OF LETTERS TO HIS CLASS.

BY CHARLES D. MEIGS, M. D.,

Professor of Midwifery, and Diseases of Women and Children, in the Jefferson Medical College, Philadelphia.

THIRD EDITION, REVISED AND ENLARGED.

*In one large and handsome octavo volume of nearly seven hundred pages.*

The gratifying appreciation of his labors, as evinced by the exhaustion of two large impressions of this work within a few years, has not been lost upon the author, who has endeavored in every way to render it worthy of the favor with which it has been received. The opportunity thus afforded for a second revision has been improved, and the work is now presented as in every way superior to its predecessors, additions and alterations having been made whenever the advance of science has rendered them desirable. The typographical execution of the work will also be found to have undergone a similar improvement; by an increase in the size of the page, the additions have been incorporated without enlarging the bulk of the volume, and the work is now confidently presented as in every way worthy the position it has acquired as the standard American text-book on the Diseases of Females.

It contains a vast amount of practical knowledge, by one who has accurately observed and retained the experience of many years, and who tells the result in a free, familiar, and pleasant manner.—*Dublin Quarterly Journal.*

There is an off-hand fervor, a glow, and a warm-heartedness infecting the effort of Dr. Meigs, which is entirely captivating, and which absolutely hurries the reader through from beginning to end. Besides, the book teems with solid instruction, and it shows the very highest evidence of ability, viz: the clearness with which the information is presented. We know of no better test of one's understanding a subject than the evidence of the power of lucidly explaining it. The most elementary, as well as the obscurest subjects, under the pencil of Prof. Meigs, are isolated and made to stand out in such bold relief, as to produce distinct impressions upon the mind and memory of the reader.—*The Charleston Med. Journal.*

Professor Meigs has enlarged and amended this great work, for such it unquestionably is, having passed the ordeal of criticism at home and abroad, but been improved thereby; for in this new edition the author has introduced real improvements, and increased the value and utility of the book immeasurably. It presents so many novel, bright, and sparkling thoughts; such an exuberance of new ideas on almost every page, that we confess ourselves to have become enamored with the book and its author; and cannot withhold our congratulations from our Philadelphia conferees, that such a teacher is in their service. We regret that our limits will not allow of a more extended notice of this work, but must content ourselves with thus commending it as worthy of diligent perusal by physicians as well as students, who are seeking to be thoroughly instructed in the important practical subjects of which it treats.—*N. Y. Med. Gazette.*

~~~~~

A TREATISE ON THE DISEASES AND SPECIAL HYGIENE OF FEMALES.

BY COLOMBAT DE L'ISÈRE.

TRANSLATED FROM THE FRENCH, WITH ADDITIONS.

BY CHARLES D. MEIGS, M. D.,

Professor of Midwifery, and Diseases of Women and Children, in the Jefferson Medical College, Philadelphia.

A NEW AND REVISED EDITION.

In one octavo volume of over seven hundred pages, with numerous illustrations.



Bond's Placenta Forceps.

New and Improved Edition—(Now Ready.)

A PRACTICAL TREATISE ON
INFLAMMATION OF THE UTERUS,
 Its Cervix and Appendages;
 AND ON ITS CONNECTION WITH UTERINE DISEASE.
 BY HENRY BENNET, M. D.

Fourth
 American,
 from
 the Third
 and
 Revised
 London
 Edition.



Virgin Uterus.



Post Partum Uterus.

In
 one neat
 octavo
 volume, of
 over
 400 pages,
 with
 wood-cuts.

This edition has been carefully revised and altered, and various additions have been made, which render it more complete, and, if possible, more worthy of the high appreciation in which it is held by the medical profession throughout the world. A copy should be in the possession of every physician.—*Charleston Med. Journal and Review*, March, 1854.

So through the whole work we have evidence of added thought and study of new facts obtained, and of clearer views of treatment. The work is now a standard treatise. Most of our readers are undoubtedly familiar with its previous editions, and we can only call the attention of those who have not recognized these new views of uterine pathology to their real importance.—*Buffalo Medical Journal*, April, 1854.

The present revised and enlarged edition must be acknowledged the best treatise of the kind in the English language, and we know of none in the French or German that are equally valuable. It certainly is no small compliment to Dr. Bennet, that his treatise is published in the latter countries, and that a fourth edition has been called for and issued in this country. It must be admitted that Dr. Bennet's writings have served to mark a new era in the literature, if not in the pathological doctrines of this department of Medicine. In point of general accuracy and practical value, there certainly is no other treatise that equals this. No physician's library can be complete without the incomparably excellent work of Mr. Bennet.—*N. Y. Journal of Medicine*, May, 1854.

New Edition—Now Ready.

THE CAUSES AND TREATMENT OF ABORTION AND STERILITY;

Being the Result of an Extended Practical Inquiry into the Physiological and Morbid Conditions of the Uterus. By JAMES WHITEHEAD, F. R. C. S., &c. Second American Edition. In one octavo volume, extra cloth, of 368 pages.

A PRACTICAL TREATISE ON THE DISEASES PECULIAR TO WOMEN,


Illustrated by Cases derived from Hospital and Private Practice. By SAMUEL ASHWELL, M. D. With Additions by PAUL BECK GODDARD, M. D. Second American Edition. In one octavo volume of 520 pages.

CLINICAL MIDWIFERY;

Comprising the Histories of Five Hundred and Forty-five Cases of Difficult, Prematural, and Complicated Labor, with Commentaries. By ROBERT LEE, M. D., F. R. S., &c. From the Second London Edition. In one royal 12mo. volume, extra cloth, of 238 pages.

ON PARTURITION,

AND THE PRINCIPLES AND PRACTICE OF OBSTETRICS. By W. TYLER SMITH, M. D., Lecturer on Obstetrics in the Hunterian School of Medicine. In one large royal 12mo. volume of 400 pages.



Ricord's Speculum.

ON THE DISEASES OF WOMEN; INCLUDING DISEASES OF PREGNANCY AND CHILDBED.

BY FLEETWOOD CHURCHILL, M. D., M. R. I. A., &c.

A New American Edition, revised by the Author.

WITH NOTES AND ADDITIONS, BY D. FRANCIS CONDIE, M. D.

In one large octavo volume of nearly seven hundred pages.

From the Author's Preface.

In reviewing this edition, at the request of my American publishers, I have inserted several new sections and chapters, and I have added, I believe, all the information we have derived from recent researches; in addition to which the publishers have been fortunate enough to secure the services of an able and highly esteemed editor in Dr. Condie.

Former editions of this work have been noticed in previous numbers of the Journal. The sentiments of high commendation expressed in those notices, have only to be repeated in this; not from the fact that the profession at large are not aware of the high merits which this work really possesses, but from a desire to see the principles and doctrines therein contained more generally recognized, and more universally carried out in practice.—*N. Y. Journal of Medicine.*

We know of no author who deserves that approbation, on "the diseases of females," to the same extent that Dr. Churchill does. His, indeed, is the only thorough treatise we know of on the subject; and it may be commended to practitioners and students as a masterpiece in its particular department. The former editions of this work have been commended strongly in this journal, and they have won their way to an extended, and a well-deserved popularity. This fifth edition, before us, is well calculated to maintain Dr. Churchill's high reputation. It was revised and enlarged by the author, for his American publishers, and it seems to us that there is scarcely any species of

desirable information on its subjects that may not be found in this work.—*The Western Journal of Medicine and Surgery.*

We are gratified to announce a new and revised edition of Dr. Churchill's valuable work on the diseases of females. We have ever regarded it as one of the very best works on the subjects embraced within its scope, in the English language; and the present edition, enlarged and revised by the author, renders it still more entitled to the confidence of the profession. The valuable notes of Prof. Huston have been retained, and contribute, in no small degree, to enhance the value of the work. It is a source of congratulation that the publishers have permitted the author to be, in this instance, his own editor, thus securing all the revision which an author alone is capable of making.—*The Western Lancet.*

As a comprehensive manual for students, or a work of reference for practitioners, we only speak with common justice when we say that it surpasses any other that has ever issued on the same subject from the British press.—*The Dublin Quarterly Journal.*

~~~~~  
BY THE SAME AUTHOR.

## ON THE DISEASES OF INFANTS AND CHILDREN.

*In one large and handsome volume, of over six hundred pages.*

The present volume will sustain the reputation acquired by the author from his previous works. The reader will find in it full and judicious directions for the management of infants at birth, and a compendious, but clear account of the diseases to which children are liable, and the most successful mode of treating them. We must not close this notice without calling attention to the author's style, which is perspicuous and polished to a degree, we regret to say, not generally characteristic of medical works. We recommend the work of Dr. Churchill most cordially, both to students and practitioners, as a valuable and reliable guide in the treatment of the diseases of children.—*Am. Journ. of the Med. Sciences.*

We regard this volume as possessing more claims to completeness than any other of the kind with which we are acquainted. Most cordially and earnestly, therefore, do we commend it to our professional brethren, and we feel assured that the stamp of their approbation will in due time be impressed upon it. After an attentive perusal of its contents, we hesitate not to say, that it is one of the most comprehensive ever written upon the diseases of children, and that, for copiousness of reference, extent of research, and perspicuity of detail, it is scarcely to be equalled, and not to be excelled, in any language.—*Dublin Quarterly Journal.*

BY THE SAME AUTHOR.

ESSAYS ON THE PUERPERAL FEVER, AND OTHER DISEASES PECULIAR TO WOMEN. Selected from the writings of British Authors previous to the close of the Eighteenth Century. In one neat octavo volume, of about four hundred and fifty pages.

New and Improved Edition—(Just Issued.)

A PRACTICAL TREATISE ON THE  
DISEASES OF CHILDREN.

BY D. FRANCIS CONDIE, M. D.

Fourth Edition, Revised and Augmented.

In one large octavo volume of nearly seven hundred and fifty pages.

From the Author's Preface.

The demand for another edition has afforded the author an opportunity of again subjecting the entire treatise to a careful revision, and of incorporating in it every important observation recorded since the appearance of the last edition, in reference to the pathology and therapeutics of the several diseases of which it treats.

Every species of hypothetical reasoning has, as much as possible, been avoided. The author has endeavored throughout the work to confine himself to a simple statement of well-ascertained pathological facts, and plain therapeutical directions—his chief desire being to render it what its title imports it to be, A PRACTICAL TREATISE ON THE DISEASES OF CHILDREN.

We feel persuaded that the American medical profession will soon regard it not only as a very good, but as the VERY BEST "Practical Treatise on the Diseases of Children."—*American Medical Journal*.

One of the best works upon the diseases of children in the English language.—*Western Lancet*.

We pronounced the first edition to be the best work

on the diseases of children in the English language, and, notwithstanding all that has been published, we still regard it in that light.—*Medical Examiner*.

Taken as a whole, in our judgment, Dr. Condie's Treatise is the one from the perusal of which the practitioner in this country will rise with the greatest satisfaction.—*Western Journ. of Medicine and Surgery*

New and Revised Edition—(Now Ready.)

LECTURES ON THE  
DISEASES OF INFANCY AND CHILDHOOD.

BY CHARLES WEST, M. D.,

Physician to the Hospital for Sick Children, &c.

SECOND AMERICAN, FROM THE SECOND AND ENLARGED LONDON EDITION.

In one octavo volume of nearly five hundred pages.

In the preparation of the second edition of these Lectures, the whole work has been carefully revised. A few formulæ have been introduced and a minute alphabetical index has been appended, while additions, amounting altogether to fifty pages, have been made wherever I felt that more extended observation, or more careful reflection had enabled me to supply some of those deficiencies which I am well aware, are still far too numerous. The work now contains the result of 640 observations, and 199 post-mortem examinations, chiefly made among 16,276 children who came under my notice during the ten years of my connection with the Children's Infirmary in Lambeth.—*Author's Preface to the Second Edition*.

Dr. West has placed the profession under deep obligation by this able, thorough, and finished work upon a subject which almost daily taxes to the utmost the skill of the general practitioner. He has with singular felicity threaded his way through all the tortuous labyrinths of the difficult subject he has undertaken to elucidate, and has in many of the darkest corners left a light, for the benefit of succeeding travellers, which will never be extinguished. Not the least cap-

titivating feature in this admirable performance is its easy, conversational style, which acquires force from its very simplicity, and leaves an impression upon the memory, of the truths it conveys, as clear and refreshing as its own purity. The author's position secured him extraordinary facilities for the investigation of children's diseases, and his powers of observation and discrimination have enabled him to make the most of these great advantages.—*Nashville Medical Journal*.

WORKS BY WILLIAM P. DEWEES, M. D.

Twelfth Edition—(Just Issued.)

A COMPREHENSIVE SYSTEM OF MIDWIFERY.

ILLUSTRATED BY OCCASIONAL CASES AND MANY ENGRAVINGS.

In one octavo volume of six hundred pages.

Tenth Edition—(Just Issued.)

A TREATISE ON THE PHYSICAL AND MEDICAL TREATMENT OF CHILDREN.

In one octavo volume of five hundred and forty-eight pages.

Tenth Edition—(Just Issued.)

A TREATISE ON THE DISEASES OF FEMALES.

In one octavo volume of five hundred and thirty-two pages, with plates.

Much Improved Edition—(Now Ready.)

# A UNIVERSAL FORMULARY;

CONTAINING THE  
METHODS OF PREPARING AND ADMINISTERING  
OFFICINAL AND OTHER MEDICINES.

THE WHOLE ADAPTED TO PHYSICIANS AND PHARMACEUTISTS,

BY ROBERT E. GRIFFITH, M. D.,

SECOND EDITION, THOROUGHLY REVISED, WITH NUMEROUS ADDITIONS,

BY ROBERT P. THOMAS, M. D.,


Professor of Materia Medica in the Philadelphia College of Pharmacy.

In one large and handsome octavo volume, of 650 pages.

With Illustrations.

The speedy exhaustion of a large edition, and demand for a second, sufficiently show the position which this work has so rapidly attained as an authoritative and convenient book of reference for the physician and pharmacist. The opportunity thus afforded for its improvement has not been neglected. In its revision, Prof. Thomas (to whom this task has been confided in consequence of the death of the author), has spared no labor, in the hope of rendering it the most complete and correct work on the subject as yet presented to the profession. All the newly introduced articles of the *Materia Medica* have been inserted, such formulæ as had escaped the attention of the author have been added, numerous illustrative wood-cuts have been introduced, and the whole has been most carefully read and examined, to insure the absolute correctness, so indispensable in a work of this nature. The amount of these additions may be estimated from the fact that not only has the page been considerably enlarged, but the volume has also been increased by about seventy-five pages, while the arrangement of the formulæ and the general typographical execution will be found to have undergone great improvement. To the practitioner, its copious collection of all the forms and combinations of the articles of the *Pharmacopœia* render it an invaluable book of reference, while its very complete embodiment of officinal preparations of all kinds, derived from all sources, American, English, and Continental, make it an indispensable assistant to the apothecary.

Besides the FORMULARY, this work contains Tables of Weights and Measures, Specific Gravity, Temperature for certain Pharmaceutical Operations, Hydrometrical Equivalents, Specific Gravities of some of the Preparations of the *Pharmacopœias*, Relation between different Thermometrical Scales, Explanation of Abbreviations used in Formulæ, Vocabulary of Words used in Prescriptions, Observations on the Management of the Sick Room, Doses of Medicines, Rules for the Administration of Medicines, Management of Convalescence and Relapses, Dietetic Preparations not included in the Formulary, List of Incompatibles, Posological Table, Table of Pharmaceutical Names which differ in the *Pharmacopœias*, Officinal Preparations and Directions, and Poisons. The vast body of information contained throughout the volume is rendered easy of reference by three extended Indexes, one of Diseases and their Remedies, one of Pharmaceutical and Botanical Names, and a complete general Index.



Displacement Apparatus.

Dr. Griffith's Formulary is worthy of recommendation, not only on account of the care which has been bestowed on it by its estimable author, but for its general accuracy, and the richness of its details.—*Medical Examiner*.

A very useful work, and a most complete compendium on the subject of *materia medica*. We know of no work in our language, or any other, so comprehensive in all its details.—*London Lancet*.

Pre-eminent among the best and most useful compilations of the present day will be found the work

before us, which can have been produced only at a very great cost of thought and labor. We are not cognizant of the existence of a parallel work. Its value will be apparent to our readers from the sketch given of its contents. We strongly recommend it to all who are engaged either in practical medicine, or more exclusively with its literature.—*London Med. Gazette*.

A valuable acquisition to the medical practitioner, and a useful book of reference to the apothecary on numerous occasions.—*Amer. Journal of Pharmacy*.

**A PRIZE ESSAY ON THE USE OF ALCOHOLIC LIQUORS IN HEALTH AND DISEASE.** By WILLIAM B. CARPENTER, M. D., author of "Human Physiology," &c. New edition, with a Preface by D. F. CONDIE, M. D., and Explanations of Scientific Words. In one neat 12mo. volume. (*Now Ready*)

This new edition has been prepared with a view to an extended circulation of this important little work, which is universally recognized as the best exponent of the laws of physiology and pathology applied to the subject of intoxicating liquors, in a form suited both for the profession and the public. To secure a wider dissemination of its doctrines the publishers have done up copies in flexible cloth, suitable for mailing, which will be forwarded through the post-office, free, on receipt of fifty cents. Societies and others supplied in quantities for distribution at a liberal deduction.



Now Complete.

THE ELEMENTS OF  
MATERIA MEDICA AND THERAPEUTICS.

BY JONATHAN PEREIRA, M. D., F. R. S., &c.

Third American Edition, Enlarged and Improved by the Author.

INCLUDING

NOTICES OF MOST OF THE MEDICINAL SUBSTANCES IN USE IN THE CIVILIZED WORLD,

AND FORMING AN

ENCYCLOPÆDIA OF MATERIA MEDICA.

EDITED BY JOSEPH CARSON, M. D.,

Professor of Materia Medica and Pharmacy in the University of Pennsylvania, &c.

In two very large and handsome octavo volumes of 2100 pages on small type, with over four hundred and fifty illustrations; strongly bound in leather, with raised bands

VOLUME I.—Lately issued, contains the Inorganic Materia Medica, of 838 pages, with 145 cuts.

VOLUME II.—Now ready, embraces the Organic Materia Medica, and forms a very large octavo volume of 1250 pages, with two plates and three hundred handsome wood-cuts.



Buds of *Cavaens Sarsaparilla*.

The present edition of this valuable and standard work will enhance in every respect its well-deserved reputation. The care bestowed upon its revision by the author may be estimated by the fact that its size has been increased by about five hundred pages. These additions have extended to every portion of the work, and embrace not only the materials afforded by the recent editions of the Pharmacopœias, but also all the important information accessible to the care and industry of the author in treatises, essays, memoirs, monographs, and from correspondents in various parts of the globe. In this manner, the work comprises the most recent and reliable information respecting all the articles of the Materia Medica, their natural and commercial history, chemical and therapeutical properties, preparation, uses, doses, and modes of administration, brought up to the present time, with a completeness not to be met with elsewhere. A considerable portion of the work which preceded the remainder in London, has also enjoyed the advantage of a further revision by the author expressly for this country, and in addition to this the editor, Professor Carson, has made whatever additions appeared desirable to adapt it thoroughly to the U. S. Pharmacopœia, and to the wants of the American profession. An equal improvement will likewise be observable in every department of its mechanical execution. It is printed from new type, on good white paper, with a greatly extended and improved series of illustrations.

Gentlemen who have the first volume are recommended to complete their copies without delay. The first volume will no longer be sold separate.

The work is in every way improved and enlarged, being, in fact, a complete encyclopedia of materia medica, superior to any work on the subject hitherto published.—*Charleston Med. Journal and Review*.

The third edition of his "Elements of Materia Medica," is by far the most elaborate treatise in the English language, and will, while medical literature is cherished, continue a monument alike honorable to his genius, as to his learning and industry.—*American Journal of Pharmacy*, March, 1854.

When we remember that Philology, Natural History, Botany, Chemistry, Physics, and the Microscope, are all brought forward to elucidate the subject, one cannot fail to see that the reader has here a work worthy of the name of an encyclopedia of materia medica. Our own opinion of its merits is that of its editors, and also that of the whole profession, both of this and foreign countries, namely, "that in copiousness of details, in extent, variety, and accuracy of information, and in lucid explanation of difficult and recondite subjects, it surpasses all other works on materia me-

dica hitherto published." The illustrations have been increased, and this edition by Dr. Carson cannot well be regarded in any other light than that of a treasure which should be found in the library of every physician.—*New York Journal of Med. Science*, March, 1854.

None the less welcome, because so long expected, comes this second volume of the great treasury of pharmacological learning. When Dr. Pereira commenced the compilation of this superb work, he appears to have been resolved not only to exhaust the entire field of medical literature in regard to his topic, but to use every aid that could be furnished by botany, chemistry, and the microscope, and in addition, to cultivate a thorough knowledge of the Drug trade in all its various ramifications. Whatever Dr. P. asserts positively may be relied on implicitly. There never was a writer more cautious in his statements or who would cling to a difficult question more tenaciously until he had mastered it.—*N. J. Med. Reporter*, April, 1854.

It should be in the library and on the table of every practitioner.—*Dublin Quarterly Medical Journal*.

PEREIRA'S MATERIA MEDICA.  
(Specimen Wood-cut.)



Various Species of *Copaifera*. (Copaiva)

New and Improved Edition—(Just Issued.)

## GENERAL THERAPEUTICS AND MATERIA MEDICA; ADAPTED FOR A MEDICAL TEXT-BOOK.

BY ROBLEY DUNGLISON, M. D.,

Professor of Institutes of Medicine in Jefferson Medical College, Philadelphia.

With  
one hundred  
and  
eighty-seven  
illustrations.  
Fifth edition,  
Revised  
and  
Improved.  
In two very  
handsome  
octavo volumes,  
of  
1100 pages.



*Ipomoea purga (Jalap)*

The new editions of the United States Pharmacopœia and those of London and Dublin, have rendered necessary a thorough revision of this work. In accomplishing this, the author has spared no pains in rendering it a complete exponent of all that is new and reliable, both in the departments of Therapeutics and Materia Medica. The book has thus been somewhat enlarged, and a like improvement will be found in every department of its mechanical execution. As a convenient text-book for the student, therefore, containing within a moderate compass a satisfactory résumé of its important subject, it is again presented as even more worthy than heretofore of the very great favor which it has received.

In this work of Dr. Dunglison, we recognize the same untiring industry in the collection and embodying of facts on the several subjects of which he treats, that has heretofore distinguished him, and we cheerfully point to these volumes as two of the most interesting that we know of. In noticing the additions to this, the fourth edition, there is very little in the periodical or annual literature of the profession, published in the interval which has elapsed since the issue of the first, that has escaped the careful search of the author. As a book for reference, it is invaluable.—*Charleston Medical Journal and Review*.

It may be said to be the work now upon the subjects upon which it treats.—*Western Lancet*.

As a text-book for students, for whom it is particularly designed, we know of none superior to it.—*St. Louis Medical and Surgical Journal*.

It purports to be a new edition, but it is rather a new book, so greatly has it been improved, both in the amount and quality of the matter which it contains.—*N. O. Medical and Surgical Journal*.

We bespeak for this edition, from the profession, an increase of patronage over any of its former ones, on account of its increased merit.—*N. Y. Journal of Medicine*.

We consider this work unequalled.—*Boston Medical and Surgical Journal*.

BY THE SAME AUTHOR.

## NEW REMEDIES; WITH FORMULÆ FOR THEIR ADMINISTRATION.

Sixth Edition, with Extensive Additions.

In one very large octavo volume of over seven hundred and fifty pages.

One of the most useful of the author's works.—*Southern Medical and Surgical Journal*.

This well-known and standard book has now reached its sixth edition, and has been enlarged and improved by the introduction of all the recent gifts to therapeutics which the last few years have so richly produced, including the anæsthetic agents, &c. This elaborate and useful volume should be found in every medical library, for as a book of reference, for physicians, it is unsurpassed by any other work in existence, and the double index for diseases and for reme-

dis, will be found greatly to enhance its value.—*New York Medical Gazette*.

The great learning of the author, and his remarkable industry in pushing his researches into every source whence information is derivable, has enabled him to throw together an extensive mass of facts and statements, accompanied by full reference to authorities; which last feature renders the work practically valuable to investigators who desire to examine the original papers.—*The American Journal of Pharmacy*.

**MATERIA MEDICA AND THERAPEUTICS.**

INCLUDING THE PREPARATIONS OF  
**THE PHARMACOPŒIAS OF LONDON, EDINBURGH, DUBLIN, AND OF THE UNITED STATES.**  
 WITH MANY NEW MEDICINES.

BY  
**J. FORBES ROYLE, M.D., F.R.S., &c.**

EDITED BY **JOSEPH CARSON, M.D.,**  
 Professor of Materia Medica and Pharmacy in the University of Pennsylvania.

In one very  
 handsome octavo volume  
 of nearly  
 seven hundred pages, with about  
 one hundred beautiful  
 illustrations.

This work is, indeed, a most valuable one, and will fill up an important vacancy that existed between Dr. Pereira's most learned and complete system of Materia Medica, and the class of productions on the other extreme, which are necessarily imperfect from their small extent.—*British and Foreign Medical Review.*



*Althaea officinalis.*

New and Enlarged Edition to 1854—(Now Ready.)

**THE MEDICAL FORMULARY.**

Being a Collection of Prescriptions, derived from the Writings and Practice of many of the most eminent Physicians of America and Europe. Together with the usual Dietetic Preparations and Antidotes for Poisons. To which is added an Appendix, on the Endermic use of Medicines, and on the use of Ether and Chloroform. The whole accompanied with a few brief Pharmacœutic and Medical Observations. By **BENJAMIN ELLIS, M. D.** Tenth edition, revised and much extended, by **ROBERT P. THOMAS, M. D.,** Professor of Materia Medica in the Philadelphia College of Pharmacy, &c. In one neat octavo volume, extra cloth, of three hundred pages.

This work has received a very complete revision at the hands of the editor, who has made whatever alterations and additions the progress of medical and pharmaceutical science has rendered advisable, introducing fully the new remedial agents, and revising the whole by the latest improvements of the Pharmacopœia. To accommodate these additions, the size of the page has been increased, and the volume itself considerably enlarged, while every effort has been made to secure the typographical accuracy which has so long merited the confidence of the profession.

It will prove particularly useful to students and young practitioners, as the most important prescriptions employed in modern practice, which lie scattered through our medical literature, are here collected and conveniently arranged for reference.—*Charleston Med. Journal and Review.*

After an examination of the new matter and the

alterations, we believe the reputation of the work built up by the author, and the late distinguished editor, will continue to flourish under the auspices of the present editor, who has the industry and accuracy, and, we would say, conscientiousness requisite for the responsible task.—*American Journal of Pharmacy,* March, 1854.

**THE THREE KINDS OF COD-LIVER OIL,** comparatively considered, with their Chemical and Therapeutic Properties. By **D. L. DE JONGH, M. D.** Translated, with an Appendix and Cases, by **EDWARD CAREY, M. D.** To which is added an article on the subject from "Dun- glison on New Remedies." In one small 12mo. volume, extra cloth.

**PREPARING.**

**ELEMENTS OF GENERAL AND SPECIAL THERAPEUTICS.** By **ALFRED STILLE, M. D.** In one large and handsome volume.

In one  
very large  
and  
handsome  
octavo volume,  
of over  
1000 pages,  
strongly bound  
in leather.



With  
two  
hundred  
and  
thirteen  
engravings  
on  
wood.

*Cucumis Colocynthis.*

## A DISPENSATORY.

Or Commentary on the Pharmacopias of Great Britain and the United States;

COMPRISING

THE NATURAL HISTORY, DESCRIPTION, CHEMISTRY, PHARMACY, ACTION, USES, AND DOSES OF  
THE ARTICLES OF THE MATERIA MEDICA.

BY ROBERT V. CHRISTISON, M. D., V. P. R. S. E., & Co.

Second Edition, Revised and Improved.

With a Supplement containing the most important New Remedies.

WITH COPIOUS ADDITIONS, BY R. EGLESFELD GRIFFITH, M. D.

It is not needful that we should compare it with the other pharmacopias extant, which enjoy and merit the confidence of the profession: it is enough to say that it appears to us as perfect as a Dispensatory, in the present state of pharmaceutical science, could be made. If it omits any details pertaining to this branch of knowledge which the student has a right to expect

in such a work, we confess the omission has escaped our scrutiny. We cordially recommend this work to such of our readers as are in need of a Dispensatory. They cannot make choice of a better.—*Western Journal of Medicine and Surgery.*

There is not in any language a more complete and perfect Treatise.—*New York Annalist.*

## MEDICAL BOTANY;

OR, DESCRIPTIONS OF THE MORE IMPORTANT PLANTS USED IN MEDICINE.

With their History, Properties, and Modes of Administration.

BY R. EGLESFELD GRIFFITH, M. D.

WITH THREE HUNDRED AND THIRTY-FOUR ILLUSTRATIONS.

In one large and handsome 8vo. volume, of over seven hundred pages.

One of the few books which supply a positive deficiency in our medical literature.—*Western Lancet.*

We hope the day is not distant when this work will not only be a text-book in every medical school and college in the Union, but find a place in the library of every private practitioner.—*N. Y. Journal of Medicine.*

One of the greatest acquisitions to American medical literature. It should by all means be introduced, at the very earliest period, into our medical schools, and occupy a place in the library of every physician in the land.—*Southwestern Medical Advocate.*

Admirably calculated for the physician and student; we have seen no work which promises greater advantage to the profession.—*N. O. Med. and Surg. Journal.*



*Datura Stramonium.*

# PRACTICAL PHARMACY;

THE ARRANGEMENTS, APPARATUS, AND MANIPULATIONS OF THE PHARMACEUTICAL SHOP AND LABORATORY.

BY FRANCIS MOHR, PH. D.,

Assessor Pharmaciae of the Royal Prussian College of Medicine, Coblenz;

AND THEOPHILUS REDWOOD,

Professor of Chemistry and Pharmacy to the Pharmaceutical Society of Great Britain.


EDITED, WITH EXTENSIVE ADDITIONS,

BY WILLIAM PROCTER, JR.,

Professor of Pharmacy in the Philadelphia College of Pharmacy.

Illustrated by five hundred Engravings on Wood.

In one large and very handsome octavo volume, of nearly six hundred pages.



Apparatus for Spreading Plasters.

This work will be found of great value, not only to apothecaries, but also to physicians who live at a distance from competent pharmacutists. Comprehending, as it does, all the manipulations and operations of preparing and dispensing medicines, from the extraction of a stopper or the tying of a cork, to the most complicated and delicate processes of pharmacy, illustrated at every step with a profusion of engravings, it may be regarded as an indispensable assistant to the druggist and country practitioner, while its very moderate price places it within the reach of all.

It is a book, however, which will be in the hands of almost every one who is much interested in pharmaceutical operations, as we know of no other publication so well calculated to fill a void long felt.—*Medical Examiner*.

This work is very full and complete, and details, in a style uncommonly clear and lucid, not only the

more complicated and difficult processes, but those not less important ones, the most simple and common.—*Buffalo Medical Journal*.

The country practitioner who is obliged to dispense his own medicines, will find it a most valuable assistant.—*Monthly Journal and Retrospect*.

SYNOPSIS OF THE COURSE OF LECTURES ON MATERIA MEDICA AND PHARMACY, delivered in the University of Pennsylvania. By JOSEPH CARSON, M. D., Professor of Materia Medica and Pharmacy in the University of Pennsylvania. In one very neat octavo volume, of 208 pages.

A DISPENSATORY AND THERAPEUTICAL REMEMBRANCER, Comprising the entire lists of Materia Medica, with every Practical Formula contained in the three British Pharmacopœias. By JOHN MAYNE, M. D., M. R. C. S. Edited, with the addition of the Formulæ of the United States Pharmacopœia, by R. EGLERFELD GRIFFITH, M. D. In one 12mo. volume, extra cloth, of over 300 large pages.

NOW READY.

# HANDBOOK OF CHEMISTRY,

## THEORETICAL, PRACTICAL, AND TECHNICAL.

BY F. A. ABEL AND C. L. BLOXAM.

WITH A PREFACE BY A. W. HOFMANN, M. D.

And Numerous Illustrations on Wood.

*In one large and handsome octavo volume of nearly seven hundred large pages.**Preparation of Carbonic Oxide.*

The object of this work is to present such a general view of PRACTICAL CHEMISTRY as shall enable the student to acquire, with the least possible expenditure of time, a knowledge which will either suffice for the ordinary applications of chemistry to the useful arts, or will serve as a sound basis for the education of the professional chemist. It will thus be seen to fill a void among chemical text-books, there being no modern work in the language presenting a complete view of chemical science in this aspect. The innumerable and continually multiplying applications of the science in medicine and the useful arts, will therefore render this a work for daily reference by those who wish to understand their profession thoroughly, or to keep on a level with the practical improvements of the day.

A very condensed summary of the contents is subjoined:—

**INTRODUCTION.**—Specific Gravity—Definitions of Chemical Terms—Equivalents—Affinity—Nomenclature and Notation—Phenomena relating to the Physical condition of Bodies.

**CHEMICAL MANIPULATION.**—Apparatus for the Preparation of Gases with the aid of Heat—Method of fitting up Apparatus for the Generation of Gases—Apparatus for the Generation of Gases at ordinary Temperatures—Arrangements for Purifying Gases—Apparatus for Collecting Gases—Combustion of Gases—Combustion in Gases—Transference of Gases—Measurement of Gases—Solution of Gases—Condensation of Gases—Distillation and Sublimation—Disintegration—Solution—Infusion—Digestion—Filtration—Evaporation—Crystallization—Ignition—Fusion—Sources of Heat in the Laboratory—Use of the Blowpipe—Gas-blowing.

**ELEMENTARY CHEMISTRY.**—Classification—Oxygen—Nitrogen—Chlorine—Bromine—Iodine—Fluorine—Sulphur—Selenium—Phosphorus—Carbon—Boron—Silicon—General Considerations on the Metals—Potassium—Sodium—Lithium—Ammonium—Barium—Strontium—Calcium—Magnesium—Aluminum—Glucinum—Thorium—Yttrium—Erbium—Terbium—Cerium—Lanthanum—Didymium—Zirconium—Chromium—Uranium—Iron—Manganese—Zinc—Nickel—Cobalt—Vanadium—

Cadmium—Copper—Bismuth—Gold—Platinum—Palladium—Rhodium—Iridium—Osmium—Ruthenium—Tin—Antimony—Arsenic—Tungsten—Molybdenum—Tellurium—Titanium—Mercury—Lead—Silver—Tantalum.

**ANALYTICAL CHEMISTRY.**—Definitions of Analysis—Analytical Classification of Metals—Reactions of the Oxides of Metals—Analytical Classification of Inorganic Acids—Reactions of Inorganic Acids—Analytical Classification of Organic Acids—Reactions of Organic Acids—Systematic Course of Qualitative Analysis—Quantitative Analysis—Operations in Quantitative Analysis—Special Methods of Quantitative Analysis; common salt, phosphate of soda, heavy spar, Rochelle salt, marble, limestones, alum, chrome-iron-ore, clay, wavelite, iron-ores, manganeseiferous spathic iron-ore, sulphate of copper, calamine, copper pyrites, tartar-emetic, pewter, fusible alloy, type-metal, brass, bronze, gun-metal, German silver, Britannia metal, standard gold, standard silver, amalgams, spels—cobalt, valuation of manganese ores—Chlorimetry—Alkalimetry—Acidimetry—Analysis of Gunpowder, glass, cast-iron, ultramarine—Separation of chlorine, bromine, and iodine—Analysis of Mineral waters—Soils—Ashes of Vegetables—Ashes of Animals.

**MEDICAL CHEMISTRY;** for the Use of Students and the Profession; being a Manual of the Science, with its Applications to Toxicology, Physiology, Therapeutics, Hygiene, &c. &c. By D. PEREIRA GARDNER, M. D. In one neat volume, royal 12mo., of 400 pages, with numerous wood-cuts.

New and Improved Edition, to 1854—(Now Ready.)

# ELEMENTARY CHEMISTRY;

THEORETICAL AND PRACTICAL.

BY GEORGE FOWNES, F. R. S.,  
Late Professor of Practical Chemistry in University College, London.


EDITED, WITH ADDITIONS,

BY ROBERT BRIDGES, M. D.,  
Professor of Chemistry in the Philadelphia College of Pharmacy, &c.


A NEW AMERICAN, FROM THE LAST AND REVISED LONDON EDITION.



Uric Acid Calculus.



Oxalate of Lime Calculus.



Apparatus for Liquefying Carbonic Acid.

With one hundred and eighty-one Illustrations.

In one large royal 12mo. volume, of over 550 pages, clearly printed on small type.

The lamented death of the author has caused the revision of this edition to pass into the hands of those distinguished chemists, H. Bence Jones and A. W. Hofmann, who have fully sustained its reputation by the additions which they have made, more especially in the portion devoted to Organic Chemistry, considerably increasing the size of the volume. This labor has been so thoroughly performed, that the American Editor has found but little to add, his notes consisting chiefly of such matters as the rapid advance of the science has rendered necessary, or of investigations which had apparently been overlooked by the author's friends.

The volume is therefore again presented as an exponent of the most advanced state of chemical science, and as not unworthy a continuation of the marked favor which it has received as an elementary text-book.

We know of no better text-book, especially in the difficult department of organic chemistry, upon which it is particularly full and satisfactory. We would recommend it to preceptors as a capital "office book" for their students who are beginners in Chemistry. It is copiously illustrated with excellent wood-cuts, and altogether admirably "got up."—*N. J. Med. Reporter*, March, 1854.

A standard manual, which has long enjoyed the reputation of embodying much knowledge in a small space. The author has achieved the difficult task of condensation with masterly tact. His book is concise without being dry, and brief without being too dogmatical or general.—*Virginia Med. and Surg. Journal*.

The work of Dr. Fownes has long been before the public, and its merits have been fully appreciated as the best text-book on Chemistry now in existence. We do not, of course, place it in a rank superior to the works of Brande, Graham, Turner, Gregory, or Gmelin, but we say that, as a work for students, it is preferable to any of them.—*London Journ. of Medicine*.

We unhesitatingly recommend it to medical students.—*N. W. Med. and Surg. Journal*, February, 1854.

One of the best elementary works on Chemistry accessible to the American and English student.—*N. Y. Journal of Medicine*, March, 1854.

We can safely commend it as one of the most reliable and comprehensive though brief works on this important and interesting department of medical science. No man is better qualified than the late distinguished professor of Practical Chemistry in University College, London, to write the book that was needed for students on this subject, and none, in our opinion, have succeeded so well.—*Ky. Medical Reporter*, March, 1854.

This is one of the most popular text-books with medical students, who are good judges of merit. It contains enough of Physics, Elementary Chemistry, Organic Chemistry, and general principles, to make it peculiarly adapted to their wants.—*Southern Medical and Surgical Journal*, Augusta.



New and Enlarged Edition.

Second American  
from  
an entirely  
Revised  
and  
greatly enlarged  
English  
edition.  
With  
numerous  
engravings  
on wood.

*Preparation of Ammonia.*

**ELEMENTS OF CHEMISTRY;**  
INCLUDING  
THE APPLICATIONS OF THE SCIENCE IN THE ARTS.  
BY THOMAS GRAHAM, F. R. S.

EDITED, WITH NOTES, BY ROBERT BRIDGES, M. D.

PART I. (lately issued), large 8vo., 430 pages, 185 Illustrations.

PART II. completing the work. (Preparing.)

A PRACTICAL HANDBOOK  
OF  
MEDICAL CHEMISTRY.

BY JOHN E. BOWMAN,  
Demonstrator of Chemistry in King's College, London, &c.

With numerous Illustrations on Wood.

In one  
neat royal 12mo. volume, of about  
three hundred pages.

*Mucus and Epithelium in Urine.*

BY THE SAME AUTHOR.

AN INTRODUCTION TO PRACTICAL CHEMISTRY;  
INCLUDING ANALYSIS.

In  
one neat  
royal 12mo. volume  
of 300 pages,  
with  
about 100  
illustrations.

*Reduction of Metallic Oxides.*

New and Improved Edition—(Just Issued.)

## MEDICAL JURISPRUDENCE.

BY ALFRED S. TAYLOR, M. D., F. R. S.,

Lecturer on Medical Jurisprudence and Chemistry in Guy's Hospital, &c.

THIRD AMERICAN, FROM THE FOURTH AND IMPROVED ENGLISH EDITION.

With Notes and References to American Decisions,

BY EDWARD HARTSHORNE, M. D.

In one large octavo volume of about seven hundred pages.

In the preparation of the English edition, from which this has been printed, the author has found it necessary to revise the whole of the chapters, as well as to make numerous alterations and additions, together with references to many recent cases of importance. A Glossary has also been added for the convenience of those whose studies have not been directed specially to this subject. The notes of the American editor embrace the additions formerly made by Dr. Griffith, who revised the work on its first appearance in this country, together with such new matter as his experience and the progress of the science have shown to be advisable. The work may therefore be regarded as fully on a level with the most recent discoveries, and worthy of the reputation which it has acquired as a complete and compendious guide for the physician and lawyer.

We know of no work on Medical Jurisprudence which contains in the same space anything like the same amount of valuable matter.—*N. Y. Journal of Medicine.*

The American editor has appended several important facts, the whole constituting by far the best, most reliable, and interesting treatise on Medical Jurisprudence, and one that we cannot too strongly recommend to all who desire to become acquainted with the true and correct exposition of this department of medical literature.—*Northern Lancet.*

No work upon the subject can be put into the hands of students either of law or medicine which will engage them more closely or profitably; and none could be offered to the busy practitioner of either calling, for the purpose of casual or hasty reference, that would be more likely to afford the aid desired. We therefore recommend it as the best and safest manual for daily use.—*American Journal of Medical Sciences.*

We have heretofore had reason to refer to it in terms of commendation, and need now only state that in the edition before us, the author has completely revised the whole work, making many additions and

alterations, and brought it fully up to the present state of knowledge. The task of the American editor has been to present all the important facts and cases that have recently occurred in our own country, bearing on the subjects treated of. No better work can be placed in the hands of the physician or jurist.—*St. Louis Medical and Surgical Journal.*

So well is this work known to the members both of the medical and legal professions, and so highly is it appreciated by them, that it cannot be necessary for us to say a word in its commendation: its having already reached a fourth edition being the best possible testimony in its favor. The author has obviously subjected the entire work to a very careful revision.—*British and Foreign Med.-Chirurgy. Review.*

It is not excess of praise to say that the volume before us is the very best treatise extant on Medical Jurisprudence. In saying this, we do not wish to be understood as detracting from the merits of the excellent works of Beck, Ryan, Trull, Guy, and others; but in interest and value we think it must be conceded that Taylor is superior to anything that has preceded it.—*N. W. Medical and Surgical Journal.*

BY THE SAME AUTHOR.

## ON POISONS,

IN RELATION TO MEDICAL JURISPRUDENCE AND MEDICINE.

EDITED, WITH NOTES AND ADDITIONS,

BY R. E. GRIFFITH, M. D.

In one large octavo volume of nearly seven hundred pages.

The most elaborate work on the subject that our literature possesses.—*British and Foreign Medico-Chirurgical Review.*

It contains a vast body of facts, which embrace all that is important in toxicology, all that is necessary to the guidance of the medical jurist, and all that can be desired by the lawyer.—*Medico-Chirurgy. Review.*

One of the most practical and trustworthy works on Poisons in our language.—*Western Journ. of Med.*

It is, so far as our knowledge extends, incomparably the best upon the subject; in the highest degree creditable to the author, entirely trustworthy, and indispensable to the student and practitioner.—*N. Y. Analyst.*

LETTERS TO A CANDID INQUIRER ON ANIMAL MAGNETISM. Description and Analysis of the Phenomena. Details of Facts and Cases. By WILLIAM GREGORY, F. R. S. E., Professor of Chemistry in the University of Edinburgh, &c. In one neat volume, royal 12mo., extra cloth.

HUMAN HEALTH; or, the Influence of Atmosphere and Locality, Change of Air and Climate, Seasons, Food, Clothing, Bathing, Exercise, Sleep, &c. &c., on Healthy Man; constituting Elements of Hygiene.

By ROWLEY DUNGLISON, M. D., Professor of the Institutes of Medicine in the Jefferson Medical College, Philadelphia. Second edition, with many modifications and additions. In one octavo volume, of 464 pages.

THE LAWS OF HEALTH IN RELATION TO MIND AND BODY. A Series of Letters from an old Practitioner to a Patient. By LEONEL JOHN BRALL, M. R. C. S., &c. In one handsome volume, royal 12mo., extra cloth.

Just Issued.

# THE BOOK OF NATURE

AN ELEMENTARY INTRODUCTION TO  
 PHYSICS, MINERALOGY,  
 ASTRONOMY, GEOLOGY,  
 CHEMISTRY, BOTANY,

BY FRIEDRICH SCHOEDLER,

Professor of the Natural Sciences at Worms,


First American Edition, with a Glossary and other Additions and Improvements

TRANSLATED FROM THE SIXTH GERMAN EDITION


BY HENRY MEDLOCK, F. C. S.

Profusely  
 Illustrated  
 with nearly  
 seven hundred  
 beautiful  
 engravings on  
 wood.

In one large and  
 handsome  
 crown octavo  
 volume,  
 of 692 pages,  
 extra cloth.



Machinery of a Watch (I)



Fecundation of Plants (Botany).



INAMUS rymond

**THE BOOK OF NATURE—(Continued.)**

To accommodate those who desire to use the separate portions of this work, the publishers have prepared an edition in parts, as follows, which may be had singly, neatly done up in flexible cloth, at very reasonable prices.

|                                |                                    |                                                |                                   |
|--------------------------------|------------------------------------|------------------------------------------------|-----------------------------------|
| <b>Natural Philosophy,</b>     | 114 pages, with 149 illustrations. | <b>Botany,</b>                                 | 98 pages, with 176 illustrations. |
| <b>Astronomy,</b>              | 64 " " 51 "                        | <b>Zoology and Physiology,</b>                 | 106 pages, with 84 "              |
| <b>Chemistry,</b>              | 110 " " 48 "                       | <b>Introduction, Glossary, Index, &amp;c.,</b> | 96 pages.                         |
| <b>Mineralogy and Geology,</b> | 104 pages, with 167 "              |                                                |                                   |

The necessity of some acquaintance with the Natural and Physical Sciences is now so universally admitted in all thorough education, while the circle of facts and principles embraced in the study has enlarged so rapidly, that a compendious Manual like the *BOOK OF NATURE* cannot fail to supply a want frequently felt and expressed by a large and growing class.

The reputation of the present volume in England and Germany, where repeated editions have been rapidly called for, is sufficient proof of the author's success in condensing and popularizing the principles of his numerous subjects. The publishers therefore would merely state that, in reproducing the work, they have spared no pains to render it even better adapted to the American student. It has been passed through the press under the care of a competent editor, who has corrected such errors as had escaped the attention of the English translator, and has made whatever additions appeared necessary to bring it completely on a level with the existing state of science. These will be found principally in the sections on Botany and Geology; especially the latter, in which references have been made to the numerous and systematic Government surveys of the several States, and the whole adapted to the nomenclature and systems generally used in this country. A copious Glossary has been appended, and numerous additional illustrations have been introduced wherever the elucidation of the text appeared to render them desirable.

It is therefore confidently presented as an excellent Manual for the private student, an indispensable work for District, School, and Township Libraries, or as a complete and thorough Class-book for collegiate and academical use.

This volume, as its title shows, covers nearly all the sciences, and embodies a vast amount of information for instruction. No other work that we have seen presents the reader with so wide a range of elementary knowledge, with so full illustrations, at so cheap a rate.—*Sullivan's Journal*.

Written with remarkable clearness, and scrupulously correct in its details.—*Mining Journal*.

The most complete and copious *résumé* of the sciences that has ever been offered to the public in a form so usable and cheap.—*Glasgow Citizen*.

It is utterly impossible in the space we have at com-

mand, to make our readers acquainted with the full value of this admirable volume.—*Glasgow Herald*.

Dr. Schoedler's work, as described by Liebig, is at once the most useful and beautiful book of the class to which it belongs. It is a valuable addition to our scientific literature. The wood-cuts and illustrations are numerous, and remarkably well executed.—*Economist*.

His expositions are most lucid. There are few who will not follow him with pleasure as well as with profit through his masterly exposition of the principles and primary laws of science. It should certainly be made a class-book in schools.—*Critic*.

**ASPECTS OF NATURE**

IN DIFFERENT LANDS AND DIFFERENT CLIMATES. With Scientific Elucidations. By ALEXANDER VON HUMBOLDT. Translated by Mrs. SABINE. In one large and handsome royal 12mo. volume, extra cloth.

**AMERICAN ORNITHOLOGY.**

By CHARLES BONAPARTE, Prince of Canino. In four very handsome folio volumes, half bound, with splendid colored plates.

**ZOOLOGICAL RECREATIONS.**

By W. T. BRODERIP, F. R. S. From the second London edition. In one neat royal 12mo. volume, extra cloth.

**CHEMISTRY OF THE FOUR SEASONS;**

SPRING, SUMMER, AUTUMN, AND WINTER. By THOMAS GRIFFITH. In one handsome volume, large royal 12mo., extra cloth, with numerous illustrations.

**ETHNOGRAPHY AND PHILOLOGY**

OF THE UNITED STATES EXPLORING EXPEDITION. By HORATIO HALE. In one large imperial 4to. volume, beautifully printed, and strongly bound in extra cloth.

THE PHYSICAL SCIENCES—(Now Complete.)  
 HANDBOOKS OF  
 NATURAL PHILOSOPHY AND ASTRONOMY.

BY DIONYSIUS LARDNER, D. C. L.,

Formerly Professor of Natural Philosophy and Astronomy in University College, London.

This valuable work is now complete, consisting of three volumes or courses, as follows:—

**First Course.**

**Mechanics, Hydrostatics, Hydraulics, Pneumatics, Sound, and Optics.**

In one large royal 12mo. volume, of 750 pages, with 424 illustrations.

**Second Course.**

**Heat, Magnetism, Common Electricity, and Voltaic Electricity.**

In one volume, royal 12mo., of four hundred and fifty pages, with 244 illustrations.

**Third Course.**

**Astronomy and Meteorology.**

In one volume, royal 12mo., of nearly 800 pages, with 37 plates, and over 200 illustrations.

These volumes  
 can be had  
 either separately or  
 in uniform sets,  
 containing in all about  
 two thousand pages,  
 with neatly  
 one thousand illustrations  
 on steel and wood.



*Aurora Borealis.*

To accommodate those who desire separate treatises on the leading departments of Natural Philosophy, the First Course may also be had divided in three portions, as follows:—

PART I.—MECHANICS.

“ II.—HYDROSTATICS, HYDRAULICS, PNEUMATICS, and SOUND.

“ III.—OPTICS.

It will thus be seen that this work furnishes either a complete course of instruction on these subjects, or separate treatises on all the different branches of Natural Science.

From among a vast number of recommendatory notices the publishers submit a few.

Nothing could be better for the purposes intended.—  
*British Quarterly Review.*

The “First Course” of Natural Philosophy has been already noticed by us. The “Second” merits equal praise. Dr. Lardner’s professional abilities are well known in our colleges and academies; and it is a great felicity to have, from such a source, complete scientific manuals. These handbooks are prepared with a direct view to the wants of those who wish to obtain a knowledge of physical science, without minute mathematical details. While this does not impair the work for the scholar, it brings it within the reach of the less favored, but more aspiring classes of the community; a more benevolent exercise of elevated talent cannot be imagined.—*Southern Methodist Quarterly Review.*

This “Handbook,” then, is an honest addition to our scientific literature. It is not the hasty production of a man who has been skimming the surface of the pool, and fudging the hem of the garment, but

the matured production of an accomplished mathematician, who has devoted his life to the study of physics, who has soared high and dived deep, who has traversed the rough places as well as the smooth, and than whom there can be no safer and more intelligent guide through those pleasant paths which belong to popular science.—*London Literary Gazette.*

The author, celebrated for his skill and learning in Physical and Mechanical Science, has composed this work for “the satisfaction of those who desire to obtain a knowledge of the elements of Physics, without pursuing them through their mathematical consequences and details.” On a cursory examination, we should think this volume admirably suited to carry out the author’s intentions. The principles are stated in precise and terse language, and the illustrations are happily given.—*N. Y. Churchman.*

One of the best popular scientific works that has yet appeared.—*English Journal of Education.*

## OUTLINES OF ASTRONOMY.

BY SIR JOHN F. W. HERSCHEL, BART., F. R. S., &c.

A new Edition.

WITH NUMEROUS PLATES AND WOOD-CUTS.

In one very handsome crown octavo volume, of over five hundred and fifty pages.

We now take leave of this remarkable work, which we hold to be, beyond a doubt, the greatest and most remarkable of the works in which the laws of astronomy and the appearance of the heavens are described to those who are not mathematicians nor observers, and recalled to those who are. It is the reward of men who can descend from the advancement of knowledge to care for its diffusion, that their works are essential to all, that they become the manuals of the proficient as well as the text-books of the learner.—*Athenæum*.

There is perhaps no book in the English language on the subject, which, whilst it contains so many of the facts of Astronomy (which it attempts to explain with as little technical language as possible), is so attractive in its style, and so clear and forcible in its illustrations.—*Evangelical Review*.

Probably no book ever written upon any science, embraces within so small a compass an entire epitome of everything known within all its various departments, practical, theoretical, and physical.—*Examiner*.

## THE GEOLOGICAL OBSERVER.

BY SIR HENRY T. DE LA BECHE, C. B., F. R. S., &c.

WITH OVER THREE HUNDRED WOOD-CUTS.

In one large and handsome octavo volume, of seven hundred pages.

The object which the author has proposed to himself in this work has been to afford a general view of the chief points of the science, such as existing observations would lead us to infer were established, to show how the correctness of such observations may be tested, and to sketch the direction in which they may apparently be extended. To accomplish this, he has entered into a minute detail of the results of geological investigations throughout the earth's surface, and, without confining himself to a mere detail of facts, he has endeavored to explain the causes, as well as the results of geological action. The assistance of numerous diagrams and illustrations has also been resorted to, to make manifest to the student the various forms of geological phenomena, and to guide him in the observations which he may desire to make for himself. To exemplify the manner in which this plan has been carried out, a very brief summary of the contents is subjoined.

- |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p>I. Introduction.<br/>                 II. Decomposition of Rocks.<br/>                 III. Removal of the Parts of Rocks by Water.<br/>                 IV. Lacustrine Deposits.<br/>                 V. Action of the Sea on Coasts.<br/>                 VI. Distribution and Deposit of Sediment in Tideless Seas.<br/>                 VII. Distribution and Deposit of Sediment in Tidal Seas.<br/>                 VIII. Chemical Deposits in Seas.<br/>                 IX. Preservation of the Remains of existing Life in Mineral Matter.<br/>                 X. Transportation of Mineral Matter by Ice.<br/>                 XI. Ossiferous Caves and Osseous Breccia.<br/>                 XII. Volcanoes and their Products.<br/>                 XIII. Salses, or Mud Volcanoes.<br/>                 XIV. Earthquakes.<br/>                 XV. Quiet Rise and Subsidence of Land.</p> | <p>XVI. Sunk (Submarine) Forests, and Raised Beaches.<br/>                 XVII. Temperature of the Earth.<br/>                 XVIII. Mode of Accumulation of Detrital and Fossiliferous Rocks.<br/>                 XIX. Igneous Products of Earlier Date than those of Modern Volcanoes.<br/>                 XX. Consolidation and Adjustment of the Component Parts of Rocks.<br/>                 XXI. Bending, Contortion, and Fracture of Bedded Rocks.<br/>                 XXII. Filling of Fissures and other Cavities with Mineral Matter.<br/>                 XXIII. Partial Removal and Denudation of Rocks.<br/>                 XXIV. Geological Maps and Sections.<br/>                 Additional Notices.</p> |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|



Tabular Iceberg.

NEW AND IMPROVED EDITION—(Jus

**PHYSICAL GEOGRAPHY**

BY MARY SOMERVILLE

A New American, from the Third and Revised Edition

WITH NOTES AND A GLOSSARY

BY W. S. W. RUSCHENBERGER, M. D.,

*In one large and handsome royal 12mo. volume, of nearly*

The subject of Physical Geography is one of which the acknowledgment of its introduction into all systems of education which pretend to drill the scholar into a mechanical knowledge of the names of territorial divisions of the earth's surface. A want is now felt of structure of the globe, externally and internally, and of the causes of land and water, forest and desert, heat and cold, tides, currents, and physical phenomena occurring around us, which have so direct an influence on the human race. This is all summed up in "Physical Geography," the resumé of all that is known on the natural history and present habits of the human race—the practical application of the principles which are elucidated by the scientific observer. This vast and interesting subject is so ably grappled by Mrs. Somerville, who in the present volume has set forth in a popular yet condensed account of the globe, in its relations to geological forces, its configuration and divisions into land and water, its meteorology, mineral productions, vegetation, and animal life, tracing the causes at work, and their influence on plants, animals, and man. This, taken in conjunction with ordinary political geography, leads to a mere catalogue of names and boundaries, and, in addition to the vast amount of information imparted, tends to impress the whole more strongly on the mind.

Eulogy is unnecessary with regard to a work like the present, which has passed through three editions on each side of the Atlantic within the space of a few years. We therefore only consider it necessary to state that the last London edition has been revised at the hands of the author, who introduced whatever improvements the advance of science rendered desirable, and that the present issue has been subjected to a careful examination on the part of the editor, to adapt it more speedily to the requirements of the age. In this revision care has been exercised in both the text and the glossary to obtain a work of this nature, and in its present improved and enlarged state, and in view of the increase of price, it is confidently presented by the publishers as in every respect of the striking favor with which it has been everywhere received.

**PHYSICAL ATLAS OF NATURAL PHILOSOPHY**

For the use of Colleges, Academies, and Families. By ALEXANDER K. BIRD, F. G. S., &c. In one large volume, imperial quarto, handsomely illustrated with twenty-six plates, engraved and colored in the best style. Together with twelve pages of descriptive letter-press, and a very copious Index.

A work which should be in every family and every school-room, for it affords a clear and concise view of the physical world. By the ingenious arrangement adopted by the author, it makes clear the connection between observation relative to the present condition of the earth, arrangement of Geology, Hydrography, Meteorology, and Natural History. The work is filled with a body of important information, nowhere else to be found condensed into so small a compass, while a very full Index renders the whole easy of reference.


**ELEMENTS OF NATURAL PHILOSOPHY**

BEING AN EXPERIMENTAL INTRODUCTION TO THE STUDY OF NATURAL PHILOSOPHY. Illustrated with over three hundred wood-cuts. By GOLDING BIRD, M. D., F. R. S., &c. From the Third and Improved London edition. 12mo. volume, of over 400 very large pages, on small type.

We are astonished to find that there is room in so small a book for even the bare recital of so many subjects. Where everything is treated succinctly, great judgment and much time are needed in making a selection and winnowing the wheat from the chaff. Dr. Bird has no need to partition as a shield against the world, and continues to be the language of this wide range of subjects. *American Review*, Apr.

**ELEMENTS OF PHYSIOLOGY**

OR, NATURAL PHILOSOPHY, GENERAL AND MEDICAL, IN PLAIN, OR NON-TECHNICAL LANGUAGE. By WILLIAM ARNOTT, M. D. With about two hundred illustrations.

*Electro-Magnetic Machine.*

## PRINCIPLES OF PHYSICS AND METEOROLOGY.

BY J. MULLER,

Professor of Physics at the University of Freiburg, &c.

Revised, and Illustrated

*With over five hundred beautiful Engravings on Wood, and two colored Plates.*

In one very handsome octavo volume, of nearly 650 pages.

In the work before us, the first thing which strikes us is the profusion of wood-cuts beyond what we should have imagined any publisher could furnish at the price—and very good cuts. The matter of the work is sound, and ranges over the subjects well. The manner is popular, and the author, though a mathematician, introduces only formulæ to look at now and then, without algebraic processes. We strongly recommend this work, particularly for the libraries of schools, mechanics' institutes, and the like. The present volume contains excellent reading and reference for those who are to have but one book on the subject.—*Athenæum*.

The plan adopted by Muller is simple; it reminds us of the excellent and popular treatise published many years since by Dr. Arnott, but it takes a much wider range of subjects. Like it, all the necessary

explanations are given in clear and concise language, without more than an occasional reference to mathematics; and the treatise is most abundantly illustrated with well-executed wood engravings.

The author has actually contrived to comprise in about five hundred pages, including the space occupied by illustrations, Mechanics, the laws of Motion, Acoustics, Light, Magnetism, Electricity, Galvanism, Electro-Magnetism, Heat, and Meteorology.

Medical practitioners and students, even if they have the means to procure, have certainly not the time to study an elaborate treatise in every branch of science; and the question therefore is, simply, whether they are to remain wholly ignorant of such subjects, or to make a profitable use of the labors of those who have the happy art of saying or suggesting much in a small space.

### ZOOPHYTES AND CORALS.

By JAMES D. DANA. In one large imperial quarto volume, extra cloth, with wood-cuts. (U. S. Exploring Expedition.)

Also, AN ATLAS to the above, in one imperial folio volume, half bound in morocco, with sixty-one large plates, exquisitely colored after nature.

*By the same Author.*

THE STRUCTURE AND CLASSIFICATION OF ZOOPHYTES. In one imperial quarto volume, extra cloth.



**PRINCIPLES OF THE  
MECHANICS OF MACHINERY AND ENGINEERING.**

**BY PROFESSOR JULIUS WEISBACH.**


TRANSLATED AND EDITED BY PROFESSOR GORDON, OF GLASGOW.

*First American Edition, with Additions,*

**BY PROFESSOR WALTER R. JOHNSON.**

*In two very handsome octavo volumes, of nearly 900 pages, with about 900 superb Illustrations.*

*Handsomely and strongly bound in extra cloth.*



*Jonval's Turbine.*

These volumes contain an immense amount of practical information necessary to the mechanical, architectural, and civil engineer; the whole thoroughly brought up to the most advanced state of scientific investigation. The following rapid summary of the contents will show the manner in which the subjects are successively brought forward.

**Volume I.**

- SECT. I. Mathematical Science of Motion.
- SECT. II. Mechanics in the Physical Science of Motion in general.
- SECT. III. Statics of Rigid Bodies.
- SECT. IV. Dynamics of Rigid Bodies.
- SECT. V. Statics of Fluid Bodies.
- SECT. VI. Dynamics of Fluid Bodies.

**Volume II.**

- SECTION I. Application of Mechanics in Building.
- DIVISION II. Application of Mechanics to Machinery.
- SECTION II. Of Moving Powers and their Effects.

These subjects, treated in all their ramifications and practical applications to Machines, Water-powers, Dams, Bridges, Water-wheels, Turbines, Roofs, Walls, Arches, Windmills, Scales, Steam-boilers, Water-engines, &c. &c., present points of primary importance to every one engaged in machine-work of every description, building, or engineering; while the comprehension of the whole is greatly facilitated by the profusion of clear and beautifully executed illustrations which are interspersed throughout the work.

**THE YOUNG MILLWRIGHT,**

**AND MILLER'S GUIDE.** Illustrated by twenty-eight Descriptive Plates. By OLIVER EVANS, with Additions and Corrections by THOMAS P. JONES. With a Description of an Improved Merchant Flour-mill, by C. and O. EVANS. Fourteenth edition. In one handsome octavo volume, strongly bound in leather.

**TECHNOLOGY;**  
OR,  
**CHEMISTRY APPLIED TO THE ARTS AND TO MANUFACTURES.**

BY DR. F. KNAPP.

Professor at the University of Giessen.

EDITED, WITH NUMEROUS NOTES AND ADDITIONS,  
BY DR. EDMUND RONALDS AND DR. THOMAS RICHARDSON.

*First American Edition, with Notes and Additions,*

BY PROF. WALTER R. JOHNSON.

In two  
very handsome  
octavo  
volumes  
strongly and  
handsomely  
bound in  
extra cloth.



*Sulphuric Acid Still.*

Containing  
about  
one thousand  
pages  
and nearly  
five hundred  
splendid  
illustrations on  
wood.

*Summary of Contents of Volume I.*

Heat—Coals, Relative Values of Fuels, Applications of Heat—Modes of effecting Illumination, Illuminating Power of different Materials—Manufacture of Salt—Soda—Sulphuric Acid, Potashes—Borax and Boracic Acid—Saltpetre—Gunpowder—Gun Cotton—Nitric Acid—Soap—Oil—Varnish—Lampblack—Coke—Patent fuel—Application of Heat—Ven-

tilation—Consumption of Smoke—Lamps—Candles—Gas.

*Summary of Contents of Volume II.*

Manufacture of Glass—Bottle Glass—Cylinder Glass—Crown Glass—Plate Glass—Flint Glass—Alum—Copperas—China-ware—Stoneware—Bricks—Lime—Gypsum—Magnesia.

These numerous and important branches of manufacture are treated practically, in all their details and processes, illustrated profusely with a vast number of engravings, representing fully the machinery employed, and the various stages of the processes. Every one, therefore, engaged in the manufacture of Pottery, Porcelain, Soap, Bricks, Gas, Gas Fixtures, Lamps, Chemicals of Various Descriptions, Gunpowder, Lampblack, Salt, Glass, &c. &c., will find in these volumes details of all the most recent and approved modes now followed in England and on the Continent, presenting a vast body of useful and indispensable information, nowhere else combined in the same compass.

**ELEMENTS OF OPTICS.**

By SIR DAVID BREWSTER. With Notes and Additions by A. D. BACHE, LL. D. In one 12mo. volume, half bound, with numerous wood-cuts.

Skinner's Complete Edition of Youatt on the Horse.  
A BOOK FOR EVERY FARMER AND COUNTRY GENTLEMAN.

## THE HORSE.

BY WILLIAM YOUATT.

A NEW EDITION, WITH NUMEROUS ILLUSTRATIONS.

TOGETHER WITH A  
GENERAL HISTORY OF THE HORSE; A DISSERTATION ON THE AMERICAN TROTTER HORSE;  
HOW TRAINED AND JOCKEYED; AN ACCOUNT OF HIS REMARKABLE PERFORMANCES;

AND  
AN ESSAY ON THE ASS AND THE MULE.

BY J. S. SKINNER,

Assistant Postmaster-General, and Editor of the Turf Register.

*In one octavo volume, of nearly 450 large pages, with numerous illustrations.*

This edition of Youatt's well-known and standard work on the Management, Diseases, and Treatment of the Horse, has already obtained such a wide circulation throughout the country, that the Publishers need say nothing to attract to it the attention and confidence of all who keep Horses or are interested in their improvement.

*By the same Author.*

## THE DOG.

EDITED, WITH ADDITIONS, BY E. J. LEWIS, M. D.

WITH NUMEROUS VERY HANDSOME PLATES AND WOOD-CUTS.

*In one very handsome volume, crown octavo, done up in beautiful crimson cloth, gilt stamps.*



*The Beagle.*

This volume will be found to contain a vast amount of information, nowhere else accessible, and of the utmost interest and importance to all who keep dogs. The anatomy, diseases, treatment, breeding, and breaking of that animal are described fully and clearly, together with the peculiarities of the various races and varieties; the whole illustrated in the finest style, and presented at a remarkably low price.

## A DICTIONARY OF MODERN GARDENING.


By GEORGE WILLIAM JOHNSON, Esq., Fellow of the Horticultural Society of India, &c. &c.


With 180 wood-cuts. Edited, with numerous Additions, by DAVID LANDRETH, of Philadelphia.

In one large royal 12mo. volume, of nearly 650 double-column pages, small type.

In the American edition, many modifications and additions have been made, so as to render the work a complete and satisfactory book of reference upon every subject connected with modern gardening in its most extended sense; while great care has been exercised in adapting it to the practice of every section of this country.









**LANE MEDICAL LIBRARY**

To avoid fine, this book should be returned  
on or before the date last stamped below.

|  |  |  |
|--|--|--|
|  |  |  |
|--|--|--|





